Differential Equations & Linear Algebra

FOURTH

HENRY EDWARDS DAVID E. PENNEY DAVID T. CALVIS



DIFFERENTIAL EQUATIONS & LINEAR ALGEBRA

Fourth Edition

C. Henry Edwards David E. Penney

The University of Georgia

David T. Calvis

Baldwin Wallace University



Director, Courseware Portfolio Management: Deirdre Lynch Executive Editor: Jeff Weidenaar Editorial Assistant: Jennifer Snyder Managing Producer: Karen Wernholm Content Producer: Tamela Ambush Producer: Nicholas Sweenv Field Marketing Manager: Evan St. Cyr Product Marketing Manager: Yvonne Vannatta Marketing Assistant: Jon Bryant Senior Author Support/Technology Specialist: Joe Vetere Manager, Rights Management: Gina M. Cheselka Manufacturing Buyer: Carol Melville, LSC Communications Composition: Dennis Kletzing, Kletzing Typesetting Corp. Cover Design: Studio Montage Cover Image: Thegoodly/Getty Images Photo Credits: Icon/Logo, Rashadashurov/Fotolia; Screenshots from Texas Instruments Incorporated. Courtesy of Texas Instruments Incorporated; p. 45, Screenshot by Wolfram Alpha LLC.

Copyright ©2018, 2010, 2005 by Pearson Education, Inc. or its affiliates. All Rights Reserved. Printed in the United States of America. This publication is protected by copyright, and permission should be obtained from the publisher prior to any prohibited reproduction, storage in a retrieval system, or transmission in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise. For information regarding permissions, request forms and the appropriate contacts within the Pearson Rights & Permissions Department, please visit www.pearsoned.com/permissions/.

PEARSON, ALWAYS LEARNING is an exclusive trademark owned by Pearson Education, Inc. or its affiliates in the United States and/or other countries.

Unless otherwise indicated herein, any third-party trademarks that may appear in this work are the property of their respective owners and any references to third-party trademarks, logos or other trade dress are for demonstrative or descriptive purposes only. Such references are not intended to imply any sponsorship, endorsement, authorization, or promotion of Pearson's products by the owners of such marks, or any relationship between the owner and Pearson Education, Inc. or its affiliates, authors, licensees or distributors.

Library of Congress Cataloging-in-Publication Data

- Names: Edwards, C. Henry (Charles Henry), 1937– | Penney, David E. | Calvis, David
- Title: Differential equations & linear algebra / C. Henry Edwards, David E. Penney, The University of Georgia; with the assistance of David Calvis, Baldwin-Wallace College.
- Description: Fourth edition. | Boston : Pearson, [2018] | Includes bibliographical references and index.
- Identifies: LCCN 2016030491 | ISBN 9780134497181 (hardcover) | ISBN 013449718X (hardcover)
- Subjects: LCSH: Differential equations. | Algebras, Linear.
- Classification: LCC QA372 .E34 2018 | DDC 515/.35--dc23
- LC record available at https://lccn.loc.gov/2016030491

1 16



ISBN 13: 978-0-13-449718-1 ISBN 10: 0-13-449718-X

CONTENTS

Application Modules vi Preface ix

CHAPTER

First-Order Differential Equations

1.1 Differential Equations and Mathematical Models 1

1

- 1.2 Integrals as General and Particular Solutions 10
- 1.3 Slope Fields and Solution Curves 17
- 1.4 Separable Equations and Applications 30
- 1.5 Linear First-Order Equations 46
- 1.6 Substitution Methods and Exact Equations 58

CHAPTER 2

Mathematical Models and Numerical Methods 75

- 2.1 Population Models 75
- 2.2 Equilibrium Solutions and Stability 87
- 2.3 Acceleration-Velocity Models 94
- 2.4 Numerical Approximation: Euler's Method 106
- 2.5 A Closer Look at the Euler Method 117
- 2.6 The Runge–Kutta Method 127

CHAPTER 3

Linear Systems and Matrices 138

- 3.1 Introduction to Linear Systems 147
- 3.2 Matrices and Gaussian Elimination 146
- 3.3 Reduced Row-Echelon Matrices 156
- 3.4 Matrix Operations 164
- 3.5 Inverses of Matrices 175
- 3.6 Determinants 188
- 3.7 Linear Equations and Curve Fitting 203

CHAPTER

Vector Spaces 211

- 4.1 The Vector Space R³ 211
- 4.2 The Vector Space Rⁿ and Subspaces 221
- 4.3 Linear Combinations and Independence of Vectors 228
- 4.4 Bases and Dimension for Vector Spaces 235
- 4.5 Row and Column Spaces 242
- 4.6 Orthogonal Vectors in Rⁿ 250
- 4.7 General Vector Spaces 257

CHAPTER

Higher-Order Linear Differential Equations 265

- 5.1 Introduction: Second-Order Linear Equations 265
- 5.2 General Solutions of Linear Equations 279
- 5.3 Homogeneous Equations with Constant Coefficients 291
- 5.4 Mechanical Vibrations 302
- 5.5 Nonhomogeneous Equations and Undetermined Coefficients 314
- 5.6 Forced Oscillations and Resonance 327

CHAPTER 6

Eigenvalues and Eigenvectors 339

- 6.1 Introduction to Eigenvalues 339
- 6.2 Diagonalization of Matrices 347
- 6.3 Applications Involving Powers of Matrices 354

CHAPTER 7

Linear Systems of Differential Equations 365

- 7.1 First-Order Systems and Applications 365
- 7.2 Matrices and Linear Systems 375
- 7.3 The Eigenvalue Method for Linear Systems 385
- 7.4 A Gallery of Solution Curves of Linear Systems 398
- 7.5 Second-Order Systems and Mechanical Applications 424
- 7.6 Multiple Eigenvalue Solutions 437
- 7.7 Numerical Methods for Systems 454

CHAPTER

Matrix Exponential Methods 469

- 8.1 Matrix Exponentials and Linear Systems 469
- 8.2 Nonhomogeneous Linear Systems 482
- 8.3 Spectral Decomposition Methods 490

CHAPTER **O**

9.1

Nonlinear Systems and Phenomena 503

- Stability and the Phase Plane 503
- 9.2 Linear and Almost Linear Systems 514
- 9.3 Ecological Models: Predators and Competitors 526
- 9.4 Nonlinear Mechanical Systems 539

CHAPTER 10

Laplace Transform Methods 557

- 10.1 Laplace Transforms and Inverse Transforms 557
- 10.2 Transformation of Initial Value Problems 567
- 10.3 Translation and Partial Fractions 578
- 10.4 Derivatives, Integrals, and Products of Transforms 587
- 10.5 Periodic and Piecewise Continuous Input Functions 594

CHAPTER 11

Power Series Methods 604

- 11.1 Introduction and Review of Power Series 604
- 11.2 Power Series Solutions 616
- 11.3 Frobenius Series Solutions 627
- 11.4 Bessel Functions 642

References for Further Study 652 Appendix A: Existence and Uniqueness of Solutions 654 Appendix B: Theory of Determinants 668 Answers to Selected Problems 677 Index 733

APPLICATION MODULES

The modules listed below follow the indicated sections in the text. Most provide computing projects that illustrate the corresponding text sections. Many of these modules are enhanced by the supplementary material found at the new Expanded Applications website, which can be accessed by visiting **goo.g1/BXB9k4**. For more information about the Expanded Applications, please review the Principal Features of this Revision section of the preface.

- 1.3 Computer-Generated Slope Fields and Solution Curves
- 1.4 The Logistic Equation
- 1.5 Indoor Temperature Oscillations
- 1.6 Computer Algebra Solutions
- 2.1 Logistic Modeling of Population Data
- 2.3 Rocket Propulsion
- 2.4 Implementing Euler's Method
- 2.5 Improved Euler Implementation
- 2.6 Runge-Kutta Implementation
- 3.2 Automated Row Operations
- 3.3 Automated Row Reduction
- 3.5 Automated Solution of Linear Systems
- 5.1 Plotting Second-Order Solution Families
- 5.2 Plotting Third-Order Solution Families
- 5.3 Approximate Solutions of Linear Equations
- 5.5 Automated Variation of Parameters
- 5.6 Forced Vibrations
- 7.1 Gravitation and Kepler's Laws of Planetary Motion
- 7.3 Automatic Calculation of Eigenvalues and Eigenvectors
- 7.4 Dynamic Phase Plane Graphics
- 7.5 Earthquake-Induced Vibrations of Multistory Buildings
- 7.6 Defective Eigenvalues and Generalized Eigenvectors
- 7.7 Comets and Spacecraft
- 8.1 Automated Matrix Exponential Solutions
- 8.2 Automated Variation of Parameters
- 9.1 Phase Plane Portraits and First-Order Equations
- 9.2 Phase Plane Portraits of Almost Linear Systems
- 9.3 Your Own Wildlife Conservation Preserve
- 9.4 The Rayleigh, van der Pol, and FitzHugh-Nagumo Equations

- 10.1 Computer Algebra Transforms and Inverse Transforms
- 10.2 Transforms of Initial Value Problems
- 10.3 Damping and Resonance Investigations
- 10.5 Engineering Functions
- 11.2 Automatic Computation of Series Coefficients
- 11.3 Automating the Frobenius Series Method

This page intentionally left blank

The evolution of the present text is based on experience teaching introductory differential equations and linear algebra with an emphasis on conceptual ideas and the use of applications and projects to involve students in active problem-solving experiences. Technical computing environments like *Maple*, *Mathematica*, MAT-LAB, and Python are widely available and are now used extensively by practicing engineers and scientists. This change in professional practice motivates a shift from the traditional concentration on manual symbolic methods to coverage also of qualitative and computer-based methods that employ numerical computation and graphical visualization to develop greater conceptual understanding. A bonus of this more comprehensive approach is accessibility to a wider range of more realistic applications of differential equations.

Both the conceptual and the computational aspects of such a course depend heavily on the perspective and techniques of linear algebra. Consequently, the study of differential equations and linear algebra in tandem reinforces the learning of both subjects. In this book we therefore have combined core topics in elementary differential equations with those concepts and methods of elementary linear algebra that are needed for a contemporary introduction to differential equations.

Principal Features of This Revision

This 4th edition is the most comprehensive and wide-ranging revision in the history of this text.

We have enhanced the exposition, as well as added graphics, in numerous sections throughout the book. We have also inserted new applications, including biological. Moreover we have exploited throughout the new interactive computer technology that is now available to students on devices ranging from desktop and laptop computers to smartphones and graphing calculators. While the text continues to use standard computer algebra systems such as *Mathematica, Maple*, and MATLAB, we have now added the Wolfram | Alpha website. In addition, this is the first edition of this book to feature Python, a computer platform that is freely available on the internet and which is gaining in popularity as an all-purpose scientific computing environment.

However, with a single exception of a new section inserted in Chapter 7 (noted below), the class-tested table of contents of the book remains unchanged. Therefore, instructors notes and syllabi will not require revision to continue teaching with this new edition.

A conspicuous feature of this edition is the insertion of about 80 new computergenerated figures, many of them illustrating interactive computer applications with slider bars or touchpad controls that can be used to change initial values or parameters in a differential equation, and immediately see in real time the resulting changes in the structure of its solutions. Some illustrations of the revisions and updating in this edition:

New Exposition In a number of sections, we have added new text and graphics to enhance student understanding of the subject matter. For instance, see the new introductory treatments of separable equations in Section 1.4 (page 30), of linear equations in Section 1.5 (page 46), and of isolated critical points in Sections 9.1 (page 503) and 9.2 (page 514). Also we have updated the examples and accompanying graphics in Sections 2.4–2.6, 7.3, and 7.7 to illustrate modern calculator technology.

New Interactive Technology and Graphics New figures throughout the text illustrate the capability that modern computing technology platforms offer to vary initial conditions and other parameters interactively. These figures are accompanied by detailed instructions that allow students to recreate the figures and make full use of the interactive features. For example, Section 7.4 includes the figure shown, a *Mathematica*-drawn phase plane diagram for a linear system of the form $\mathbf{x}' = \mathbf{A}\mathbf{x}$; after putting the accompanying code into *Mathematica*, the user can immediately

see the effect of changing the initial condition by clicking and dragging the "locator point" initially set at (4, 2).

Similarly, the application module for Section 5.1 now offers MATLAB and TI-Nspire graphics with interactive slider bars that vary the coefficients of a linear differential equation. The Section 11.2 application module features a new MATLAB graphic in which the user can vary the order of a series solution of an initial value problem, again immediately displaying the resulting graphical change in the corresponding approximate solution.



New Mathematica graphic in Section 7.4

New Content The single entirely new section for this edition is Section 7.4, which is devoted to the construction of a "gallery" of phase plane portraits illustrating all the possible geometric behaviors of solutions of the 2-dimensional linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. In motivation and preparation for the detailed study of eigenvalueeigenvector methods in subsequent sections of Chapter 7 (which then follow in the same order as in the previous edition), Section 7.4 shows how the particular arrangements of eigenvalues and eigenvectors of the coefficient matrix A correspond to identifiable patterns-"fingerprints," so to speak-in the phase plane portrait of the system. The resulting gallery is shown in the two pages of phase plane portraits in Figure 7.4.16 (pages 417-418) at the end of the section. The new 7.4 application module (on dynamic phase plane portraits, page 421) shows how students can use interactive computer systems to bring to life this gallery by allowing initial conditions, eigenvalues, and even eigenvectors to vary in real time. This dynamic approach is then illustrated with several new graphics inserted in the remainder of Chapter 7.

Finally, for a new biological application, see the application module for Section 9.4, which now includes a substantial investigation (page 551) of the nonlinear FitzHugh–Nagumo equations of neuroscience, which were introduced to model the behavior of neurons in the nervous system.

New Topical Headings Many of the examples and problems are now organized under headings that make the topic easy to see at a glance. This not only adds to the readability of the book, but it also makes it easier to choose in-class examples and homework problems. For instance, most of the text examples in Section 1.4 are

now labelled by topic, and the same is true of the wealth of problems following this section.

New Expanded Applications Website The effectiveness of the application modules located throughout the text is greatly enhanced by the supplementary material found at the new Expanded Applications website. Nearly all of the application mod-

ules in the text are marked with and a unique "tiny URL"—a web address that leads directly to an Expanded Applications page containing a wealth of electronic resources supporting that module. Typical Expanded Applications materials include an enhanced and expanded PDF version of the text with further discussion or additional applications, together with computer files in a variety of platforms, including *Mathematica*, *Maple*, MATLAB, and in some cases Python and/or TI calculator. These files provide all code appearing in the text as well as equivalent versions in other platforms, allowing students to immediately use the material in the Application Module on the computing platform of their choice. In addition to the URLs dispersed throughout the text, the Expanded Applications can be accessed by going to the Expanded Applications home page through this URL: goo.g1/BXB9k4. Note that when you enter URLs for the Extended Applications, take care to distinguish the following characters:

1 =lowercase L 1 =one I =uppercase I 0 =zero

```
o = uppercase O
```

Features of This Text

Computing Features The following features highlight the flavor of computing technology that distinguishes much of our exposition.

- Almost 600 *computer-generated figures* show students vivid pictures of direction fields, solution curves, and phase plane portraits that bring symbolic solutions of differential equations to life.
- About three dozen *application modules* follow key sections throughout the text. Most of these applications outline technology investigations that can be carried out using a variety of popular technical computing systems and which seek to actively engage students in the application of new technology. These modules are accompanied by the new Expanded Applications website previously detailed, which provides explicit code for nearly all of the applications in a number of popular technology platforms.
- The early introduction of numerical solution techniques in Chapter 2 (on mathematical models and numerical methods) allows for a fresh numerical emphasis throughout the text. Here and in Chapter 7, where numerical techniques for systems are treated, a concrete and tangible flavor is achieved by the inclusion of numerical algorithms presented in parallel fashion for systems ranging from graphing calculators to MATLAB and Python.

Modeling Features Mathematical modeling is a goal and constant motivation for the study of differential equations. For a small sample of the range of applications in this text, consider the following questions:

- What explains the commonly observed time lag between indoor and outdoor daily temperature oscillations? (Section 1.5)
- What makes the difference between doomsday and extinction in alligator populations? (Section 2.1)

- How do a unicycle and a car react differently to road bumps? (Sections 5.6 and 7.5)
- Why might an earthquake demolish one building and leave standing the one next door? (Section 7.5)
- How can you predict the time of next perihelion passage of a newly observed comet? (Section 7.7)
- What determines whether two species will live harmoniously together or whether competition will result in the extinction of one of them and the survival of the other? (Section 9.3)

Organization and Content This text reshapes the usual sequence of topics to accommodate new technology and new perspectives. For instance:

- After a precis of first-order equations in Chapter 1 (though with the coverage of certain traditional symbolic methods streamlined a bit), Chapter 2 offers an early introduction to mathematical modeling, stability and qualitative properties of differential equations, and numerical methods—a combination of topics that frequently are dispersed later in an introductory course.
- Chapters 3 (Linear Systems and Matrices), 4 (Vector Spaces), and 6 (Eigenvalues and Eigenvectors) provide concrete and self-contained coverage of the elementary linear algebra concepts and techniques that are needed for the solution of linear differential equations and systems. Chapter 4 includes sections 4.5 (row and column spaces) and 4.6 (orthogonal vectors in Rⁿ). Chapter 6 concludes with applications of diagonalizable matrices and a proof of the Cayley–Hamilton theorem for such matrices.
- Chapter 5 exploits the linear algebra of Chapters 3 and 4 to present efficiently the theory and solution of single linear differential equations. Chapter 7 is based on the eigenvalue approach to linear systems, and includes (in Section 7.6) the Jordan normal form for matrices and its application to the general Cayley–Hamilton theorem. This chapter includes an unusual number of applications (ranging from railway cars to earthquakes) of the various cases of the eigenvalue method, and concludes in Section 7.7 with numerical methods for systems.
- Chapter 8 is devoted to matrix exponentials with applications to linear systems of differential equations. The spectral decomposition method of Section 8.3 offers students an especially concrete approach to the computation of matrix exponentials.
- Chapter 9 exploits linear methods for the investigation of nonlinear systems and phenomena, and ranges from phase plane analysis to applications involving ecological and mechanical systems.
- Chapters 10 treats Laplace transform methods for the solution of constantcoefficient linear differential equations with a goal of handling the piecewise continuous and periodic forcing functions that are common in physical applications. Chapter 11 treats power series methods with a goal of discussing Bessel's equation with sufficient detail for the most common elementary applications.

This edition of the text also contains over 1800 end-of-section exercises, including a wealth of application problems. The Answers to Selected Problems section (page 677) includes answers to most odd-numbered problems plus a good many even-numbered ones, as well as about 175 computer-generated graphics to enhance its value as a learning aid. **Instructor's Solutions Manual** (0-13-449825-9) is available for instructors to download at Pearson's Instructor Resource Center (pearsonhighered.com/irc). This manual provides worked-out solutions for most of the problems in the book.

Student's Solutions Manual (0-13-449814-3) contains solutions for most of the odd-numbered problems.

Both manuals have been reworked extensively for this edition with improved explanations and more details inserted in the solutions of many problems.

Acknowledgments

In preparing this revision we profited greatly from the advice and assistance of the following very capable and perceptive reviewers:

Anthony Aidoo, Eastern Connecticut State University Miklos Bona, University of Florida Elizabeth Bradley, University of Louisville Mark Bridger, Northeastern University Raymond A. Claspadle, University of Memphis Gregory Davis, University of Wisconsin, Green Bay Sigal Gottlieb, University of Massachusetts, Dartmouth Zoran Grujic, University of Virginia Grant Gustafson, University of Utah Semion Gutman, University of Oklahoma Richard Jardine, Keene State College Yang Kuang, Arizona State University Dening Li, West Virginia University Carl Lutzer, Rochester Institute of Technology Francisco Sayas-Gonzalez, University of Delaware Morteza Shafii-Mousavi, Indiana University, South Bend Brent Solie, Knox College Ifran Ul-Haq, University of Wisconsin, Platteville Luther White, University of Oklahoma Hong-Ming Yin, Washington State University

We are grateful to our editor, Jeff Weidenaar, for advice and numerous suggestions that enhanced and shaped this revision; to Jennifer Snyder for her counsel and coordination of the editorial process; to Tamela Ambush and Julie Kidd for their supervision of the production process; and to Joe Vetere for his assistance with technical aspects of production of the supplementary manuals. It is a pleasure to (once again) credit Dennis Kletzing and his extraordinary TeXpertise for the attractive presentation of the text and graphics in this book (and its predecessors over the past decade).

Henry Edwards h.edwards@mindspring.com

David Calvis dcalvis@bw.edu This page intentionally left blank

First-Order Differential Equations

1.1 Differential Equations and Mathematical Models

The laws of the universe are written in the language of mathematics. Algebra is sufficient to solve many static problems, but the most interesting natural phenomena involve change and are described by equations that relate changing quantities.

Because the derivative dx/dt = f'(t) of the function f is the rate at which the quantity x = f(t) is changing with respect to the independent variable t, it is natural that equations involving derivatives are frequently used to describe the changing universe. An equation relating an unknown function and one or more of its derivatives is called a **differential equation**.

Example 1 The differential equation

$$\frac{dx}{dt} = x^2 + t^2$$

involves both the unknown function x(t) and its first derivative x'(t) = dx/dt. The differential equation

$$\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 7y = 0$$

involves the unknown function y of the independent variable x and the first two derivatives y' and y'' of y.

The study of differential equations has three principal goals:

- **1.** To discover the differential equation that describes a specified physical situation.
- **2.** To find—either exactly or approximately—the appropriate solution of that equation.
- **3.** To interpret the solution that is found.

In algebra, we typically seek the unknown *numbers* that satisfy an equation such as $x^3 + 7x^2 - 11x + 41 = 0$. By contrast, in solving a differential equation, we

are challenged to find the unknown *functions* y = y(x) for which an identity such as y'(x) = 2xy(x)—that is, the differential equation

$$\frac{dy}{dx} = 2xy$$

—holds on some interval of real numbers. Ordinarily, we will want to find *all* solutions of the differential equation, if possible.

If C is a constant and Example 2

$$y(x) = Ce^{x^2},\tag{1}$$

then

$$\frac{dy}{dx} = C\left(2xe^{x^2}\right) = (2x)\left(Ce^{x^2}\right) = 2xy.$$

Thus every function y(x) of the form in Eq. (1) satisfies—and thus is a solution of—the differential equation

$$\frac{dy}{dx} = 2xy \tag{2}$$

for all x. In particular, Eq. (1) defines an *infinite* family of different solutions of this differential equation, one for each choice of the arbitrary constant C. By the method of separation of variables (Section 1.4) it can be shown that every solution of the differential equation in (2) is of the form in Eq. (1).

Differential Equations and Mathematical Models

The following three examples illustrate the process of translating scientific laws and principles into differential equations. In each of these examples the independent variable is time t, but we will see numerous examples in which some quantity other than time is the independent variable.

Newton's law of cooling may be stated in this way: The *time rate of change* Rate of cooling (the rate of change with respect to time t) of the temperature T(t) of a body is proportional to the difference between T and the temperature A of the surrounding medium (Fig. 1.1.1). That is.

$$\frac{dT}{dt} = -k(T - A),\tag{3}$$

where k is a positive constant. Observe that if T > A, then dT/dt < 0, so the temperature is a decreasing function of t and the body is cooling. But if T < A, then dT/dt > 0, so that T is increasing.

Thus the physical law is translated into a differential equation. If we are given the values of k and A, we should be able to find an explicit formula for T(t), and then—with the aid of this formula-we can predict the future temperature of the body.

Draining tank Torricelli's law implies that the *time rate of change* of the volume V of water in a draining tank (Fig. 1.1.2) is proportional to the square root of the depth y of water in the tank:

$$\frac{dV}{dt} = -k\sqrt{y},\tag{4}$$

where k is a constant. If the tank is a cylinder with vertical sides and cross-sectional area A. then V = Ay, so $dV/dt = A \cdot (dy/dt)$. In this case Eq. (4) takes the form

$$\frac{dy}{dt} = -h\sqrt{y},\tag{5}$$

FIGURE 1.1.2. Torricelli's law of draining, Eq. (4), describes the draining of a water tank.

where h = k/A is a constant.



FIGURE 1.1.1. Newton's law of

cooling, Eq. (3), describes the cooling

Population growth The *time rate of change* of a population P(t) with constant birth and death rates is, in many simple cases, proportional to the size of the population. That is,

$$\frac{dP}{dt} = kP,\tag{6}$$

where k is the constant of proportionality.

Let us discuss Example 5 further. Note first that each function of the form

$$P(t) = Ce^{kt} \tag{7}$$

is a solution of the differential equation

$$\frac{dP}{dt} = kP$$

in (6). We verify this assertion as follows:

$$P'(t) = Cke^{kt} = k\left(Ce^{kt}\right) = kP(t)$$

for all real numbers t. Because substitution of each function of the form given in (7) into Eq. (6) produces an identity, all such functions are solutions of Eq. (6).

Thus, even if the value of the constant k is known, the differential equation dP/dt = kP has *infinitely many* different solutions of the form $P(t) = Ce^{kt}$, one for each choice of the "arbitrary" constant C. This is typical of differential equations. It is also fortunate, because it may allow us to use additional information to select from among all these solutions a particular one that fits the situation under study.

Example 6 Population growth Suppose that $P(t) = Ce^{kt}$ is the population of a colony of bacteria at time *t*, that the population at time t = 0 (hours, h) was 1000, and that the population doubled after 1 h. This additional information about P(t) yields the following equations:

$$1000 = P(0) = Ce^{0} = C,$$

$$2000 = P(1) = Ce^{k}.$$

It follows that C = 1000 and that $e^k = 2$, so $k = \ln 2 \approx 0.693147$. With this value of k the differential equation in (6) is

$$\frac{dP}{dt} = (\ln 2)P \approx (0.693147)P.$$

Substitution of $k = \ln 2$ and C = 1000 in Eq. (7) yields the particular solution

$$P(t) = 1000e^{(\ln 2)t} = 1000(e^{\ln 2})^t = 1000 \cdot 2^t$$
 (because $e^{\ln 2} = 2$)

that satisfies the given conditions. We can use this particular solution to predict future populations of the bacteria colony. For instance, the predicted number of bacteria in the population after one and a half hours (when t = 1.5) is

~ / ~

$$P(1.5) = 1000 \cdot 2^{3/2} \approx 2828.$$

The condition P(0) = 1000 in Example 6 is called an **initial condition** because we frequently write differential equations for which t = 0 is the "starting time." Figure 1.1.3 shows several different graphs of the form $P(t) = Ce^{kt}$ with $k = \ln 2$. The graphs of all the infinitely many solutions of dP/dt = kP in fact fill the entire two-dimensional plane, and no two intersect. Moreover, the selection of any one point P_0 on the *P*-axis amounts to a determination of P(0). Because exactly one solution passes through each such point, we see in this case that an initial condition $P(0) = P_0$ determines a unique solution agreeing with the given data.



FIGURE 1.1.3. Graphs of $P(t) = Ce^{kt}$ with $k = \ln 2$.

Mathematical Models

Our brief discussion of population growth in Examples 5 and 6 illustrates the crucial process of *mathematical modeling* (Fig. 1.1.4), which involves the following:

- **1.** The formulation of a real-world problem in mathematical terms; that is, the construction of a mathematical model.
- 2. The analysis or solution of the resulting mathematical problem.
- **3.** The interpretation of the mathematical results in the context of the original real-world situation—for example, answering the question originally posed.



FIGURE 1.1.4. The process of mathematical modeling.

In the population example, the real-world problem is that of determining the population at some future time. A **mathematical model** consists of a list of variables (P and t) that describe the given situation, together with one or more equations relating these variables (dP/dt = kP, $P(0) = P_0$) that are known or are assumed to hold. The mathematical analysis consists of solving these equations (here, for P as a function of t). Finally, we apply these mathematical results to attempt to answer the original real-world question.

As an example of this process, think of first formulating the mathematical model consisting of the equations dP/dt = kP, P(0) = 1000, describing the bacteria population of Example 6. Then our mathematical analysis there consisted of solving for the solution function $P(t) = 1000e^{(\ln 2)t} = 1000 \cdot 2^t$ as our mathematical result. For an interpretation in terms of our real-world situation—the actual bacteria population—we substituted t = 1.5 to obtain the predicted population of $P(1.5) \approx 2828$ bacteria after 1.5 hours. If, for instance, the bacteria population is growing under ideal conditions of unlimited space and food supply, our prediction may be quite accurate, in which case we conclude that the mathematical model is adequate for studying this particular population.

On the other hand, it may turn out that no solution of the selected differential equation accurately fits the actual population we're studying. For instance, for *no* choice of the constants *C* and *k* does the solution $P(t) = Ce^{kt}$ in Eq. (7) accurately describe the actual growth of the human population of the world over the past few centuries. We must conclude that the differential equation dP/dt = kP is inadequate for modeling the world population—which in recent decades has "leveled off" as compared with the steeply climbing graphs in the upper half (P > 0) of Fig. 1.1.3. With sufficient insight, we might formulate a new mathematical model including a perhaps more complicated differential equation, one that takes into account such factors as a limited food supply and the effect of increased population on birth and death rates. With the formulation of this new mathematical model, we may attempt to traverse once again the diagram of Fig. 1.1.4 in a counterclockwise manner. If we can solve the new differential equation, we get new solution functions to com-

pare with the real-world population. Indeed, a successful population analysis may require refining the mathematical model still further as it is repeatedly measured against real-world experience.

But in Example 6 we simply ignored any complicating factors that might affect our bacteria population. This made the mathematical analysis quite simple, perhaps unrealistically so. A satisfactory mathematical model is subject to two contradictory requirements: It must be sufficiently detailed to represent the real-world situation with relative accuracy, yet it must be sufficiently simple to make the mathematical analysis practical. If the model is so detailed that it fully represents the physical situation, then the mathematical analysis may be too difficult to carry out. If the model is too simple, the results may be so inaccurate as to be useless. Thus there is an inevitable tradeoff between what is physically realistic and what is mathematically possible. The construction of a model that adequately bridges this gap between realism and feasibility is therefore the most crucial and delicate step in the process. Ways must be found to simplify the model mathematically without sacrificing essential features of the real-world situation.

Mathematical models are discussed throughout this book. The remainder of this introductory section is devoted to simple examples and to standard terminology used in discussing differential equations and their solutions.

Examples and Terminology

Example 7	If C is a constant and $y(x) = 1/(C - x)$, then					
		$\frac{dy}{dx} = \frac{1}{(C-x)^2} = y^2$				
	if $x \neq C$. Thus	$y(x) = \frac{1}{C - x}$	(8)			

defines a solution of the differential equation

$$\frac{dy}{dx} = y^2 \tag{9}$$

on any interval of real numbers not containing the point x = C. Actually, Eq. (8) defines a *one-parameter family* of solutions of $dy/dx = y^2$, one for each value of the arbitrary constant or "parameter" C. With C = 1 we get the particular solution

$$y(x) = \frac{1}{1-x}$$

that satisfies the initial condition y(0) = 1. As indicated in Fig. 1.1.5, this solution is continuous on the interval $(-\infty, 1)$ but has a vertical asymptote at x = 1.

Example 8 Verify that the function $y(x) = 2x^{1/2} - x^{1/2} \ln x$ satisfies the differential equation

$$4x^2y'' + y = 0 (10)$$

for all
$$x > 0$$
.

Solution First we compute the derivatives

$$y'(x) = -\frac{1}{2}x^{-1/2}\ln x$$
 and $y''(x) = \frac{1}{4}x^{-3/2}\ln x - \frac{1}{2}x^{-3/2}$.

Then substitution into Eq. (10) yields

$$4x^{2}y'' + y = 4x^{2}\left(\frac{1}{4}x^{-3/2}\ln x - \frac{1}{2}x^{-3/2}\right) + 2x^{1/2} - x^{1/2}\ln x = 0$$

if x is positive, so the differential equation is satisfied for all x > 0.

The fact that we can write a differential equation is not enough to guarantee that it has a solution. For example, it is clear that the differential equation

$$(y')^2 + y^2 = -1 \tag{11}$$

has *no* (real-valued) solution, because the sum of nonnegative numbers cannot be negative. For a variation on this theme, note that the equation

$$(y')^2 + y^2 = 0 (12)$$

obviously has only the (real-valued) solution $y(x) \equiv 0$. In our previous examples any differential equation having at least one solution indeed had infinitely many.

The **order** of a differential equation is the order of the highest derivative that appears in it. The differential equation of Example 8 is of second order, those in Examples 2 through 7 are first-order equations, and

$$y^{(4)} + x^2 y^{(3)} + x^5 y = \sin x$$

is a fourth-order equation. The most general form of an *n*th-order differential equation with independent variable x and unknown function or dependent variable y = y(x) is

$$F\left(x, y, y', y'', \dots, y^{(n)}\right) = 0,$$
(13)

where *F* is a specific real-valued function of n + 2 variables.

Our use of the word *solution* has been until now somewhat informal. To be precise, we say that the continuous function u = u(x) is a **solution** of the differential equation in (13) **on the interval** *I* provided that the derivatives $u', u'', \ldots, u^{(n)}$ exist on *I* and

$$F\left(x, u, u', u'', \dots, u^{(n)}\right) = 0$$

for all x in I. For the sake of brevity, we may say that u = u(x) satisfies the differential equation in (13) on I.

Remark Recall from elementary calculus that a differentiable function on an open interval is necessarily continuous there. This is why only a continuous function can qualify as a (differentiable) solution of a differential equation on an interval.

Continued Figure 1.1.5 shows the two "connected" branches of the graph y = 1/(1 - x). The left-hand branch is the graph of a (continuous) solution of the differential equation $y' = y^2$ that is defined on the interval $(-\infty, 1)$. The right-hand branch is the graph of a *different* solution of the differential equation that is defined (and continuous) on the different interval $(1, \infty)$. So the single formula y(x) = 1/(1 - x) actually defines two different solutions (with different domains of definition) of the same differential equation $y' = y^2$.

Example 9 If A and B are constants and

$$y(x) = A\cos 3x + B\sin 3x,$$
(14)

then two successive differentiations yield

$$y'(x) = -3A\sin 3x + 3B\cos 3x, y''(x) = -9A\cos 3x - 9B\sin 3x = -9y(x)$$

for all x. Consequently, Eq. (14) defines what it is natural to call a *two-parameter family* of solutions of the second-order differential equation

$$y'' + 9y = 0 (15)$$

on the whole real number line. Figure 1.1.6 shows the graphs of several such solutions.



FIGURE 1.1.5. The solution of $y' = y^2$ defined by y(x) = 1/(1-x).



FIGURE 1.1.6. The three solutions $y_1(x) = 3\cos 3x$, $y_2(x) = 2\sin 3x$, and $y_3(x) = -3\cos 3x + 2\sin 3x$ of the differential equation y'' + 9y = 0.

≻

Although the differential equations in (11) and (12) are exceptions to the general rule, we will see that an *n*th-order differential equation ordinarily has an *n*-parameter family of solutions—one involving *n* different arbitrary constants or parameters.

In both Eqs. (11) and (12), the appearance of y' as an implicitly defined function causes complications. For this reason, we will ordinarily assume that any differential equation under study can be solved explicitly for the highest derivative that appears; that is, that the equation can be written in the so-called *normal form*

$$y^{(n)} = G\left(x, y, y', y'', \dots, y^{(n-1)}\right),$$
(16)

where G is a real-valued function of n + 1 variables. In addition, we will always seek only real-valued solutions unless we warn the reader otherwise.

All the differential equations we have mentioned so far are **ordinary** differential equations, meaning that the unknown function (dependent variable) depends on only a *single* independent variable. If the dependent variable is a function of two or more independent variables, then partial derivatives are likely to be involved; if they are, the equation is called a **partial** differential equation. For example, the temperature u = u(x, t) of a long thin uniform rod at the point x at time t satisfies (under appropriate simple conditions) the partial differential equation

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2},$$

where k is a constant (called the *thermal diffusivity* of the rod). In Chapters 1 through 8 we will be concerned only with *ordinary* differential equations and will refer to them simply as differential equations.

In this chapter we concentrate on *first-order* differential equations of the form

$$\frac{dy}{dx} = f(x, y). \tag{17}$$

We also will sample the wide range of applications of such equations. A typical mathematical model of an applied situation will be an **initial value problem**, consisting of a differential equation of the form in (17) together with an **initial condi-**tion $y(x_0) = y_0$. Note that we call $y(x_0) = y_0$ an initial condition whether or not $x_0 = 0$. To **solve** the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0$$
 (18)

means to find a differentiable function y = y(x) that satisfies both conditions in Eq. (18) on some interval containing x_0 .

Example 10 Given the solution y(x) = 1/(C - x) of the differential equation $dy/dx = y^2$ discussed in Example 7, solve the initial value problem

$$\frac{dy}{dx} = y^2, \quad y(1) = 2.$$

Solution We need only find a value of C so that the solution y(x) = 1/(C - x) satisfies the initial condition y(1) = 2. Substitution of the values x = 1 and y = 2 in the given solution yields

$$2 = y(1) = \frac{1}{C - 1}$$



FIGURE 1.1.7. The solutions of $y' = y^2$ defined by y(x) = 2/(3 - 2x).

so 2C - 2 = 1, and hence $C = \frac{3}{2}$. With this value of C we obtain the desired solution

$$y(x) = \frac{1}{\frac{3}{2} - x} = \frac{2}{3 - 2x}$$

Figure 1.1.7 shows the two branches of the graph y = 2/(3 - 2x). The left-hand branch is the graph on $(-\infty, \frac{3}{2})$ of the solution of the given initial value problem $y' = y^2$, y(1) = 2. The right-hand branch passes through the point (2, -2) and is therefore the graph on $(\frac{3}{2}, \infty)$ of the solution of the different initial value problem $y' = y^2$, y(2) = -2.

The central question of greatest immediate interest to us is this: If we are given a differential equation known to have a solution satisfying a given initial condition, how do we actually *find* or *compute* that solution? And, once found, what can we do with it? We will see that a relatively few simple techniques—separation of variables (Section 1.4), solution of linear equations (Section 1.5), elementary substitution methods (Section 1.6)—are enough to enable us to solve a variety of first-order equations having impressive applications.

1.1 **Problems**

In Problems 1 through 12, verify by substitution that each given function is a solution of the given differential equation. Throughout these problems, primes denote derivatives with respect to x.

1. $y' = 3x^2$; $y = x^3 + 7$ 2. y' + 2y = 0; $y = 3e^{-2x}$ 3. y'' + 4y = 0; $y_1 = \cos 2x$, $y_2 = \sin 2x$ 4. y'' = 9y; $y_1 = e^{3x}$, $y_2 = e^{-3x}$ 5. $y' = y + 2e^{-x}$; $y = e^x - e^{-x}$ 6. y'' + 4y' + 4y = 0; $y_1 = e^{-2x}$, $y_2 = xe^{-2x}$ 7. y'' - 2y' + 2y = 0; $y_1 = e^x \cos x$, $y_2 = e^x \sin x$ 8. $y'' + y = 3\cos 2x$, $y_1 = \cos x - \cos 2x$, $y_2 = \sin x - \cos 2x$ 9. $y' + 2xy^2 = 0$; $y = \frac{1}{1 + x^2}$ 10. $x^2y'' + xy' - y = \ln x$; $y_1 = x - \ln x$, $y_2 = \frac{1}{2} - \ln x$

10.
$$x^{2}y'' + 5xy' + 4y = 0; y_{1} = \frac{1}{x^{2}}, y_{2} = \frac{\ln x}{x^{2}}$$

11. $x^{2}y'' + 5xy' + 4y = 0; y_{1} = \frac{1}{x^{2}}, y_{2} = \frac{\ln x}{x^{2}}$
12. $x^{2}y'' - xy' + 2y = 0; y_{1} = x \cos(\ln x), y_{2} = x \sin(\ln x)$

In Problems 13 through 16, substitute $y = e^{rx}$ into the given differential equation to determine all values of the constant r for which $y = e^{rx}$ is a solution of the equation.

13. 3y' = 2y**14.** 4y'' = y**15.** y'' + y' - 2y = 0**16.** 3y'' + 3y' - 4y = 0

In Problems 17 through 26, first verify that y(x) satisfies the given differential equation. Then determine a value of the constant C so that y(x) satisfies the given initial condition. Use a computer or graphing calculator (if desired) to sketch several typical solutions of the given differential equation, and highlight the one that satisfies the given initial condition.

17.
$$y' + y = 0$$
; $y(x) = Ce^{-x}$, $y(0) = 2$
18. $y' = 2y$; $y(x) = Ce^{2x}$, $y(0) = 3$
19. $y' = y + 1$; $y(x) = Ce^{x} - 1$, $y(0) = 5$

20. $y' = x - y; y(x) = Ce^{-x} + x - 1, y(0) = 10$ **21.** $y' + 3x^2y = 0; y(x) = Ce^{-x^3}, y(0) = 7$ **22.** $e^y y' = 1; y(x) = \ln(x + C), y(0) = 0$ **23.** $x\frac{dy}{dx} + 3y = 2x^5; y(x) = \frac{1}{4}x^5 + Cx^{-3}, y(2) = 1$ **24.** $xy' - 3y = x^3; y(x) = x^3(C + \ln x), y(1) = 17$ **25.** $y' = 3x^2(y^2 + 1); y(x) = \tan(x^3 + C), y(0) = 1$ **26.** $y' + y \tan x = \cos x; y(x) = (x + C)\cos x, y(\pi) = 0$

In Problems 27 through 31, a function y = g(x) is described by some geometric property of its graph. Write a differential equation of the form dy/dx = f(x, y) having the function g as its solution (or as one of its solutions).

- **27.** The slope of the graph of g at the point (x, y) is the sum of x and y.
- **28.** The line tangent to the graph of g at the point (x, y) intersects the x-axis at the point (x/2, 0).
- **29.** Every straight line normal to the graph of *g* passes through the point (0, 1). Can you *guess* what the graph of such a function *g* might look like?
- **30.** The graph of g is normal to every curve of the form $y = x^2 + k$ (k is a constant) where they meet.
- **31.** The line tangent to the graph of g at (x, y) passes through the point (-y, x).

Differential Equations as Models

In Problems 32 through 36, write—in the manner of Eqs. (3) through (6) of this section—a differential equation that is a mathematical model of the situation described.

- **32.** The time rate of change of a population *P* is proportional to the square root of *P*.
- **33.** The time rate of change of the velocity v of a coasting motorboat is proportional to the square of v.
- **34.** The acceleration dv/dt of a Lamborghini is proportional to the difference between 250 km/h and the velocity of the car.

- **35.** In a city having a fixed population of P persons, the time rate of change of the number N of those persons who have heard a certain rumor is proportional to the number of those who have not yet heard the rumor.
- **36.** In a city with a fixed population of P persons, the time rate of change of the number N of those persons infected with a certain contagious disease is proportional to the product of the number who have the disease and the number who do not.

In Problems 37 through 42, determine by inspection at least one solution of the given differential equation. That is, use your knowledge of derivatives to make an intelligent guess. Then test your hypothesis.

37.	y'' = 0	38.	y' = y
39.	$xy' + y = 3x^2$	40.	$(y')^2 + y^2 = 1$
41.	$y' + y = e^x$	42.	y'' + y = 0

Problems 43 through 46 concern the differential equation

$$\frac{dx}{dt} = kx^2$$

where k is a constant.

- **43.** (a) If k is a constant, show that a general (one-parameter) solution of the differential equation is given by x(t) = 1/(C kt), where C is an arbitrary constant.
 - (b) Determine by inspection a solution of the initial value problem $x' = kx^2$, x(0) = 0.
- **44.** (a) Assume that k is positive, and then sketch graphs of solutions of $x' = kx^2$ with several typical positive values of x(0).
 - (**b**) How would these solutions differ if the constant *k* were negative?
- **45.** Suppose a population P of rodents satisfies the differential equation $dP/dt = kP^2$. Initially, there are P(0) =



FIGURE 1.1.8. Graphs of solutions of the equation $dy/dx = y^2$.

2 rodents, and their number is increasing at the rate of dP/dt = 1 rodent per month when there are P = 10 rodents. Based on the result of Problem 43, how long will it take for this population to grow to a hundred rodents? To a thousand? What's happening here?

- **46.** Suppose the velocity v of a motorboat coasting in water satisfies the differential equation $dv/dt = kv^2$. The initial speed of the motorboat is v(0) = 10 meters per second (m/s), and v is decreasing at the rate of 1 m/s^2 when v = 5 m/s. Based on the result of Problem 43, long does it take for the velocity of the boat to decrease to 1 m/s^2 . To $\frac{1}{10}$ m/s? When does the boat come to a stop?
- **47.** In Example 7 we saw that y(x) = 1/(C x) defines a one-parameter family of solutions of the differential equation $dy/dx = y^2$. (a) Determine a value of *C* so that y(10) = 10. (b) Is there a value of *C* such that y(0) = 0? Can you nevertheless find by inspection a solution of $dy/dx = y^2$ such that y(0) = 0? (c) Figure 1.1.8 shows typical graphs of solutions of the form y(x) = 1/(C x). Does it appear that these solution curves fill the entire xy-plane? Can you conclude that, given any point (a, b) in the plane, the differential equation $dy/dx = y^2$ has exactly one solution y(x) satisfying the condition y(a) = b?
- **48.** (a) Show that $y(x) = Cx^4$ defines a one-parameter family of differentiable solutions of the differential equation xy' = 4y (Fig. 1.1.9). (b) Show that

$$y(x) = \begin{cases} -x^4 & \text{if } x < 0\\ x^4 & \text{if } x \ge 0 \end{cases}$$

defines a differentiable solution of xy' = 4y for all x, but is not of the form $y(x) = Cx^4$. (c) Given any two real numbers a and b, explain why—in contrast to the situation in part (c) of Problem 47—there exist infinitely many differentiable solutions of xy' = 4y that all satisfy the condition y(a) = b.



FIGURE 1.1.9. The graph $y = Cx^4$ for various values of *C*.

.2 Integrals as General and Particular Solutions

The first-order equation dy/dx = f(x, y) takes an especially simple form if the right-hand-side function f does not actually involve the dependent variable y, so

$$\frac{dy}{dx} = f(x). \tag{1}$$

In this special case we need only integrate both sides of Eq. (1) to obtain

$$y(x) = \int f(x) \, dx + C. \tag{2}$$

This is a **general solution** of Eq. (1), meaning that it involves an arbitrary constant C, and for every choice of C it is a solution of the differential equation in (1). If G(x) is a particular antiderivative of f—that is, if $G'(x) \equiv f(x)$ —then

$$y(x) = G(x) + C.$$
(3)

The graphs of any two such solutions $y_1(x) = G(x) + C_1$ and $y_2(x) = G(x) + C_2$ on the same interval *I* are "parallel" in the sense illustrated by Figs. 1.2.1 and 1.2.2. There we see that the constant *C* is geometrically the vertical distance between the two curves y(x) = G(x) and y(x) = G(x) + C.

To satisfy an initial condition $y(x_0) = y_0$, we need only substitute $x = x_0$ and $y = y_0$ into Eq. (3) to obtain $y_0 = G(x_0) + C$, so that $C = y_0 - G(x_0)$. With this choice of *C*, we obtain the **particular solution** of Eq. (1) satisfying the initial value problem

$$\frac{dy}{dx} = f(x), \quad y(x_0) = y_0$$

We will see that this is the typical pattern for solutions of first-order differential equations. Ordinarily, we will first find a *general solution* involving an arbitrary constant *C*. We can then attempt to obtain, by appropriate choice of *C*, a *particular solution* satisfying a given initial condition $y(x_0) = y_0$.

Remark As the term is used in the previous paragraph, a *general solution* of a first-order differential equation is simply a one-parameter family of solutions. A natural question is whether a given general solution contains *every* particular solution of the differential equation. When this is known to be true, we call it **the** general solution of the differential equation. For example, because any two antiderivatives of the same function f(x) can differ only by a constant, it follows that every solution of Eq. (1) is of the form in (2). Thus Eq. (2) serves to define **the** general solution of (1).

Example 1 Solve the in

>

$$\frac{dy}{dx} = 2x + 3, \quad y(1) = 2.$$

Solutio

$$y(x) = \int (2x+3) \, dx = x^2 + 3x + C.$$

Figure 1.2.3 shows the graph $y = x^2 + 3x + C$ for various values of *C*. The particular solution we seek corresponds to the curve that passes through the point (1, 2), thereby satisfying the initial condition

$$y(1) = (1)^2 + 3 \cdot (1) + C = 2.$$

It follows that C = -2, so the desired particular solution is

$$y(x) = x^2 + 3x - 2.$$







FIGURE 1.2.2. Graphs of $y = \sin x + C$ for various values of *C*.

1.2 INTEGRALS AS GENERAL AND PARTICULAR SOLUTIONS



FIGURE 1.2.3. Solution curves for the differential equation in Example 1.

Second-order equations. The observation that the special first-order equation dy/dx = f(x) is readily solvable (provided that an antiderivative of f can be found) extends to second-order differential equations of the special form

$$\frac{d^2y}{dx^2} = g(x),\tag{4}$$

in which the function g on the right-hand side involves neither the dependent variable y nor its derivative dy/dx. We simply integrate once to obtain

$$\frac{dy}{dx} = \int y''(x) \, dx = \int g(x) \, dx = G(x) + C_1,$$

where G is an antiderivative of g and C_1 is an arbitrary constant. Then another integration yields

$$y(x) = \int y'(x) \, dx = \int [G(x) + C_1] \, dx = \int G(x) \, dx + C_1 x + C_2$$

where C_2 is a second arbitrary constant. In effect, the second-order differential equation in (4) is one that can be solved by solving successively the *first-order* equations

$$\frac{dv}{dx} = g(x)$$
 and $\frac{dy}{dx} = v(x)$.

Velocity and Acceleration

Direct integration is sufficient to allow us to solve a number of important problems concerning the motion of a particle (or *mass point*) in terms of the forces acting on it. The motion of a particle along a straight line (the *x*-axis) is described by its **position function**

$$x = f(t) \tag{5}$$

giving its x-coordinate at time t. The velocity of the particle is defined to be

$$v(t) = f'(t);$$
 that is, $v = \frac{dx}{dt}.$ (6)

Its **acceleration** a(t) is a(t) = v'(t) = x''(t); in Leibniz notation,

$$a = \frac{dv}{dt} = \frac{d^2x}{dt^2}.$$
(7)

Equation (6) is sometimes applied either in the indefinite integral form $x(t) = \int v(t) dt$ or in the definite integral form

$$x(t) = x(t_0) + \int_{t_0}^t v(s) \, ds,$$

which you should recognize as a statement of the fundamental theorem of calculus (precisely because dx/dt = v).

Newton's *second law of motion* says that if a force F(t) acts on the particle and is directed along its line of motion, then

$$ma(t) = F(t);$$
 that is, $F = ma,$ (8)

where *m* is the mass of the particle. If the force *F* is known, then the equation x''(t) = F(t)/m can be integrated twice to find the position function x(t) in terms of two constants of integration. These two arbitrary constants are frequently determined by the **initial position** $x_0 = x(0)$ and the **initial velocity** $v_0 = v(0)$ of the particle.

Constant acceleration. For instance, suppose that the force F, and therefore the acceleration a = F/m, are *constant*. Then we begin with the equation

$$\frac{dv}{dt} = a \quad (a \text{ is a constant}) \tag{9}$$

and integrate both sides to obtain

$$v(t) = \int a \, dt = at + C_1.$$

We know that $v = v_0$ when t = 0, and substitution of this information into the preceding equation yields the fact that $C_1 = v_0$. So

$$v(t) = \frac{dx}{dt} = at + v_0.$$
⁽¹⁰⁾

A second integration gives

$$x(t) = \int v(t) dt = \int (at + v_0) dt = \frac{1}{2}at^2 + v_0t + C_2,$$

and the substitution t = 0, $x = x_0$ gives $C_2 = x_0$. Therefore,

$$x(t) = \frac{1}{2}at^2 + v_0t + x_0.$$
(11)

Thus, with Eq. (10) we can find the velocity, and with Eq. (11) the position, of the particle at any time *t* in terms of its *constant* acceleration *a*, its initial velocity v_0 , and its initial position x_0 .

Example 2 Lunar lander A lunar lander is falling freely toward the surface of the moon at a speed of 450 meters per second (m/s). Its retrorockets, when fired, provide a constant deceleration of 2.5 meters per second per second (m/s^2) (the gravitational acceleration produced by the moon is assumed to be included in the given deceleration). At what height above the lunar surface should the retrorockets be activated to ensure a "soft touchdown" (v = 0 at impact)? We denote by x(t) the height of the lunar lander above the surface, as indicated in Fig. 1.2.4. We let t = 0 denote the time at which the retrorockets should be fired. Then $v_0 = -450$ (m/s, negative because the height x(t) is decreasing), and a = +2.5, because an upward thrust increases the velocity v (although it decreases the *speed* |v|). Then Eqs. (10) and (11) become

$$v(t) = 2.5t - 450 \tag{12}$$

(13)

$$x(t) = 1.25t^2 - 450t + x_0,$$

where x_0 is the height of the lander above the lunar surface at the time t = 0 when the retrorockets should be activated.

From Eq. (12) we see that v = 0 (soft touchdown) occurs when t = 450/2.5 = 180 s (that is, 3 minutes); then substitution of t = 180, x = 0 into Eq. (13) yields

$$x_0 = 0 - (1.25)(180)^2 + 450(180) = 40,500$$

meters—that is, $x_0 = 40.5$ km $\approx 25\frac{1}{6}$ miles. Thus the retrorockets should be activated when the lunar lander is 40.5 kilometers above the surface of the moon, and it will touch down softly on the lunar surface after 3 minutes of decelerating descent.

and

Lunar surface

FIGURE 1.2.4. The lunar lander of Example 2.

Physical Units

Numerical work requires units for the measurement of physical quantities such as distance and time. We sometimes use ad hoc units—such as distance in miles or kilometers and time in hours—in special situations (such as in a problem involving an auto trip). However, the foot-pound-second (fps) and meter-kilogram-second (mks) unit systems are used more generally in scientific and engineering problems. In fact, fps units are commonly used only in the United States (and a few other countries), while mks units constitute the standard international system of scientific units.

	fps units mks ur		
Force	pound (lb)	newton (N)	
Mass	slug	kilogram (kg)	
Distance	foot (ft)	meter (m)	
Time	second (s)	second (s)	
g	32 ft/s^2	9.8 m/s^2	

The last line of this table gives values for the gravitational acceleration g at the surface of the earth. Although these approximate values will suffice for most examples and problems, more precise values are 9.7805 m/s² and 32.088 ft/s² (at sea level at the equator).

Both systems are compatible with Newton's second law F = ma. Thus 1 N is (by definition) the force required to impart an acceleration of 1 m/s² to a mass of 1 kg. Similarly, 1 slug is (by definition) the mass that experiences an acceleration of 1 ft/s² under a force of 1 lb. (We will use mks units in all problems requiring mass units and thus will rarely need slugs to measure mass.)

Inches and centimeters (as well as miles and kilometers) also are commonly used in describing distances. For conversions between fps and mks units it helps to remember that

1 in. = 2.54 cm (exactly) and 1 lb
$$\approx$$
 4.448 N.

For instance,

1 ft = 12 in.
$$\times 2.54 \frac{\text{cm}}{\text{in.}} = 30.48 \text{ cm},$$

and it follows that

1 mi = 5280 ft × 30.48
$$\frac{\text{cm}}{\text{ft}}$$
 = 160934.4 cm \approx 1.609 km.

Thus a posted U.S. speed limit of 50 mi/h means that—in international terms—the legal speed limit is about $50 \times 1.609 \approx 80.45$ km/h.

Vertical Motion with Gravitational Acceleration

The weight W of a body is the force exerted on the body by gravity. Substitution of a = g and F = W in Newton's second law F = ma gives

$$W = mg \tag{14}$$

for the weight W of the mass m at the surface of the earth (where $g \approx 32 \text{ ft/s}^2 \approx 9.8 \text{ m/s}^2$). For instance, a mass of m = 20 kg has a weight of $W = (20 \text{ kg})(9.8 \text{ m/s}^2) = 196 \text{ N}$. Similarly, a mass m weighing 100 pounds has mks weight

$$W = (100 \text{ lb})(4.448 \text{ N/lb}) = 444.8 \text{ N},$$

so its mass is

$$m = \frac{W}{g} = \frac{444.8 \text{ N}}{9.8 \text{ m/s}^2} \approx 45.4 \text{ kg}.$$

To discuss vertical motion it is natural to choose the y-axis as the coordinate system for position, frequently with y = 0 corresponding to "ground level." If we choose the *upward* direction as the positive direction, then the effect of gravity on a vertically moving body is to decrease its height and also to decrease its velocity v = dy/dt. Consequently, if we ignore air resistance, then the acceleration a = dv/dt of the body is given by

$$\frac{dv}{dt} = -g. \tag{15}$$

This acceleration equation provides a starting point in many problems involving vertical motion. Successive integrations (as in Eqs. (10) and (11)) yield the velocity and height formulas

$$v(t) = -gt + v_0 \tag{16}$$

$$y(t) = -\frac{1}{2}gt^2 + v_0t + y_0.$$
 (17)

Here, y_0 denotes the initial (t = 0) height of the body and v_0 its initial velocity.

Projectile motion

(a) Suppose that a ball is thrown straight upward from the ground $(y_0 = 0)$ with initial velocity $v_0 = 96$ (ft/s, so we use g = 32 ft/s² in fps units). Then it reaches its maximum height when its velocity (Eq. (16)) is zero,

$$v(t) = -32t + 96 = 0$$

and thus when t = 3 s. Hence the maximum height that the ball attains is

$$y(3) = -\frac{1}{2} \cdot 32 \cdot 3^2 + 96 \cdot 3 + 0 = 144$$
 (ft)

(with the aid of Eq. (17)).

(b) If an arrow is shot straight upward from the ground with initial velocity $v_0 = 49$ (m/s, so we use g = 9.8 m/s² in mks units), then it returns to the ground when

$$y(t) = -\frac{1}{2} \cdot (9.8)t^2 + 49t = (4.9)t(-t + 10) = 0,$$

and thus after 10 s in the air.

A Swimmer's Problem

Figure 1.2.5 shows a northward-flowing river of width w = 2a. The lines $x = \pm a$ represent the banks of the river and the *y*-axis its center. Suppose that the velocity v_R at which the water flows increases as one approaches the center of the river, and indeed is given in terms of distance *x* from the center by

$$v_R = v_0 \left(1 - \frac{x^2}{a^2} \right). \tag{18}$$

You can use Eq. (18) to verify that the water does flow the fastest at the center, where $v_R = v_0$, and that $v_R = 0$ at each riverbank.



FIGURE 1.2.5. A swimmer's problem (Example 4).

Example 3

1.2 INTEGRALS AS GENERAL AND PARTICULAR SOLUTIONS

Suppose that a swimmer starts at the point (-a, 0) on the west bank and swims due east (relative to the water) with constant speed v_S . As indicated in Fig. 1.2.5, his velocity vector (relative to the riverbed) has horizontal component v_S and vertical component v_R . Hence the swimmer's direction angle α is given by

$$\tan \alpha = \frac{v_R}{v_S}.$$

Because $\tan \alpha = dy/dx$, substitution using (18) gives the differential equation

$$\frac{dy}{dx} = \frac{v_0}{v_S} \left(1 - \frac{x^2}{a^2} \right) \tag{19}$$

for the swimmer's trajectory y = y(x) as he crosses the river.

River crossing Suppose that the river is 1 mile wide and that its midstream velocity is $v_0 = 9$ mi/h. If the swimmer's velocity is $v_S = 3$ mi/h, then Eq. (19) takes the form

$$\frac{dy}{dx} = 3(1-4x^2).$$

Integration yields

$$y(x) = \int (3 - 12x^2) \, dx = 3x - 4x^3 + C$$

for the swimmer's trajectory. The initial condition $y\left(-\frac{1}{2}\right) = 0$ yields C = 1, so

$$y(x) = 3x - 4x^3 + 1$$

Then

$$y\left(\frac{1}{2}\right) = 3\left(\frac{1}{2}\right) - 4\left(\frac{1}{2}\right)^3 + 1 = 2,$$

so the swimmer drifts 2 miles downstream while he swims 1 mile across the river.

1.2 **Problems**

In Problems 1 through 10, find a function y = f(x) satisfying the given differential equation and the prescribed initial condition.

1. $\frac{dy}{dx} = 2x + 1; y(0) = 3$ 2. $\frac{dy}{dx} = (x - 2)^2; y(2) = 1$ 3. $\frac{dy}{dx} = \sqrt{x}; y(4) = 0$ 4. $\frac{dy}{dx} = \frac{1}{x^2}; y(1) = 5$ 5. $\frac{dy}{dx} = \frac{1}{\sqrt{x + 2}}; y(2) = -1$ 6. $\frac{dy}{dx} = x\sqrt{x^2 + 9}; y(-4) = 0$ 7. $\frac{dy}{dx} = \frac{10}{x^2 + 1}; y(0) = 0$ 8. $\frac{dy}{dx} = \cos 2x; y(0) = 1$ 9. $\frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}}; y(0) = 0$ 10. $\frac{dy}{dx} = xe^{-x}; y(0) = 1$ In Problems 11 through 18, find the position function x(t) of a moving particle with the given acceleration a(t), initial position $x_0 = x(0)$, and initial velocity $v_0 = v(0)$.

11. $a(t) = 50, v_0 = 10, x_0 = 20$ **12.** $a(t) = -20, v_0 = -15, x_0 = 5$ **13.** $a(t) = 3t, v_0 = 5, x_0 = 0$ **14.** $a(t) = 2t + 1, v_0 = -7, x_0 = 4$ **15.** $a(t) = 4(t + 3)^2, v_0 = -1, x_0 = 1$ **16.** $a(t) = \frac{1}{\sqrt{t+4}}, v_0 = -1, x_0 = 1$ **17.** $a(t) = \frac{1}{(t+1)^3}, v_0 = 0, x_0 = 0$ **18.** $a(t) = 50 \sin 5t, v_0 = -10, x_0 = 8$

Velocity Given Graphically

In Problems 19 through 22, a particle starts at the origin and travels along the x-axis with the velocity function v(t) whose graph is shown in Figs. 1.2.6 through 1.2.9. Sketch the graph of the resulting position function x(t) for $0 \le t \le 10$.



FIGURE 1.2.9. Graph of the velocity function v(t) of Problem 22.

Problems 23 through 28 explore the motion of projectiles under constant acceleration or deceleration.

- **23.** What is the maximum height attained by the arrow of part (b) of Example 3?
- **24.** A ball is dropped from the top of a building 400 ft high. How long does it take to reach the ground? With what speed does the ball strike the ground?
- **25.** The brakes of a car are applied when it is moving at 100 km/h and provide a constant deceleration of 10 meters per second per second (m/s^2) . How far does the car travel before coming to a stop?
- 26. A projectile is fired straight upward with an initial velocity of 100 m/s from the top of a building 20 m high and falls to the ground at the base of the building. Find (a) its maximum height above the ground; (b) when it passes the top of the building; (c) its total time in the air.
- **27.** A ball is thrown straight downward from the top of a tall building. The initial speed of the ball is 10 m/s. It strikes the ground with a speed of 60 m/s. How tall is the building?
- **28.** A baseball is thrown straight downward with an initial speed of 40 ft/s from the top of the Washington Monument (555 ft high). How long does it take to reach the ground, and with what speed does the baseball strike the ground?
- **29.** Variable acceleration A diesel car gradually speeds up so that for the first 10 s its acceleration is given by

$$\frac{dv}{dt} = (0.12)t^2 + (0.6)t \quad (\text{ft/s}^2).$$

If the car starts from rest ($x_0 = 0$, $v_0 = 0$), find the distance it has traveled at the end of the first 10 s and its velocity at that time.

Problems 30 through 32 explore the relation between the speed of an auto and the distance it skids when the brakes are applied.

- **30.** A car traveling at 60 mi/h (88 ft/s) skids 176 ft after its brakes are suddenly applied. Under the assumption that the braking system provides constant deceleration, what is that deceleration? For how long does the skid continue?
- **31.** The skid marks made by an automobile indicated that its brakes were fully applied for a distance of 75 m before it came to a stop. The car in question is known to have a constant deceleration of 20 m/s^2 under these conditions. How fast—in km/h—was the car traveling when the brakes were first applied?
- **32.** Suppose that a car skids 15 m if it is moving at 50 km/h when the brakes are applied. Assuming that the car has the same constant deceleration, how far will it skid if it is moving at 100 km/h when the brakes are applied?

Problems 33 and 34 explore vertical motion on a planet with gravitational acceleration different than the Earth's.

- **33.** On the planet Gzyx, a ball dropped from a height of 20 ft hits the ground in 2 s. If a ball is dropped from the top of a 200-ft-tall building on Gzyx, how long will it take to hit the ground? With what speed will it hit?
- **34.** A person can throw a ball straight upward from the surface of the earth to a maximum height of 144 ft. How high could this person throw the ball on the planet Gzyx of Problem 33?
- **35.** Velocity in terms of height A stone is dropped from rest at an initial height *h* above the surface of the earth. Show that the speed with which it strikes the ground is $v = \sqrt{2gh}$.
- **36.** Varying gravitational acceleration Suppose a woman has enough "spring" in her legs to jump (on earth) from the ground to a height of 2.25 feet. If she jumps straight upward with the same initial velocity on the moon—where the surface gravitational acceleration is (approximately) 5.3 ft/s^2 —how high above the surface will she rise?
- 37. At noon a car starts from rest at point A and proceeds at constant acceleration along a straight road toward point B. If the car reaches B at 12:50 P.M. with a velocity of 60 mi/h, what is the distance from A to B?
- **38.** At noon a car starts from rest at point *A* and proceeds with constant acceleration along a straight road toward point *C*, 35 miles away. If the constantly accelerated car arrives at *C* with a velocity of 60 mi/h, at what time does it arrive at *C*?
- **39.** River crossing If a = 0.5 mi and $v_0 = 9$ mi/h as in Example 4, what must the swimmer's speed v_S be in order that he drifts only 1 mile downstream as he crosses the river?
- **40.** River crossing Suppose that a = 0.5 mi, $v_0 = 9$ mi/h, and $v_S = 3$ mi/h as in Example 4, but that the velocity of the river is given by the fourth-degree function

$$v_R = v_0 \left(1 - \frac{x^4}{a^4} \right)$$

rather than the quadratic function in Eq. (18). Now find how far downstream the swimmer drifts as he crosses the river.

- **41. Interception of bomb** A bomb is dropped from a helicopter hovering at an altitude of 800 feet above the ground. From the ground directly beneath the helicopter, a projectile is fired straight upward toward the bomb, exactly 2 seconds after the bomb is released. With what initial velocity should the projectile be fired in order to hit the bomb at an altitude of exactly 400 feet?
- **42.** Lunar lander A spacecraft is in free fall toward the surface of the moon at a speed of 1000 mph (mi/h). Its retrorockets, when fired, provide a constant deceleration of 20,000 mi/h². At what height above the lunar surface should the astronauts fire the retrorockets to insure a soft touchdown? (As in Example 2, ignore the moon's gravitational field.)
- **43.** Solar wind Arthur Clarke's *The Wind from the Sun* (1963) describes Diana, a spacecraft propelled by the solar wind. Its aluminized sail provides it with a constant acceleration of $0.001g = 0.0098 \text{ m/s}^2$. Suppose this spacecraft starts from rest at time t = 0 and simultaneously fires a projectile (straight ahead in the same direction) that travels at one-tenth of the speed $c = 3 \times 10^8 \text{ m/s}$ of light. How long will it take the spacecraft to catch up with the projectile, and how far will it have traveled by then?
- **44.** Length of skid A driver involved in an accident claims he was going only 25 mph. When police tested his car, they found that when its brakes were applied at 25 mph, the car skidded only 45 feet before coming to a stop. But the driver's skid marks at the accident scene measured 210 feet. Assuming the same (constant) deceleration, determine the speed he was actually traveling just prior to the accident.
- **45.** Kinematic formula Use Eqs. (10) and (11) to show that $v(t)^2 v_0^2 = 2a[x(t) x_0]$ for all t when the acceleration a = dv/dt is constant. Then use this "kinematic formula"—commonly presented in introductory physics courses—to confirm the result of Example 2.

1.3 Slope Fields and Solution Curves

Consider a differential equation of the form

$$\frac{dy}{dx} = f(x, y) \tag{1}$$

where the right-hand function f(x, y) involves both the independent variable x and the dependent variable y. We might think of integrating both sides in (1) with respect to x, and hence write $y(x) = \int f(x, y(x)) dx + C$. However, this approach does not lead to a solution of the differential equation, because the indicated integral involves the *unknown* function y(x) itself, and therefore cannot be evaluated explicitly. Actually, there exists *no* straightforward procedure by which a general differential equation can be solved explicitly. Indeed, the solutions of such a simple-looking differential equation as $y' = x^2 + y^2$ cannot be expressed in terms of the ordinary

Example 1

elementary functions studied in calculus textbooks. Nevertheless, the graphical and numerical methods of this and later sections can be used to construct *approximate* solutions of differential equations that suffice for many practical purposes.

Slope Fields and Graphical Solutions

There is a simple geometric way to think about solutions of a given differential equation y' = f(x, y). At each point (x, y) of the xy-plane, the value of f(x, y) determines a slope m = f(x, y). A solution of the differential equation is simply a differentiable function whose graph y = y(x) has this "correct slope" at each point (x, y(x)) through which it passes—that is, y'(x) = f(x, y(x)). Thus a **solution curve** of the differential equation y' = f(x, y)—the graph of a solution of the equation—is simply a curve in the xy-plane whose tangent line at each point (x, y) has slope m = f(x, y). For instance, Fig. 1.3.1 shows a solution curve of the differential equation y' = x - y together with its tangent lines at three typical points.



FIGURE 1.3.1. A solution curve for the differential equation y' = x - y together with tangent lines having

- slope $m_1 = x_1 y_1$ at the point (x_1, y_1) ;
- slope $m_2 = x_2 y_2$ at the point (x_2, y_2) ; and
- slope $m_3 = x_3 y_3$ at the point (x_3, y_3) .

This geometric viewpoint suggests a graphical method for constructing approximate solutions of the differential equation y' = f(x, y). Through each of a representative collection of points (x, y) in the plane we draw a short line segment having the proper slope m = f(x, y). All these line segments constitute a **slope field** (or a **direction field**) for the equation y' = f(x, y).

Figures 1.3.2 (a)–(d) show slope fields and solution curves for the differential equation

$$\frac{ly}{lx} = ky \tag{2}$$

with the values k = 2, 0.5, -1, and -3 of the parameter k in Eq. (2). Note that each slope field yields important qualitative information about the set of all solutions of the differential equation. For instance, Figs. 1.3.2(a) and (b) suggest that each solution y(x) approaches $\pm \infty$ as $x \to +\infty$ if k > 0, whereas Figs. 1.3.2(c) and (d) suggest that $y(x) \to 0$ as $x \to +\infty$ if k < 0. Moreover, although the sign of k determines the *direction* of increase or decrease of y(x), its absolute value |k| appears to determine the *rate of change* of y(x). All this is apparent from slope fields like those in Fig. 1.3.2, even without knowing that the general solution of Eq. (2) is given explicitly by $y(x) = Ce^{kx}$.

A slope field suggests visually the general shapes of solution curves of the differential equation. Through each point a solution curve should proceed in such

1.3 SLOPE FIELDS AND SOLUTION CURVES 9



FIGURE 1.3.2(a) Slope field and solution curves for y' = 2y.



FIGURE 1.3.2(d) Slope field and solution curves for y' = -3y.



FIGURE 1.3.2(b) Slope field and solution curves for y' = (0.5)y.



FIGURE 1.3.2(c) Slope field and solution curves for y' = -y.

$x \setminus y$	-4	-3	-2	-1	0	1	2	3	4
-4	0	-1	-2	-3	-4	-5	-6	-7	-8
-3	1	0	-1	-2	-3	-4	-5	-6	-7
-2	2	1	0	-1	-2	-3	-4	-5	-6
-1	3	2	1	0	-1	-2	-3	-4	-5
0	4	3	2	1	0	-1	-2	-3	-4
1	5	4	3	2	1	0	-1	-2	-3
2	6	5	4	3	2	1	0	-1	-2
3	7	6	5	4	3	2	1	0	-1
4	8	7	6	5	4	3	2	1	0

FIGURE 1.3.3. Values of the slope y' = x - y for $-4 \le x$, $y \le 4$.

a direction that its tangent line is nearly parallel to the nearby line segments of the slope field. Starting at any initial point (a, b), we can attempt to sketch freehand an approximate solution curve that threads its way through the slope field, following the visible line segments as closely as possible.

Example 2 Construct a slope field for the differential equation y' = x - y and use it to sketch an approximate solution curve that passes through the point (-4, 4).

Solution Figure 1.3.3 shows a table of slopes for the given equation. The numerical slope m = x - y appears at the intersection of the horizontal *x*-row and the vertical *y*-column of the table. If you inspect the pattern of upper-left to lower-right diagonals in this table, you can see that it was easily and quickly constructed. (Of course, a more complicated function f(x, y) on the right-hand side of the differential equation would necessitate more complicated calculations.) Figure 1.3.4 shows the corresponding slope field, and Fig. 1.3.5 shows an approximate solution curve sketched through the point (-4, 4) so as to follow this slope field as closely as possible. At each point it appears to proceed in the direction indicated by the nearby line segments of the slope field.

Although a spreadsheet program (for instance) readily constructs a table of slopes as in Fig. 1.3.3, it can be quite tedious to plot by hand a sufficient number of slope segments as in Fig. 1.3.4. However, most computer algebra systems include commands for quick and ready construction of slope fields with as many line segments as desired; such commands are illustrated in the application material for this section. The more line segments are constructed, the more accurately solution curves can be visualized and sketched. Figure 1.3.6 shows a "finer" slope field for





FIGURE 1.3.4. Slope field for y' = x - ycorresponding to the table of slopes in Fig. 1.3.3.

FIGURE 1.3.5. The solution curve through (-4, 4).



FIGURE 1.3.6. Slope field and typical solution curves for y' = x - y.

the differential equation y' = x - y of Example 2, together with typical solution curves treading through this slope field.

If you look closely at Fig. 1.3.6, you may spot a solution curve that appears to be a straight line! Indeed, you can verify that the linear function y = x - 1 is a solution of the equation y' = x - y, and it appears likely that the other solution curves approach this straight line as an asymptote as $x \to +\infty$. This inference illustrates the fact that a slope field can suggest tangible information about solutions that is not at all evident from the differential equation itself. Can you, by tracing the appropriate solution curve in this figure, infer that $y(3) \approx 2$ for the solution y(x) of the initial value problem y' = x - y, y(-4) = 4?

Applications of Slope Fields

The next two examples illustrate the use of slope fields to glean useful information in physical situations that are modeled by differential equations. Example 3 is based on the fact that a baseball moving through the air at a moderate speed v (less than about 300 ft/s) encounters air resistance that is approximately proportional to v. If the baseball is thrown straight downward from the top of a tall building or from a hovering helicopter, then it experiences both the downward acceleration of gravity and an upward acceleration of air resistance. If the y-axis is directed downward, then the ball's velocity v = dy/dt and its gravitational acceleration g = 32 ft/s² are both positive, while its acceleration due to air resistance is negative. Hence its total acceleration is of the form

$$\frac{dv}{dt} = g - kv. \tag{3}$$

A typical value of the air resistance proportionality constant might be k = 0.16.



FIGURE 1.3.7. Slope field and typical solution curves for v' = 32 - 0.16v.

Suppose you throw a baseball straight downward from a helicopter hov-Falling baseball ering at an altitude of 3000 feet. You wonder whether someone standing on the ground below could conceivably catch it. In order to estimate the speed with which the ball will land, you can use your laptop's computer algebra system to construct a slope field for the differential equation

$$\frac{dv}{dt} = 32 - 0.16v.$$
 (4)

The result is shown in Fig. 1.3.7, together with a number of solution curves corresponding to different values of the initial velocity v(0) with which you might throw the baseball downward. Note that all these solution curves appear to approach the horizontal line v = 200as an asymptote. This implies that-however you throw it-the baseball should approach the *limiting velocity* v = 200 ft/s instead of accelerating indefinitely (as it would in the absence of any air resistance). The handy fact that 60 mi/h = 88 ft/s yields

$$v = 200\frac{\text{ft}}{\text{s}} \times \frac{60 \text{ mi/h}}{88 \text{ ft/s}} \approx 136.36 \frac{\text{mi}}{\text{h}}.$$

Perhaps a catcher accustomed to 100 mi/h fastballs would have some chance of fielding this speeding ball.

Comment If the ball's initial velocity is v(0) = 200, then Eq. (4) gives v'(0) = 32 - 200(0.16)(200) = 0, so the ball experiences *no* initial acceleration. Its velocity therefore remains unchanged, and hence $v(t) \equiv 200$ is a constant "equilibrium solution" of the differential equation. If the initial velocity is greater than 200, then the initial acceleration given by Eq. (4) is negative, so the ball slows down as it falls. But if the initial velocity is less than 200, then the initial acceleration given by (4) is positive, so the ball speeds up as it falls. It therefore seems quite reasonable that, because of air resistance, the baseball will approach a limiting velocity of 200 ft/s-whatever initial velocity it starts with. You might like to verify that—in the absence of air resistance—this ball would hit the ground at over 300 mi/h.

In Section 2.1 we will discuss in detail the logistic differential equation

$$\frac{dP}{dt} = kP(M-P) \tag{5}$$

(6)

that often is used to model a population P(t) that inhabits an environment with *carrying capacity M*. This means that M is the maximum population that this environment can sustain on a long-term basis (in terms of the maximum available food, for instance).

Limiting population If we take k = 0.0004 and M = 150, then the logistic equation in (5) takes the form $\frac{dP}{dt} = 0.0004P(150 - P) = 0.06P - 0.0004P^2.$



FIGURE 1.3.8. Slope field and typical solution curves for $P' = 0.06P - 0.0004P^2.$

The positive term 0.06P on the right in (6) corresponds to natural growth at a 6% annual rate (with time t measured in years). The negative term $-0.0004P^2$ represents the inhibition of growth due to limited resources in the environment.

Figure 1.3.8 shows a slope field for Eq. (6), together with a number of solution curves corresponding to possible different values of the initial population P(0). Note that all these solution curves appear to approach the horizontal line P = 150 as an asymptote. This implies that—whatever the initial population—the population P(t) approaches the *limiting popula*tion P = 150 as $t \to \infty$.

Comment If the initial population is P(0) = 150, then Eq. (6) gives

$$P'(0) = 0.0004(150)(150 - 150) = 0,$$

so the population experiences no initial (instantaneous) change. It therefore remains unchanged, and hence $P(t) \equiv 150$ is a constant "equilibrium solution" of the differential equation. If the initial population is greater than 150, then the initial rate of change given by (6) is negative, so the population immediately begins to decrease. But if the initial population is less than 150, then the initial rate of change given by (6) is positive, so the population immediately begins to increase. It therefore seems quite reasonable to conclude that the population will approach a limiting value of 150-whatever the (positive) initial population.

Existence and Uniqueness of Solutions

Before one spends much time attempting to solve a given differential equation, it is wise to know that solutions actually exist. We may also want to know whether there is only one solution of the equation satisfying a given initial condition—that is, whether its solutions are unique.
Example 5 (a) **Failure of existence** The initial value problem

$$y' = \frac{1}{x}, \quad y(0) = 0$$
 (7)

has *no* solution, because no solution $y(x) = \int (1/x) dx = \ln |x| + C$ of the differential equation is defined at x = 0. We see this graphically in Fig. 1.3.9, which shows a direction field and some typical solution curves for the equation y' = 1/x. It is apparent that the indicated direction field "forces" all solution curves near the y-axis to plunge downward so that none can pass through the point (0, 0).



FIGURE 1.3.9. Direction field and typical solution curves for the equation y' = 1/x.

FIGURE 1.3.10. Direction field and two different solution curves for the initial value problem $y' = 2\sqrt{y}$, y(0) = 0.

(b) Failure of uniqueness On the other hand, you can readily verify that the initial value problem

$$y' = 2\sqrt{y}, \quad y(0) = 0$$
 (8)

has the *two* different solutions $y_1(x) = x^2$ and $y_2(x) \equiv 0$ (see Problem 27). Figure 1.3.10 shows a direction field and these two different solution curves for the initial value problem in (8). We see that the curve $y_1(x) = x^2$ threads its way through the indicated direction field, whereas the differential equation $y' = 2\sqrt{y}$ specifies slope y' = 0 along the *x*-axis $y_2(x) = 0$.

Example 5 illustrates the fact that, before we can speak of "the" solution of an initial value problem, we need to know that it has *one and only one* solution. Questions of existence and uniqueness of solutions also bear on the process of mathematical modeling. Suppose that we are studying a physical system whose behavior is completely determined by certain initial conditions, but that our proposed mathematical model involves a differential equation *not* having a unique solution satisfying those conditions. This raises an immediate question as to whether the mathematical model adequately represents the physical system.

The theorem stated below implies that the initial value problem y' = f(x, y), y(a) = b has one and only one solution defined near the point x = a on the x-axis, provided that both the function f and its partial derivative $\partial f/\partial y$ are continuous near the point (a, b) in the xy-plane. Methods of proving existence and uniqueness theorems are discussed in the Appendix.



FIGURE 1.3.11. The rectangle *R* and *x*-interval *I* of Theorem 1, and the solution curve y = y(x) through the point (a, b).



FIGURE 1.3.12. The solution curve through the initial point (0, 1) leaves the rectangle *R* before it reaches the right side of *R*.

THEOREM 1 Existence and Uniqueness of Solutions

Suppose that both the function f(x, y) and its partial derivative $D_y f(x, y)$ are continuous on some rectangle *R* in the *xy*-plane that contains the point (a, b) in its interior. Then, for some open interval *I* containing the point *a*, the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(a) = b \tag{9}$$

has one and only one solution that is defined on the interval I. (As illustrated in Fig. 1.3.11, the solution interval I may not be as "wide" as the original rectangle R of continuity; see Remark 3 below.)

Remark 1 In the case of the differential equation dy/dx = -y of Example 1 and Fig. 1.3.2(c), both the function f(x, y) = -y and the partial derivative $\partial f/\partial y = -1$ are continuous everywhere, so Theorem 1 implies the existence of a unique solution for any initial data (a, b). Although the theorem ensures existence only on some open interval containing x = a, each solution $y(x) = Ce^{-x}$ actually is defined for all x.

Remark 2 In the case of the differential equation $dy/dx = 2\sqrt{y}$ of Example 5(b) and Eq. (8), the function $f(x, y) = 2\sqrt{y}$ is continuous wherever y > 0, but the partial derivative $\partial f/\partial y = 1/\sqrt{y}$ is discontinuous when y = 0, and hence at the point (0, 0). This is why it is possible for there to exist two different solutions $y_1(x) = x^2$ and $y_2(x) \equiv 0$, each of which satisfies the initial condition y(0) = 0.

Remark 3 In Example 7 of Section 1.1 we examined the especially simple differential equation $dy/dx = y^2$. Here we have $f(x, y) = y^2$ and $\partial f/\partial y = 2y$. Both of these functions are continuous everywhere in the *xy*-plane, and in particular on the rectangle -2 < x < 2, 0 < y < 2. Because the point (0, 1) lies in the interior of this rectangle, Theorem 1 guarantees a unique solution—necessarily a continuous function—of the initial value problem

$$\frac{y}{x} = y^2, \quad y(0) = 1$$
 (10)

on some open x-interval containing a = 0. Indeed this is the solution

 $\frac{d}{d}$

)

$$v(x) = \frac{1}{1-x}$$

that we discussed in Example 7. But y(x) = 1/(1-x) is discontinuous at x = 1, so our unique continuous solution does not exist on the entire interval -2 < x < 2. Thus the solution interval I of Theorem 1 may not be as wide as the rectangle R where f and $\partial f/\partial y$ are continuous. Geometrically, the reason is that the solution curve provided by the theorem may leave the rectangle—wherein solutions of the differential equation are guaranteed to exist—before it reaches the one or both ends of the interval (see Fig. 1.3.12).

The following example shows that, if the function f(x, y) and/or its partial derivative $\partial f/\partial y$ fail to satisfy the continuity hypothesis of Theorem 1, then the initial value problem in (9) may have *either* no solution *or* many—even infinitely many—solutions.

Example 6

Consider the first-order differential equation

$$x\frac{dy}{dx} = 2y.$$
 (11)

Applying Theorem 1 with f(x, y) = 2y/x and $\partial f/\partial y = 2/x$, we conclude that Eq. (11) must have a unique solution near any point in the *xy*-plane where $x \neq 0$. Indeed, we see immediately by substitution in (11) that

v

$$(x) = Cx^2 \tag{12}$$



FIGURE 1.3.13. There are infinitely many solution curves through the point (0, 0), but no solution curves through the point (0, b) if $b \neq 0$.

satisfies Eq. (11) for any value of the constant C and for all values of the variable x. In particular, the initial value problem

$$x\frac{dy}{dx} = 2y, \quad y(0) = 0$$
 (13)

has infinitely many different solutions, whose solution curves are the parabolas $y = Cx^2$ illustrated in Fig. 1.3.13. (In case C = 0 the "parabola" is actually the x-axis y = 0.)

Observe that all these parabolas pass through the origin (0, 0), but none of them passes through any other point on the *y*-axis. It follows that the initial value problem in (13) has infinitely many solutions, but the initial value problem

$$x\frac{dy}{dx} = 2y, \quad y(0) = b \tag{14}$$

has no solution if $b \neq 0$.

Finally, note that through any point off the *y*-axis there passes only one of the parabolas $y = Cx^2$. Hence, if $a \neq 0$, then the initial value problem

$$x\frac{dy}{dx} = 2y, \quad y(a) = b \tag{15}$$

has a unique solution on any interval that contains the point x = a but not the origin x = 0. In summary, the initial value problem in (15) has

- a unique solution near (a, b) if $a \neq 0$;
- no solution if a = 0 but $b \neq 0$;
- infinitely many solutions if a = b = 0.

Still more can be said about the initial value problem in (15). Consider a typical initial point off the *y*-axis—for instance the point (-1, 1) indicated in Fig. 1.3.14. Then for any value of the constant *C* the function defined by

$$y(x) = \begin{cases} x^2 & \text{if } x \le 0, \\ Cx^2 & \text{if } x > 0 \end{cases}$$
(16)



FIGURE 1.3.14. There are infinitely many solution curves through the point (1, -1).

is continuous and satisfies the initial value problem

$$x\frac{dy}{dx} = 2y, \quad y(-1) = 1.$$
 (17)

For a particular value of *C*, the solution curve defined by (16) consists of the left half of the parabola $y = x^2$ and the right half of the parabola $y = Cx^2$. Thus the unique solution curve near (-1, 1) branches at the origin into the infinitely many solution curves illustrated in Fig. 1.3.14.

We therefore see that Theorem 1 (if its hypotheses are satisfied) guarantees uniqueness of the solution near the initial point (a, b), but a solution curve through (a, b) may eventually branch elsewhere so that uniqueness is lost. Thus a solution may exist on a larger interval than one on which the solution is unique. For instance, the solution $y(x) = x^2$ of the initial value problem in (17) exists on the whole x-axis, but this solution is unique only on the negative x-axis $-\infty < x < 0$.

1.3 **Problems**

In Problems 1 through 10, we have provided the slope field of the indicated differential equation, together with one or more solution curves. Sketch likely solution curves through the additional points marked in each slope field.







FIGURE 1.3.16.











FIGURE 1.3.19.



















FIGURE 1.3.24.

A more detailed version of Theorem 1 says that, if the function f(x, y) is continuous near the point (a, b), then at least one solution of the differential equation y' = f(x, y) exists on some open interval I containing the point x = a and, moreover, that if in addition the partial derivative $\partial f/\partial y$ is continuous near (a, b), then this solution is unique on some (perhaps smaller) interval J. In Problems 11 through 20, determine whether existence of at least one solution of the given initial value problem is thereby guaranteed and, if so, whether uniqueness of that solution is guaranteed.

11.
$$\frac{dy}{dx} = 2x^2y^2; \quad y(1) = -1$$

12. $\frac{dy}{dx} = x \ln y; \quad y(1) = 1$
13. $\frac{dy}{dx} = \sqrt[3]{y}; \quad y(0) = 1$
14. $\frac{dy}{dx} = \sqrt[3]{y}; \quad y(0) = 0$
15. $\frac{dy}{dx} = \sqrt{x - y}; \quad y(2) = 2$
16. $\frac{dy}{dx} = \sqrt{x - y}; \quad y(2) = 1$
17. $y\frac{dy}{dx} = x - 1; \quad y(0) = 1$
18. $y\frac{dy}{dx} = x - 1; \quad y(1) = 0$
19. $\frac{dy}{dx} = \ln(1 + y^2); \quad y(0) = 0$
20. $\frac{dy}{dx} = x^2 - y^2; \quad y(0) = 1$

In Problems 21 and 22, first use the method of Example 2 to construct a slope field for the given differential equation. Then sketch the solution curve corresponding to the given initial condition. Finally, use this solution curve to estimate the desired value of the solution y(x). **21.** y' = x + y, y(0) = 0; y(-4) = ?**22.** y' = y - x, y(4) = 0; y(-4) = ?

Problems 23 and 24 are like Problems 21 and 22, but now use a computer algebra system to plot and print out a slope field for the given differential equation. If you wish (and know how), you can check your manually sketched solution curve by plotting it with the computer.

23.
$$y' = x^2 + y^2 - 1$$
, $y(0) = 0$; $y(2) = ?$

24. $y' = x + \frac{1}{2}y^2$, y(-2) = 0; y(2) = ?

25. Falling parachutist You bail out of the helicopter of Example 3 and pull the ripcord of your parachute. Now k = 1.6 in Eq. (3), so your downward velocity satisfies the initial value problem

$$\frac{dv}{dt} = 32 - 1.6v, \quad v(0) = 0.$$

In order to investigate your chances of survival, construct a slope field for this differential equation and sketch the appropriate solution curve. What will your limiting velocity be? Will a strategically located haystack do any good? How long will it take you to reach 95% of your limiting velocity?

26. Deer population Suppose the deer population P(t) in a small forest satisfies the logistic equation

$$\frac{dP}{dt} = 0.0225P - 0.0003P^2.$$

Construct a slope field and appropriate solution curve to answer the following questions: If there are 25 deer at time t = 0 and t is measured in months, how long will it take the number of deer to double? What will be the limiting deer population?

The next seven problems illustrate the fact that, if the hypotheses of Theorem 1 are not satisfied, then the initial value problem y' = f(x, y), y(a) = b may have either no solutions, finitely many solutions, or infinitely many solutions.

27. (a) Verify that if *c* is a constant, then the function defined piecewise by

$$y(x) = \begin{cases} 0 & \text{for } x \leq c, \\ (x-c)^2 & \text{for } x > c \end{cases}$$

satisfies the differential equation $y' = 2\sqrt{y}$ for all x (including the point x = c). Construct a figure illustrating the fact that the initial value problem $y' = 2\sqrt{y}$, y(0) = 0 has infinitely many different solutions. (b) For what values of b does the initial value problem $y' = 2\sqrt{y}$, y(0) = b have (i) no solution, (ii) a unique solution that is defined for all x?

28. Verify that if k is a constant, then the function $y(x) \equiv kx$ satisfies the differential equation xy' = y for all x. Construct a slope field and several of these straight line solution curves. Then determine (in terms of a and b) how many different solutions the initial value problem xy' = y, y(a) = b has—one, none, or infinitely many.

29. Verify that if c is a constant, then the function defined piecewise by

$$y(x) = \begin{cases} 0 & \text{for } x \leq c, \\ (x-c)^3 & \text{for } x > c \end{cases}$$

satisfies the differential equation $y' = 3y^{2/3}$ for all x. Can you also use the "left half" of the cubic $y = (x - c)^3$ in piecing together a solution curve of the differential equation? (See Fig. 1.3.25.) Sketch a variety of such solution curves. Is there a point (a, b) of the xy-plane such that the initial value problem $y' = 3y^{2/3}$, y(a) = b has either no solution or a unique solution that is defined for all x? Reconcile your answer with Theorem 1.



FIGURE 1.3.25. A suggestion for Problem 29.

30. Verify that if c is a constant, then the function defined piecewise by

$$y(x) = \begin{cases} +1 & \text{if } x \leq c, \\ \cos(x-c) & \text{if } c < x < c + \pi, \\ -1 & \text{if } x \geq c + \pi \end{cases}$$

satisfies the differential equation $y' = -\sqrt{1-y^2}$ for all x. (Perhaps a preliminary sketch with c = 0 will be helpful.) Sketch a variety of such solution curves. Then determine (in terms of a and b) how many different solutions the initial value problem $y' = -\sqrt{1-y^2}$, y(a) = b has.

- **31.** Carry out an investigation similar to that in Problem 30, except with the differential equation $y' = +\sqrt{1-y^2}$. Does it suffice simply to replace $\cos(x-c)$ with $\sin(x-c)$ in piecing together a solution that is defined for all x?
- **32.** Verify that if c > 0, then the function defined piecewise by

$$y(x) = \begin{cases} 0 & \text{if } x^2 \leq c, \\ (x^2 - c)^2 & \text{if } x^2 > c \end{cases}$$

satisfies the differential equation $y' = 4x\sqrt{y}$ for all x. Sketch a variety of such solution curves for different values of c. Then determine (in terms of a and b) how many different solutions the initial value problem $y' = 4x\sqrt{y}$, y(a) = b has.

28 CHAPTER 1 FIRST-ORDER DIFFERENTIAL EQUATIONS

33. If $c \neq 0$, verify that the function defined by y(x) = x/(cx - 1) (with the graph illustrated in Fig. 1.3.26) satisfies the differential equation $x^2y' + y^2 = 0$ if $x \neq 1/c$. Sketch a variety of such solution curves for different values of *c*. Also, note the constant-valued function $y(x) \equiv 0$ that does not result from any choice of the constant *c*. Finally, determine (in terms of *a* and *b*) how many different solutions the initial value problem $x^2y' + y^2 = 0$, y(a) = b has.



FIGURE 1.3.26. Slope field for $x^2y' + y^2 = 0$ and graph of a solution y(x) = x/(cx - 1).

- 34. (a) Use the direction field of Problem 5 to estimate the values at x = 1 of the two solutions of the differential equation y' = y x + 1 with initial values y(-1) = -1.2 and y(-1) = -0.8.
 - (b) Use a computer algebra system to estimate the values at x = 3 of the two solutions of this differential equation with initial values y(-3) = -3.01 and y(-3) = -2.99.

The lesson of this problem is that small changes in initial conditions can make big differences in results.

- **35.** (a) Use the direction field of Problem 6 to estimate the values at x = 2 of the two solutions of the differential equation y' = x y + 1 with initial values y(-3) = -0.2 and y(-3) = +0.2.
 - (b) Use a computer algebra system to estimate the values at x = 2 of the two solutions of this differential equation with initial values y(-3) = -0.5 and y(-3) = +0.5.

The lesson of this problem is that big changes in initial conditions may make only small differences in results.

1.3 Application Computer-Generated Slope Fields and Solution Curves

Go to goo.gl/QjGAkl to download this application's computing resources including *Maple/Mathematica/*MATLAB. Widely available computer algebra systems and technical computing environments include facilities to automate the construction of slope fields and solution curves, as do some graphing calculators (see Figs. 1.3.27–29).

The Expanded Applications site at the URL indicated in the margin includes a discussion of *Maple*TM, *Mathematica*TM, and MATLABTM resources for the investigation of differential equations. For instance, the *Maple* command

```
with(DEtools):
DEplot(diff(y(x),x)=sin(x-y(x)), y(x), x=-5..5, y=-5..5);
```

and the Mathematica command

VectorPlot[{1, Sin[x-y]}, {x, -5, 5}, {y, -5, 5}]

produce slope fields similar to the one shown in Fig. 1.3.29. Figure 1.3.29 itself was generated with the MATLAB program dfield [John Polking and David Arnold, *Or*dinary Differential Equations Using MATLAB, 3rd edition, Hoboken, NJ: Pearson, 2003] that is freely available for educational use (math.rice.edu/~dfield). This web site also provides a stand-alone Java version of dfield that can be used in a web browser. When a differential equation is entered in the dfield setup menu (Fig. 1.3.30), you can (with mouse button clicks) plot both a slope field and the solution curve (or curves) through any desired point (or points). Another freely available and user-friendly MATLAB-based ODE package with impressive graphical capabilities is Iode (www.math.uiuc.edu/iode).

Modern technology platforms offer even further interactivity by allowing the user to vary initial conditions and other parameters "in real time." *Mathematica*'s **Manipulate** command was used to generate Fig. 1.3.31, which shows three particular solutions of the differential equation $dy/dx = \sin(x - y)$. The solid curve corresponds to the initial condition y(1) = 0. As the "locator point" initially at



FIGURE 1.3.27. TI-84 Plus CE[™] graphing calculator and TI-Nspire[™] CX CAS handheld. Screenshot from Texas Instruments Incorporated. Courtesy of Texas Instruments Incorporated.



FIGURE 1.3.28. Slope field and solution curves for the differential equation

 $\frac{dy}{dx} = \sin(x - y)$ with initial points (0, b), b = -3, -1, -2, 0, 2, 4 and window -5 $\leq x, y \leq 5$ on a TI-89 graphing calculator.

(1,0) is dragged—by mouse or touchpad—to the point (0, 3) or (2, -2), the solution curve immediately follows, resulting in the dashed curves shown. The TI-NspireTM CX CAS has similar capability; indeed, as Fig. 1.3.28 appears on the Nspire display, each of the initial points (0, *b*) can be dragged to different locations using the Nspire's touchpad, with the corresponding solution curves being instantly redrawn.

DFIELD Setup				- 🗆 ×
File Edit Window				
	The differential	equation.		
$y ' = \int \sin(x \cdot y)$				
The independent variable is x	Ť			
Parameters: =			=	
	The display v	window.		
The minimum value of x =	-5	The minimum value of y =		-5
The maximum value of x =	5	The maximum value of y = 5		
Quit	Revert		Proceed	

FIGURE 1.3.30. MATLAB **dfield** setup to construct slope field and solution curves for $y' = \sin(x - y)$.

Use a graphing calculator or computer system in the following investigations. You might warm up by generating the slope fields and some solution curves for Problems 1 through 10 in this section.

INVESTIGATION A: Plot a slope field and typical solution curves for the differential equation $dy/dx = \sin(x - y)$, but with a larger window than that of Fig. 1.3.29. With $-10 \le x \le 10, -10 \le y \le 10$, for instance, a number of apparent straight line solution curves should be visible, especially if your display allows you to drag the initial point interactively from upper left to lower right.

- (a) Substitute y = ax + b in the differential equation to determine what the coefficients *a* and *b* must be in order to get a solution. Are the results consistent with what you see on the display?
- (b) A computer algebra system gives the general solution

$$y(x) = x - 2 \tan^{-1} \left(\frac{x - 2 - C}{x - C} \right).$$



FIGURE 1.3.29. Computergenerated slope field and solution curves for the differential equation $y' = \sin(x - y)$.



FIGURE 1.3.31. Interactive *Mathematica* solution of the differential equation $y' = \sin(x - y)$. The "locator point" corresponding to the initial condition y(1) = 0 can be dragged to any other point in the display, causing the solution curve to be automatically redrawn.

Plot this solution with selected values of the constant *C* to compare the resulting solution curves with those indicated in Fig. 1.3.28. Can you see that *no* value of *C* yields the linear solution $y = x - \pi/2$ corresponding to the initial condition $y(\pi/2) = 0$? Are there any values of *C* for which the corresponding solution curves lie close to this straight line solution curve?

INVESTIGATION B: For your own personal investigation, let n be the *smallest* digit in your student ID number that is greater than 1, and consider the differential equation

$$\frac{dy}{dx} = \frac{1}{n}\cos(x - ny)$$

- (a) First investigate (as in part (a) of Investigation A) the possibility of straight line solutions.
- (b) Then generate a slope field for this differential equation, with the viewing window chosen so that you can picture some of these straight lines, plus a sufficient number of nonlinear solution curves that you can formulate a conjecture about what happens to y(x) as x → +∞. State your inference as plainly as you can. Given the initial value y(0) = y₀, try to predict (perhaps in terms of y₀) how y(x) behaves as x → +∞.
- (c) A computer algebra system gives the general solution

$$y(x) = \frac{1}{n} \left[x + 2 \tan^{-1} \left(\frac{1}{x - C} \right) \right].$$

Can you make a connection between this symbolic solution and your graphically generated solution curves (straight lines or otherwise)?

1.4 Separable Equations and Applications

In the preceding sections we saw that if the function f(x, y) does not involve the variable y, then solving the first-order differential equation

$$\frac{dy}{dx} = f(x, y) \tag{1}$$

is a matter of simply finding an antiderivative. For example, the general solution of

$$\frac{dy}{dx} = -6x\tag{2}$$

is given by

$$y(x) = \int -6x \, dx = -3x^2 + C.$$

If instead f(x, y) does involve the dependent variable y, then we can no longer solve the equation merely by integrating both sides: The differential equation

$$\frac{dy}{dx} = -6xy \tag{3a}$$

differs from Eq. (2) only in the factor y appearing on the right-hand side, but this is enough to prevent us from using the same approach to solve Eq. (3a) that was successful with Eq. (2).

1.4 SEPARABLE EQUATIONS AND APPLICATIONS31

And yet, as we will see throughout the remainder of this chapter, differential equations like (3a) often can, in fact, be solved by methods which are based on the idea of "integrating both sides." The idea behind these techniques is to rewrite the given equation in a form that, while equivalent to the given equation, allows both sides to be integrated directly, thus leading to the solution of the original differential equation.

The most basic of these methods, *separation of variables*, can be applied to Eq. (3a). First, we note that the right-hand function f(x, y) = -6xy can be viewed as the *product* of two expressions, one involving only the independent variable x, and the other involving only the dependent variable y:

$$\frac{dy}{dx} = (-6x) \cdot y. \tag{3b}$$

Next, we informally break up the derivative dy/dx into the "free-floating" differentials dx and dy—a notational convenience that leads to correct results, as we will see below—and then multiply by dx and divide by y in Eq. (3b), leading to

$$\frac{dy}{y} = -6x \, dx. \tag{3c}$$

Equation (3c) is an equivalent version of the original differential equation in (3a), but with the variables x and y separated (that is, by the equal sign), and this is what allows us to integrate both sides. The left-hand side is integrated with respect to y (with no "interference" from the variable x), and vice versa for the right-hand side. This leads to

$$\int \frac{dy}{y} = \int -6x \, dx,$$

$$\ln|y| = -3x^2 + C.$$
(4)

or

This gives the general solution of Eq. (3a) *implicitly*, and a family of solution curves is shown in Fig. 1.4.1.

In this particular case we can go on to solve for *y* to give the *explicit* general solution

$$y(x) = \pm e^{-3x^2 + C} = \pm e^{-3x^2} e^C = A e^{-3x^2},$$
(5)

where A represents the constant $\pm e^{C}$, which can take on any nonzero value. If we impose an initial condition on Eq. (3a), say y(0) = 7, then in Eq. (5) we find that A = 7, yielding the particular solution

$$y(x) = 7e^{-3x^2}$$

which is the upper emphasized solution curve shown in Fig. 1.4.1. In the same way, the initial condition y(0) = -4 leads to the particular solution

$$v(x) = -4e^{-3x^2},$$

which is the lower emphasized solution curve shown in Fig. 1.4.1.

To complete this example, we note that whereas the constant A in Eq. (5) is nonzero, taking A = 0 in (5) leads to $y(x) \equiv 0$, and this is, in fact, a solution of the given differential equation (3a). Thus Eq. (5) actually provides a solution of (3a) for *all* values of the constant A, including A = 0. Why did the method of separation of variables fail to capture all solutions of Eq. (3a)? The reason is that in the step



FIGURE 1.4.1. Slope field and solution curves for y' = -6xy.

in which we actually separated the variables, that is, in passing from Eq. (3b) to (3c), we *divided by y*, thus (tacitly) assuming that $y \neq 0$. As a result, our general solution (5), with its restriction that $A \neq 0$, "missed" the particular solution $y(x) \equiv 0$ corresponding to A = 0. Such solutions are known as *singular solutions*, and we say more about them—together with implicit and general solutions—below.

In general, the first-order differential equation (1) is called **separable** provided that f(x, y) can be written as the product of a function of x and a function of y:

$$\frac{dy}{dx} = f(x, y) = g(x)k(y) = \frac{g(x)}{h(y)},$$

where h(y) = 1/k(y). In this case the variables x and y can be *separated*—isolated on opposite sides of an equation—by writing informally the equation

$$h(y) \, dy = g(x) \, dx$$

which we understand to be concise notation for the differential equation

$$h(y)\frac{dy}{dx} = g(x).$$
 (6)

(In the preceding example, $h(y) = \frac{1}{y}$ and g(x) = -6x.) As illustrated above, we can solve this type of differential equation simply by integrating both sides with respect to *x*:

$$\int h(y(x)) \frac{dy}{dx} dx = \int g(x) dx + C;$$

equivalently,

$$\int h(y) \, dy = \int g(x) \, dx + C. \tag{7}$$

All that is required is that the antiderivatives

$$H(y) = \int h(y) \, dy$$
 and $G(x) = \int g(x) \, dx$

can be found. To see that Eqs. (6) and (7) are equivalent, note the following consequence of the chain rule:

$$D_x[H(y(x))] = H'(y(x))y'(x) = h(y)\frac{dy}{dx} = g(x) = D_x[G(x)],$$

which in turn is equivalent to

$$H(y(x)) = G(x) + C,$$
(8)

because two functions have the same derivative on an interval if and only if they differ by a constant on that interval.

$$\frac{dy}{dx} = \frac{4 - 2x}{3y^2 - 5}.$$
(9)

1.4 SEPARABLE EQUATIONS AND APPLICATIONS33

Solution Because



FIGURE 1.4.2. Slope field and solution curves for $y' = (4-2x)/(3y^2-5)$ in Example 1.

$$\frac{4-2x}{3y^2-5} = (4-2x) \cdot \frac{1}{3y^2-5} = g(x)k(y)$$

is the product of a function that depends only on x, and one that depends only on y, Eq. (9) is separable, and thus we can proceed in much the same way as in Eq. (3a). Before doing so, however, we note an important feature of Eq. (9) not shared by Eq. (3a): The function $k(y) = \frac{1}{3y^2 - 5}$ is not defined for all values of y. Indeed, setting $3y^2 - 5$ equal to zero shows that k(y), and thus $\frac{dy}{dx}$ itself, becomes infinite as y approaches either of $\pm \sqrt{\frac{5}{3}}$. Because an infinite slope corresponds to a vertical line segment, we would therefore expect the line segments in the slope field for this differential equation to be "standing straight up" along the two horizontal lines $y = \pm \sqrt{\frac{5}{3}} \approx \pm 1.29$; as Fig. 1.4.2 shows (where these two lines are dashed), this is indeed what we find.

What this means for the differential equation (9) is that *no solution curve of this equation can cross either of the horizontal lines* $y = \pm \sqrt{\frac{5}{3}}$, simply because along these lines $\frac{dy}{dx}$ is undefined. Effectively, then, these lines divide the plane into three regions—defined by the conditions $y > \sqrt{\frac{5}{3}}, -\sqrt{\frac{5}{3}} < y < \sqrt{\frac{5}{3}}$, and $y < -\sqrt{\frac{5}{3}}$ —with all solution curves of Eq. (9) remaining confined to one of these regions.

With this in mind, the general solution of the differential equation in Eq. (9) is easy to find, at least in implicit form. Separating variables and integrating both sides leads to

$$\int 3y^2 - 5 \, dy = \int 4 - 2x \, dx,$$

$$y^3 - 5y = 4x - x^2 + C.$$
 (10)

and thus

Note that unlike Eq. (4), the general solution in Eq. (10) cannot readily be solved for y; thus we cannot directly plot the solution curves of Eq. (9) in the form y = y(x), as we would like. However, what we *can* do is rearrange Eq. (10) so that the constant C is isolated on the right-hand side:

$$y^{3} - 5y - (4x - x^{2}) = C.$$
 (11)

This shows that the solution curves of the differential equation in Eq. (9) are contained in the level curves (also known as contours) of the function

$$F(x, y) = y^{3} - 5y - (4x - x^{2}).$$
(12)

Because no particular solution curve of Eq. (9) can cross either of the lines $y = \pm \sqrt{\frac{5}{3}}$ despite the fact that the level curves of F(x, y) freely do so—the particular solution curves of Eq. (9) are those *portions* of the level curves of F(x, y) which avoid the lines $y = \pm \sqrt{\frac{5}{3}}$.

For example, suppose that we wish to solve the initial value problem

$$\frac{dy}{dx} = \frac{4-2x}{3y^2-5}, \quad y(1) = 3.$$
(13)

Substituting x = 1 and y = 3 into our general solution (10) gives C = 9. Therefore our desired solution curve lies on the level curve

$$y^3 - 5y - (4x - x^2) = 9$$
(14)

of F(x, y); Fig. 1.4.2 shows this and other level curves of F(x, y). However, because the solution curve of the initial value problem (13) must pass through the point (1, 3), which lies above the line $y = \sqrt{\frac{5}{3}}$ in the *xy*-plane, our desired solution curve is restricted to that portion of the level curve (14) which satisfies $y > \sqrt{\frac{5}{3}}$. (In Fig. 1.4.2 the solution curve of the initial value problem (13) is drawn more heavily than the remainder of the level curve (14).) In the same way, Figure 1.4.2 also shows the particular solutions of Eq. (9) subject to the initial



 $f(y) = y^3 - 5y - 9.$



FIGURE 1.4.4. Slope field and solution curves for y' = -x/y.

conditions y(1) = 0 and y(1) = -2. In each of these cases, the curve corresponding to the desired particular solution is only a piece of a larger level curve of the function F(x, y). (Note that in fact, some of the level curves of F themselves consist of two pieces.)

Finally, despite the difficulty of solving Eq. (14) for y by algebraic means, we can nonetheless "solve" for y in the sense that, when a specific value of x is substituted in (14), we can attempt to solve numerically for y. For instance, taking x = 4 yields the equation

$$f(y) = y^3 - 5y - 9 = 0$$

Fig. 1.4.3 shows the graph of f. Using technology we can solve for the single real root $y \approx 2.8552$, thus yielding the value $y(4) \approx 2.8552$ for the solution of the initial value problem (13). By repeating this process for other values of x, we can create a table (like the one shown below) of corresponding x- and y-values for the solution of (13); such a table serves effectively as a "numerical solution" of this initial value problem.

x	-1	0	1	2	3	4	5	6	
у	2.5616	2.8552	3	3.0446	3	2.8552	2.5616	1.8342	

Implicit, General, and Singular Solutions

The equation K(x, y) = 0 is commonly called an **implicit solution** of a differential equation if it is satisfied (on some interval) by some solution y = y(x) of the differential equation. But note that a particular solution y = y(x) of K(x, y) = 0 may or may not satisfy a given initial condition. For example, differentiation of $x^2 + y^2 = 4$ yields

$$x + y\frac{dy}{dx} = 0,$$

so $x^2 + y^2 = 4$ is an implicit solution of the differential equation x + yy' = 0. But only the first of the two explicit solutions

$$y(x) = +\sqrt{4-x^2}$$
 and $y(x) = -\sqrt{4-x^2}$

satisfies the initial condition y(0) = 2 (Fig. 1.4.4).

Remark 1 You should not assume that every possible algebraic solution y = y(x) of an implicit solution satisfies the same differential equation. For instance, if we multiply the implicit solution $x^2 + y^2 - 4 = 0$ by the factor (y - 2x), then we get the new implicit solution

$$(y-2x)(x^2+y^2-4) = 0$$

that yields (or "contains") not only the previously noted explicit solutions $y = +\sqrt{4-x^2}$ and $y = -\sqrt{4-x^2}$ of the differential equation x + yy' = 0, but also the additional function y = 2x that does *not* satisfy this differential equation.

Remark 2 Similarly, solutions of a given differential equation can be either gained or lost when it is multiplied or divided by an algebraic factor. For instance, consider the differential equation

$$(y-2x)y\frac{dy}{dx} = -x(y-2x)$$
 (15)

having the obvious solution y = 2x. But if we divide both sides by the common factor (y - 2x), then we get the previously discussed differential equation

$$y\frac{dy}{dx} = -x$$
, or $x + y\frac{dy}{dx} = 0$, (16)

of which y = 2x is *not* a solution. Thus we "lose" the solution y = 2x of Eq. (15) upon its division by the factor (y - 2x); alternatively, we "gain" this new solution when we multiply Eq. (16) by (y - 2x). Such elementary algebraic operations to simplify a given differential equation before attempting to solve it are common in practice, but the possibility of loss or gain of such "extraneous solutions" should be kept in mind.



FIGURE 1.4.5. The general solution curves $y = (x - C)^2$ and the singular solution curve y = 0 of the differential equation $(y')^2 = 4y$.

A solution of a differential equation that contains an "arbitrary constant" (like the constant C appearing in Eqs. (4) and (10)) is commonly called a **general solution** of the differential equation; any particular choice of a specific value for C yields a single particular solution of the equation.

The argument preceding Example 1 actually suffices to show that *every* particular solution of the differential equation h(y)y' = g(x) in (6) satisfies the equation H(y(x)) = G(x) + C in (8). Consequently, it is appropriate to call (8) not merely *a* general solution of (6), but *the* general solution of (6).

In Section 1.5 we shall see that every particular solution of a *linear* first-order differential equation is contained in its general solution. By contrast, it is common for a nonlinear first-order differential equation to have both a general solution involving an arbitrary constant C and one or several particular solutions that cannot be obtained by selecting a value for C. These exceptional solutions are frequently called **singular solutions**. In Problem 30 we ask you to show that the general solution of the differential equation $(y')^2 = 4y$ yields the family of parabolas $y = (x - C)^2$ illustrated in Fig. 1.4.5, and to observe that the constant-valued function $y(x) \equiv 0$ is a singular solution that cannot be obtained from the general solution by any choice of the arbitrary constant C.

Example 2

Find all solutions of the differential equation

$$\frac{dy}{dx} = 6x(y-1)^{2/3}.$$

Solution Separation of variables gives



FIGURE 1.4.6. General and singular solution curves for $y' = 6x(y-1)^{2/3}$.

$$\int \frac{1}{3(y-1)^{2/3}} dy = \int 2x \, dx;$$
$$(y-1)^{1/3} = x^2 + C;$$
$$y(x) = 1 + (x^2 + C)^3$$

Positive values of the arbitrary constant *C* give the solution curves in Fig. 1.4.6 that lie above the line y = 1, whereas negative values yield those that dip below it. The value C = 0 gives the solution $y(x) = 1 + x^6$, but *no* value of *C* gives the singular solution $y(x) \equiv 1$ that was lost when the variables were separated. Note that the two different solutions $y(x) \equiv 1$ and $y(x) = 1 + (x^2 - 1)^3$ both satisfy the initial condition y(1) = 1. Indeed, the whole singular solution curve y = 1 consists of points where the solution is not unique and where the function $f(x, y) = 6x(y - 1)^{2/3}$ is not differentiable.

Natural Growth and Decay

The differential equation

>

$$\frac{dx}{dt} = kx \quad (k \text{ a constant}) \tag{17}$$

serves as a mathematical model for a remarkably wide range of natural phenomena—any involving a quantity whose time rate of change is proportional to its current size. Here are some examples.

POPULATION GROWTH: Suppose that P(t) is the number of individuals in a population (of humans, or insects, or bacteria) having *constant* birth and death rates β and δ (in births or deaths per individual per unit of time). Then, during a short time interval Δt , approximately $\beta P(t) \Delta t$ births and $\delta P(t) \Delta t$ deaths occur, so the change in P(t) is given approximately by

$$\Delta P \approx (\beta - \delta) P(t) \,\Delta t,$$

and therefore

$$\frac{dP}{dt} = \lim_{\Delta t \to 0} \frac{\Delta P}{\Delta t} = kP,$$
(18)

where $k = \beta - \delta$.

COMPOUND INTEREST: Let A(t) be the number of dollars in a savings account at time t (in years), and suppose that the interest is *compounded continuously* at an annual interest rate r. (Note that 10% annual interest means that r = 0.10.) Continuous compounding means that during a short time interval Δt , the amount of interest added to the account is approximately $\Delta A = rA(t) \Delta t$, so that

$$\frac{dA}{dt} = \lim_{\Delta t \to 0} \frac{\Delta A}{\Delta t} = rA.$$
(19)

RADIOACTIVE DECAY: Consider a sample of material that contains N(t) atoms of a certain radioactive isotope at time t. It has been observed that a constant fraction of those radioactive atoms will spontaneously decay (into atoms of another element or into another isotope of the same element) during each unit of time. Consequently, the sample behaves exactly like a population with a constant death rate and no births. To write a model for N(t), we use Eq. (18) with N in place of P, with k > 0 in place of δ , and with $\beta = 0$. We thus get the differential equation

$$\frac{dN}{dt} = -kN.$$
(20)

The value of k depends on the particular radioactive isotope.

The key to the method of *radiocarbon dating* is that a constant proportion of the carbon atoms in any living creature is made up of the radioactive isotope ¹⁴C of carbon. This proportion remains constant because the fraction of ¹⁴C in the atmosphere remains almost constant, and living matter is continuously taking up carbon from the air or is consuming other living matter containing the same constant ratio of ¹⁴C atoms to ordinary ¹²C atoms. This same ratio permeates all life, because organic processes seem to make no distinction between the two isotopes.

The ratio of 14 C to normal carbon remains constant in the atmosphere because, although 14 C is radioactive and slowly decays, the amount is continuously replenished through the conversion of 14 N (ordinary nitrogen) to 14 C by cosmic rays bombarding the upper atmosphere. Over the long history of the planet, this decay and replenishment process has come into nearly steady state.

Of course, when a living organism dies, it ceases its metabolism of carbon and the process of radioactive decay begins to deplete its ¹⁴C content. There is no replenishment of this ¹⁴C, and consequently the ratio of ¹⁴C to normal carbon begins to drop. By measuring this ratio, the amount of time elapsed since the death of the organism can be estimated. For such purposes it is necessary to measure the **decay constant** k. For ¹⁴C, it is known that $k \approx 0.0001216$ if t is measured in years.

(Matters are not as simple as we have made them appear. In applying the technique of radiocarbon dating, extreme care must be taken to avoid contaminating the sample with organic matter or even with ordinary fresh air. In addition, the cosmic ray levels apparently have not been constant, so the ratio of ¹⁴C in the atmosphere has varied over the past centuries. By using independent methods of dating samples, researchers in this area have compiled tables of correction factors to enhance the accuracy of this process.)

1.4 SEPARABLE EQUATIONS AND APPLICATIONS37

DRUG ELIMINATION: In many cases the amount A(t) of a certain drug in the bloodstream, measured by the excess over the natural level of the drug, will decline at a rate proportional to the current excess amount. That is,

$$\frac{dA}{dt} = -\lambda A,\tag{21}$$

where $\lambda > 0$. The parameter λ is called the **elimination constant** of the drug.

The Natural Growth Equation

The prototype differential equation dx/dt = kx with x(t) > 0 and k a constant (either negative or positive) is readily solved by separating the variables and integrating:

$$\int \frac{1}{x} dx = \int k dt;$$
$$\ln x = kt + C$$

$$\ln x = kt + C.$$

FIGURE 1.4.7. Natural growth.

х x_0 $x = x_0 \ e^{kt}$ (k < 0)

FIGURE 1.4.8. Natural decay.

Then we solve for *x*:

$$e^{\ln x} = e^{kt+C}; \quad x = x(t) = e^C e^{kt} = A e^{kt}.$$

Because C is a constant, so is $A = e^{C}$. It is also clear that $A = x(0) = x_{0}$, so the particular solution of Eq. (17) with the initial condition $x(0) = x_0$ is simply

$$x(t) = x_0 e^{kt}. (22)$$

Because of the presence of the natural exponential function in its solution, the differential equation

$$\frac{x}{t} = kx \tag{23}$$

is often called the exponential or natural growth equation. Figure 1.4.7 shows a typical graph of x(t) in the case k > 0; the case k < 0 is illustrated in Fig. 1.4.8.

Example 3 **World population** According to data listed at **www.census.gov**, the world's total population reached 6 billion persons in mid-1999, and was then increasing at the rate of about 212 thousand persons each day. Assuming that natural population growth at this rate continues, we want to answer these questions:

(a) What is the annual growth rate k?

(b) What will be the world population at the middle of the 21st century?

(c) How long will it take the world population to increase tenfold—thereby reaching the 60 billion that some demographers believe to be the maximum for which the planet can provide adequate food supplies?

Solution

(a) We measure the world population P(t) in billions and measure time in years. We take t = 0 to correspond to (mid) 1999, so $P_0 = 6$. The fact that P is increasing by 212,000, or 0.000212 billion, persons per day at time t = 0 means that

$$P'(0) = (0.000212)(365.25) \approx 0.07743$$

billion per year. From the natural growth equation P' = kP with t = 0 we now obtain

$$k = \frac{P'(0)}{P(0)} \approx \frac{0.07743}{6} \approx 0.0129.$$

Thus the world population was growing at the rate of about 1.29% annually in 1999. This value of k gives the world population function

$$P(t) = 6e^{0.0129t}$$



(b) With t = 51 we obtain the prediction

$$P(51) = 6e^{(0.0129)(51)} \approx 11.58$$
 (billion)

for the world population in mid-2050 (so the population will almost have doubled in the just over a half-century since 1999).

(c) The world population should reach 60 billion when

$$60 = 6e^{0.0129t}$$
; that is, when $t = \frac{\ln 10}{0.0129} \approx 178$;

and thus in the year 2177.

Note Actually, the rate of growth of the world population is expected to slow somewhat during the next half-century, and the best current prediction for the 2050 population is "only" 9.1 billion. A simple mathematical model cannot be expected to mirror precisely the complexity of the real world.

The decay constant of a radioactive isotope is often specified in terms of another empirical constant, the *half-life* of the isotope, because this parameter is more convenient. The **half-life** τ of a radioactive isotope is the time required for *half* of it to decay. To find the relationship between k and τ , we set $t = \tau$ and $N = \frac{1}{2}N_0$ in the equation $N(t) = N_0 e^{-kt}$, so that $\frac{1}{2}N_0 = N_0 e^{-k\tau}$. When we solve for τ , we find that

$$\tau = \frac{\ln 2}{k}.$$
(24)

For example, the half-life of ¹⁴C is $\tau \approx (\ln 2)/(0.0001216)$, approximately 5700 years.

Radiometric dating A specimen of charcoal found at Stonehenge turns out to contain 63% as much ¹⁴C as a sample of present-day charcoal of equal mass. What is the age of the sample?

Solution

We take t = 0 as the time of the death of the tree from which the Stonehenge charcoal was made and N_0 as the number of ¹⁴C atoms that the Stonehenge sample contained then. We are given that $N = (0.63)N_0$ now, so we solve the equation $(0.63)N_0 = N_0e^{-kt}$ with the value k = 0.0001216. Thus we find that

$$t = -\frac{\ln(0.63)}{0.0001216} \approx 3800 \text{ (years)}.$$

Thus the sample is about 3800 years old. If it has any connection with the builders of Stonehenge, our computations suggest that this observatory, monument, or temple—whichever it may be—dates from 1800 B.C. or earlier.

Cooling and Heating

>

According to Newton's law of cooling (Eq. (3) of Section 1.1), the time rate of change of the temperature T(t) of a body immersed in a medium of constant temperature A is proportional to the difference A - T. That is,

$$\frac{dT}{dt} = k(A - T), \tag{25}$$

where *k* is a positive constant. This is an instance of the linear first-order differential equation with constant coefficients:

$$\frac{dx}{dt} = ax + b. \tag{26}$$

It includes the exponential equation as a special case (b = 0) and is also easy to solve by separation of variables.

- **Example 5** Cooling A 4-lb roast, initially at 50°F, is placed in a 375°F oven at 5:00 P.M. After 75 minutes it is found that the temperature T(t) of the roast is 125°F. When will the roast be 150°F (medium rare)?
 - Solution We take time t in minutes, with t = 0 corresponding to 5:00 P.M. We also assume (somewhat unrealistically) that at any instant the temperature T(t) of the roast is uniform throughout. We have T(t) < A = 375, T(0) = 50, and T(75) = 125. Hence

$$\frac{dT}{dt} = k(375 - T);$$

$$\int \frac{1}{375 - T} dT = \int k \, dt;$$

$$-\ln(375 - T) = kt + C;$$

$$375 - T = Be^{-kt}.$$

Now T(0) = 50 implies that B = 325, so $T(t) = 375 - 325e^{-kt}$. We also know that T = 125 when t = 75. Substitution of these values in the preceding equation yields

$$k = -\frac{1}{75} \ln\left(\frac{250}{325}\right) \approx 0.0035.$$

Hence we finally solve the equation

$$150 = 375 - 325e^{(-0.0035)t}$$

for $t = -[\ln(225/325)]/(0.0035) \approx 105$ (min), the total cooking time required. Because the roast was placed in the oven at 5:00 P.M., it should be removed at about 6:45 P.M.

Torricelli's Law

Suppose that a water tank has a hole with area a at its bottom, from which water is leaking. Denote by y(t) the depth of water in the tank at time t, and by V(t) the volume of water in the tank then. It is plausible—and true, under ideal conditions—that the velocity of water exiting through the hole is

$$v = \sqrt{2gy},\tag{27}$$

which is the velocity a drop of water would acquire in falling freely from the surface of the water to the hole (see Problem 35 of Section 1.2). One can derive this formula beginning with the assumption that the sum of the kinetic and potential energy of the system remains constant. Under real conditions, taking into account the constriction of a water jet from an orifice, $v = c\sqrt{2gy}$, where *c* is an empirical constant between 0 and 1 (usually about 0.6 for a small continuous stream of water). For simplicity we take c = 1 in the following discussion.

As a consequence of Eq. (27), we have

$$\frac{dV}{dt} = -av = -a\sqrt{2gy};$$
(28a)

equivalently,

$$\frac{dV}{dt} = -k\sqrt{y} \quad \text{where} \quad k = a\sqrt{2g}. \tag{28b}$$

This is a statement of Torricelli's law for a draining tank. Let A(y) denote the horizontal cross-sectional area of the tank at height y. Then, applied to a thin horizontal

slice of water at height \overline{y} with area $A(\overline{y})$ and thickness $d\overline{y}$, the integral calculus method of cross sections gives

$$V(y) = \int_0^y A(\overline{y}) \, d\overline{y}$$

The fundamental theorem of calculus therefore implies that dV/dy = A(y) and hence that

$$\frac{dV}{dt} = \frac{dV}{dy} \cdot \frac{dy}{dt} = A(y)\frac{dy}{dt}.$$
(29)

From Eqs. (28) and (29) we finally obtain

$$A(y)\frac{dy}{dt} = -a\sqrt{2gy} = -k\sqrt{y},$$
(30)

an alternative form of Torricelli's law.

From the right triangle in Fig. 1.4.9, we see that

Draining tank A hemispherical bowl has top radius 4 ft and at time t = 0 is full of water. At that moment a circular hole with diameter 1 in. is opened in the bottom of the tank. How long will it take for all the water to drain from the tank?

Solution

Example 6

$$A(y) = \pi r^{2} = \pi \left[16 - (4 - y)^{2} \right] = \pi (8y - y^{2})$$

With g = 32 ft/s², Eq. (30) becomes

$$\pi (8y - y^2) \frac{dy}{dt} = -\pi \left(\frac{1}{24}\right)^2 \sqrt{2 \cdot 32y};$$
$$\int (8y^{1/2} - y^{3/2}) \, dy = -\int \frac{1}{72} \, dt;$$
$$\frac{16}{3} y^{3/2} - \frac{2}{5} y^{5/2} = -\frac{1}{72}t + C.$$

Now y(0) = 4, so

$$C = \frac{16}{3} \cdot 4^{3/2} - \frac{2}{5} \cdot 4^{5/2} = \frac{448}{15}.$$

The tank is empty when y = 0, thus when

$$t = 72 \cdot \frac{448}{15} \approx 2150$$
 (s);

that is, about 35 min 50 s. So it takes slightly less than 36 min for the tank to drain.

In the case of an upright cylindrical tank with constant cross-sectional area *A*, Torricelli's law in Eq. (30) takes the form

$$\frac{dy}{dt} = -c\sqrt{y}$$

with c = k/A. With initial condition y(0) = 0 we routinely separate variables and integrate to get

$$y = \frac{1}{4}c^2t^2$$

However, this formula if taken at face value implies that y > 0 if t > 0. But if the tank is empty at time t = 0 as prescribed by the initial condition y(0) = 0, then certainly the tank remains empty thereafter, so $y(t) \equiv 0$ for t > 0.

To see what is going on here, note that the right-hand side function $f(t, y) = -c\sqrt{y}$ in our differential equation $y' = -c\sqrt{y}$ does *not* satisfy the condition that $\partial f/\partial y$ be continuous at (0, 0), so the existence-uniqueness theorem of Section 1.3 does *not* guarantee uniqueness of a solution near t = 0. Indeed, we note the two different but physically meaningful solutions

$$y_1(t) \equiv 0$$
 for all t, and $y_2(t) = \begin{cases} \frac{1}{4}c^2t^2 & \text{for } t < 0, \\ 0 & \text{for } t \ge 0 \end{cases}$



FIGURE 1.4.9. Draining a hemispherical tank.

Example 7

of the initial value problem $y' = -c\sqrt{y}$, y(0) = 0. The constant solution $y_1(t) \equiv 0$ corresponds to a tank that always has been and always will be empty, while $y_2(t)$ corresponds to a tank draining while t < 0 that empties precisely at time t = 0 and remains empty thereafter.

Thus this example provides a concrete physical situation described by an initial value problem with non-unique solutions.

1.4 Problems

Find general solutions (implicit if necessary, explicit if convenient) of the differential equations in Problems 1 through 18. Primes denote derivatives with respect to x.

- 1. $\frac{dy}{dx} + 2xy = 0$ 3. $\frac{dy}{dx} = y \sin x$ 5. $2\sqrt{x} \frac{dy}{dx} = \sqrt{1-y^2}$ 7. $\frac{dy}{dx} = (64xy)^{1/3}$ 9. $(1-x^2)\frac{dy}{dx} = 2y$ 10. $(1+x)^2\frac{dy}{dx} = (1+y)^2$ 11. $y' = xy^3$ 12. $yy' = x(y^2+1)$ 13. $y^3\frac{dy}{dx} = (y^4+1)\cos x$ 14. $\frac{dy}{dx} = \frac{1+\sqrt{x}}{1+\sqrt{y}}$ 15. $\frac{dy}{dx} = \frac{(x-1)y^5}{x^2(2y^3-y)}$ 16. $(x^2+1)(\tan y)y' = x$ 17. y' = 1+x+y+xy (Suggestion: Factor the right-hand
- **17.** y' = 1 + x + y + xy (Suggestion: Factor the right-han side.)
- **18.** $x^2y' = 1 x^2 + y^2 x^2y^2$

Find explicit particular solutions of the initial value problems in Problems 19 through 28.

19. $\frac{dy}{dx} = ye^x$, y(0) = 2e20. $\frac{dy}{dx} = 3x^2(y^2 + 1)$, y(0) = 121. $2y\frac{dy}{dx} = \frac{x}{\sqrt{x^2 - 16}}$, y(5) = 222. $\frac{dy}{dx} = 4x^3y - y$, y(1) = -323. $\frac{dy}{dx} + 1 = 2y$, y(1) = 124. $\frac{dy}{dx} = y \cot x$, $y(\frac{1}{2}\pi) = \frac{1}{2}\pi$ 25. $x\frac{dy}{dx} - y = 2x^2y$, y(1) = 126. $\frac{dy}{dx} = 2xy^2 + 3x^2y^2$, y(1) = -127. $\frac{dy}{dx} = 6e^{2x-y}$, y(0) = 028. $2\sqrt{x}\frac{dy}{dx} = \cos^2 y$, $y(4) = \pi/4$

Problems 29 through 32 explore the connections among general and singular solutions, existence, and uniqueness.

- **29.** (a) Find a general solution of the differential equation $dy/dx = y^2$. (b) Find a singular solution that is not included in the general solution. (c) Inspect a sketch of typical solution curves to determine the points (a, b) for which the initial value problem $y' = y^2$, y(a) = b has a unique solution.
- **30.** Solve the differential equation $(dy/dx)^2 = 4y$ to verify the general solution curves and singular solution curve that are illustrated in Fig. 1.4.5. Then determine the points (a, b) in the plane for which the initial value problem $(y')^2 = 4y$, y(a) = b has (a) no solution, (b) infinitely many solutions that are defined for all x, (c) on some neighborhood of the point x = a, only finitely many solutions.
- **31.** Discuss the difference between the differential equations $(dy/dx)^2 = 4y$ and $dy/dx = 2\sqrt{y}$. Do they have the same solution curves? Why or why not? Determine the points (a, b) in the plane for which the initial value problem $y' = 2\sqrt{y}$, y(a) = b has (**a**) no solution, (**b**) a unique solution, (**c**) infinitely many solutions.
- 32. Find a general solution and any singular solutions of the differential equation $dy/dx = y\sqrt{y^2 1}$. Determine the points (a, b) in the plane for which the initial value problem $y' = y\sqrt{y^2 1}$, y(a) = b has (a) no solution, (b) a unique solution, (c) infinitely many solutions.
- **33. Population growth** A certain city had a population of 25,000 in 1960 and a population of 30,000 in 1970. Assume that its population will continue to grow exponentially at a constant rate. What population can its city planners expect in the year 2000?
- **34.** Population growth In a certain culture of bacteria, the number of bacteria increased sixfold in 10 h. How long did it take for the population to double?
- **35.** Radiocarbon dating Carbon extracted from an ancient skull contained only one-sixth as much ¹⁴C as carbon extracted from present-day bone. How old is the skull?
- **36.** Radiocarbon dating Carbon taken from a purported relic of the time of Christ contained 4.6×10^{10} atoms of 14 C per gram. Carbon extracted from a present-day specimen of the same substance contained 5.0×10^{10} atoms of 14 C per gram. Compute the approximate age of the relic. What is your opinion as to its authenticity?
- **37.** Continuously compounded interest Upon the birth of their first child, a couple deposited \$5000 in an account that pays 8% interest compounded continuously. The interest payments are allowed to accumulate. How much will the account contain on the child's eighteenth birth-day?

42 CHAPTER 1 FIRST-ORDER DIFFERENTIAL EQUATIONS

- **38.** Continuously compounded interest Suppose that you discover in your attic an overdue library book on which your grandfather owed a fine of 30 cents 100 years ago. If an overdue fine grows exponentially at a 5% annual rate compounded continuously, how much would you have to pay if you returned the book today?
- **39. Drug elimination** Suppose that sodium pentobarbital is used to anesthetize a dog. The dog is anesthetized when its bloodstream contains at least 45 milligrams (mg) of sodium pentobarbitol per kilogram of the dog's body weight. Suppose also that sodium pentobarbitol is eliminated exponentially from the dog's bloodstream, with a half-life of 5 h. What single dose should be administered in order to anesthetize a 50-kg dog for 1 h?
- **40. Radiometric dating** The half-life of radioactive cobalt is 5.27 years. Suppose that a nuclear accident has left the level of cobalt radiation in a certain region at 100 times the level acceptable for human habitation. How long will it be until the region is again habitable? (Ignore the probable presence of other radioactive isotopes.)
- **41. Isotope formation** Suppose that a mineral body formed in an ancient cataclysm—perhaps the formation of the earth itself—originally contained the uranium isotope 238 U (which has a half-life of 4.51×10^9 years) but no lead, the end product of the radioactive decay of 238 U. If today the ratio of 238 U atoms to lead atoms in the mineral body is 0.9, when did the cataclysm occur?
- **42.** Radiometric dating A certain moon rock was found to contain equal numbers of potassium and argon atoms. Assume that all the argon is the result of radioactive decay of potassium (its half-life is about 1.28×10^9 years) and that one of every nine potassium atom disintegrations yields an argon atom. What is the age of the rock, measured from the time it contained only potassium?
- **43.** Cooling A pitcher of buttermilk initially at 25° C is to be cooled by setting it on the front porch, where the temperature is 0° C. Suppose that the temperature of the buttermilk has dropped to 15° C after 20 min. When will it be at 5° C?
- **44.** Solution rate When sugar is dissolved in water, the amount *A* that remains undissolved after *t* minutes satisfies the differential equation dA/dt = -kA (k > 0). If 25% of the sugar dissolves after 1 min, how long does it take for half of the sugar to dissolve?
- **45.** Underwater light intensity The intensity I of light at a depth of x meters below the surface of a lake satisfies the differential equation dI/dx = (-1.4)I. (a) At what depth is the intensity half the intensity I_0 at the surface (where x = 0)? (b) What is the intensity at a depth of 10 m (as a fraction of I_0)? (c) At what depth will the intensity be 1% of that at the surface?
- **46.** Barometric pressure and altitude The barometric pressure p (in inches of mercury) at an altitude x miles above sea level satisfies the initial value problem dp/dx = (-0.2)p, p(0) = 29.92. (a) Calculate the barometric pressure at 10,000 ft and again at 30,000 ft. (b) Without prior

conditioning, few people can survive when the pressure drops to less than 15 in. of mercury. How high is that?

- **47. Spread of rumor** A certain piece of dubious information about phenylethylamine in the drinking water began to spread one day in a city with a population of 100,000. Within a week, 10,000 people had heard this rumor. Assume that the rate of increase of the number who have heard the rumor is proportional to the number who have not yet heard it. How long will it be until half the population of the city has heard the rumor?
- **48.** Isotope formation According to one cosmological theory, when uranium was first generated in the early evolution of the universe following the "big bang," the isotopes 235 U and 238 U were produced in equal amounts. Given the half-lives of 4.51×10^9 years for 238 U and 7.10×10^8 years for 235 U, calculate the length of time required to reach the present distribution of 137.7 atoms of 238 U for each atom of 235 U.
- **49.** Cooling A cake is removed from an oven at 210°F and left to cool at room temperature, which is 70°F. After 30 min the temperature of the cake is 140°F. When will it be 100°F?
- **50.** Pollution increase The amount A(t) of atmospheric pollutants in a certain mountain valley grows naturally and is tripling every 7.5 years.
 - (a) If the initial amount is 10 pu (pollutant units), write a formula for A(t) giving the amount (in pu) present after t years.
 - (b) What will be the amount (in pu) of pollutants present in the valley atmosphere after 5 years?
 - (c) If it will be dangerous to stay in the valley when the amount of pollutants reaches 100 pu, how long will this take?
- **51. Radioactive decay** An accident at a nuclear power plant has left the surrounding area polluted with radioactive material that decays naturally. The initial amount of radioactive material present is 15 su (safe units), and 5 months later it is still 10 su.
 - (a) Write a formula giving the amount A(t) of radioactive material (in su) remaining after t months.
 - (b) What amount of radioactive material will remain after 8 months?
 - (c) How long—total number of months or fraction thereof—will it be until A = 1 su, so it is safe for people to return to the area?
- **52. Growth of languages** There are now about 3300 different human "language families" in the whole world. Assume that all these are derived from a single original language and that a language family develops into 1.5 language families every 6 thousand years. About how long ago was the single original human language spoken?
- **53. Growth of languages** Thousands of years ago ancestors of the Native Americans crossed the Bering Strait from Asia and entered the western hemisphere. Since then, they have fanned out across North and South America. The single language that the original Native Americans spoke has

since split into many Indian "language families." Assume (as in Problem 52) that the number of these language families has been multiplied by 1.5 every 6000 years. There are now 150 Native American language families in the western hemisphere. About when did the ancestors of today's Native Americans arrive?

Torricelli's Law

Problems 54 through 64 illustrate the application of Torricelli's law.

- 54. A tank is shaped like a vertical cylinder; it initially contains water to a depth of 9 ft, and a bottom plug is removed at time t = 0 (hours). After 1 h the depth of the water has dropped to 4 ft. How long does it take for all the water to drain from the tank?
- **55.** Suppose that the tank of Problem 54 has a radius of 3 ft and that its bottom hole is circular with radius 1 in. How long will it take the water (initially 9 ft deep) to drain completely?
- 56. At time t = 0 the bottom plug (at the vertex) of a full conical water tank 16 ft high is removed. After 1 h the water in the tank is 9 ft deep. When will the tank be empty?
- 57. Suppose that a cylindrical tank initially containing V_0 gallons of water drains (through a bottom hole) in T minutes. Use Torricelli's law to show that the volume of water in the tank after $t \leq T$ minutes is $V = V_0 [1 (t/T)]^2$.
- **58.** A water tank has the shape obtained by revolving the curve $y = x^{4/3}$ around the *y*-axis. A plug at the bottom is removed at 12 noon, when the depth of water in the tank is 12 ft. At 1 P.M. the depth of the water is 6 ft. When will the tank be empty?
- 59. A water tank has the shape obtained by revolving the parabola x² = by around the y-axis. The water depth is 4 ft at 12 noon, when a circular plug in the bottom of the tank is removed. At 1 P.M. the depth of the water is 1 ft.
 (a) Find the depth y(t) of water remaining after t hours.
 (b) When will the tank be empty? (c) If the initial radius of the top surface of the water is 2 ft, what is the radius of the circular hole in the bottom?
- **60.** A cylindrical tank with length 5 ft and radius 3 ft is situated with its axis horizontal. If a circular bottom hole with a radius of 1 in. is opened and the tank is initially half full of water, how long will it take for the liquid to drain completely?
- **61.** A spherical tank of radius 4 ft is full of water when a circular bottom hole with radius 1 in. is opened. How long will be required for all the water to drain from the tank?
- **62.** Suppose that an initially full hemispherical water tank of radius 1 m has its flat side as its bottom. It has a bottom hole of radius 1 cm. If this bottom hole is opened at 1 P.M., when will the tank be empty?
- **63.** Consider the initially full hemispherical water tank of Example 8, except that the radius *r* of its circular bottom hole is now unknown. At 1 P.M. the bottom hole is opened and at 1:30 P.M. the depth of water in the tank is 2 ft. (a) Use Torricelli's law in the form $dV/dt = -(0.6)\pi r^2 \sqrt{2gy}$

(taking constriction into account) to determine when the tank will be empty. (b) What is the radius of the bottom hole?

64. A 12 h water clock is to be designed with the dimensions shown in Fig. 1.4.10, shaped like the surface obtained by revolving the curve y = f(x) around the *y*-axis. What should this curve be, and what should the radius of the circular bottom hole be, in order that the water level will fall at the *constant* rate of 4 inches per hour (in./h)?



FIGURE 1.4.10. The clepsydra.

- **65.** Time of death Just before midday the body of an apparent homicide victim is found in a room that is kept at a constant temperature of 70°F. At 12 noon the temperature of the body is 80°F and at 1 P.M. it is 75°F. Assume that the temperature of the body at the time of death was 98.6°F and that it has cooled in accord with Newton's law. What was the time of death?
- **66.** Snowplow problem Early one morning it began to snow at a constant rate. At 7 A.M. a snowplow set off to clear a road. By 8 A.M. it had traveled 2 miles, but it took two more hours (until 10 A.M.) for the snowplow to go an additional 2 miles. (a) Let t = 0 when it began to snow, and let x denote the distance traveled by the snowplow at time t. Assuming that the snowplow clears snow from the road at a constant rate (in cubic feet per hour, say), show that

$$k\frac{dx}{dt} = \frac{1}{t}$$

where k is a constant. (b) What time did it start snowing? (*Answer:* 6 A.M.)

- **67. Snowplow problem** A snowplow sets off at 7 A.M. as in Problem 66. Suppose now that by 8 A.M. it had traveled 4 miles and that by 9 A.M. it had moved an additional 3 miles. What time did it start snowing? This is a more difficult snowplow problem because now a transcendental equation must be solved numerically to find the value of *k*. (*Answer:* 4:27 A.M.)
- **68. Brachistochrone** Figure 1.4.11 shows a bead sliding down a frictionless wire from point P to point Q. The

44 CHAPTER 1 FIRST-ORDER DIFFERENTIAL EQUATIONS

brachistochrone problem asks what shape the wire should be in order to minimize the bead's time of descent from P to Q. In June of 1696, John Bernoulli proposed this problem as a public challenge, with a 6-month deadline (later extended to Easter 1697 at George Leibniz's request). Isaac Newton, then retired from academic life and serving as Warden of the Mint in London, received Bernoulli's challenge on January 29, 1697. The very next day he communicated his own solution—the curve of minimal descent time is an arc of an inverted cycloid—to the Royal Society of London. For a modern derivation of this result, suppose the bead starts from rest at the origin P and let y = y(x) be the equation of the desired curve in a coordinate system with the y-axis pointing downward. Then a mechanical analogue of Snell's law in optics implies that

$$\frac{\sin \alpha}{v} = \text{constant},$$
 (i)

where α denotes the angle of deflection (from the vertical) of the tangent line to the curve—so $\cot \alpha = y'(x)$ (why?)—and $v = \sqrt{2gy}$ is the bead's velocity when it has descended a distance y vertically (from KE = $\frac{1}{2}mv^2 = mgy = -PE$).



FIGURE 1.4.11. A bead sliding down a wire—the brachistochrone problem.

(a) First derive from Eq. (i) the differential equation

$$\frac{dy}{dx} = \sqrt{\frac{2a - y}{y}},$$
 (ii)

where a is an appropriate positive constant.

(b) Substitute $y = 2a \sin^2 t$, $dy = 4a \sin t \cos t dt$ in (ii) to derive the solution

$$x = a(2t - \sin 2t), \quad y = a(1 - \cos 2t)$$
 (iii)

for which t = y = 0 when x = 0. Finally, the substitution of $\theta = 2t$ in (iii) yields the standard parametric equations $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ of the cycloid that is generated by a point on the rim of a circular wheel of radius *a* as it rolls along the *x*-axis. [See Example 5 in Section 9.4 of Edwards and Penney, *Calculus: Early Transcendentals*, 7th edition, Hoboken, NJ: Pearson, 2008.]

69. Hanging cable Suppose a uniform flexible cable is suspended between two points $(\pm L, H)$ at equal heights located symmetrically on either side of the *x*-axis (Fig. 1.4.12). Principles of physics can be used to show that the shape y = y(x) of the hanging cable satisfies the differential equation

$$a\frac{d^2y}{dx^2} = \sqrt{1 + \left(\frac{dy}{dx}\right)^2},$$

where the constant $a = T/\rho$ is the ratio of the cable's tension T at its lowest point x = 0 (where y'(0) = 0) and its (constant) linear density ρ . If we substitute v = dy/dx, $dv/dx = d^2y/dx^2$ in this second-order differential equation, we get the first-order equation

$$a\frac{dv}{dx} = \sqrt{1+v^2}.$$

Solve this differential equation for $y'(x) = v(x) = \sinh(x/a)$. Then integrate to get the shape function

$$y(x) = a \cosh\left(\frac{x}{a}\right) + C$$

of the hanging cable. This curve is called a *catenary*, from the Latin word for *chain*.



FIGURE 1.4.12. The catenary.

1.4 Application The Logistic Equation



As in Eq. (7) of this section, the solution of a separable differential equation reduces to the evaluation of two indefinite integrals. It is tempting to use a symbolic algebra system for this purpose. We illustrate this approach using the *logistic differential equation*

$$\frac{dx}{dt} = ax - bx^2 \tag{1}$$

that models a population x(t) with births (per unit time) proportional to x and deaths proportional to x^2 . Here we concentrate on the solution of Eq. (1) and defer discussion of population applications to Section 2.1.

If a = 0.01 and b = 0.0001, for instance, Eq. (1) is

$$\frac{dx}{dt} = (0.01)x - (0.0001)x^2 = \frac{x}{10000}(100 - x).$$
(2)

Separation of variables leads to

$$\int \frac{1}{x(100-x)} dx = \int \frac{1}{10000} dt = \frac{t}{10000} + C.$$
 (3)

We can evaluate the integral on the left by using the Maple command

int(1/(x*(100 - x)), x);

the Mathematica command

or the MATLAB command

syms x; int(1/(x*(100 - x)))

Alternatively, we could use the freely available Wolfram|Alpha system (www.wolframalpha.com); the query

integrate 1/(x*(100 - x))

produces the output shown in Fig. 1.4.14.

Any computer algebra system gives a result of the form

$$\frac{1}{100}\ln x - \frac{1}{100}\ln(x - 100) = \frac{t}{10000} + C \tag{4}$$

equivalent to the graphing calculator result shown in Fig. 1.4.13.

You can now apply the initial condition $x(0) = x_0$, combine logarithms, and finally exponentiate to solve Eq. (4) for the particular solution

$$x(t) = \frac{100x_0 e^{t/100}}{100 - x_0 + x_0 e^{t/100}}$$
(5)

of Eq. (2). The slope field and solution curves shown in Fig. 1.4.15 suggest that, whatever is the initial value x_0 , the solution x(t) approaches 100 as $t \to +\infty$. Can you use Eq. (5) to verify this conjecture?

INVESTIGATION: For your own personal logistic equation, take a = m/n and b = 1/n in Eq. (1), with *m* and *n* being the *largest* two distinct digits (in either order) in your student ID number.

- (a) First generate a slope field for your differential equation and include a sufficient number of solution curves that you can see what happens to the population as t → +∞. State your inference plainly.
- (b) Next use a computer algebra system to solve the differential equation symbolically; then use the symbolic solution to find the limit of x(t) as $t \to +\infty$. Was your graphically based inference correct?
- (c) Finally, state and solve a numerical problem using the symbolic solution. For instance, how long does it take x to grow from a selected initial value x₀ to a given target value x₁?



FIGURE 1.4.15. Slope field and solution curves for $x' = (0.01)x - (0.0001)x^2$.



FIGURE 1.4.13. TI-89 screen showing the integral in Eq. (3).



 $\frac{1}{100}$ (log(x) - log(100 - x)) + constant

FIGURE 1.4.14. Wolfram|Alpha display showing the integral in Eq. (3). Screenshot of Wolfram|Alpha output. Used by permission of WolframAlpha LLC.

1.5 Linear First-Order Equations

We turn now to another important method for solving first-order differential equations that rests upon the idea of "integrating both sides." In Section 1.4 we saw that the first step in solving a separable differential equation is to multiply and/or divide both sides of the equation by whatever is required in order to separate the variables. For instance, to solve the equation

$$\frac{dy}{dx} = 2xy \quad (y > 0),\tag{1}$$

we divide both sides by y (and, so to speak, multiply by the differential dx) to get

$$\frac{dy}{y} = 2x \, dx.$$

Integrating both sides then gives the general solution $\ln y = x^2 + C$.

There is another way to approach the differential equation in (1), however, which—while leading to the same general solution—opens the door not only to the solution method discussed in this section, but to other methods of solving differential equations as well. What is common to all these methods is the idea that if a given equation is difficult to solve, then perhaps *multiplying* both sides of the equation by a suitably chosen function of x and/or y may result in an equivalent equation that can be solved more easily. Thus, in Eq. (1), rather than *divide* both sides by y, we could instead *multiply* both sides by the factor 1/y. (Of course algebraically these two are the same, but we are highlighting the fact that often the crucial first step in solving a differential equation is to multiply both of its sides by the "right" function.) Applying this to Eq. (1) (while leaving dx in place) gives

$$\frac{1}{v} \cdot \frac{dy}{dx} = 2x.$$
(2)

The significance of Eq. (2) is that, unlike Eq. (1), *both sides are recognizable as a derivative*. By the chain rule, the left-hand side of Eq. (2) can be written as

$$\frac{1}{y} \cdot \frac{dy}{dx} = D_x(\ln y),$$

and of course the right hand side of Eq. (2) is $D_x(x^2)$. Thus each side of Eq. (2) can be viewed as a derivative with respect to x:

$$D_x(\ln y) = D_x(x^2)$$

Integrating both sides with respect to x gives the same general solution $\ln y = x^2 + C$ that we found before.

We were able to solve the differential equation in Eq. (1), then, by first multiplying both of its sides by a factor—known as an *integrating factor*—chosen so that both sides of the resulting equation could be recognized as a derivative. Solving the equation then becomes simply a matter of integrating both sides. More generally, an **integrating factor** for a differential equation is a function $\rho(x, y)$ such that multiplication of each side of the differential equation by $\rho(x, y)$ yields an equation in which each side is recognizable as a derivative. In some cases integrating factors involve both of the variables x and y; however, our second solution of Eq. (1) was based on the integrating factor $\rho(y) = 1/y$, which depends only on y. Our goal in this section is to show how integrating factors can be used to solve a broad and important category of first-order differential equations.

A linear first-order equation is a differential equation of the form

$$\frac{dy}{dx} + P(x)y = Q(x).$$
(3)

We assume that the coefficient functions P(x) and Q(x) are continuous on some interval on the *x*-axis. (Can you see that the differential equation in Eq. (1), in addition to being separable, is also linear? Is *every* separable equation also linear?) Assuming that the necessary antiderivatives can be found, the general linear equation in (3) can always be solved by multiplying by the integrating factor

$$\rho(x) = e^{\int P(x) \, dx}.\tag{4}$$

The result is

 \succ

$$e^{\int P(x) \, dx} \frac{dy}{dx} + P(x) e^{\int P(x) \, dx} y = Q(x) e^{\int P(x) \, dx}.$$
(5)

Because

$$D_x\left[\int P(x)\,dx\right] = P(x),$$

the left-hand side is the derivative of the *product* $y(x) \cdot e^{\int P(x) dx}$, so Eq. (5) is equivalent to

$$D_x\left[y(x)\cdot e^{\int P(x)\,dx}\right] = Q(x)e^{\int P(x)\,dx}.$$

Integration of both sides of this equation gives

$$y(x)e^{\int P(x)\,dx} = \int \left(Q(x)e^{\int P(x)\,dx}\right)\,dx + C.$$

Finally, solving for y, we obtain the general solution of the linear first-order equation in (3):

$$y(x) = e^{-\int P(x) dx} \left[\int \left(Q(x) e^{\int P(x) dx} \right) dx + C \right].$$
(6)

This formula should **not** be memorized. In a specific problem it generally is simpler to use the *method* by which we developed the formula. That is, in order to solve an equation that can be written in the form in Eq. (3) with the coefficient functions P(x) and Q(x) displayed explicitly, you should attempt to carry out the following steps.

METHOD: SOLUTION OF LINEAR FIRST-ORDER EQUATIONS

- **1.** Begin by calculating the integrating factor $\rho(x) = e^{\int P(x) dx}$.
- **2.** Then multiply both sides of the differential equation by $\rho(x)$.
- **3.** Next, recognize the left-hand side of the resulting equation as the derivative of a product:

$$D_x \left[\rho(x) y(x) \right] = \rho(x) Q(x).$$

4. Finally, integrate this equation,

$$\rho(x)y(x) = \int \rho(x)Q(x)\,dx + C,$$

then solve for *y* to obtain the general solution of the original differential equation.

Remark 1 Given an initial condition $y(x_0) = y_0$, you can (as usual) substitute $x = x_0$ and $y = y_0$ into the general solution and solve for the value of *C* yielding the particular solution that satisfies this initial condition.

Remark 2 You need not supply explicitly a constant of integration when you find the integrating factor $\rho(x)$. For if we replace

$$\int P(x) dx$$
 with $\int P(x) dx + K$

in Eq. (4), the result is

$$\rho(x) = e^{K + \int P(x) \, dx} = e^K e^{\int P(x) \, dx}$$

But the constant factor e^K does not affect materially the result of multiplying both sides of the differential equation in (3) by $\rho(x)$, so we might as well take K = 0. You may therefore choose for $\int P(x) dx$ any convenient antiderivative of P(x), without bothering to add a constant of integration.

Example 1 Solve the initial value problem

$$\frac{dy}{dx} - y = \frac{11}{8}e^{-x/3}, \quad y(0) = -1.$$

Solution

Here we have
$$P(x) \equiv -1$$
 and $Q(x) = \frac{11}{8}e^{-x/3}$, so the integrating factor is

$$\rho(x) = e^{\int (-1) \, dx} = e^{-x}$$

Multiplication of both sides of the given equation by e^{-x} yields

$$e^{-x}\frac{dy}{dx} - e^{-x}y = \frac{11}{8}e^{-4x/3},\tag{7}$$

which we recognize as

$$\frac{d}{dx}\left(e^{-x}y\right) = \frac{11}{8}e^{-4x/3}$$

Hence integration with respect to x gives

$$e^{-x}y = \int \frac{11}{8}e^{-4x/3} dx = -\frac{33}{32}e^{-4x/3} + C,$$

and multiplication by e^x gives the general solution

$$y(x) = Ce^{x} - \frac{33}{32}e^{-x/3}.$$
(8)

Substitution of x = 0 and y = -1 now gives $C = \frac{1}{32}$, so the desired particular solution is

$$y(x) = \frac{1}{32}e^x - \frac{33}{32}e^{-x/3} = \frac{1}{32}\left(e^x - 33e^{-x/3}\right).$$

Remark Figure 1.5.1 shows a slope field and typical solution curves for Eq. (7), including the one passing through the point (0, -1). Note that some solutions grow rapidly in the positive direction as *x* increases, while others grow rapidly in the negative direction. The behavior of a given solution curve is determined by its initial condition $y(0) = y_0$. The two types of behavior are separated by the particular solution $y(x) = -\frac{33}{32}e^{-x/3}$ for which C = 0 in Eq. (8), so $y_0 = -\frac{33}{32}$ for the solution curve that is dashed in Fig. 1.5.1. If $y_0 > -\frac{33}{32}$, then C > 0 in Eq. (8), so the term e^x eventually dominates the behavior of y(x), and hence $y(x) \to +\infty$ as $x \to +\infty$. But if $y_0 < -\frac{33}{32}$, then C < 0, so both terms in y(x) are negative and therefore $y(x) \to -\infty$ as $x \to +\infty$. Thus the initial condition $y_0 = -\frac{33}{32}$ is *critical* in the sense that solutions that start above $-\frac{33}{32}$ on the *y*-axis grow in the positive direction, while solutions that start lower than $-\frac{33}{32}$ grow in the negative direction as $x \to +\infty$. The interpretation of a mathematical model often hinges on finding such a critical condition that separates one kind of behavior of a solution from a different kind of behavior.



FIGURE 1.5.1. Slope field and solution curves for $y' = y + \frac{11}{8}e^{-x/3}$.

Example 2 Find a general solution of

$$(x^2 + 1)\frac{dy}{dx} + 3xy = 6x.$$
 (9)

Solution

After division of both sides of the equation by
$$x^2 + 1$$
, we recognize the result

$$\frac{dy}{dx} + \frac{3x}{x^2 + 1}y = \frac{6x}{x^2 + 1}$$

as a first-order linear equation with $P(x) = 3x/(x^2 + 1)$ and $Q(x) = 6x/(x^2 + 1)$. Multiplication by

$$\rho(x) = \exp\left(\int \frac{3x}{x^2 + 1} \, dx\right) = \exp\left(\frac{3}{2}\ln(x^2 + 1)\right) = (x^2 + 1)^{3/2}$$

yields

$$(x^{2}+1)^{3/2} \frac{dy}{dx} + 3x(x^{2}+1)^{1/2}y = 6x(x^{2}+1)^{1/2}$$

and thus

$$D_x\left[(x^2+1)^{3/2}y\right] = 6x(x^2+1)^{1/2}.$$

Integration then yields

$$(x^{2}+1)^{3/2}y = \int 6x(x^{2}+1)^{1/2} dx = 2(x^{2}+1)^{3/2} + C.$$

Multiplication of both sides by $(x^2 + 1)^{-3/2}$ gives the general solution

$$y(x) = 2 + C(x^2 + 1)^{-3/2}.$$
 (10)

Remark Figure 1.5.2 shows a slope field and typical solution curves for Eq. (9). Note that, as
$$x \to +\infty$$
, all other solution curves approach the constant solution curve $y(x) \equiv 2$ that corresponds to $C = 0$ in Eq. (10). This constant solution can be described as an *equilibrium solution* of the differential equation, because $y(0) = 2$ implies that $y(x) = 2$ for all x (and thus the value of the solution remains forever where it starts). More generally, the word "equilibrium" connotes "unchanging," so by an equilibrium solution of a differential equation is meant a constant solution $y(x) \equiv c$, for which it follows that $y'(x) \equiv 0$. Note that substitution of $y' = 0$ in the differential equation (9) yields $3xy = 6x$, so it follows that $y = 2$ if $x \neq 0$. Hence we see that $y(x) \equiv 2$ is the only equilibrium solution of this differential equation, as seems visually obvious in Fig. 1.5.2.

A Closer Look at the Method

The preceding derivation of the solution in Eq. (6) of the linear first-order equation y' + Py = Q bears closer examination. Suppose that the coefficient functions P(x) and Q(x) are continuous on the (possibly unbounded) open interval *I*. Then the antiderivatives

$$\int P(x) dx$$
 and $\int \left(Q(x) e^{\int P(x) dx} \right) dx$

exist on *I*. Our derivation of Eq. (6) shows that *if* y = y(x) is a solution of Eq. (3) on *I*, *then* y(x) is given by the formula in Eq. (6) for some choice of the constant *C*. Conversely, you may verify by direct substitution (Problem 31) that the function y(x) given in Eq. (6) satisfies Eq. (3). Finally, given a point x_0 of *I* and any number y_0 , there is—as previously noted—a unique value of *C* such that $y(x_0) = y_0$. Consequently, we have proved the following existence-uniqueness theorem.



FIGURE 1.5.2. Slope field and solution curves for the differential equation in Eq. (9).

THEOREM 1 The Linear First-Order Equation

If the functions P(x) and Q(x) are continuous on the open interval *I* containing the point x_0 , then the initial value problem

has a unique solution y(x) on I, given by the formula in Eq. (6) with an appropriate value of C.

Remark 1 Theorem 1 gives a solution on the *entire* interval *I* for a *linear* differential equation, in contrast with Theorem 1 of Section 1.3, which guarantees only a solution on a possibly smaller interval.

Remark 2 Theorem 1 tells us that every solution of Eq. (3) is included in the general solution given in Eq. (6). Thus a *linear* first-order differential equation has *no* singular solutions.

Remark 3 The appropriate value of the constant C in Eq. (6)—as needed to solve the initial value problem in Eq. (11)—can be selected "automatically" by writing

$$\rho(x) = \exp\left(\int_{x_0}^x P(t) dt\right),$$

$$y(x) = \frac{1}{\rho(x)} \left[y_0 + \int_{x_0}^x \rho(t) Q(t) dt\right].$$
(12)

The indicated limits x_0 and x effect a choice of indefinite integrals in Eq. (6) that guarantees in advance that $\rho(x_0) = 1$ and that $y(x_0) = y_0$ (as you can verify directly by substituting $x = x_0$ in Eq. (12)).

Example 3 Solve the initial value problem

$$x^{2}\frac{dy}{dx} + xy = \sin x, \quad y(1) = y_{0}.$$
 (13)

Solution Division by x^2 gives the linear first-order equation

$$\frac{dy}{dx} + \frac{1}{x}y = \frac{\sin x}{x^2}$$

with P(x) = 1/x and $Q(x) = (\sin x)/x^2$. With $x_0 = 1$ the integrating factor in (12) is

$$\rho(x) = \exp\left(\int_{1}^{x} \frac{1}{t} dt\right) = \exp(\ln x) = x,$$

so the desired particular solution is given by

$$y(x) = \frac{1}{x} \left[y_0 + \int_1^x \frac{\sin t}{t} \, dt \right].$$
 (14)

,

In accord with Theorem 1, this solution is defined on the whole positive *x*-axis.

Comment In general, an integral such as the one in Eq. (14) would (for given x) need to be approximated numerically—using Simpson's rule, for instance—to find the value y(x) of the solution at x. In this case, however, we have the sine integral function

$$\operatorname{Si}(x) = \int_0^x \frac{\sin t}{t} \, dt$$



FIGURE 1.5.3. Typical solution curves defined by Eq. (15).



FIGURE 1.5.4. The single-tank mixture problem.

which appears with sufficient frequency in applications that its values have been tabulated. A good set of tables of special functions is Abramowitz and Stegun, *Handbook of Mathematical Functions* (New York: Dover, 1965). Then the particular solution in Eq. (14) reduces to

$$y(x) = \frac{1}{x} \left[y_0 + \int_0^x \frac{\sin t}{t} dt - \int_0^1 \frac{\sin t}{t} dt \right] = \frac{1}{x} \left[y_0 + \operatorname{Si}(x) - \operatorname{Si}(1) \right].$$
(15)

The sine integral function is available in most scientific computing systems and can be used to plot typical solution curves defined by Eq. (15). Figure 1.5.3 shows a selection of solution curves with initial values $y(1) = y_0$ ranging from $y_0 = -3$ to $y_0 = 3$. It appears that on each solution curve, $y(x) \rightarrow 0$ as $x \rightarrow +\infty$, and this is in fact true because the sine integral function is bounded.

In the sequel we will see that it is the exception—rather than the rule—when a solution of a differential equation can be expressed in terms of elementary functions. We will study various devices for obtaining good approximations to the values of the nonelementary functions we encounter. In Chapter 2 we will discuss numerical integration of differential equations in some detail.

Mixture Problems

As a first application of linear first-order equations, we consider a tank containing a solution—a mixture of solute and solvent—such as salt dissolved in water. There is both inflow and outflow, and we want to compute the *amount* x(t) of solute in the tank at time t, given the amount $x(0) = x_0$ at time t = 0. Suppose that solution with a concentration of c_i grams of solute per liter of solution flows into the tank at the constant rate of r_i liters per second, and that the solution in the tank—kept thoroughly mixed by stirring—flows out at the constant rate of r_o liters per second.

To set up a differential equation for x(t), we estimate the change Δx in x during the brief time interval $[t, t + \Delta t]$. The amount of solute that flows into the tank during Δt seconds is $r_i c_i \Delta t$ grams. To check this, note how the cancellation of dimensions checks our computations:

$$\left(r_i \frac{\text{liters}}{\text{second}}\right) \left(c_i \frac{\text{grams}}{\text{liter}}\right) (\Delta t \text{ seconds})$$

yields a quantity measured in grams.

The amount of solute that flows out of the tank during the same time interval depends on the concentration $c_o(t)$ of solute in the solution at time t. But as noted in Fig. 1.5.4, $c_o(t) = x(t)/V(t)$, where V(t) denotes the volume (not constant unless $r_i = r_o$) of solution in the tank at time t. Then

$$\Delta x = \{\text{grams input}\} - \{\text{grams output}\} \approx r_i c_i \Delta t - r_o c_o \Delta t$$

We now divide by Δt :

$$\frac{\Delta x}{\Delta t} \approx r_i c_i - r_o c_o.$$

Finally, we take the limit as $\Delta t \rightarrow 0$; if all the functions involved are continuous and x(t) is differentiable, then the error in this approximation also approaches zero, and we obtain the differential equation

$$\frac{dx}{dt} = r_i c_i - r_o c_o, \tag{16}$$

in which r_i , c_i , and r_o are constants, but c_o denotes the variable concentration

$$c_o(t) = \frac{x(t)}{V(t)} \tag{17}$$

of solute in the tank at time t. Thus the amount x(t) of solute in the tank satisfies the differential equation

$$\frac{dx}{dt} = r_i c_i - \frac{r_o}{V} x. \tag{18}$$

If $V_0 = V(0)$, then $V(t) = V_0 + (r_i - r_o)t$, so Eq. (18) is a linear first-order differential equation for the amount x(t) of solute in the tank at time t.

Important Equation (18) need not be committed to memory. It is the *process* we used to obtain that equation—examination of the behavior of the system over a short time interval $[t, t + \Delta t]$ —that you should strive to understand, because it is a very useful tool for obtaining all sorts of differential equations.

Remark It was convenient for us to use g/L mass/volume units in deriving Eq. (18). But any other consistent system of units can be used to measure amounts of solute and volumes of solution. In the following example we measure both in cubic kilometers.

Example 4 Mixture problem Assume that Lake Erie has a volume of 480 km³ and that its rate of inflow (from Lake Huron) and outflow (to Lake Ontario) are both 350 km³ per year. Suppose that at the time t = 0 (years), the pollutant concentration of Lake Erie—caused by past industrial pollution that has now been ordered to cease—is five times that of Lake Huron. If the outflow henceforth is perfectly mixed lake water, how long will it take to reduce the pollution concentration in Lake Erie to twice that of Lake Huron?

Solution Here we have

 $V = 480 \text{ (km}^3\text{)},$ $r_i = r_o = r = 350 \text{ (km}^3/\text{yr}),$ $c_i = c \text{ (the pollutant concentration of Lake Huron), and}$ $x_0 = x(0) = 5cV,$

and the question is this: When is x(t) = 2cV? With this notation, Eq. (18) is the separable equation

$$\frac{dx}{dt} = rc - \frac{r}{V}x,\tag{19}$$

which we rewrite in the linear first-order form

$$\frac{dx}{dt} + px = q \tag{20}$$

with constant coefficients p = r/V, q = rc, and integrating factor $\rho = e^{pt}$. You can either solve this equation directly or apply the formula in (12). The latter gives

$$x(t) = e^{-pt} \left[x_0 + \int_0^t q e^{pt} dt \right] = e^{-pt} \left[x_0 + \frac{q}{p} \left(e^{pt} - 1 \right) \right]$$

= $e^{-rt/V} \left[5cV + \frac{rc}{r/V} \left(e^{rt/V} - 1 \right) \right];$
 $x(t) = cV + 4cVe^{-rt/V}.$ (21)

To find when x(t) = 2cV, we therefore need only solve the equation

$$cV + 4cVe^{-rt/V} = 2cV$$
 for $t = \frac{V}{r}\ln 4 = \frac{480}{350}\ln 4 \approx 1.901$ (years).

- **Example 5** Mixture problem A 120-gallon (gal) tank initially contains 90 lb of salt dissolved in 90 gal of water. Brine containing 2 lb/gal of salt flows into the tank at the rate of 4 gal/min, and the well-stirred mixture flows out of the tank at the rate of 3 gal/min. How much salt does the tank contain when it is full?
 - Solution The interesting feature of this example is that, due to the differing rates of inflow and outflow, the volume of brine in the tank increases steadily with V(t) = 90 + t gallons. The change Δx in the amount x of salt in the tank from time t to time $t + \Delta t$ (minutes) is given by

$$\Delta x \approx (4)(2) \Delta t - 3\left(\frac{x}{90+t}\right) \Delta t,$$

so our differential equation is

$$\frac{dx}{dt} + \frac{3}{90+t}x = 8.$$

An integrating factor is

$$\rho(x) = \exp\left(\int \frac{3}{90+t} \, dt\right) = e^{3\ln(90+t)} = (90+t)^3,$$

which gives

$$D_t \left[(90+t)^3 x \right] = 8(90+t)^3;$$

(90+t)³x = 2(90+t)⁴ + C

Substitution of x(0) = 90 gives $C = -(90)^4$, so the amount of salt in the tank at time t is

$$x(t) = 2(90+t) - \frac{90^4}{(90+t)^3}.$$

The tank is full after 30 min, and when t = 30, we have

$$x(30) = 2(90 + 30) - \frac{90^4}{120^3} \approx 202 \text{ (lb)}$$

of salt in the tank.

1.5 Problems

Find general solutions of the differential equations in Problems 1 through 25. If an initial condition is given, find the corresponding particular solution. Throughout, primes denote derivatives with respect to x.

1.
$$y' + y = 2$$
, $y(0) = 0$
3. $y' + 3y = 2xe^{-3x}$
5. $xy' + 2y = 3x$, $y(1) = 5$
6. $xy' + 5y = 7x^2$, $y(2) = 5$
7. $2xy' + y = 10\sqrt{x}$
8. $3xy' + y = 12x$
9. $xy' - y = x$, $y(1) = 7$
10. $2xy' - 3y = 9x^3$
11. $xy' + y = 3xy$, $y(1) = 0$
12. $xy' + 3y = 2x^5$, $y(2) = 1$
13. $y' + y = e^x$, $y(0) = 1$
14. $xy' - 3y = x^3$, $y(1) = 10$
15. $y' + 2xy = x$, $y(0) = -2$
16. $y' = (1 - y)\cos x$, $y(\pi) = 2$
17. $(1 + x)y' + y = \cos x$, $y(0) = 1$
18. $xy' = 2y + x^3 \cos x$

19. $y' + y \cot x = \cos x$ **20.** y' = 1 + x + y + xy, y(0) = 0 **21.** $xy' = 3y + x^4 \cos x, y(2\pi) = 0$ **22.** $y' = 2xy + 3x^2 \exp(x^2), y(0) = 5$ **23.** $xy' + (2x - 3)y = 4x^4$ **24.** $(x^2 + 4)y' + 3xy = x, y(0) = 1$ **25.** $(x^2 + 1)\frac{dy}{dx} + 3x^3y = 6x \exp\left(-\frac{3}{2}x^2\right), y(0) = 1$

Solve the differential equations in Problems 26 through 28 by regarding y as the independent variable rather than x.

26.
$$(1 - 4xy^2)\frac{dy}{dx} = y^3$$

27. $(x + ye^y)\frac{dy}{dx} = 1$
28. $(1 + 2xy)\frac{dy}{dx} = 1 + y^2$

29. Express the general solution of dy/dx = 1 + 2xy in terms of the **error function**

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt.$$

54 CHAPTER 1 FIRST-ORDER DIFFERENTIAL EQUATIONS

30. Express the solution of the initial value problem

$$2x\frac{dy}{dx} = y + 2x\cos x, \quad y(1) = 0$$

as an integral as in Example 3 of this section.

Problems 31 and 32 illustrate—for the special case of firstorder linear equations—techniques that will be important when we study higher-order linear equations in Chapter 3.

31. (a) Show that

$$v_c(x) = Ce^{-\int P(x) dx}$$

is a general solution of dy/dx + P(x)y = 0. (b) Show that

$$y_{p}(x) = e^{-\int P(x) dx} \left[\int \left(Q(x) e^{\int P(x) dx} \right) dx \right]$$

is a particular solution of dy/dx + P(x)y = Q(x). (c) Suppose that $y_c(x)$ is any general solution of dy/dx + P(x)y = 0 and that $y_p(x)$ is any particular solution of dy/dx + P(x)y = Q(x). Show that $y(x) = y_c(x) + y_p(x)$ is a general solution of dy/dx + P(x)y = Q(x).

32. (a) Find constants A and B such that $y_p(x) = A \sin x + B \cos x$ is a solution of $dy/dx + y = 2 \sin x$. (b) Use the result of part (a) and the method of Problem 31 to find the general solution of $dy/dx + y = 2 \sin x$. (c) Solve the initial value problem $dy/dx + y = 2 \sin x$, y(0) = 1.

Mixture Problems

Problems 33 through 37 illustrate the application of linear first-order differential equations to mixture problems.

- **33.** A tank contains 1000 liters (L) of a solution consisting of 100 kg of salt dissolved in water. Pure water is pumped into the tank at the rate of 5 L/s, and the mixture—kept uniform by stirring— is pumped out at the same rate. How long will it be until only 10 kg of salt remains in the tank?
- **34.** Consider a reservoir with a volume of 8 billion cubic feet (ft^3) and an initial pollutant concentration of 0.25%. There is a daily inflow of 500 million ft^3 of water with a pollutant concentration of 0.05% and an equal daily outflow of the well-mixed water in the reservoir. How long will it take to reduce the pollutant concentration in the reservoir to 0.10%?
- **35.** Rework Example 4 for the case of Lake Ontario, which empties into the St. Lawrence River and receives inflow from Lake Erie (via the Niagara River). The only differences are that this lake has a volume of 1640 km³ and an inflow-outflow rate of 410 km³/year.
- 36. A tank initially contains 60 gal of pure water. Brine containing 1 lb of salt per gallon enters the tank at 2 gal/min, and the (perfectly mixed) solution leaves the tank at 3 gal/min; thus the tank is empty after exactly 1 h. (a) Find the amount of salt in the tank after t minutes. (b) What is the maximum amount of salt ever in the tank?

- **37.** A 400-gal tank initially contains 100 gal of brine containing 50 lb of salt. Brine containing 1 lb of salt per gallon enters the tank at the rate of 5 gal/s, and the well-mixed brine in the tank flows out at the rate of 3 gal/s. How much salt will the tank contain when it is full of brine?
- **38.** Two tanks Consider the *cascade* of two tanks shown in Fig. 1.5.5, with $V_1 = 100$ (gal) and $V_2 = 200$ (gal) the volumes of brine in the two tanks. Each tank also initially contains 50 lb of salt. The three flow rates indicated in the figure are each 5 gal/min, with pure water flowing into tank 1. (a) Find the amount x(t) of salt in tank 1 at time *t*. (b) Suppose that y(t) is the amount of salt in tank 2 at time *t*. Show first that

$$\frac{dy}{dt} = \frac{5x}{100} - \frac{5y}{200},$$

and then solve for y(t), using the function x(t) found in part (a). (c) Finally, find the maximum amount of salt ever in tank 2.



FIGURE 1.5.5. A cascade of two tanks.

- **39.** Two tanks Suppose that in the cascade shown in Fig. 1.5.5, tank 1 initially contains 100 gal of pure ethanol and tank 2 initially contains 100 gal of pure water. Pure water flows into tank 1 at 10 gal/min, and the other two flow rates are also 10 gal/min. (a) Find the amounts x(t) and y(t) of ethanol in the two tanks at time $t \ge 0$. (b) Find the maximum amount of ethanol ever in tank 2.
- **40.** Multiple tanks A multiple cascade is shown in Fig. 1.5.6. At time t = 0, tank 0 contains 1 gal of ethanol and 1 gal of water; all the remaining tanks contain 2 gal of pure water each. Pure water is pumped into tank 0 at 1 gal/min, and the varying mixture in each tank is pumped into the one below it at the same rate. Assume, as usual, that the mixtures are kept perfectly uniform by stirring. Let $x_n(t)$ denote the amount of ethanol in tank *n* at time *t*.



FIGURE 1.5.6. A multiple cascade.

(a) Show that $x_0(t) = e^{-t/2}$. (b) Show by induction on *n* that

$$x_n(t) = \frac{t^n e^{-t/2}}{n! 2^n}$$
 for $n \ge 0$.

(c) Show that the maximum value of $x_n(t)$ for n > 0 is $M_n = x_n(2n) = n^n e^{-n}/n!$. (d) Conclude from Stirling's approximation $n! \approx n^n e^{-n} \sqrt{2\pi n}$ that $M_n \approx (2\pi n)^{-1/2}$.

- **41.** Retirement savings A 30-year-old woman accepts an engineering position with a starting salary of \$30,000 per year. Her salary S(t) increases exponentially, with $S(t) = 30e^{t/20}$ thousand dollars after *t* years. Meanwhile, 12% of her salary is deposited continuously in a retirement account, which accumulates interest at a continuous annual rate of 6%. (a) Estimate ΔA in terms of Δt to derive the differential equation satisfied by the amount A(t) in her retirement account after *t* years. (b) Compute A(40), the amount available for her retirement at age 70.
- **42.** Falling hailstone Suppose that a falling hailstone with density $\delta = 1$ starts from rest with negligible radius r = 0. Thereafter its radius is r = kt (k is a constant) as it grows by accretion during its fall. Use Newton's second law—according to which the net force F acting on a possibly variable mass m equals the time rate of change dp/dt of its momentum p = mv—to set up and solve the initial value problem

$$\frac{d}{dt}(mv) = mg, \quad v(0) = 0,$$

where *m* is the variable mass of the hailstone, v = dy/dt is its velocity, and the positive *y*-axis points downward. Then show that dv/dt = g/4. Thus the hailstone falls as though it were under *one-fourth* the influence of gravity. 43. Figure 1.5.7 shows a slope field and typical solution curves for the equation y' = x - y. (a) Show that every solution curve approaches the straight line y = x - 1 as x → +∞. (b) For each of the five values y₁ = 3.998, 3.999, 4.000, 4.001, and 4.002, determine the initial value y₀ (accurate to four decimal places) such that y(5) = y₁ for the solution satisfying the initial condition y(-5) = y₀.



FIGURE 1.5.7. Slope field and solution curves for y' = x - y.

44. Figure 1.5.8 shows a slope field and typical solution curves for the equation y' = x + y. (a) Show that every solution curve approaches the straight line y = -x - 1 as x → -∞. (b) For each of the five values y₁ = -10, -5, 0, 5, and 10, determine the initial value y₀ (accurate to five decimal places) such that y(5) = y₁ for the solution satisfying the initial condition y(-5) = y₀.



FIGURE 1.5.8. Slope field and solution curves for y' = x + y.

Polluted Reservoir

Problems 45 and 46 deal with a shallow reservoir that has a one-square-kilometer water surface and an average water depth of 2 meters. Initially it is filled with fresh water, but at time t = 0 water contaminated with a liquid pollutant begins flowing into the reservoir at the rate of 200 thousand cubic meters per month. The well-mixed water in the reservoir flows out at the same rate. Your first task is to find the amount x(t) of pollutant (in millions of liters) in the reservoir after t months.

45. The incoming water has a pollutant concentration of c(t) = 10 liters per cubic meter (L/m³). Verify that the graph of x(t) resembles the steadily rising curve in

56 CHAPTER 1 FIRST-ORDER DIFFERENTIAL EQUATIONS

Fig. 1.5.9, which approaches asymptotically the graph of the equilibrium solution $x(t) \equiv 20$ that corresponds to the reservoir's long-term pollutant content. How long does it take the pollutant concentration in the reservoir to reach 10 L/m³?

46. The incoming water has pollutant concentration $c(t) = 10(1 + \cos t) \text{ L/m}^3$ that varies between 0 and 20, with an average concentration of 10 L/m³ and a period of oscillation of slightly over $6\frac{1}{4}$ months. Does it seem predictable that the lake's pollutant content should ultimately oscillate periodically about an average level of 20 million liters? Verify that the graph of x(t) does, indeed, resemble the oscillatory curve shown in Fig. 1.5.9. How long does

it take the pollutant concentration in the reservoir to reach 10 L/m^3 ?



FIGURE 1.5.9. Graphs of solutions in Problems 45 and 46.

1.5 Application Indoor Temperature Oscillations

Go to goo.gl/QVuenz to download this application's computing resources including *Maple/Mathematica*/MATLAB/ Python. For an interesting applied problem that involves the solution of a linear differential equation, consider indoor temperature oscillations that are driven by outdoor temperature oscillations of the form

$$A(t) = a_0 + a_1 \cos \omega t + b_1 \sin \omega t. \tag{1}$$

If $\omega = \pi/12$, then these oscillations have a period of 24 hours (so that the cycle of outdoor temperatures repeats itself daily) and Eq. (1) provides a realistic model for the temperature outside a house on a day when no change in the overall day-to-day weather pattern is occurring. For instance, for a typical July day in Athens, Georgia with a minimum temperature of 70°F when t = 4 (4 A.M.) and a maximum of 90°F when t = 16 (4 P.M.), we would take

$$A(t) = 80 - 10\cos\omega(t - 4) = 80 - 5\cos\omega t - 5\sqrt{3}\sin\omega t.$$
 (2)

We derived Eq. (2) by using the identity $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$ to get $a_0 = 80$, $a_1 = -5$, and $b_1 = -5\sqrt{3}$ in Eq. (1).

If we write Newton's law of cooling (Eq. (3) of Section 1.1) for the corresponding indoor temperature u(t) at time t, but with the outside temperature A(t) given by Eq. (1) instead of a constant ambient temperature A, we get the linear first-order differential equation

$$\frac{du}{dt} = -k(u - A(t));$$

that is,

$$\frac{du}{dt} + ku = k(a_0 + a_1 \cos \omega t + b_1 \sin \omega t)$$
(3)

with coefficient functions $P(t) \equiv k$ and Q(t) = kA(t). Typical values of the proportionality constant k range from 0.2 to 0.5 (although k might be greater than 0.5 for a poorly insulated building with open windows, or less than 0.2 for a well-insulated building with tightly sealed windows).

SCENARIO: Suppose that our air conditioner fails at time $t_0 = 0$ one midnight, and we cannot afford to have it repaired until payday at the end of the month. We therefore want to investigate the resulting indoor temperatures that we must endure for the next several days.

Begin your investigation by solving Eq. (3) with the initial condition $u(0) = u_0$ (the indoor temperature at the time of the failure of the air conditioner). You may want to use the integral formulas in 49 and 50 of the endpapers, or possibly a computer algebra system. You should get the solution

$$u(t) = a_0 + c_0 e^{-kt} + c_1 \cos \omega t + d_1 \sin \omega t,$$
(4)

where

$$c_{0} = u_{0} - a_{0} - \frac{k^{2}a_{1} - k\omega b_{1}}{k^{2} + \omega^{2}},$$
$$c_{1} = \frac{k^{2}a_{1} - k\omega b_{1}}{k^{2} + \omega^{2}}, \quad d_{1} = \frac{k\omega a_{1} + k^{2}b_{1}}{k^{2} + \omega^{2}}$$

with $\omega = \pi/12$.

With $a_0 = 80$, $a_1 = -5$, $b_1 = -5\sqrt{3}$ (as in Eq. (2)), $\omega = \pi/12$, and k = 0.2 (for instance), this solution reduces (approximately) to

$$u(t) = 80 + e^{-t/5} \left(u_0 - 82.3351 \right) + (2.3351) \cos \frac{\pi t}{12} - (5.6036) \sin \frac{\pi t}{12}.$$
 (5)

Observe first that the "damped" exponential term in Eq. (5) approaches zero as $t \to +\infty$, leaving the long-term "steady periodic" solution

$$u_{\rm sp}(t) = 80 + (2.3351)\cos\frac{\pi t}{12} - (5.6036)\sin\frac{\pi t}{12}.$$
 (6)

Consequently, the long-term indoor temperatures oscillate every 24 hours around the same average temperature 80°F as the average outdoor temperature.

Figure 1.5.10 shows a number of solution curves corresponding to possible initial temperatures u_0 ranging from 65°F to 95°F. Observe that—whatever the initial temperature—the indoor temperature "settles down" within about 18 hours to a periodic daily oscillation. But the amplitude of temperature variation is less indoors than outdoors. Indeed, using the trigonometric identity mentioned earlier, Eq. (6) can be rewritten (verify this!) as

$$u(t) = 80 - (6.0707) \cos\left(\frac{\pi t}{12} - 1.9656\right)$$
$$= 80 - (6.0707) \cos\frac{\pi}{12}(t - 7.5082). \tag{7}$$

Do you see that this implies that the indoor temperature varies between a minimum of about 74°F and a maximum of about 86°F?

Finally, comparison of Eqs. (2) and (7) indicates that the indoor temperature lags behind the outdoor temperature by about $7.5082 - 4 \approx 3.5$ hours, as illustrated in Fig. 1.5.11. Thus the temperature inside the house continues to rise until about 7:30 P.M. each evening, so the hottest part of the day inside is early evening rather than late afternoon (as outside).



FIGURE 1.5.10. Solution curves given by Eq. (5) with $u_0 = 65, 68, 71, \dots, 92, 95$.



FIGURE 1.5.11. Comparison of indoor and outdoor temperature oscillations.
For a personal problem to investigate, carry out a similar analysis using average July daily maximum/minimum figures for your own locale and a value of k appropriate to your own home. You might also consider a winter day instead of a summer day. (What is the winter-summer difference for the indoor temperature problem?) You may wish to explore the use of available technology both to solve the differential equation and to graph its solution for the indoor temperature in comparison with the outdoor temperature.

1.6 Substitution Methods and Exact Equations

The first-order differential equations we have solved in the previous sections have all been either separable or linear. But many applications involve differential equations that are neither separable nor linear. In this section we illustrate (mainly with examples) substitution methods that sometimes can be used to transform a given differential equation into one that we already know how to solve.

For instance, the differential equation

$$\frac{dy}{dx} = f(x, y),\tag{1}$$

with dependent variable y and independent variable x, may contain a conspicuous combination

$$v = \alpha(x, y) \tag{2}$$

of x and y that suggests itself as a new independent variable v. Thus the differential equation

$$\frac{dy}{dx} = (x+y+3)^2$$

practically demands the substitution v = x + y + 3 of the form in Eq. (2).

If the substitution relation in Eq. (2) can be solved for

$$v = \beta(x, v), \tag{3}$$

then application of the chain rule—regarding v as an (unknown) function of x—yields

$$\frac{dy}{dx} = \frac{\partial\beta}{\partial x}\frac{dx}{dx} + \frac{\partial\beta}{\partial v}\frac{dv}{dx} = \beta_x + \beta_v\frac{dv}{dx},\tag{4}$$

where the partial derivatives $\partial \beta / \partial x = \beta_x(x, v)$ and $\partial \beta / \partial v = \beta_v(x, v)$ are *known* functions of x and v. If we substitute the right-hand side in (4) for dy/dx in Eq. (1) and then solve for dv/dx, the result is a *new* differential equation of the form

$$\frac{dv}{dx} = g(x, v) \tag{5}$$

with *new* dependent variable v. If this new equation is either separable or linear, then we can apply the methods of preceding sections to solve it.

If v = v(x) is a solution of Eq. (5), then $y = \beta(x, v(x))$ will be a solution of the original Eq. (1). The trick is to select a substitution such that the transformed Eq. (5) is one we can solve. Even when possible, this is not always easy; it may require a fair amount of ingenuity or trial and error.

$$\frac{dy}{dx} = (x+y+3)^2$$

1.6 SUBSTITUTION METHODS AND EXACT EQUATIONS59

Solution As indicated earlier, let's try the substitution

$$v = x + y + 3$$
; that is, $y = v - x - 3$.

Then

$$\frac{dy}{dx} = \frac{dv}{dx} - 1,$$

so the transformed equation is

$$\frac{lv}{lx} = 1 + v^2.$$

This is a separable equation, and we have no difficulty in obtaining its solution

$$x = \int \frac{dv}{1 + v^2} = \tan^{-1} v + C.$$

So $v = \tan(x - C)$. Because v = x + y + 3, the general solution of the original equation $dy/dx = (x + y + 3)^2$ is $x + y + 3 = \tan(x - C)$; that is,

$$y(x) = \tan(x - C) - x - 3.$$

Remark Figure 1.6.1 shows a slope field and typical solution curves for the differential equation of Example 1. We see that, although the function $f(x, y) = (x + y + 3)^2$ is continuously differentiable for all x and y, each solution is continuous only on a bounded interval. In particular, because the tangent function is continuous on the open interval $(-\pi/2, \pi/2)$, the particular solution with arbitrary constant value C is continuous on the interval where $-\pi/2 < x - C < \pi/2$; that is, $C - \pi/2 < x < C + \pi/2$. This situation is fairly typical of non-linear differential equations, in contrast with linear differential equations, whose solutions are continuous wherever the coefficient functions in the equation are continuous.

Example 1 illustrates the fact that any differential equation of the form

$$\frac{dy}{dx} = F(ax + by + c) \tag{6}$$

can be transformed into a separable equation by use of the substitution v = ax + by + c (see Problem 55). The paragraphs that follow deal with other classes of first-order equations for which there are standard substitutions that are known to succeed.

Homogeneous Equations

A **homogeneous** first-order differential equation is one that can be written in the form

$$\frac{dy}{dx} = F\left(\frac{y}{x}\right).\tag{7}$$

If we make the substitutions

>

$$v = \frac{y}{x}, \quad y = vx, \quad \frac{dy}{dx} = v + x\frac{dv}{dx},$$
 (8)

then Eq. (7) is transformed into the *separable* equation

$$x\frac{dv}{dx} = F(v) - v$$

Thus every homogeneous first-order differential equation can be reduced to an integration problem by means of the substitutions in (8).



FIGURE 1.6.1. Slope field and solution curves for $y' = (x + y + 3)^2$.

Remark A dictionary definition of "homogeneous" is "of a similar kind or nature." Consider a differential equation of the form

$$Ax^m y^n \frac{dy}{dx} = Bx^p y^q + Cx^r y^s, \qquad (*)$$

whose polynomial coefficient functions are "homogeneous" in the sense that each of their terms has the same total degree, m + n = p + q = r + s = K. If we divide each side of (*) by x^{K} , then the result—because $x^{m}y^{n}/x^{m+n} = (y/x)^{n}$, and so forth—is the equation

$$A\left(\frac{y}{x}\right)^n \frac{dy}{dx} = B\left(\frac{y}{x}\right)^q + C\left(\frac{y}{x}\right)^s,$$

which evidently can be written (by another division) in the form of Eq. (7). More generally, a differential equation of the form P(x, y)y' = Q(x, y) with polynomial coefficients P and Q is homogeneous if the terms in these polynomials all have the same total degree K. The differential equation in the following example is of this form with K = 2.

Example 2

Solve the differential equation

$$2xy\frac{dy}{dx} = 4x^2 + 3y^2$$

Solution This equation is neither separable nor linear, but we recognize it as a homogeneous equation by writing it in the form

$$\frac{dy}{dx} = \frac{4x^2 + 3y^2}{2xy} = 2\left(\frac{x}{y}\right) + \frac{3}{2}\left(\frac{y}{x}\right).$$

The substitutions in (8) then take the form

$$y = vx$$
, $\frac{dy}{dx} = v + x \frac{dv}{dx}$, $v = \frac{y}{x}$, and $\frac{1}{v} = \frac{x}{y}$.

 $v + x\frac{dv}{dx} = \frac{2}{v} + \frac{3}{2}v,$

These yield

and hence

$$x\frac{dv}{dx} = \frac{2}{v} + \frac{v}{2} = \frac{v^2 + 4}{2v};$$
$$\int \frac{2v}{v^2 + 4} \, dv = \int \frac{1}{x} \, dx;$$
$$\ln(v^2 + 4) = \ln|x| + \ln C.$$

We apply the exponential function to both sides of the last equation to obtain

$$v^{2} + 4 = C|x|;$$

 $\frac{y^{2}}{x^{2}} + 4 = C|x|;$
 $y^{2} + 4x^{2} = kx^{3}.$

Note that the left-hand side of this equation is necessarily nonnegative. It follows that
$$k > 0$$
 in the case of solutions that are defined for $x > 0$, while $k < 0$ for solutions where $x < 0$.
Indeed, the family of solution curves illustrated in Fig. 1.6.2 exhibits symmetry about both coordinate axes. Actually, there are positive-valued and negative-valued solutions of the forms $y(x) = \pm \sqrt{kx^3 - 4x^2}$ that are defined for $x > 4/k$ if the constant k is positive, and for $x < 4/k$ if k is negative.



FIGURE 1.6.2. Slope field and solution curves for $2xyy' = 4x^2 + 3y^2$.

Example 3 Solve the in

Solve the initial value problem

$$x\frac{dy}{dx} = y + \sqrt{x^2 - y^2}, \quad y(x_0) = 0,$$

where $x_0 > 0$.

Solution

We divide both sides by x and find that

$$\frac{dy}{dx} = \frac{y}{x} + \sqrt{1 - \left(\frac{y}{x}\right)^2},$$

so we make the substitutions in (8); we get

$$v + x\frac{dv}{dx} = v + \sqrt{1 - v^2};$$
$$\int \frac{1}{\sqrt{1 - v^2}} dv = \int \frac{1}{x} dx;$$
$$\sin^{-1} v = \ln x + C.$$

We need not write $\ln |x|$ because x > 0 near $x = x_0 > 0$. Now note that $v(x_0) = y(x_0)/x_0 = 0$, so $C = \sin^{-1} 0 - \ln x_0 = -\ln x_0$. Hence

$$v = \frac{y}{x} = \sin\left(\ln x - \ln x_0\right) = \sin\left(\ln \frac{x}{x_0}\right),$$

and therefore

>

$$y(x) = x \sin\left(\ln\frac{x}{x_0}\right)$$

is the desired particular solution. Figure 1.6.3 shows some typical solution curves. Because of the radical in the differential equation, these solution curves are confined to the indicated triangular region $x \ge |y|$. You can check that the boundary lines y = x and y = -x (for x > 0) are singular solution curves that consist of points of tangency with the solution curves found earlier.

Bernoulli Equations

A first-order differential equation of the form

$$\frac{dy}{dx} + P(x)y = Q(x)y^n \tag{9}$$

is called a **Bernoulli equation**. If either n = 0 or n = 1, then Eq. (9) is linear. Otherwise, as we ask you to show in Problem 56, the substitution

$$v = v^{1-n} \tag{10}$$

transforms Eq. (9) into the linear equation

$$\frac{dv}{dx} + (1-n)P(x)v = (1-n)Q(x).$$

Rather than memorizing the form of this transformed equation, it is more efficient to make the substitution in Eq. (10) explicitly, as in the following examples.



FIGURE 1.6.3. Solution curves for $xy' = y + \sqrt{x^2 - y^2}$.

Example 4

If we rewrite the homogeneous equation $2xyy' = 4x^2 + 3y^2$ of Example 2 in the form

$$\frac{dy}{dx} - \frac{3}{2x}y = \frac{2x}{y},$$

we see that it is also a Bernoulli equation with P(x) = -3/(2x), Q(x) = 2x, n = -1, and 1 - n = 2. Hence we substitute

$$v = y^2$$
, $y = v^{1/2}$, and $\frac{dy}{dx} = \frac{dy}{dv}\frac{dv}{dx} = \frac{1}{2}v^{-1/2}\frac{dv}{dx}$.

This gives

$$\frac{1}{2}v^{-1/2}\frac{dv}{dx} - \frac{3}{2x}v^{1/2} = 2xv^{-1/2}.$$

Then multiplication by $2v^{1/2}$ produces the linear equation

$$\frac{dv}{dx} - \frac{3}{x}v = 4x$$

with integrating factor $\rho = e^{\int (-3/x) dx} = x^{-3}$. So we obtain

$$D_{x}(x^{-3}v) = \frac{4}{x^{2}};$$

$$x^{-3}v = -\frac{4}{x} + C;$$

$$x^{-3}y^{2} = -\frac{4}{x} + C;$$

$$y^{2} = -4x^{2} + Cx^{3}.$$

Example 5 The

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

is neither separable nor linear nor homogeneous, but it is a Bernoulli equation with $n = \frac{4}{3}$, $1 - n = -\frac{1}{3}$. The substitutions

$$v = y^{-1/3}$$
, $y = v^{-3}$, and $\frac{dy}{dx} = \frac{dy}{dv}\frac{dv}{dx} = -3v^{-4}\frac{dv}{dx}$

transform it into

$$-3xv^{-4}\frac{dv}{dx} + 6v^{-3} = 3xv^{-4}.$$

Division by $-3xv^{-4}$ yields the linear equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

with integrating factor $\rho = e^{\int (-2/x) dx} = x^{-2}$. This gives

$$D_x(x^{-2}v) = -\frac{1}{x^2}; \quad x^{-2}v = \frac{1}{x} + C; \quad v = x + Cx^2;$$

and finally,

$$y(x) = \frac{1}{(x+Cx^2)^3}.$$

Example 6 The equation

$$2xe^{2y}\frac{dy}{dx} = 3x^4 + e^{2y}$$
(11)

1.6 SUBSTITUTION METHODS AND EXACT EQUATIONS63

is neither separable, nor linear, nor homogeneous, nor is it a Bernoulli equation. But we observe that y appears only in the combinations e^{2y} and $D_x(e^{2y}) = 2e^{2y}y'$. This prompts the substitution

$$v = e^{2y}, \quad \frac{dv}{dx} = 2e^{2y}\frac{dy}{dx}$$

that transforms Eq. (11) into the linear equation $xv'(x) = 3x^4 + v(x)$; that is,

$$\frac{dv}{dx} - \frac{1}{x}v = 3x^3$$

After multiplying by the integrating factor $\rho = 1/x$, we find that

$$\frac{1}{x}v = \int 3x^2 \, dx = x^3 + C, \quad \text{so} \quad e^{2y} = v = x^4 + Cx,$$

and hence

$$y(x) = \frac{1}{2} \ln \left| x^4 + Cx \right|.$$



Suppose that an airplane departs from the point (a, 0) located due east of its intended destination—an airport located at the origin (0, 0). The plane travels with constant speed v_0 relative to the wind, which is blowing due north with constant speed w. As indicated in Fig. 1.6.4, we assume that the plane's pilot maintains its heading directly toward the origin.

Figure 1.6.5 helps us derive the plane's velocity components relative to the ground. They are

$$\frac{dx}{dt} = -v_0 \cos \theta = -\frac{v_0 x}{\sqrt{x^2 + y^2}},$$
$$\frac{dy}{dt} = -v_0 \sin \theta + w = -\frac{v_0 y}{\sqrt{x^2 + y^2}} + w$$

Hence the trajectory y = f(x) of the plane satisfies the differential equation

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{1}{v_0 x} \left(v_0 y - w \sqrt{x^2 + y^2} \right).$$
(12)

If we set

$$k = \frac{w}{v_0},\tag{13}$$

the ratio of the windspeed to the plane's airspeed, then Eq. (12) takes the homogeneous form

$$\frac{dy}{dx} = \frac{y}{x} - k \left[1 + \left(\frac{y}{x}\right)^2 \right]^{1/2}.$$
(14)

The substitution y = xv, y' = v + xv' then leads routinely to

$$\int \frac{dv}{\sqrt{1+v^2}} = -\int \frac{k}{x} \, dx. \tag{15}$$

By trigonometric substitution, or by consulting a table for the integral on the left, we find that

$$\ln\left(v + \sqrt{1 + v^2}\right) = -k\ln x + C,$$
(16)



FIGURE 1.6.4. The airplane headed for the origin.



FIGURE 1.6.5. The components of the velocity vector of the airplane.

and the initial condition v(a) = y(a)/a = 0 yields

1

$$C = k \ln a. \tag{17}$$

As we ask you to show in Problem 68, the result of substituting (17) in Eq. (16) and then solving for v is

$$y = \frac{1}{2} \left[\left(\frac{x}{a}\right)^{-k} - \left(\frac{x}{a}\right)^{k} \right].$$
(18)

Because y = xv, we finally obtain

$$y(x) = \frac{a}{2} \left[\left(\frac{x}{a}\right)^{1-k} - \left(\frac{x}{a}\right)^{1+k} \right]$$
(19)

for the equation of the plane's trajectory.

Note that only in the case k < 1 (that is, $w < v_0$) does the curve in Eq. (19) pass through the origin, so that the plane reaches its destination. If $w = v_0$ (so that k = 1), then Eq. (19) takes the form $y(x) = \frac{1}{2}a(1 - x^2/a^2)$, so the plane's trajectory approaches the point (0, a/2) rather than (0, 0). The situation is even worse if $w > v_0$ (so k > 1)—in this case it follows from Eq. (19) that $y \to +\infty$ as $x \to 0$. The three cases are illustrated in Fig. 1.6.6.

Flight trajectory If a = 200 mi, $v_0 = 500 \text{ mi/h}$, and w = 100 mi/h, then $k = w/v_0 = \frac{1}{5}$, so the plane will succeed in reaching the airport at (0, 0). With these values, Eq. (19) yields

$$y(x) = 100 \left[\left(\frac{x}{200} \right)^{4/5} - \left(\frac{x}{200} \right)^{6/5} \right].$$
 (20)

Now suppose that we want to find the maximum amount by which the plane is blown off course during its trip. That is, what is the maximum value of y(x) for $0 \le x \le 200$? Differentiation of the function in Eq. (20) yields

 $\frac{dy}{dx} = \frac{1}{2} \left[\frac{4}{5} \left(\frac{x}{200} \right)^{-1/5} - \frac{6}{5} \left(\frac{x}{200} \right)^{1/5} \right],$

and we readily solve the equation y'(x) = 0 to obtain $(x/200)^{2/5} = \frac{2}{3}$. Hence

$$y_{\text{max}} = 100 \left[\left(\frac{2}{3} \right)^2 - \left(\frac{2}{3} \right)^3 \right] = \frac{400}{27} \approx 14.81.$$

Thus the plane is blown almost 15 mi north at one point during its westward journey. (The graph of the function in Eq. (20) is the one used to construct Fig. 1.6.4. The vertical scale there is exaggerated by a factor of 4.)

Exact Differential Equations

We have seen that a general solution y(x) of a first-order differential equation is often defined implicitly by an equation of the form

$$F(x, y(x)) = C, \tag{21}$$

where *C* is a constant. On the other hand, given the identity in (21), we can recover the original differential equation by differentiating each side with respect to *x*. Provided that Eq. (21) implicitly defines *y* as a differentiable function of *x*, this gives the original differential equation in the form

$$\frac{\partial F}{\partial x} + \frac{\partial F}{\partial y}\frac{dy}{dx} = 0$$



FIGURE 1.6.6. The three cases $w < v_0$ (plane velocity exceeds wind velocity), $w = v_0$ (equal velocities), and $w > v_0$ (wind is greater).

Example 7

Solution

1.6 SUBSTITUTION METHODS AND EXACT EQUATIONS65

that is,

 \succ

>

$$M(x, y) + N(x, y)\frac{dy}{dx} = 0,$$
 (22)

where $M(x, y) = F_x(x, y)$ and $N(x, y) = F_y(x, y)$.

It is sometimes convenient to rewrite Eq. (22) in the more symmetric form

$$M(x, y) dx + N(x, y) dy = 0,$$
(23)

called its **differential** form. The general first-order differential equation y' = f(x, y) can be written in this form with M = f(x, y) and $N \equiv -1$. The preceding discussion shows that, *if* there exists a function F(x, y) such that

$$\frac{\partial F}{\partial x} = M$$
 and $\frac{\partial F}{\partial y} = N$,

then the equation

$$F(x, y) = C$$

implicitly defines a general solution of Eq. (23). In this case, Eq. (23) is called an **exact differential equation**—the differential

$$dF = F_x \, dx + F_y \, dy$$

of F(x, y) is exactly M dx + N dy.

Natural questions are these: How can we determine whether the differential equation in (23) is exact? And if it is exact, how can we find the function F such that $F_x = M$ and $F_y = N$? To answer the first question, let us recall that if the mixed second-order partial derivatives F_{xy} and F_{yx} are continuous on an open set in the xy-plane, then they are equal: $F_{xy} = F_{yx}$. If Eq. (23) is exact and M and N have continuous partial derivatives, it then follows that

$$\frac{\partial M}{\partial y} = F_{xy} = F_{yx} = \frac{\partial N}{\partial x}.$$

Thus the equation

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \tag{24}$$

is a *necessary condition* that the differential equation M dx + N dy = 0 be exact. That is, if $M_y \neq N_x$, then the differential equation in question is not exact, so we need not attempt to find a function F(x, y) such that $F_x = M$ and $F_y = N$ —there is no such function.

Example 8 The differential equation

$$y^3 \, dx + 3x y^2 \, dy = 0 \tag{25}$$

is exact because we can immediately see that the function $F(x, y) = xy^3$ has the property that $F_x = y^3$ and $F_y = 3xy^2$. Thus a general solution of Eq. (25) is

$$xy^3 = C;$$

if you prefer, $y(x) = kx^{-1/3}$.

But suppose that we divide each term of the differential equation in Example 8 by y^2 to obtain

$$y \, dx + 3x \, dy = 0.$$
 (26)

This equation is not exact because, with M = y and N = 3x, we have

$$\frac{\partial M}{\partial y} = 1 \neq 3 = \frac{\partial N}{\partial x}.$$

Hence the necessary condition in Eq. (24) is not satisfied.

We are confronted with a curious situation here. The differential equations in (25) and (26) are essentially equivalent, and they have exactly the same solutions, yet one is exact and the other is not. In brief, whether a given differential equation is exact or not is related to the precise form M dx + N dy = 0 in which it is written.

Theorem 1 tells us that (subject to differentiability conditions usually satisfied in practice) the necessary condition in (24) is also a *sufficient* condition for exactness. In other words, if $M_y = N_x$, then the differential equation M dx + N dy = 0 is exact.

THEOREM 1 Criterion for Exactness

Suppose that the functions M(x, y) and N(x, y) are continuous and have continuous first-order partial derivatives in the open rectangle R: a < x < b, c < y < d. Then the differential equation

$$M(x, y) dx + N(x, y) dy = 0$$
 (23)

is exact in R if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$
 (24)

at each point of *R*. That is, there exists a function F(x, y) defined on *R* with $\partial F/\partial x = M$ and $\partial F/\partial y = N$ if and only if Eq. (24) holds on *R*.

Proof: We have seen already that it is necessary for Eq. (24) to hold if Eq. (23) is to be exact. To prove the converse, we must show that if Eq. (24) holds, then we can construct a function F(x, y) such that $\partial F/\partial x = M$ and $\partial F/\partial y = N$. Note first that, for *any* function g(y), the function

$$F(x, y) = \int M(x, y) \, dx + g(y) \tag{27}$$

satisfies the condition $\partial F/\partial x = M$. (In Eq. (27), the notation $\int M(x, y) dx$ denotes an antiderivative of M(x, y) with respect to x.) We plan to choose g(y) so that

$$N = \frac{\partial F}{\partial y} = \left(\frac{\partial}{\partial y} \int M(x, y) \, dx\right) + g'(y)$$

as well; that is, so that

$$g'(y) = N - \frac{\partial}{\partial y} \int M(x, y) \, dx.$$
⁽²⁸⁾

To see that there *is* such a function of *y*, it suffices to show that the right-hand side in Eq. (28) is a function of *y* alone. We can then find g(y) by integrating with respect

1.6 SUBSTITUTION METHODS AND EXACT EQUATIONS67

to y. Because the right-hand side in Eq. (28) is defined on a rectangle, and hence on an interval as a function of x, it suffices to show that its derivative with respect to x is identically zero. But

$$\frac{\partial}{\partial x} \left(N - \frac{\partial}{\partial y} \int M(x, y) \, dx \right) = \frac{\partial N}{\partial x} - \frac{\partial}{\partial x} \frac{\partial}{\partial y} \int M(x, y) \, dx$$
$$= \frac{\partial N}{\partial x} - \frac{\partial}{\partial y} \frac{\partial}{\partial x} \int M(x, y) \, dx$$
$$= \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = 0$$

by hypothesis. So we can, indeed, find the desired function g(y) by integrating Eq. (28). We substitute this result in Eq. (27) to obtain

$$F(x, y) = \int M(x, y) \, dx + \int \left(N(x, y) - \frac{\partial}{\partial y} \int M(x, y) \, dx \right) dy \tag{29}$$

as the desired function with $F_x = M$ and $F_y = N$.

Instead of memorizing Eq. (29), it is usually better to solve an exact equation M dx + N dy = 0 by carrying out the process indicated by Eqs. (27) and (28). First we integrate M(x, y) with respect to x and write

$$F(x, y) = \int M(x, y) \, dx + g(y),$$

thinking of the function g(y) as an "arbitrary constant of integration" as far as the variable x is concerned. Then we determine g(y) by imposing the condition that $\partial F/\partial y = N(x, y)$. This yields a general solution in the implicit form F(x, y) = C.

xample 9 Solve the differential equation

$$(6xy - y^3) dx + (4y + 3x^2 - 3xy^2) dy = 0.$$
(30)

Solution

Let
$$M(x, y) = 6xy - y^3$$
 and $N(x, y) = 4y + 3x^2 - 3xy^2$. The given equation is exact because

$$\frac{\partial M}{\partial y} = 6x - 3y^2 = \frac{\partial N}{\partial x}.$$

Integrating $\partial F/\partial x = M(x, y)$ with respect to x, we get

$$F(x, y) = \int (6xy - y^3) \, dx = 3x^2y - xy^3 + g(y).$$

Then we differentiate with respect to y and set $\partial F/\partial y = N(x, y)$. This yields

$$\frac{\partial F}{\partial y} = 3x^2 - 3xy^2 + g'(y) = 4y + 3x^2 - 3xy^2,$$

and it follows that g'(y) = 4y. Hence $g(y) = 2y^2 + C_1$, and thus

$$F(x, y) = 3x^2y - xy^3 + 2y^2 + C_1.$$

Therefore, a general solution of the differential equation is defined implicitly by the equation

$$3x^2y - xy^3 + 2y^2 = C (31)$$

(we have absorbed the constant C_1 into the constant C).



FIGURE 1.6.7. Slope field and solution curves for the exact equation in Example 9.

Remark Figure 1.6.7 shows a rather complicated structure of solution curves for the differential equation of Example 9. The solution satisfying a given initial condition $y(x_0) = y_0$ is defined implicitly by Eq. (31), with *C* determined by substituting $x = x_0$ and $y = y_0$ in the equation. For instance, the particular solution satisfying y(0) = 1 is defined implicitly by the equation $3x^2y - xy^3 + 2y^2 = 2$. The other two special points in the figure—at (0,0) and near (0.75, 2.12)—are ones where both coefficient functions in Eq. (30) vanish, so the theorem of Section 1.3 does not guarantee a unique solution.

Reducible Second-Order Equations

A second-order differential equation involves the second derivative of the unknown function y(x), and thus has the general form

$$F(x, y, y', y'') = 0.$$
 (32)

If *either* the dependent variable *y* or the independent variable *x* is missing from a second-order equation, then it is easily reduced by a simple substitution to a first-order equation that may be solvable by the methods of this chapter.

Dependent variable y missing. If y is missing, then Eq. (32) takes the form

$$F(x, y', y'') = 0.$$
 (33)

Then the substitution

$$p = y' = \frac{dy}{dx}, \quad y'' = \frac{dp}{dx} \tag{34}$$

results in the first-order differential equation

$$F(x, p, p') = 0.$$

If we can solve this equation for a general solution $p(x, C_1)$ involving an arbitrary constant C_1 , then we need only write

$$y(x) = \int y'(x) \, dx = \int p(x, C_1) \, dx + C_2$$

to get a solution of Eq. (33) that involves two arbitrary constants C_1 and C_2 (as is to be expected in the case of a second-order differential equation).

Example 10 Solution

Solve the equation xy'' + 2y' = 6x in which the dependent variable y is missing. The substitution defined in (34) gives the first-order equation

$$x\frac{dp}{dx} + 2p = 6x;$$
 that is, $\frac{dp}{dx} + \frac{2}{x}p = 6$

25 20 15 10 5 3 0 -5 -10-15-20-25 -2-1 0 1

FIGURE 1.6.8. Solution curves of the form $y(x) = x^2 - \frac{C_1}{x}$ for $C_1 = 0, \pm 3, \pm 10, \pm 20, \pm 35, \pm 60, \pm 100.$

Observing that the equation on the right here is linear, we multiply by its integrating factor $\rho = \exp\left(\int (2/x) dx\right) = e^{2\ln x} = x^2$ and get

$$D_x(x^2 p) = 6x^2,$$

$$x^2 p = 2x^3 + C_1,$$

$$p = \frac{dy}{dx} = 2x + \frac{C_1}{x^2}$$

A final integration with respect to x yields the general solution

$$y(x) = x^2 - \frac{C_1}{x} + C_2$$

of the second-order equation xy'' + 2y' = 6x. Solution curves with $C_1 = 0$ but $C_2 \neq 0$ are simply vertical translates of the parabola $y = x^2$ (for which $C_1 = C_2 = 0$). Figure 1.6.8 shows this parabola and some typical solution curves with $C_2 = 0$ but $C_1 \neq 0$. Solution curves with C_1 and C_2 both nonzero are vertical translates of those (other than the parabola) shown in Fig. 1.6.8.

1.6 SUBSTITUTION METHODS AND EXACT EQUATIONS69

Independent variable x missing. If x is missing, then Eq. (32) takes the form

$$F(y, y', y'') = 0.$$
 (35)

Then the substitution

$$p = y' = \frac{dy}{dx}, \quad y'' = \frac{dp}{dx} = \frac{dp}{dy}\frac{dy}{dx} = p\frac{dp}{dy}$$
(36)

results in the *first-order* differential equation

$$F\left(y, p, p\frac{dp}{dy}\right) = 0$$

for p as a function of y. If we can solve this equation for a general solution $p(y, C_1)$ involving an arbitrary constant C_1 , then (assuming that $y' \neq 0$) we need only write

$$x(y) = \int \frac{dx}{dy} \, dy = \int \frac{1}{dy/dx} \, dy = \int \frac{1}{p} \, dy = \int \frac{dy}{p(y, C_1)} + C_2$$

If the final integral $P = \int (1/p) dy$ can be evaluated, the result is an implicit solution $x(y) = P(y, C_1) + C_2$ of our second-order differential equation.

Example 11 Solve the equation $yy'' = (y')^2$ in which the independent variable x is missing.

Solution We assume temporarily that y and y' are both nonnegative, and then point out at the end that this restriction is unnecessary. The substitution defined in (36) gives the first-order equation

$$yp\frac{dp}{dy} = p^2$$

Then separation of variables gives

$$\int \frac{dp}{p} = \int \frac{dy}{y},$$

ln $p = \ln y + C$ (because $y > 0$ and $p = y' > 0$),
 $p = C_1 y$

where $C_1 = e^C$. Hence

$$\frac{dx}{dy} = \frac{1}{p} = \frac{1}{C_1 y},$$
$$C_1 x = \int \frac{dy}{y} = \ln y + C_2$$

The resulting general solution of the second-order equation $yy'' = (y')^2$ is

$$y(x) = \exp(C_1 x - C_2) = A e^{Bx},$$

where $A = e^{-C_2}$ and $B = C_1$. Despite our temporary assumptions, which imply that the constants *A* and *B* are both positive, we readily verify that $y(x) = Ae^{Bx}$ satisfies $yy'' = (y')^2$ for *all* real values of *A* and *B*. With B = 0 and different values of *A*, we get all horizontal lines in the plane as solution curves. The upper half of Fig. 1.6.9 shows the solution curves obtained with A = 1 (for instance) and different positive values of *B*. With A = -1 these solution curves are reflected in the *x*-axis, and with negative values of *B* they are reflected in the *y*-axis. In particular, we see that we get solutions of $yy'' = (y')^2$, allowing both positive and negative possibilities for both *y* and *y'*.



FIGURE 1.6.9. The solution curves $y = Ae^{Bx}$ with B = 0 and A = 0, ± 1 are the horizontal lines $y = 0, \pm 1$. The exponential curves with B > 0 and $A = \pm 1$ are in color, those with B < 0 and $A = \pm 1$ are black.

1.6 Problems

Find general solutions of the differential equations in Problems 1 through 30. Primes denote derivatives with respect to x throughout.

1. (x + y)y' = x - y**2.** $2xyy' = x^2 + 2y^2$ 4. (x - y)y' = x + y3. $xy' = y + 2\sqrt{xy}$ 5. x(x + y)y' = y(x - y)6. (x + 2y)y' = y8. $x^2 y' = xy + x^2 e^{y/x}$ 7. $xy^2y' = x^3 + y^3$ 9. $x^2y' = xy + y^2$ 11. $(x^2 - y^2)y' = 2xy$ 10. $xyy' = x^2 + 3y^2$ 12. $xyy' = y^2 + x\sqrt{4x^2 + y^2}$ 13. $xy' = y + \sqrt{x^2 + y^2}$ 14. $yy' + x = \sqrt{x^2 + y^2}$ **15.** x(x + y)y' + y(3x + y) = 0**16.** $y' = \sqrt{x + y + 1}$ 17. $y' = (4x + y)^2$ **19.** $x^2 y' + 2xy = 5y^3$ **18.** (x + y)y' = 1**20.** $y^2 y' + 2xy^3 = 6x$ **21.** $y' = y + y^3$ **23.** $xy' + 6y = 3xy^{4/3}$ **22.** $x^2 y' + 2xy = 5y^4$ **24.** $2xy' + y^3e^{-2x} = 2xy$ **25.** $y^2(xy'+y)(1+x^4)^{1/2} = x$ **26.** $3y^2y' + y^3 = e^{-x}$ **27.** $3xy^2y' = 3x^4 + y^3$ **28.** $xe^{y}y' = 2(e^{y} + x^{3}e^{2x})$ **29.** $(2x \sin y \cos y)y' = 4x^2 + \sin^2 y$ **30.** $(x + e^y)y' = xe^{-y} - 1$

In Problems 31 through 42, verify that the given differential equation is exact; then solve it.

31.
$$(2x + 3y) dx + (3x + 2y) dy = 0$$

32. $(4x - y) dx + (6y - x) dy = 0$
33. $(3x^2 + 2y^2) dx + (4xy + 6y^2) dy = 0$
34. $(2xy^2 + 3x^2) dx + (2x^2y + 4y^3) dy = 0$
35. $\left(x^3 + \frac{y}{x}\right) dx + (y^2 + \ln x) dy = 0$
36. $(1 + ye^{xy}) dx + (2y + xe^{xy}) dy = 0$
37. $(\cos x + \ln y) dx + \left(\frac{x}{y} + e^y\right) dy = 0$
38. $(x + \tan^{-1} y) dx + \frac{x + y}{1 + y^2} dy = 0$
39. $(3x^2y^3 + y^4) dx + (3x^3y^2 + y^4 + 4xy^3) dy = 0$
40. $(e^x \sin y + \tan y) dx + (e^x \cos y + x \sec^2 y) dy = 0$
41. $\left(\frac{2x}{y} - \frac{3y^2}{x^4}\right) dx + \left(\frac{2y}{x^3} - \frac{x^2}{y^2} + \frac{1}{\sqrt{y}}\right) dy = 0$
42. $\frac{2x^{5/2} - 3y^{5/3}}{2x^{5/2}y^{2/3}} dx + \frac{3y^{5/3} - 2x^{5/2}}{3x^{3/2}y^{5/3}} dy = 0$

Find a general solution of each reducible second-order differential equation in Problems 43–54. Assume x, y and/or y'positive where helpful (as in Example 11).

43.
$$xy'' = y'$$
 44. $yy'' + (y')^2 = 0$

- **45.** y'' + 4y = 0 **46.** xy'' + y' = 4x
- **47.** $y'' = (y')^2$

49. $yy'' + (y')^2 = yy'$

51. $y'' = 2y(y')^3$

50. $y'' = (x + y')^2$

48. $x^2 y'' + 3x y' = 2$

52. $y^3 y'' = 1$

- **53.** y'' = 2yy' **54.** $yy'' = 3(y')^2$
- **55.** Show that the substitution v = ax + by + c transforms the differential equation dy/dx = F(ax + by + c) into a separable equation.
- 56. Suppose that $n \neq 0$ and $n \neq 1$. Show that the substitution $v = y^{1-n}$ transforms the Bernoulli equation $dy/dx + P(x)y = Q(x)y^n$ into the linear equation

$$\frac{dv}{dx} + (1-n)P(x)v(x) = (1-n)Q(x).$$

- 57. Show that the substitution $v = \ln y$ transforms the differential equation $dy/dx + P(x)y = Q(x)(y \ln y)$ into the linear equation dv/dx + P(x) = Q(x)v(x).
- 58. Use the idea in Problem 57 to solve the equation

$$x\frac{dy}{dx} - 4x^2y + 2y\ln y = 0.$$

59. Solve the differential equation

$$\frac{dy}{dx} = \frac{x - y - 1}{x + y + 3}$$

by finding *h* and *k* so that the substitutions x = u + h, y = v + k transform it into the homogeneous equation

$$\frac{dv}{du} = \frac{u-v}{u+v}.$$

60. Use the method in Problem 59 to solve the differential equation

$$\frac{dy}{dx} = \frac{2y - x + 7}{4x - 3y - 18}.$$

- **61.** Make an appropriate substitution to find a solution of the equation $dy/dx = \sin(x y)$. Does this general solution contain the linear solution $y(x) = x \pi/2$ that is readily verified by substitution in the differential equation?
- 62. Show that the solution curves of the differential equation

$$\frac{dy}{dx} = -\frac{y(2x^3 - y^3)}{x(2y^3 - x^3)}$$

are of the form $x^3 + y^3 = Cxy$.

63. The equation $dy/dx = A(x)y^2 + B(x)y + C(x)$ is called a **Riccati equation**. Suppose that one particular solution $y_1(x)$ of this equation is known. Show that the substitution

$$y = y_1 + \frac{1}{v}$$

transforms the Riccati equation into the linear equation

$$\frac{dv}{dx} + (B + 2Ay_1)v = -A.$$

Use the method of Problem 63 to solve the equations in Problems 64 and 65, given that $y_1(x) = x$ *is a solution of each.*

1.6 SUBSTITUTION METHODS AND EXACT EQUATIONS71

64.
$$\frac{dy}{dx} + y^2 = 1 + x^2$$

65. $\frac{dy}{dx} + 2xy = 1 + x^2 + y^2$

66. An equation of the form

$$y = xy' + g(y') \tag{37}$$

is called a **Clairaut equation**. Show that the oneparameter family of straight lines described by

$$y(x) = Cx + g(C) \tag{38}$$

is a general solution of Eq. (37).

67. Consider the Clairaut equation

$$y = xy' - \frac{1}{4}(y')^2$$

for which $g(y') = -\frac{1}{4}(y')^2$ in Eq. (37). Show that the line

$$y = Cx - \frac{1}{4}C^2$$

is tangent to the parabola $y = x^2$ at the point $\left(\frac{1}{2}C, \frac{1}{4}C^2\right)$.

Explain why this implies that $y = x^2$ is a singular solution of the given Clairaut equation. This singular solution and the one-parameter family of straight line solutions are illustrated in Fig. 1.6.10.



FIGURE 1.6.10. Solutions of the Clairaut equation of Problem 67. The "typical" straight line with equation $y = Cx - \frac{1}{4}C^2$ is tangent to the parabola at the point $(\frac{1}{2}C, \frac{1}{4}C^2)$.

- 68. Derive Eq. (18) in this section from Eqs. (16) and (17).
- **69.** Flight trajectory In the situation of Example 7, suppose that a = 100 mi, $v_0 = 400$ mi/h, and w = 40 mi/h. Now how far northward does the wind blow the airplane?
- **70. Flight trajectory** As in the text discussion, suppose that an airplane maintains a heading toward an airport at the origin. If $v_0 = 500$ mi/h and w = 50 mi/h (with the wind blowing due north), and the plane begins at the point (200, 150), show that its trajectory is described by

$$y + \sqrt{x^2 + y^2} = 2(200x^9)^{1/10}$$

- 71. River crossing A river 100 ft wide is flowing north at w feet per second. A dog starts at (100, 0) and swims at v₀ = 4 ft/s, always heading toward a tree at (0, 0) on the west bank directly across from the dog's starting point.
 (a) If w = 2 ft/s, show that the dog reaches the tree. (b) If w = 4 ft/s, show that the dog reaches instead the point on the west bank 50 ft north of the tree. (c) If w = 6 ft/s, show that the dog never reaches the west bank.
- 72. In the calculus of plane curves, one learns that the *curva-ture* κ of the curve y = y(x) at the point (x, y) is given by

$$\kappa = \frac{|y''(x)|}{[1 + y'(x)^2]^{3/2}}$$

and that the curvature of a circle of radius r is $\kappa = 1/r$. [See Example 3 in Section 11.6 of Edwards and Penney, *Calculus: Early Transcendentals*, 7th edition, Hoboken, NJ: Pearson, 2008.] Conversely, substitute $\rho = y'$ to derive a general solution of the second-order differential equation

$$ry'' = [1 + (y')^2]^{3/2}$$

(with r constant) in the form

$$(x-a)^2 + (y-b)^2 = r^2$$

Thus a circle of radius r (or a part thereof) is the *only* plane curve with constant curvature 1/r.

1.6 Application Computer Algebra Solutions

Go to goo.gl/tLcVCl to download this application's computing resources including *Maple/Mathematica/*MATLAB. Computer algebra systems typically include commands for the "automatic" solution of differential equations. But two different such systems often give different results whose equivalence is not clear, and a single system may give the solution in an overly complicated form. Consequently, computer algebra solutions of differential equations often require considerable "processing" or simplification by a human user in order to yield concrete and applicable information. Here we illustrate these issues using the interesting differential equation

$$\frac{dy}{dx} = \sin(x - y) \tag{1}$$

that appeared in the Section 1.3 Application. The Maple command

dsolve(D(y)(x) = sin(x - y(x)), y(x));

yields the simple and attractive result

$$y(x) = x - 2\tan^{-1}\left(\frac{x - 2 - C1}{x - C1}\right)$$
(2)

that was cited there. But the supposedly equivalent Mathematica command

$$DSolve[y'[x] == Sin[x - y[x]], y[x], x]$$

and the Wolfram Alpha query

y' = sin(x - y)

both yield considerably more complicated results from which—with a fair amount of effort in simplification—one can extract the quite different looking solution

$$y(x) = 2\cos^{-1}\left(\frac{2\cos\frac{x}{2} + (x-c)\left(\cos\frac{x}{2} + \sin\frac{x}{2}\right)}{\sqrt{2 + 2(x-c+1)^2}}\right).$$
 (3)

This apparent disparity is not unusual; different symbolic algebra systems, or even different versions of the same system, often yield different forms of a solution of the same differential equation. As an alternative to attempted reconciliation of such seemingly disparate results as in Eqs. (2) and (3), a common tactic is simplification of the differential equation before submitting it to a computer algebra system.

EXERCISE 1: Show that the plausible substitution v = x - y in Eq. (1) yields the separable equation

$$\frac{dv}{dx} = 1 - \sin v. \tag{4}$$

Now the *Maple* command int(1/(1 - sin(v)), v) yields

$$\int \frac{dv}{1-\sin v} = \frac{2}{1-\tan\frac{v}{2}} \tag{5}$$

(omitting the constant of integration, as symbolic computer algebra systems often do).

EXERCISE 2: Use simple algebra to deduce from Eq. (5) the integral formula

$$\int \frac{dv}{1-\sin v} = \frac{1+\tan\frac{v}{2}}{1-\tan\frac{v}{2}} + C.$$
 (6)

EXERCISE 3: Deduce from (6) that Eq. (4) has the general solution

$$v(x) = 2 \tan^{-1} \left(\frac{x - 1 + C}{x + 1 + C} \right)$$

and hence that Eq. (1) has the general solution

$$y(x) = x - 2\tan^{-1}\left(\frac{x - 1 + C}{x + 1 + C}\right).$$
(7)

$\operatorname{deSolve}(y' = \sin(x - y), x, y)$	$\frac{-\cos(x-y)}{-\sin(x-y)} = x + c$
	omyc yr 1

FIGURE 1.6.11. Implicit solution of $y' = \sin(x - y)$ generated by a TI-Nspire CX CAS.

EXERCISE 4: Finally, reconcile the forms in Eq. (2) and Eq. (7). What is the relation between the constants *C* and *C*1?

EXERCISE 5: Show that the integral in Eq. (5) yields immediately the graphing calculator implicit solution shown in Fig. 1.6.11.

INVESTIGATION: For your own personal differential equation, let p and q be two distinct nonzero digits in your student ID number, and consider the differential equation

$$\frac{dy}{dx} = \frac{1}{p}\cos(x - qy). \tag{8}$$

- (a) Find a symbolic general solution using a computer algebra system and/or some combination of the techniques listed in this project.
- (b) Determine the symbolic particular solution corresponding to several typical initial conditions of the form $y(x_0) = y_0$.
- (c) Determine the possible values of a and b such that the straight line y = ax + b is a solution curve of Eq. (8).
- (d) Plot a direction field and some typical solution curves. Can you make a connection between the symbolic solution and your (linear and nonlinear) solution curves?

Chapter 1 Summary

In this chapter we have discussed applications of and solution methods for several important types of first-order differential equations, including those that are separable (Section 1.4), linear (Section 1.5), or exact (Section 1.6). In Section 1.6 we also discussed substitution techniques that can sometimes be used to transform a given first-order differential equation into one that is either separable, linear, or exact.

Lest it appear that these methods constitute a "grab bag" of special and unrelated techniques, it is important to note that they are all versions of a single idea. Given a differential equation

$$f(x, y, y') = 0,$$
 (1)

we attempt to write it in the form

$$\frac{d}{dx}\left[G(x,y)\right] = 0.$$
(2)

It is precisely to obtain the form in Eq. (2) that we multiply the terms in Eq. (1) by an appropriate integrating factor (even if all we are doing is separating the variables). But once we have found a function G(x, y) such that Eqs. (1) and (2) are equivalent, a general solution is defined implicitly by means of the equation

$$G(x, y) = C \tag{3}$$

that one obtains by integrating Eq. (2).

Given a specific first-order differential equation to be solved, we can attack it by means of the following steps:

- Is it *separable*? If so, separate the variables and integrate (Section 1.4).
- Is it *linear*? That is, can it be written in the form

$$\frac{dy}{dx} + P(x)y = Q(x)?$$

If so, multiply by the integrating factor $\rho = \exp(\int P \, dx)$ of Section 1.5.

- Is it *exact*? That is, when the equation is written in the form M dx + N dy = 0, is $\partial M / \partial y = \partial N / \partial x$ (Section 1.6)?
- If the equation as it stands is not separable, linear, or exact, is there a plausible substitution that will make it so? For instance, is it homogeneous (Section 1.6)?

Many first-order differential equations succumb to the line of attack outlined here. Nevertheless, many more do not. Because of the wide availability of computers, numerical techniques are commonly used to *approximate* the solutions of differential equations that cannot be solved readily or explicitly by the methods of this chapter. Indeed, most of the solution curves shown in figures in this chapter were plotted using numerical approximations rather than exact solutions. Several numerical methods for the appropriate solution of differential equations will be discussed in Chapter 2.

Chapter 1 Review Problems

Find general solutions of the differential equations in Problems 1 through 30. Primes denote derivatives with respect to x.

- **2.** $xv^2 + 3v^2 x^2v' = 0$ 1. $x^3 + 3y - xy' = 0$ 3. $xy + y^2 - x^2y' = 0$ 4. $2xy^3 + e^x + (3x^2y^2 + \sin y)y' = 0$ 6. $2xy^2 + x^2y' = y^2$ 5. $3y + x^4y' = 2xy$ 8. $2xy + x^2y' = y^2$ 10. $y' = 1 + x^2 + y^2 + x^2y^2$ 7. $2x^2y + x^3y' = 1$ 9. $xy' + 2y = 6x^2\sqrt{y}$ 11. $x^2 y' = xy + 3y^2$ 12. $6xy^3 + 2y^4 + (9x^2y^2 + 8xy^3)y' = 0$ **14.** $x^{3}y' = x^{2}y - y^{3}$ **16.** $y' = x^{2} - 2xy + y^{2}$ 13. $4xy^2 + y' = 5x^4y^2$ 15. $y' + 3y = 3x^2e^{-3x}$ 17. $e^{x} + ye^{xy} + (e^{y} + xe^{yx})y' = 0$ **18.** $2x^2y - x^3y' = y^3$ **19.** $3x^5y^2 + x^3y' = 2y^2$ **20.** $xy' + 3y = 3x^{-3/2}$ **21.** $(x^2 - 1)y' + (x - 1)y = 1$ **22.** $xy' = 6y + 12x^4y^{2/3}$ **23.** $e^{y} + y \cos x + (xe^{y} + \sin x)y' = 0$
- **24.** $9x^2y^2 + x^{3/2}y' = y^2$ **25.** 2y + (x+1)y' = 3x + 3 **26.** $9x^{1/2}y^{4/3} - 12x^{1/5}y^{3/2} + (8x^{3/2}y^{1/3} - 15x^{6/5}y^{1/2})y' = 0$ **27.** $3y + x^3y^4 + 3xy' = 0$ **28.** $y + xy' = 2e^{2x}$ **29.** $(2x+1)y' + y = (2x+1)^{3/2}$ **30.** $y' = \sqrt{x+y}$

Each of the differential equations in Problems 31 through 36 is of two different types considered in this chapter—separable, linear, homogeneous, Bernoulli, exact, etc. Hence, derive general solutions for each of these equations in two different ways; then reconcile your results.

31.
$$\frac{dy}{dx} = 3(y+7)x^2$$

32. $\frac{dy}{dx} = xy^3 - xy$
33. $\frac{dy}{dx} = -\frac{3x^2 + 2y^2}{4xy}$
34. $\frac{dy}{dx} = \frac{x+3y}{y-3x}$
35. $\frac{dy}{dx} = \frac{2xy + 2x}{x^2 + 1}$
36. $\frac{dy}{dx} = \frac{\sqrt{y} - y}{\tan x}$

Mathematical Models and Numerical Methods

2.1 Population Models

In Section 1.4 we introduced the exponential differential equation dP/dt = kP, with solution $P(t) = P_0 e^{kt}$, as a mathematical model for natural population growth that occurs as a result of constant birth and death rates. Here we present a more general population model that accommodates birth and death rates that are not necessarily constant. As before, however, our population function P(t) will be a *continuous* approximation to the actual population, which of course changes only by integral increments—that is, by one birth or death at a time.

Suppose that the population changes only by the occurrence of births and deaths—there is no immigration or emigration from outside the country or environment under consideration. It is customary to track the growth or decline of a population in terms of its *birth rate* and *death rate* functions defined as follows:

- $\beta(t)$ is the number of births per unit of population per unit of time at time t;
- $\delta(t)$ is the number of deaths per unit of population per unit of time at time t.

Then the numbers of births and deaths that occur during the time interval $[t, t + \Delta t]$ is given (approximately) by

births: $\beta(t) \cdot P(t) \cdot \Delta t$, deaths: $\delta(t) \cdot P(t) \cdot \Delta t$.

Hence the change ΔP in the population during the time interval $[t, t + \Delta t]$ of length Δt is

$$\Delta P = \{\text{births}\} - \{\text{deaths}\} \approx \beta(t) \cdot P(t) \cdot \Delta t - \delta(t) \cdot P(t) \cdot \Delta t$$

so

$$\frac{\Delta P}{\Delta t} \approx \left[\beta(t) - \delta(t)\right] P(t).$$

The error in this approximation should approach zero as $\Delta t \rightarrow 0$, so—taking the limit—we get the differential equation

$$\blacktriangleright \qquad \qquad \frac{dP}{dt} = (\beta - \delta)P, \tag{1}$$

in which we write $\beta = \beta(t)$, $\delta = \delta(t)$, and P = P(t) for brevity. Equation (1) is the **general population equation**. If β and δ are constant, Eq. (1) reduces to the natural growth equation with $k = \beta - \delta$. But it also includes the possibility that β and δ are variable functions of *t*. The birth and death rates need not be known in advance; they may well depend on the unknown function P(t).

Population explosion Suppose that an alligator population numbers 100 initially, and that its death rate is $\delta = 0$ (so none of the alligators is dying). If the birth rate is $\beta = (0.0005)P$ —and thus increases as the population does—then Eq. (1) gives the initial value problem

$$\frac{dP}{dt} = (0.0005)P^2, \quad P(0) = 100$$

(with *t* in years). Then upon separating the variables we get

$$\int \frac{1}{P^2} dP = \int (0.0005) dt;$$
$$-\frac{1}{P} = (0.0005)t + C.$$

Substitution of t = 0, P = 100 gives C = -1/100, and then we readily solve for

$$P(t) = \frac{2000}{20 - t}.$$

For instance, P(10) = 2000/10 = 200, so after 10 years the alligator population has doubled. But we see that $P \to +\infty$ as $t \to 20$, so a real "population explosion" occurs in 20 years. Indeed, the direction field and solution curves shown in Fig. 2.1.1 indicate that a population explosion always occurs, whatever the size of the (positive) initial population $P(0) = P_0$. In particular, it appears that the population always becomes unbounded in a *finite* period of time.

Bounded Populations and the Logistic Equation

In situations as diverse as the human population of a nation and a fruit fly population in a closed container, it is often observed that the birth rate decreases as the population itself increases. The reasons may range from increased scientific or cultural sophistication to a limited food supply. Suppose, for example, that the birth rate β is a *linear* decreasing function of the population size *P*, so that $\beta = \beta_0 - \beta_1 P$, where β_0 and β_1 are positive constants. If the death rate $\delta = \delta_0$ remains constant, then Eq. (1) takes the form

$$\frac{dP}{dt} = (\beta_0 - \beta_1 P - \delta_0) P;$$

that is,

$$\frac{dP}{dt} = aP - bP^2,\tag{2}$$

where $a = \beta_0 - \delta_0$ and $b = \beta_1$.

If the coefficients *a* and *b* are both positive, then Eq. (2) is called the **logistic** equation. For the purpose of relating the behavior of the population P(t) to the values of the parameters in the equation, it is useful to rewrite the logistic equation in the form

$$\frac{dP}{dt} = kP(M - P),\tag{3}$$

where k = b and M = a/b are constants.



FIGURE 2.1.1. Slope field and solution curves for the equation $dP/dt = (0.0005)P^2$ in Example 1.

Example 2 Logistic model In Example 4 of Section 1.3 we explored graphically a population that is modeled by the logistic equation

$$\frac{dP}{dt} = 0.0004P(150 - P) = 0.06P - 0.0004P^2.$$
(4)

To solve this differential equation symbolically, we separate the variables and integrate. We get

$$\int \frac{dP}{P(150 - P)} = \int 0.0004 \, dt,$$

$$\frac{1}{150} \int \left(\frac{1}{P} + \frac{1}{150 - P}\right) dP = \int 0.0004 \, dt \qquad \text{[partial fractions]},$$

$$\ln|P| - \ln|150 - P| = 0.06t + C,$$

$$\frac{P}{150 - P} = \pm e^C e^{0.06t} = Be^{0.06t} \quad \text{[where } B = \pm e^C \text{]}.$$

If we substitute t = 0 and $P = P_0 \neq 150$ into this last equation, we find that $B = P_0/(150 - P_0)$. Hence

$$\frac{P}{150 - P} = \frac{P_0 e^{0.06t}}{150 - P_0}$$

Finally, this equation is easy to solve for the population

$$P(t) = \frac{150P_0}{P_0 + (150 - P_0)e^{-0.06t}}$$
(5)

at time *t* in terms of the initial population $P_0 = P(0)$. Figure 2.1.2 shows a number of solution curves corresponding to different values of the initial population ranging from $P_0 = 20$ to $P_0 = 300$. Note that all these solution curves appear to approach the horizontal line P = 150 as an asymptote. Indeed, you should be able to see directly from Eq. (5) that $\lim_{t\to\infty} P(t) = 150$, whatever the initial value $P_0 > 0$.

Limiting Populations and Carrying Capacity

The finite limiting population noted in Example 2 is characteristic of logistic populations. In Problem 32 we ask you to use the method of solution of Example 2 to show that the solution of the logistic initial value problem

$$\frac{dP}{dt} = kP(M-P), \quad P(0) = P_0 \tag{6}$$

is

$$P(t) = \frac{MP_0}{P_0 + (M - P_0)e^{-kMt}}.$$
(7)

Actual animal populations are positive valued. If $P_0 = M$, then (7) reduces to the unchanging (constant-valued) "equilibrium population" $P(t) \equiv M$. Otherwise, the behavior of a logistic population depends on whether $0 < P_0 < M$ or $P_0 > M$. If $0 < P_0 < M$, then we see from (6) and (7) that P' > 0 and

$$P(t) = \frac{MP_0}{P_0 + (M - P_0)e^{-kMt}} = \frac{MP_0}{P_0 + \{\text{pos. number}\}} < \frac{MP_0}{P_0} = M.$$

However, if $P_0 > M$, then we see from (6) and (7) that P' < 0 and

$$P(t) = \frac{MP_0}{P_0 + (M - P_0)e^{-kMt}} = \frac{MP_0}{P_0 + \{\text{neg. number}\}} > \frac{MP_0}{P_0} = M.$$



FIGURE 2.1.2. Typical solution curves for the logistic equation $P' = 0.06P - 0.0004P^2$.



FIGURE 2.1.3. Typical solution curves for the logistic equation P' = kP(M - P). Each solution curve that starts below the line P = M/2 has an inflection point on this line. (See Problem 34.)

In either case, the "positive number" or "negative number" in the denominator has absolute value less than P_0 and—because of the exponential factor—approaches 0 as $t \to +\infty$. It follows that

$$\lim_{t \to +\infty} P(t) = \frac{MP_0}{P_0 + 0} = M.$$
(8)

Thus a population that satisfies the logistic equation does *not* grow without bound like a naturally growing population modeled by the exponential equation P' = kP. Instead, it approaches the finite **limiting population** M as $t \to +\infty$. As illustrated by the typical logistic solution curves in Fig. 2.1.3, the population P(t) steadily increases and approaches M from below if $0 < P_0 < M$, but steadily decreases and approaches M from above if $P_0 > M$. Sometimes M is called the **carrying capacity** of the environment, considering it to be the maximum population that the environment can support on a long-term basis.

Example 3 Limiting population Suppose that in 1885 the population of a certain country was 50 million and was growing at the rate of 750,000 people per year at that time. Suppose also that in 1940 its population was 100 million and was then growing at the rate of 1 million per year. Assume that this population satisfies the logistic equation. Determine both the limiting population *M* and the predicted population for the year 2000.

Solution

We substitute the two given pairs of data in Eq. (3) and find that

$$0.75 = 50k(M - 50), \quad 1.00 = 100k(M - 100).$$

We solve simultaneously for M = 200 and k = 0.0001. Thus the limiting population of the country in question is 200 million. With these values of M and k, and with t = 0 corresponding to the year 1940 (in which $P_0 = 100$), we find that—according to Eq. (7)—the population in the year 2000 will be

$$P(60) = \frac{100 \cdot 200}{100 + (200 - 100)e^{-(0.0001)(200)(60)}}$$

about 153.7 million people.

Historical Note

The logistic equation was introduced (around 1840) by the Belgian mathematician and demographer P. F. Verhulst as a possible model for human population growth. In the next two examples we compare natural growth and logistic model fits to the 19th-century U.S. population census data, then compare projections for the 20th century.

Example 4 Percentage growth rate The U.S. population in 1800 was 5.308 million and in 1900 was 76.212 million. If we take $P_0 = 5.308$ (with t = 0 in 1800) in the natural growth model $P(t) = P_0 e^{rt}$ and substitute t = 100, P = 76.212, we find that

76.212 = 5.308
$$e^{100r}$$
, so $r = \frac{1}{100} \ln \frac{76.212}{5.308} \approx 0.026643$.

Thus our natural growth model for the U.S. population during the 19th century is

$$P(t) = (5.308)e^{(0.026643)t}$$
(9)

(with *t* in years and *P* in millions). Because $e^{0.026643} \approx 1.02700$, the average population growth between 1800 and 1900 was about 2.7% per year.

Example 5 Logistic modeling The U.S. population in 1850 was 23.192 million. If we take $P_0 = 5.308$ and substitute the data pairs t = 50, P = 23.192 (for 1850) and t = 100, P = 76.212 (for 1900) in the logistic model formula in Eq. (7), we get the two equations

$$\frac{(5.308)M}{5.308 + (M - 5.308)e^{-50kM}} = 23.192,$$

$$\frac{(5.308)M}{5.308 + (M - 5.308)e^{-100kM}} = 76.212$$
(10)

in the two unknowns k and M. Nonlinear systems like this ordinarily are solved numerically using an appropriate computer system. But with the right algebraic trick (Problem 36 in this section) the equations in (10) can be solved manually for k = 0.000167716, M = 188.121. Substitution of these values in Eq. (7) yields the logistic model

$$P(t) = \frac{998.546}{5.308 + (182.813)e^{-(0.031551)t}}.$$
(11)

The table in Fig. 2.1.4 compares the actual 1800–2010 U.S. census population figures with those predicted by the exponential growth model in (9) and by the logistic model in (11). Both models agree well with the 19th-century figures. But the exponential model diverges appreciably from the census data in the early decades of the 20th century, whereas the logistic model remains accurate until 1940. By the end of the 20th century the exponential model vastly overestimates the actual U.S. population; indeed it predicts a U.S. population of nearly

Year	Actual U.S. Pop.	Exponential Model	Exponential Error	Logistic Model	Logistic Error
1800	5.308	5.308	0.000	5.308	0.000
1810	7.240	6.929	0.311	7.202	0.038
1820	9.638	9.044	0.594	9.735	-0.097
1830	12.861	11.805	1.056	13.095	-0.234
1840	17.064	15.409	1.655	17.501	-0.437
1850	23.192	20.113	3.079	23.192	0.000
1860	31.443	26.253	5.190	30.405	1.038
1870	38.558	34.268	4.290	39.326	-0.768
1880	50.189	44.730	5.459	50.034	0.155
1890	62.980	58.387	4.593	62.435	0.545
1900	76.212	76.212	0.000	76.213	-0.001
1910	92.228	99.479	-7.251	90.834	1.394
1920	106.022	129.849	-23.827	105.612	0.410
1930	123.203	169.492	-46.289	119.834	3.369
1940	132.165	221.237	-89.072	132.886	-0.721
1950	151.326	288.780	-137.454	144.354	6.972
1960	179.323	376.943	-197.620	154.052	25.271
1970	203.302	492.023	-288.721	161.990	41.312
1980	226.542	642.236	-415.694	168.316	58.226
1990	248.710	838.308	-589.598	173.252	75.458
2000	281.422	1094.240	-812.818	177.038	104.384
2010	308.745	1428.307	-1119.562	179.905	128.839

FIGURE 2.1.4. Comparison of exponential growth and logistic models with U.S. census populations (in millions).



FIGURE 2.1.5. Percentage errors in the exponential and logistic population models for 1800–1950.

1.5 billion in the year 2010, over 3.6 times the actual value. The logistic model, on the other hand, underestimates the U.S. population, but with a percentage error of less than 43%.

The two models are compared in Fig. 2.1.5, where plots of their respective errors—as a *percentage* of the actual population—are shown for the 1800–1950 period. We see that the logistic model tracks the actual population reasonably well throughout this 150-year period. However, the exponential error is considerably larger during the 19th century and literally goes off the chart during the first half of the 20th century.

In order to measure the extent to which a given model fits actual data, it is customary to define the **average error** (in the model) as *the square root of the average of the squares of the individual errors* (the latter appearing in the fourth and sixth columns of the table in Fig. 2.1.4). Using only the 1800–1900 data, this definition gives 3.162 for the average error in the exponential model, while the average error in the logistic model is only 0.452. Consequently, even in 1900 we might well have anticipated that the logistic model would predict the U.S. population growth during the 20th century more accurately than the exponential model.

The moral of Examples 4 and 5 is simply that one should not expect too much of models that are based on severely limited information (such as just a pair of data points). Much of the science of *statistics* is devoted to the analysis of large "data sets" to formulate useful (and perhaps reliable) mathematical models.

More Applications of the Logistic Equation

We next describe some situations that illustrate the varied circumstances in which the logistic equation is a satisfactory mathematical model.

1. *Limited environment situation*. A certain environment can support a population of at most M individuals. It is then reasonable to expect the growth rate $\beta - \delta$ (the combined birth and death rates) to be proportional to M - P, because we may think of M - P as the potential for further expansion. Then $\beta - \delta = k(M - P)$, so that

$$\frac{dP}{dt} = (\beta - \delta)P = kP(M - P).$$

The classic example of a limited environment situation is a fruit fly population in a closed container.

2. *Competition situation.* If the birth rate β is constant but the death rate δ is proportional to *P*, so that $\delta = \alpha P$, then

$$\frac{dP}{dt} = (\beta - \alpha P)P = kP(M - P).$$

This might be a reasonable working hypothesis in a study of a cannibalistic population, in which all deaths result from chance encounters between individuals. Of course, competition between individuals is not usually so deadly, nor its effects so immediate and decisive.

3. Joint proportion situation. Let P(t) denote the number of individuals in a constant-size susceptible population M who are infected with a certain contagious and incurable disease. The disease is spread by chance encounters. Then P'(t) should be proportional to the product of the number P of individuals having the disease and the number M - P of those not having it, and therefore dP/dt = kP(M - P). Again we discover that the mathematical model is the logistic equation. The mathematical description of the spread of a rumor in a population of M individuals is identical.

Example 6 Spread of rumor Suppose that at time t = 0, 10 thousand people in a city with population M = 100 thousand people have heard a certain rumor. After 1 week the number P(t) of those who have heard it has increased to P(1) = 20 thousand. Assuming that P(t) satisfies a logistic equation, when will 80% of the city's population have heard the rumor?

Solution Substituting $P_0 = 10$ and M = 100 (thousand) in Eq. (7), we get

$$P(t) = \frac{1000}{10 + 90e^{-100kt}}.$$
(12)

Then substitution of t = 1, P = 20 gives the equation

$$20 = \frac{1000}{10 + 90e^{-100k}}$$

that is readily solved for

$$e^{-100k} = \frac{4}{9}$$
, so $k = \frac{1}{100} \ln \frac{9}{4} \approx 0.008109$.

With P(t) = 80, Eq. (12) takes the form

$$80 = \frac{1000}{10 + 90e^{-100kt}},$$

which we solve for $e^{-100kt} = \frac{1}{36}$. It follows that 80% of the population has heard the rumor when

$$t = \frac{\ln 36}{100k} = \frac{\ln 36}{\ln \frac{9}{4}} \approx 4.42$$

thus after about 4 weeks and 3 days.

Doomsday versus Extinction

Consider a population P(t) of unsophisticated animals in which females rely solely on chance encounters to meet males for reproductive purposes. It is reasonable to expect such encounters to occur at a rate that is proportional to the product of the number P/2 of males and the number P/2 of females, hence at a rate proportional to P^2 . We therefore assume that births occur at the rate kP^2 (per unit time, with k constant). The birth rate (births/time/population) is then given by $\beta = kP$. If the death rate δ is constant, then the general population equation in (1) yields the differential equation

$$\frac{dP}{dt} = kP^2 - \delta P = kP(P - M) \tag{13}$$

(where $M = \delta/k > 0$) as a mathematical model of the population.

Note that the right-hand side in Eq. (13) is the *negative* of the right-hand side in the logistic equation in (3). We will see that the constant M is now a **threshold population**, with the way the population behaves in the future depending critically on whether the initial population P_0 is less than or greater than M.

Example 7 Doomsday vs. extinction Consider an animal population P(t) that is modeled by the equation

$$\frac{dP}{dt} = 0.0004P(P-150) = 0.0004P^2 - 0.06P.$$
(14)

We want to find P(t) if (a) P(0) = 200; (b) P(0) = 100.

Solution To solve the equation in (14), we separate the variables and integrate. We get

r

$$\int \frac{dP}{P(P-150)} = \int 0.0004 \, dt,$$

$$-\frac{1}{150} \int \left(\frac{1}{P} - \frac{1}{P-150}\right) dP = \int 0.0004 \, dt \quad \text{[partial fractions]},$$

$$\ln|P| - \ln|P - 150| = -0.06t + C,$$

$$\frac{P}{P-150} = \pm e^C e^{-0.06t} = B e^{-0.06t} \quad \text{[where } B = \pm e^C \text{]}. \tag{15}$$

C

(a) Substitution of t = 0 and P = 200 into (15) gives B = 4. With this value of B we solve Eq. (15) for

$$P(t) = \frac{600e^{-0.06t}}{4e^{-0.06t} - 1}.$$
(16)

Note that, as t increases and approaches $T = \ln(4)/0.06 \approx 23.105$, the positive denominator on the right in (16) decreases and approaches 0. Consequently $P(t) \rightarrow +\infty$ as $t \rightarrow T^-$. This is a *doomsday* situation—a real population *explosion*.

(b) Substitution of t = 0 and P = 100 into (15) gives B = -2. With this value of B we solve Eq. (15) for

$$P(t) = \frac{300e^{-0.06t}}{2e^{-0.06t} + 1} = \frac{300}{2 + e^{0.06t}}.$$
(17)

Note that, as t increases without bound, the positive denominator on the right in (16) approaches $+\infty$. Consequently, $P(t) \to 0$ as $t \to +\infty$. This is an (eventual) extinction situation.

Thus the population in Example 7 either explodes or is an endangered species threatened with extinction, depending on whether or not its initial size exceeds the threshold population M = 150. An approximation to this phenomenon is sometimes observed with animal populations, such as the alligator population in certain areas of the southern United States.

Figure 2.1.6 shows typical solution curves that illustrate the two possibilities for a population P(t) satisfying Eq. (13). If $P_0 = M$ (exactly!), then the population remains constant. However, this equilibrium situation is very unstable. If P_0 exceeds M (even slightly), then P(t) rapidly increases without bound, whereas if the initial (positive) population is less than M (however slightly), then it decreases (more gradually) toward zero as $t \to +\infty$. See Problem 33.



FIGURE 2.1.6. Typical solution curves for the explosion/extinction equation P' = kP(P - M).

Problems 2.1

Separate variables and use partial fractions to solve the initial value problems in Problems 1–8. Use either the exact solution or a computer-generated slope field to sketch the graphs of several solutions of the given differential equation, and highlight the indicated particular solution.

1.
$$\frac{dx}{dt} = x - x^2, x(0) = 2$$
 2. $\frac{dx}{dt} = 10x - x^2, x(0) = 1$

3.
$$\frac{dx}{dt} = 1 - x^2, x(0) = 3$$
 4. $\frac{dx}{dt} = 9 - 4x^2, x(0) = 0$

5.
$$\frac{dx}{dt} = 3x(5-x), x(0) = 8$$

6. $\frac{dx}{dt} = 3x(x-5), x(0) = 2$

7. $\frac{dx}{dt} = 4x(7-x), x(0) = 11$

8.
$$\frac{dx}{dt} = 7x(x-13), x(0) = 17$$

- 9. Population growth The time rate of change of a rabbit population P is proportional to the square root of P. At time t = 0 (months) the population numbers 100 rabbits and is increasing at the rate of 20 rabbits per month. How many rabbits will there be one year later?
- **10. Extinction by disease** Suppose that the fish population P(t) in a lake is attacked by a disease at time t = 0, with the result that the fish cease to reproduce (so that the birth rate is $\beta = 0$ and the death rate δ (deaths per week per

fish) is thereafter proportional to $1/\sqrt{P}$. If there were initially 900 fish in the lake and 441 were left after 6 weeks, how long did it take all the fish in the lake to die?

11. Fish population Suppose that when a certain lake is stocked with fish, the birth and death rates β and δ are both inversely proportional to \sqrt{P} . (a) Show that

$$P(t) = \left(\frac{1}{2}kt + \sqrt{P_0}\right)^2,$$

where k is a constant. (b) If $P_0 = 100$ and after 6 months there are 169 fish in the lake, how many will there be after 1 year?

- **12. Population growth** The time rate of change of an alligator population *P* in a swamp is proportional to the square of *P*. The swamp contained a dozen alligators in 1988, two dozen in 1998. When will there be four dozen alligators in the swamp? What happens thereafter?
- 13. Birth rate exceeds death rate Consider a prolific breed of rabbits whose birth and death rates, β and δ, are each proportional to the rabbit population P = P(t), with β > δ.
 (a) Show that

$$P(t) = \frac{P_0}{1 - kP_0 t}, \quad k \text{ constant.}$$

Note that $P(t) \rightarrow +\infty$ as $t \rightarrow 1/(kP_0)$. This is doomsday. (b) Suppose that $P_0 = 6$ and that there are nine rabbits after ten months. When does doomsday occur?

- 14. Death rate exceeds birth rate Repeat part (a) of Problem 13 in the case $\beta < \delta$. What now happens to the rabbit population in the long run?
- 15. Limiting population Consider a population P(t) satisfying the logistic equation $dP/dt = aP bP^2$, where B = aP is the time rate at which births occur and $D = bP^2$ is the rate at which deaths occur. If the initial population is $P(0) = P_0$, and B_0 births per month and D_0 deaths per month are occurring at time t = 0, show that the limiting population is $M = B_0 P_0/D_0$.
- 16. Limiting population Consider a rabbit population P(t) satisfying the logistic equation as in Problem 15. If the initial population is 120 rabbits and there are 8 births per month and 6 deaths per month occurring at time t = 0, how many months does it take for P(t) to reach 95% of the limiting population M?
- 17. Limiting population Consider a rabbit population P(t) satisfying the logistic equation as in Problem 15. If the initial population is 240 rabbits and there are 9 births per month and 12 deaths per month occurring at time t = 0, how many months does it take for P(t) to reach 105% of the limiting population M?
- **18.** Threshold population Consider a population P(t) satisfying the extinction-explosion equation $dP/dt = aP^2 bP$, where $B = aP^2$ is the time rate at which births occur and D = bP is the rate at which deaths occur. If the initial population is $P(0) = P_0$ and B_0 births per month and D_0

deaths per month are occurring at time t = 0, show that the threshold population is $M = D_0 P_0/B_0$.

- **19.** Threshold population Consider an alligator population P(t) satisfying the extinction-explosion equation as in Problem 18. If the initial population is 100 alligators and there are 10 births per month and 9 deaths per month occurring at time t = 0, how many months does it take for P(t) to reach 10 times the threshold population M?
- **20.** Threshold population Consider an alligator population P(t) satisfying the extinction-explosion equation as in Problem 18. If the initial population is 110 alligators and there are 11 births per month and 12 deaths per month occurring at time t = 0, how many months does it take for P(t) to reach 10% of the threshold population M?
- **21.** Logistic model Suppose that the population P(t) of a country satisfies the differential equation dP/dt = kP(200 P) with *k* constant. Its population in 1960 was 100 million and was then growing at the rate of 1 million per year. Predict this country's population for the year 2020.
- **22.** Logistic model Suppose that at time t = 0, half of a "logistic" population of 100,000 persons have heard a certain rumor, and that the number of those who have heard it is then increasing at the rate of 1000 persons per day. How long will it take for this rumor to spread to 80% of the population? (*Suggestion:* Find the value of *k* by substituting P(0) and P'(0) in the logistic equation, Eq. (3).)
- **23.** Solution rate As the salt KNO₃ dissolves in methanol, the number x(t) of grams of the salt in a solution after t seconds satisfies the differential equation $dx/dt = 0.8x 0.004x^2$.
 - (a) What is the maximum amount of the salt that will ever dissolve in the methanol?
 - (b) If x = 50 when t = 0, how long will it take for an additional 50 g of salt to dissolve?
- 24. Spread of disease Suppose that a community contains 15,000 people who are susceptible to Michaud's syndrome, a contagious disease. At time t = 0 the number N(t) of people who have developed Michaud's syndrome is 5000 and is increasing at the rate of 500 per day. Assume that N'(t) is proportional to the product of the numbers of those who have caught the disease and of those who have not. How long will it take for another 5000 people to develop Michaud's syndrome?
- 25. Logistic model The data in the table in Fig. 2.1.7 are given for a certain population P(t) that satisfies the logistic equation in (3). (a) What is the limiting population M? (Suggestion: Use the approximation

$$P'(t) \approx \frac{P(t+h) - P(t-h)}{2h}$$

with h = 1 to estimate the values of P'(t) when P = 25.00and when P = 47.54. Then substitute these values in the logistic equation and solve for k and M.) (b) Use the

values of k and M found in part (a) to determine when P = 75. (Suggestion: Take t = 0 to correspond to the year 1965.)

Year	P (millions)
1964	24.63
1965	25.00
1966	25.38
:	:
2014	47.04
2015	47.54
2016	48.04

FIGURE 2.1.7. Population data for Problem 25.

- **26.** Constant death rate A population P(t) of small rodents has birth rate $\beta = (0.001)P$ (births per month per rodent) and *constant* death rate δ . If P(0) = 100 and P'(0) = 8, how long (in months) will it take this population to double to 200 rodents? (*Suggestion:* First find the value of δ .)
- **27.** Constant death rate Consider an animal population P(t) with constant death rate $\delta = 0.01$ (deaths per animal per month) and with birth rate β proportional to P. Suppose that P(0) = 200 and P'(0) = 2. (a) When is P = 1000? (b) When does doomsday occur?
- **28.** Population growth Suppose that the number x(t) (with *t* in months) of alligators in a swamp satisfies the differential equation $dx/dt = 0.0001x^2 0.01x$.
 - (a) If initially there are 25 alligators in the swamp, solve this differential equation to determine what happens to the alligator population in the long run.
 - (b) Repeat part (a), except with 150 alligators initially.
- **29.** Logistic model During the period from 1790 to 1930, the U.S. population P(t) (*t* in years) grew from 3.9 million to 123.2 million. Throughout this period, P(t) remained close to the solution of the initial value problem

$$\frac{dP}{dt} = 0.03135P - 0.0001489P^2, \quad P(0) = 3.9.$$

- (a) What 1930 population does this logistic equation predict?
- (b) What limiting population does it predict?
- (c) Has this logistic equation continued since 1930 to accurately model the U.S. population?

[This problem is based on a computation by Verhulst, who in 1845 used the 1790–1840 U.S. population data to predict accurately the U.S. population through the year 1930 (long after his own death, of course).]

30. Tumor growth A tumor may be regarded as a population of multiplying cells. It is found empirically that the "birth rate" of the cells in a tumor decreases exponentially

with time, so that $\beta(t) = \beta_0 e^{-\alpha t}$ (where α and β_0 are positive constants), and hence

$$\frac{dP}{dt} = \beta_0 e^{-\alpha t} P, \quad P(0) = P_0.$$

Solve this initial value problem for

$$P(t) = P_0 \exp\left(\frac{\beta_0}{\alpha}(1 - e^{-\alpha t})\right).$$

Observe that P(t) approaches the finite limiting population $P_0 \exp(\beta_0/\alpha)$ as $t \to +\infty$.

- **31.** Tumor growth For the tumor of Problem 30, suppose that at time t = 0 there are $P_0 = 10^6$ cells and that P(t) is then increasing at the rate of 3×10^5 cells per month. After 6 months the tumor has doubled (in size and in number of cells). Solve numerically for α , and then find the limiting population of the tumor.
- **32.** Derive the solution

$$P(t) = \frac{MP_0}{P_0 + (M - P_0)e^{-kMt}}$$

of the logistic initial value problem P' = kP(M - P), $P(0) = P_0$. Make it clear how your derivation depends on whether $0 < P_0 < M$ or $P_0 > M$.

33. (a) Derive the solution

$$P(t) = \frac{MP_0}{P_0 + (M - P_0)e^{kMt}}$$

of the extinction-explosion initial value problem P' = kP(P - M), $P(0) = P_0$.

- (b) How does the behavior of P(t) as t increases depend on whether $0 < P_0 < M$ or $P_0 > M$?
- **34.** If P(t) satisfies the logistic equation in (3), use the chain rule to show that

$$P''(t) = 2k^2 P(P - \frac{1}{2}M)(P - M)$$

Conclude that P'' > 0 if $0 < P < \frac{1}{2}M$; P'' = 0 if $P = \frac{1}{2}M$; P'' < 0 if $\frac{1}{2}M < P < M$; and P'' > 0 if P > M. In particular, it follows that any solution curve that crosses the line $P = \frac{1}{2}M$ has an inflection point where it crosses that line, and therefore resembles one of the lower S-shaped curves in Fig. 2.1.3.

35. Approach to limiting population Consider two population functions $P_1(t)$ and $P_2(t)$, both of which satisfy the logistic equation with the same limiting population M but with different values k_1 and k_2 of the constant k in Eq. (3). Assume that $k_1 < k_2$. Which population approaches M the most rapidly? You can reason *geometrically* by examining slope fields (especially if appropriate software is available), *symbolically* by analyzing the solution given in Eq. (7), or *numerically* by substituting successive values of t.

- **36.** Logistic modeling To solve the two equations in (10) for the values of k and M, begin by solving the first equation for the quantity $x = e^{-50kM}$ and the second equation for $x^2 = e^{-100kM}$. Upon equating the two resulting expressions for x^2 in terms of M, you get an equation that is readily solved for M. With M now known, either of the original equations is readily solved for k. This technique can be used to "fit" the logistic equation to any three population values P_0 , P_1 , and P_2 corresponding to equally spaced times $t_0 = 0$, t_1 , and $t_2 = 2t_1$.
- **37.** Logistic modeling Use the method of Problem 36 to fit the logistic equation to the actual U.S. population data (Fig. 2.1.4) for the years 1850, 1900, and 1950. Solve the resulting logistic equation and compare the predicted and actual populations for the years 1990 and 2000.
- **38.** Logistic modeling Fit the logistic equation to the actual U.S. population data (Fig. 2.1.4) for the years 1900,

1930, and 1960. Solve the resulting logistic equation, then compare the predicted and actual populations for the years 1980, 1990, and 2000.

39. Periodic growth rate Birth and death rates of animal populations typically are not constant; instead, they vary periodically with the passage of seasons. Find P(t) if the population P satisfies the differential equation

$$\frac{dP}{dt} = (k + b\cos 2\pi t)P$$

where t is in years and k and b are positive constants. Thus the growth-rate function $r(t) = k + b \cos 2\pi t$ varies periodically about its mean value k. Construct a graph that contrasts the growth of this population with one that has the same initial value P_0 but satisfies the natural growth equation P' = kP (same constant k). How would the two populations compare after the passage of many years?

2.1 Application Logistic Modeling of Population Data

Go to goo.gl/s3nZZ3 to download this application's computing resources including *Maple/Mathematica/*MATLAB. These investigations deal with the problem of fitting a logistic model to given population data. Thus we want to determine the numerical constants a and b so that the solution P(t) of the initial value problem

$$\frac{dP}{dt} = aP + bP^2, \quad P(0) = P_0 \tag{1}$$

approximates the given values P_0, P_1, \ldots, P_n of the population at the times $t_0 = 0$, t_1, \ldots, t_n . If we rewrite Eq. (1) (the logistic equation with kM = a and k = -b) in the form

$$\frac{1}{P}\frac{dP}{dt} = a + bP,\tag{2}$$

then we see that the points

$$\left(P(t_i), \frac{P'(t_i)}{P(t_i)}\right), \quad i = 0, 1, 2, \dots, n,$$

should all lie on the straight line with y-intercept a and slope b (as determined by the function of P on the right-hand side in Eq. (2)).

This observation provides a way to find *a* and *b*. If we can determine the approximate values of the derivatives P'_1, P'_2, \ldots corresponding to the given population data, then we can proceed with the following agenda:

- First plot the points $(P_1, P'_1/P_1)$, $(P_2, P'_2/P_2)$, ... on a sheet of graph paper with horizontal *P*-axis.
- Then use a ruler to draw a straight line that appears to approximate these points well.
- Finally, measure this straight line's *y*-intercept *a* and slope *b*.

But where are we to find the needed values of the derivative P'(t) of the (as yet) unknown function P? It is easiest to use the approximation

$$P'_{i} = \frac{P_{i+1} - P_{i-1}}{t_{i+1} - t_{i-1}}$$
(3)



suggested by Fig. 2.1.8. For instance, if we take i = 0 corresponding to the year 1790, then the U.S. population data in Fig. 2.1.9 give

$$P_1' = \frac{P_2 - P_0}{t_2 - t_0} = \frac{7.240 - 3.929}{20} \approx 0.166$$

for the slope at (t_1, P_1) corresponding to the year 1800.

INVESTIGATION A: Use Eq. (3) to verify the slope figures shown in the final column of the table in Fig. 2.1.9, then plot the points $(P_1, P'_1/P_1), \ldots, (P_{11}, P'_{11}/P_{11})$ indicated by the dots in Fig. 2.1.10. If an appropriate graphing calculator, spread-sheet, or computer program is available, use it to find the straight line y = a + bP as in (2) that best fits these points. If not, draw your own straight line approximating these points, and then measure its intercept *a* and slope *b* as accurately as you can. Next, solve the logistic equation in (1) with these numerical parameters, taking t = 0 corresponding to the year 1800. Finally, compare the predicted 20th- and 21st-century U.S. population figures with the actual data listed in Fig. 2.1.4.

Year	i	t _i	Population P _i	Slope P'_i
1790	0	-10	3.929	
1800	1	0	5.308	0.166
1810	2	10	7.240	0.217
1820	3	20	9.638	0.281
1830	4	30	12.861	0.371
1840	5	40	17.064	0.517
1850	6	50	23.192	0.719
1860	7	60	31.443	0.768
1870	8	70	38.558	0.937
1880	9	80	50.189	1.221
1890	10	90	62.980	1.301
1900	11	100	76.212	1.462
1910	12	110	92.228	





Year	World Population (billions)
1975	4.062
1980	4.440
1985	4.853
1990	5.310
1995	5.735
2000	6.127
2005	6.520
2010	6.930
2015	7.349

FIGURE 2.1.10. Points and approximating straight line for U.S. population data from 1800 to 1900.



INVESTIGATION B: Repeat Investigation A, but take t = 0 in 1900 and use only the 20th-century population data listed in Fig. 2.1.4 to create a logistic model. How well does your model predict the U.S. population in the years 1990–2010?

INVESTIGATION C: Model similarly the world population data shown in Fig. 2.1.11. The Population Division of the United Nations predicts a world population of 9.157 billion in the year 2040. What do you predict?

2.2 Equilibrium Solutions and Stability

In previous sections we have often used explicit solutions of differential equations to answer specific numerical questions. But even when a given differential equation is difficult or impossible to solve explicitly, it often is possible to extract *qualitative* information about general properties of its solutions. For example, we may be able to establish that every solution x(t) grows without bound as $t \to +\infty$, or approaches a finite limit, or is a periodic function of t. In this section we introduce—mainly by consideration of simple differential equations that *can* be solved explicitly—some of the more important qualitative questions that can sometimes be answered for equations that are difficult or impossible to solve.

Example 1



FIGURE 2.2.1. Typical solution curves for the equation of Newton's law of cooling, dx/dt = -k(x - A).

Cooling and heating Let x(t) denote the temperature of a body with initial temperature $x(0) = x_0$. At time t = 0 this body is immersed in a medium with constant temperature A. Assuming Newton's law of cooling,

$$\frac{dx}{dt} = -k(x - A) \qquad (k > 0 \quad \text{constant}), \tag{1}$$

we readily solve (by separation of variables) for the explicit solution

$$x(t) = A + (x_0 - A)e^{-kt}.$$

It follows immediately that

$$\lim_{t \to \infty} x(t) = A,$$
(2)

so the temperature of the body approaches that of the surrounding medium (as is evident to one's intuition). Note that the constant function $x(t) \equiv A$ is a solution of Eq. (1); it corresponds to the temperature of the body when it is in thermal equilibrium with the surrounding medium. In Fig. 2.2.1 the limit in (2) means that every other solution curve approaches the equilibrium solution curve x = A asymptotically as $t \to +\infty$.

x' > 0		x' < 0	
x < A	x = A	x > A	

FIGURE 2.2.2. Phase diagram for the equation dx/dt = f(x) = k(A - x).

Remark The behavior of solutions of Eq. (1) is summarized briefly by the **phase diagram** in Fig. 2.2.2—which indicates the direction (or "phase") of change in x as a function of x itself. The right-hand side f(x) = -k(x - A) = k(A - x) is positive if x < A, negative if x > A. This observation corresponds to the fact that solutions starting above the line x = A and those starting below it both approach the limiting solution $x(t) \equiv A$ as t increases (as indicated by the arrows).

In Section 2.1 we introduced the general population equation

$$\frac{dx}{dt} = (\beta - \delta)x,\tag{3}$$

where β and δ are the birth and death rates, respectively, in births or deaths per individual per unit of time. The question of whether a population x(t) is bounded or unbounded as $t \to +\infty$ is of evident interest. In many situations—like the logistic and explosion/extinction populations of Section 2.1—the birth and death rates are known functions of x. Then Eq. (3) takes the form

$$\frac{dx}{dt} = f(x). \tag{4}$$

This is an **autonomous** first-order differential equation—one in which the independent variable *t* does not appear explicitly (the terminology here stemming from the Greek word *autonomos* for "independent," e.g., of the time *t*). As in Example 1, the solutions of the equation f(x) = 0 play an important role and are called **critical points** of the autonomous differential equation dx/dt = f(x).

If x = c is a critical point of Eq. (4), then the differential equation has the constant solution $x(t) \equiv c$. A constant solution of a differential equation is sometimes called an **equilibrium solution** (one may think of a population that remains constant because it is in "equilibrium" with its environment). Thus the critical point x = c, a number, corresponds to the equilibrium solution $x(t) \equiv c$, a constant-valued function.

Example 2 illustrates the fact that the qualitative behavior (as t increases) of the solutions of an autonomous first-order equation can be described in terms of its critical points.

Example 2 Logistic equation Consider the logistic differential equation

$$\frac{dx}{dt} = kx(M-x) \tag{5}$$

(with k > 0 and M > 0). It has two critical points—the solutions x = 0 and x = M of the equation

$$f(x) = kx(M - x) = 0.$$

In Section 2.1 we discussed the logistic-equation solution

$$x(t) = \frac{Mx_0}{x_0 + (M - x_0)e^{-kMt}}$$
(6)

satisfying the initial condition $x(0) = x_0$. Note that the initial values $x_0 = 0$ and $x_0 = M$ yield the equilibrium solutions $x(t) \equiv 0$ and $x(t) \equiv M$ of Eq. (5).

We observed in Section 2.1 that if $x_0 > 0$, then $x(t) \to M$ as $t \to +\infty$. But if $x_0 < 0$, then the denominator in Eq. (6) initially is positive, but vanishes when

$$t = t_1 = \frac{1}{kM} \ln \frac{M - x_0}{-x_0} > 0.$$



FIGURE 2.2.3. Typical solution curves for the logistic equation dx/dt = kx(M - x).



FIGURE 2.2.4. Solution curves, funnel, and spout for $dx/dt = 4x - x^2$.

x' < 0	<i>x'</i> > 0		$x'\!<\!0$
<i>x</i> =	= 0	x = M	
Unst	table	Stable	

FIGURE 2.2.5. Phase diagram for the logistic equation dx/dt = f(x) = kx(M - x).

Because the numerator in (6) is negative in this case, it follows that

$$\lim_{t \to t_1^-} x(t) = -\infty \quad \text{if} \quad x_0 < 0.$$

It follows that the solution curves of the logistic equation in (5) look as illustrated in Fig. 2.2.3. Here we see graphically that every solution either approaches the equilibrium solution $x(t) \equiv M$ as t increases, or (in a visually obvious sense) diverges away from the other equilibrium solution $x(t) \equiv 0$.

Stability of Critical Points

Figure 2.2.3 illustrates the concept of *stability*. A critical point x = c of an autonomous first-order equation is said to be *stable* provided that, if the initial value x_0 is sufficiently close to c, then x(t) remains close to c for all t > 0. More precisely, the critical point c is **stable** if, for each $\epsilon > 0$, there exists $\delta > 0$ such that

$$|x_0 - c| < \delta$$
 implies that $|x(t) - c| < \epsilon$ (7)

for all t > 0. The critical point x = c is **unstable** if it is not stable.

Figure 2.2.4 shows a "wider view" of the solution curves of a logistic equation with k = 1 and M = 4. Note that the strip 3.5 < x < 4.5 enclosing the stable equilibrium curve x = 4 acts like a *funnel*—solution curves (moving from left to right) enter this strip and thereafter remain within it. By contrast, the strip -0.5 < x < 0.5 enclosing the unstable solution curve x = 0 acts like a *spout*—solution curves leave this strip and thereafter remain outside it. Thus the critical point x = M is stable, whereas the critical point x = 0 is unstable.

Remark 1 We can summarize the behavior of solutions of the logistic equation in (5)—in terms of their initial values—by means of the phase diagram shown in Fig. 2.2.5. It indicates that $x(t) \rightarrow M$ as $t \rightarrow +\infty$ if either $x_0 > M$ or $0 < x_0 < M$, whereas $x(t) \rightarrow -\infty$ as t increases if $x_0 < 0$. The fact that M is a stable critical point would be important, for instance, if we wished to conduct an experiment with a population of M bacteria. It is impossible to count precisely M bacteria for M large, but any initially positive population will approach M as t increases.

Remark 2 Related to the stability of the limiting solution M = a/b of the logistic equation

$$\frac{dx}{dt} = ax - bx^2 \tag{8}$$

is the "predictability" of M for an actual population. The coefficients a and b are unlikely to be known precisely for an actual population. But if they are replaced with close approximations a^* and b^* —derived perhaps from empirical measurements—then the approximate limiting population $M^* = a^*/b^*$ will be close to the actual limiting population M = a/b. We may therefore say that the value M of the limiting population predicted by a logistic equation not only is a stable critical point of the differential equation, but this value also is "stable" with respect to small perturbations of the constant coefficients in the equation. (Note that one of these two statements involves changes in the initial value x_0 ; the other involves changes in the coefficients a and b.)

Example 3 Explosion/extinction Consider now the explosion/extinction equation

$$\frac{dx}{dt} = kx(x - M) \tag{9}$$

of Eq. (10) in Section 2.1. Like the logistic equation, it has the two critical points x = 0 and x = M corresponding to the equilibrium solutions $x(t) \equiv 0$ and $x(t) \equiv M$. According to Problem 33 in Section 2.1, its solution with $x(0) = x_0$ is given by

$$x(t) = \frac{Mx_0}{x_0 + (M - x_0)e^{kMt}}$$
(10)



FIGURE 2.2.6. Typical solution curves for the explosion/extinction equation dx/dt = kx(x - M).

x' > 0		x' < 0	x' > 0
	x = 0	<i>x</i> =	= M
	Stable	Uns	table

FIGURE 2.2.7. Phase diagram for the explosion/extinction equation dx/dt = f(x) = kx(x - M).

(with only a single difference in sign from the logistic solution in (6)). If $x_0 < M$, then (because the coefficient of the exponential in the denominator is positive) it follows immediately from Eq. (10) that $x(t) \rightarrow 0$ as $t \rightarrow +\infty$. But if $x_0 > M$, then the denominator in (10) initially is positive, but vanishes when

$$t = t_1 = \frac{1}{kM} \ln \frac{x_0}{x_0 - M} > 0.$$

Because the numerator in (10) is positive in this case, it follows that

$$\lim_{t \to t_1^-} x(t) = +\infty \quad \text{if} \quad x_0 > M.$$

Therefore, the solution curves of the explosion/extinction equation in (9) look as illustrated in Fig. 2.2.6. A narrow band along the equilibrium curve x = 0 (as in Fig. 2.2.4) would serve as a funnel, while a band along the solution curve x = M would serve as a spout for solutions. The behavior of the solutions of Eq. (9) is summarized by the phase diagram in Fig. 2.2.7, where we see that the critical point x = 0 is stable and the critical point x = M is unstable.

Harvesting a Logistic Population

The autonomous differential equation

$$\frac{dx}{dt} = ax - bx^2 - h \tag{11}$$

(with a, b, and h all positive) may be considered to describe a logistic population with harvesting. For instance, we might think of the population of fish in a lake from which h fish per year are removed by fishing.

Example 4	Threshold/limiting population	Let us rewrite Eq. (11) in the form	
		$\frac{dx}{dt} = kx(M-x) - h,$	(12)

which exhibits the limiting population M in the case h = 0 of no harvesting. Assuming hereafter that h > 0, we can solve the quadratic equation $-kx^2 + kMx - h = 0$ for the two critical points

$$H, N = \frac{kM \pm \sqrt{(kM)^2 - 4hk}}{2k} = \frac{1}{2} \left(M \pm \sqrt{M^2 - 4h/k} \right), \tag{13}$$

assuming that the harvesting rate h is sufficiently small that $4h < kM^2$, so both roots H and N are real with 0 < H < N < M. Then we can rewrite Eq. (12) in the form

$$\frac{dx}{dt} = k(N-x)(x-H).$$
(14)

For instance, the number of critical points of the equation may change abruptly as the value of a parameter is changed. In Problem 24 we ask you to solve this equation for the solution

$$x(t) = \frac{N(x_0 - H) - H(x_0 - N)e^{-k(N - H)t}}{(x_0 - H) - (x_0 - N)e^{-k(N - H)t}}$$
(15)

in terms of the initial value $x(0) = x_0$.

Note that the exponent -k(N - H)t is negative for t > 0. If $x_0 > N$, then each of the coefficients within parentheses in Eq. (15) is positive; it follows that

If
$$x_0 > N$$
 then $x(t) \to N$ as $t \to +\infty$. (16)

In Problem 25 we ask you to deduce also from Eq. (15) that

If $H < x_0 < N$ then $x(t) \to N$ as $t \to +\infty$, whereas (17)

if
$$x_0 < H$$
 then $x(t) \to -\infty$ as $t \to t_1$ (18)

for a positive value t_1 that depends on x_0 . It follows that the solution curves of Eq. (12)—still assuming that $4h < kM^2$ —look as illustrated in Fig. 2.2.8. (Can you visualize a funnel along the line x = N and a spout along the line x = H?) Thus the constant solution $x(t) \equiv N$ is an equilibrium *limiting solution*, whereas $x(t) \equiv H$ is a *threshold solution* that separates different behaviors—the population approaches N if $x_0 > H$, while it becomes extinct because of harvesting if $x_0 < H$. Finally, the stable critical point x = N and the unstable critical point x = H are illustrated in the phase diagram in Fig. 2.2.9.



equation dx/dt = k(N-x)(x-H).

 $\frac{dx}{dt} = f(x) = k(N - x)(x - H).$

Example 5

Lake stocked with fish For a concrete application of our stability conclusions in Example 4, suppose that k = 1 and M = 4 for a logistic population x(t) of fish in a lake, measured in hundreds after t years. Without any fishing at all, the lake would eventually contain nearly 400 fish, whatever the initial population. Now suppose that h = 3, so that 300 fish are "harvested" annually (at a constant rate throughout the year). Equation (12) is then dx/dt = x(4-x) - 3, and the quadratic equation

$$-x^{2} + 4x - 3 = (3 - x)(x - 1) = 0$$

has solutions H = 1 and N = 3. Thus the threshold population is 100 fish and the (new) limiting population is 300 fish. In short, if the lake is stocked initially with more than 100 fish, then as t increases, the fish population will approach a limiting value of 300 fish. But if the lake is stocked initially with fewer than 100 fish, then the lake will be "fished out" and the fish will disappear entirely within a finite period of time.

Bifurcation and Dependence on Parameters

A biological or physical system that is modeled by a differential equation may depend crucially on the numerical values of certain coefficients or parameters that appear in the equation. For instance, the number of critical points of the equation may change abruptly as the value of a parameter is changed.

Example 6 **Critical/excessive harvesting** The differential equation

$$\frac{dx}{dt} = x(4-x) - h \tag{19}$$

(with x in hundreds) models the harvesting of a logistic population with k = 1 and limiting population M = 4 (hundred). In Example 5 we considered the case of harvesting level h = 3, and found that the new limiting population is N = 3 hundred and the threshold population is



FIGURE 2.2.10. Solution curves of the equation x' = x(4 - x) - h with critical harvesting h = 4.



FIGURE 2.2.11. Solution curves of the equation x' = x(4 - x) - h with excessive harvesting h = 5.



FIGURE 2.2.12. The parabola $(c-2)^2 = 4 - h$ is the bifurcation diagram for the differential equation x' = x(4-x) - h.

H = 1 hundred. Typical solution curves, including the equilibrium solutions $x(t) \equiv 3$ and $x(t) \equiv 1$, then look like those pictured in Fig. 2.2.8.

Now let's investigate the dependence of this picture upon the harvesting level h. According to Eq. (13) with k = 1 and M = 4, the limiting and threshold populations N and H are given by

$$H, N = \frac{1}{2} \left(4 \pm \sqrt{16 - 4h} \right) = 2 \pm \sqrt{4 - h}.$$
 (20)

If h < 4—we can consider negative values of h to describe stocking rather than harvesting the fish—then there are distinct equilibrium solutions $x(t) \equiv N$ and $x(t) \equiv H$ with N > H as in Fig. 2.2.8.

But if h = 4, then Eq. (20) gives N = H = 2, so the differential equation has only the single equilibrium solution $x(t) \equiv 2$. In this case the solution curves of the equation look like those illustrated in Fig. 2.2.10. If the initial number x_0 (in hundreds) of fish exceeds 2, then the population approaches a limiting population of 2 (hundred fish). However, any initial population $x_0 < 2$ (hundred) results in extinction with the fish dying out as a consequence of the harvesting of 4 hundred fish annually. The critical point x = 2 might therefore be described as "semistable"—it looks stable on the side x > 2 where solution curves approach the equilibrium solution $x(t) \equiv 2$ as t increases, but unstable on the side x < 2 where solution curves instead diverge away from the equilibrium solution.

If, finally, h > 4, then the quadratic equation corresponding to (20) has no real solutions and the differential equation in (19) has no equilibrium solutions. The solution curves then look like those illustrated in Fig. 2.2.11, and (whatever the initial number of fish) the population dies out as a result of the excessive harvesting.

If we imagine turning a dial to gradually increase the value of the parameter h in Eq. (19), then the picture of the solution curves changes from one like Fig. 2.2.8 with h < 4, to Fig. 2.2.10 with h = 4, to one like Fig. 2.2.11 with h > 4. Thus the differential equation has

- two critical points if *h* < 4 ;
- one critical point if h = 4;
- no critical point if h > 4.

The value h = 4—for which the qualitative nature of the solutions changes as h increases—is called a **bifurcation point** for the differential equation containing the parameter h. A common way to visualize the corresponding "bifurcation" in the solutions is to plot the **bifurcation diagram** consisting of all points (h, c), where c is a critical point of the equation x' = x(4 - x) + h. For instance, if we rewrite Eq. (20) as

$$c = 2 \pm \sqrt{4 - h},$$

 $(c - 2)^2 = 4 - h,$

where either c = N or c = H, then we get the equation of the parabola that is shown in Fig. 2.2.12. This parabola is then the bifurcation diagram for our differential equation that models a logistic fish population with harvesting at the level specified by the parameter h.

2.2 **Problems**

In Problems 1 through 12 first solve the equation f(x) = 0to find the critical points of the given autonomous differential equation dx/dt = f(x). Then analyze the sign of f(x) to determine whether each critical point is stable or unstable, and construct the corresponding phase diagram for the differential equation. Next, solve the differential equation explicitly for x(t) in terms of t. Finally, use either the exact solution or a computer-generated slope field to sketch typical solution curves for the given differential equation, and verify visually the stability of each critical point.

1.
$$\frac{dx}{dt} = x - 4$$
 2. $\frac{dx}{dt} = 3 - x$

3.
$$\frac{dx}{dt} = x^2 - 4x$$

4. $\frac{dx}{dt} = 3x - x^2$
5. $\frac{dx}{dt} = x^2 - 4$
6. $\frac{dx}{dt} = 9 - x^2$
7. $\frac{dx}{dt} = (x - 2)^2$
8. $\frac{dx}{dt} = -(3 - x)^2$
9. $\frac{dx}{dt} = x^2 - 5x + 4$
10. $\frac{dx}{dt} = 7x - x^2 - 10$
11. $\frac{dx}{dt} = (x - 1)^3$
12. $\frac{dx}{dt} = (2 - x)^3$

In Problems 13 through 18, use a computer system or graphing calculator to plot a slope field and/or enough solution curves to indicate the stability or instability of each critical point of the given differential equation. (Some of these critical points may be semistable in the sense mentioned in Example 6.)

- **13.** $\frac{dx}{dt} = (x+2)(x-2)^2$ **14.** $\frac{dx}{dt} = x(x^2-4)$ **15.** $\frac{dx}{dt} = (x^2-4)^2$ **16.** $\frac{dx}{dt} = (x^2-4)^3$ **17.** $\frac{dx}{dt} = x^2(x^2-4)$ **18.** $\frac{dx}{dt} = x^3(x^2-4)$
- 19. The differential equation $dx/dt = \frac{1}{10}x(10-x) h$ models a logistic population with harvesting at rate *h*. Determine (as in Example 6) the dependence of the number of critical points on the parameter *h*, and then construct a bifurcation diagram like Fig. 2.2.12.
- **20.** The differential equation $dx/dt = \frac{1}{100}x(x-5) + s$ models a population with stocking at rate *s*. Determine the dependence of the number of critical points *c* on the parameter *s*, and then construct the corresponding bifurcation diagram in the *sc*-plane.
- **21.** Pitchfork bifurcation Consider the differential equation $dx/dt = kx x^3$. (a) If $k \le 0$, show that the only critical value c = 0 of x is stable. (b) If k > 0, show that the critical point c = 0 is now unstable, but that the critical points $c = \pm \sqrt{k}$ are stable. Thus the qualitative nature of the solutions changes at k = 0 as the parameter k increases, and so k = 0 is a bifurcation point for the differential equation with parameter k. The plot of all points of the form (k, c) where c is a critical point of the equation $x' = kx x^3$ is the "pitchfork diagram" shown in Fig. 2.2.13.



FIGURE 2.2.13. Bifurcation diagram for $dx/dt = kx - x^3$.

22. Consider the differential equation $dx/dt = x + kx^3$ containing the parameter k. Analyze (as in Problem 21) the dependence of the number and nature of the critical points

on the value of k, and construct the corresponding bifurcation diagram.

- 23. Variable-rate harvesting Suppose that the logistic equation dx/dt = kx(M x) models a population x(t) of fish in a lake after t months during which no fishing occurs. Now suppose that, because of fishing, fish are removed from the lake at the rate of hx fish per month (with h a positive constant). Thus fish are "harvested" at a rate proportional to the existing fish population, rather than at the constant rate of Example 4. (a) If 0 < h < kM, show that the population? (b) If h ≥ kM, show that x(t) → 0 are t → +∞, so the lake is eventually fished out.
- 24. Separate variables in the logistic harvesting equation dx/dt = k(N x)(x H) and then use partial fractions to derive the solution given in Eq. (15).
- **25.** Use the alternative forms

$$x(t) = \frac{N(x_0 - H) + H(N - x_0)e^{-k(N - H)t}}{(x_0 - H) + (N - x_0)e^{-k(N - H)t}}$$
$$= \frac{H(N - x_0)e^{-k(N - H)t} - N(H - x_0)}{(N - x_0)e^{-k(N - H)t} - (H - x_0)}$$

of the solution in (15) to establish the conclusions stated in (17) and (18).

Constant-Rate Harvesting

Example 4 dealt with the case $4h > kM^2$ in the equation dx/dt = kx(M - x) - h that describes constant-rate harvesting of a logistic population. Problems 26 and 27 deal with the other cases.

26. If $4h = kM^2$, show that typical solution curves look as illustrated in Fig. 2.2.14. Thus if $x_0 \ge M/2$, then $x(t) \rightarrow M/2$ as $t \rightarrow +\infty$. But if $x_0 < M/2$, then x(t) = 0 after a finite period of time, so the lake is fished out. The critical point x = M/2 might be called *semistable*, because it looks stable from one side, unstable from the other.



FIGURE 2.2.14. Solution curves for harvesting a logistic population with $4h = kM^2$.

27. If $4h > kM^2$, show that x(t) = 0 after a finite period of time, so the lake is fished out (whatever the initial population). [*Suggestion:* Complete the square to rewrite the differential equation in the form $dx/dt = -k[(x-a)^2 + b^2]$.
Then solve explicitly by separation of variables.] The results of this and the previous problem (together with Example 4) show that $h = \frac{1}{4}kM^2$ is a critical harvesting rate for a logistic population. At any lesser harvesting rate the population approaches a limiting population *N* that is less than *M* (why?), whereas at any greater harvesting rate the population reaches extinction.

28. Alligator population This problem deals with the differential equation dx/dt = kx(x - M) - h that models the harvesting of an unsophisticated population (such as alligators). Show that this equation can be rewritten in the form dx/dt = k(x - H)(x - K), where

$$H = \frac{1}{2} \left(M + \sqrt{M^2 + 4h/k} \right) > 0,$$

$$K = \frac{1}{2} \left(M - \sqrt{M^2 + 4h/k} \right) < 0.$$

Show that typical solution curves look as illustrated in Fig. 2.2.15.

29. Consider the two differential equations

$$\frac{dx}{dt} = (x-a)(x-b)(x-c)$$
(21)

and

 $\frac{dx}{dt} = (a-x)(b-x)(c-x),$

2.3 Acceleration–Velocity Models

each having the critical points a, b, and c; suppose that a < b < c. For one of these equations, only the critical point b is stable; for the other equation, b is the only unstable critical point. Construct phase diagrams for the two equations to determine which is which. Without attempting to solve either equation explicitly, make rough sketches of typical solution curves for each. You should see two funnels and a spout in one case, two spouts and a funnel in the other.



FIGURE 2.2.15. Solution curves for harvesting a population of alligators.

In Section 1.2 we discussed vertical motion of a mass m near the surface of the earth under the influence of constant gravitational acceleration. If we neglect any effects of air resistance, then Newton's second law (F = ma) implies that the velocity v of

$$m\frac{dv}{dt} = F_G,\tag{1}$$

where $F_G = -mg$ is the (downward-directed) force of gravity, where the gravitational acceleration is $g \approx 9.8 \text{ m/s}^2$ (in mks units; $g \approx 32 \text{ ft/s}^2$ in fps units).

Example 1 No air resistance Suppose that a crossbow bolt is shot straight upward from the ground $(y_0 = 0)$ with initial velocity $v_0 = 49$ (m/s). Then Eq. (1) with g = 9.8 gives

$$\frac{dv}{dt} = -9.8$$
, so $v(t) = -(9.8)t + v_0 = -(9.8)t + 49$.

Hence the bolt's height function y(t) is given by

(22)

the mass m satisfies the equation

$$y(t) = \int \left[-(9.8)t + 49 \right] dt = -(4.9)t^2 + 49t + y_0 = -(4.9)t^2 + 49t$$

The bolt reaches its maximum height when v = -(9.8)t + 49 = 0, hence when t = 5 (s). Thus its maximum height is

$$y_{\text{max}} = y(5) = -(4.9)(5^2) + (49)(5) = 122.5 \text{ (m)}.$$

The bolt returns to the ground when y = -(4.9)t(t - 10) = 0, and thus after 10 seconds aloft.

Now we want to take account of air resistance in a problem like Example 1. The force F_R exerted by air resistance on the moving mass *m* must be added in Eq. (1), so now

$$m\frac{dv}{dt} = F_G + F_R.$$
 (2)

Newton showed in his *Principia Mathematica* that certain simple physical assumptions imply that F_R is proportional to the *square* of the velocity: $F_R = kv^2$. But empirical investigations indicate that the actual dependence of air resistance on velocity can be quite complicated. For many purposes it suffices to assume that

$$F_R = k v^p$$
,

where $1 \le p \le 2$ and the value of k depends on the size and shape of the body, as well as the density and viscosity of the air. Generally speaking, p = 1 for relatively low speeds and p = 2 for high speeds, whereas 1 for intermediate speeds. But how slow "low speed" and how fast "high speed" are depend on the same factors that determine the value of the coefficient k.

Thus air resistance is a complicated physical phenomenon. But the simplifying assumption that F_R is exactly of the form given here, with either p = 1 or p = 2, yields a tractable mathematical model that exhibits the most important qualitative features of motion with resistance.

Resistance Proportional to Velocity

Let us first consider the vertical motion of a body with mass *m* near the surface of the earth, subject to two forces: a downward gravitational force F_G and a force F_R of air resistance that is proportional to velocity (so that p = 1) and of course directed opposite the direction of motion of the body. If we set up a coordinate system with the positive *y*-direction upward and with y = 0 at ground level, then $F_G = -mg$ and

$$F_R = -kv, (3)$$

where k is a positive constant and v = dy/dt is the velocity of the body. Note that the minus sign in Eq. (3) makes F_R positive (an upward force) if the body is falling (v is negative) and makes F_R negative (a downward force) if the body is rising (v is positive). As indicated in Fig. 2.3.1, the net force acting on the body is then

$$F = F_R + F_G = -kv - mg,$$

and Newton's law of motion F = m(dv/dt) yields the equation

$$m\frac{dv}{dt} = -kv - mg$$

Thus

 \succ

$$\frac{dv}{dt} = -\rho v - g,\tag{4}$$

where $\rho = k/m > 0$. You should verify for yourself that if the positive *y*-axis were directed downward, then Eq. (4) would take the form $dv/dt = -\rho v + g$.

Equation (4) is a separable first-order differential equation, and its solution is

$$v(t) = \left(v_0 + \frac{g}{\rho}\right)e^{-\rho t} - \frac{g}{\rho}.$$
(5)



FIGURE 2.3.1. Vertical motion with air resistance.

Here, $v_0 = v(0)$ is the initial velocity of the body. Note that

$$v_{\tau} = \lim_{t \to \infty} v(t) = -\frac{g}{\rho}.$$
 (6)

Thus the speed of a body falling with air resistance does *not* increase indefinitely; instead, it approaches a *finite* limiting speed, or **terminal speed**,

$$|v_{\tau}| = \frac{g}{\rho} = \frac{mg}{k}.$$
(7)

This fact is what makes a parachute a practical invention; it even helps explain the occasional survival of people who fall without parachutes from high-flying airplanes.

We now rewrite Eq. (5) in the form

$$\frac{dy}{dt} = (v_0 - v_\tau)e^{-\rho t} + v_\tau.$$
 (8)

Integration gives

$$y(t) = -\frac{1}{\rho}(v_0 - v_\tau)e^{-\rho t} + v_\tau t + C.$$

We substitute 0 for t and let $y_0 = y(0)$ denote the initial height of the body. Thus we find that $C = y_0 + (v_0 - v_\tau)/\rho$, and so

$$y(t) = y_0 + v_\tau t + \frac{1}{\rho} (v_0 - v_\tau) (1 - e^{-\rho t}).$$
(9)

Equations (8) and (9) give the velocity v and height y of a body moving vertically under the influence of gravity and air resistance. The formulas depend on the initial height y_0 of the body, its initial velocity v_0 , and the *drag coefficient* ρ , the constant such that the acceleration due to air resistance is $a_R = -\rho v$. The two equations also involve the terminal velocity v_{τ} defined in Eq. (6).

For a person descending with the aid of a parachute, a typical value of ρ is 1.5, which corresponds to a terminal speed of $|v_{\tau}| \approx 21.3$ ft/s, or about 14.5 mi/h. With an unbuttoned overcoat flapping in the wind in place of a parachute, an unlucky skydiver might increase ρ to perhaps as much as 0.5, which gives a terminal speed of $|v_{\tau}| \approx 65$ ft/s, about 44 mi/h. See Problems 10 and 11 for some parachute-jump computations.

Example 2 Velocity-proportional resistance We again consider a bolt shot straight upward with initial velocity $v_0 = 49$ m/s from a crossbow at ground level. But now we take air resistance into account, with $\rho = 0.04$ in Eq. (4). We ask how the resulting maximum height and time aloft compare with the values found in Example 1.

Solution

We substitute
$$y_0 = 0$$
, $v_0 = 49$, and $v_\tau = -g/\rho = -245$ in Eqs. (5) and (9), and obtain
 $v(t) = 294e^{-t/25} = 245$

$$y(t) = 7350 - 245t - 7350e^{-t/25}$$

To find the time required for the bolt to reach its maximum height (when v = 0), we solve the equation

$$v(t) = 294e^{-t/25} - 245 = 0$$

for $t_m = 25 \ln(294/245) \approx 4.558$ (s). Its maximum height is then $y_{\text{max}} = v(t_m) \approx 108.280$ meters (as opposed to 122.5 meters without air resistance). To find when the bolt strikes the ground, we must solve the equation

$$y(t) = 7350 - 245t - 7350e^{-t/25} = 0.$$

Using Newton's method, we can begin with the initial guess $t_0 = 10$ and carry out the iteration $t_{n+1} = t_n - y(t_n)/y'(t_n)$ to generate successive approximations to the root. Or we can simply use the **Solve** command on a calculator or computer. We find that the bolt is in the air for $t_f \approx 9.411$ seconds (as opposed to 10 seconds without air resistance). It hits the ground with a reduced speed of $|v(t_f)| \approx 43.227$ m/s (as opposed to its initial velocity of 49 m/s).

Thus the effect of air resistance is to decrease the bolt's maximum height, the total time spent aloft, and its final impact speed. Note also that the bolt now spends more time in descent ($t_{\rm f} - t_m \approx 4.853$ s) than in ascent ($t_m \approx 4.558$ s).

Resistance Proportional to Square of Velocity

Now we assume that the force of air resistance is proportional to the *square* of the velocity:

$$F_R = \pm k v^2, \tag{10}$$

with k > 0. The choice of signs here depends on the direction of motion, which the force of resistance always opposes. Taking the positive y-direction as upward, $F_R < 0$ for upward motion (when v > 0) while $F_R > 0$ for downward motion (when v < 0). Thus the sign of F_R is always opposite that of v, so we can rewrite Eq. (10) as

$$F_R = -kv|v|. \tag{10'}$$

Then Newton's second law gives

$$m\frac{dv}{dt} = F_G + F_R = -mg - kv|v|;$$

that is,

>

$$\frac{dv}{dt} = -g - \rho v |v|, \tag{11}$$

where $\rho = k/m > 0$. We must discuss the cases of upward and downward motion separately.

UPWARD MOTION: Suppose that a projectile is launched straight upward from the initial position y_0 with initial velocity $v_0 > 0$. Then Eq. (11) with v > 0 gives the differential equation

$$\frac{dv}{dt} = -g - \rho v^2 = -g \left(1 + \frac{\rho}{g} v^2\right).$$
(12)

In Problem 13 we ask you to make the substitution $u = v \sqrt{\rho/g}$ and apply the familiar integral

$$\int \frac{1}{1+u^2} \, du = \tan^{-1} u + C$$

to derive the projectile's velocity function

$$v(t) = \sqrt{\frac{g}{\rho}} \tan\left(C_1 - t\sqrt{\rho g}\right) \quad \text{with} \quad C_1 = \tan^{-1}\left(v_0\sqrt{\frac{\rho}{g}}\right). \tag{13}$$

Because $\int \tan u \, du = -\ln |\cos u| + C$, a second integration (see Problem 14) yields the position function

$$y(t) = y_0 + \frac{1}{\rho} \ln \left| \frac{\cos \left(C_1 - t \sqrt{\rho g} \right)}{\cos C_1} \right|.$$
 (14)

DOWNWARD MOTION: Suppose that a projectile is launched (or dropped) straight downward from the initial position y_0 with initial velocity $v_0 \leq 0$. Then Eq. (11) with v < 0 gives the differential equation

$$\frac{dv}{dt} = -g + \rho v^2 = -g \left(1 - \frac{\rho}{g} v^2\right).$$
(15)

In Problem 15 we ask you to make the substitution $u = v \sqrt{\rho/g}$ and apply the integral

$$\int \frac{1}{1-u^2} \, du = \tanh^{-1} u + C$$

to derive the projectile's velocity function

$$v(t) = \sqrt{\frac{g}{\rho}} \tanh\left(C_2 - t\sqrt{\rho g}\right) \quad \text{with} \qquad C_2 = \tanh^{-1}\left(v_0\sqrt{\frac{\rho}{g}}\right). \tag{16}$$

Because $\int \tanh u \, du = \ln |\cosh u| + C$, another integration (Problem 16) yields the position function

$$y(t) = y_0 - \frac{1}{\rho} \ln \left| \frac{\cosh\left(C_2 - t\sqrt{\rho g}\right)}{\cosh C_2} \right|.$$
(17)

(Note the analogy between Eqs. (16) and (17) and Eqs. (13) and (14) for upward motion.)

If $v_0 = 0$, then $C_2 = 0$, so $v(t) = -\sqrt{g/\rho} \tanh\left(t\sqrt{\rho g}\right)$. Because

$$\lim_{x \to \infty} \tanh x = \lim_{x \to \infty} \frac{\sinh x}{\cosh x} = \lim_{x \to \infty} \frac{\frac{1}{2}(e^x - e^{-x})}{\frac{1}{2}(e^x + e^{-x})} = 1,$$

it follows that in the case of downward motion the body approaches the terminal speed

$$v_{\tau}| = \sqrt{\frac{g}{\rho}} \tag{18}$$

(as compared with $|v_{\tau}| = g/\rho$ in the case of downward motion with linear resistance described by Eq. (4)).

Example 3 Square-proportional resistance We consider once more a bolt shot straight upward with initial velocity $v_0 = 49$ m/s from a crossbow at ground level, as in Example 2. But now we assume air resistance proportional to the square of the velocity, with $\rho = 0.0011$ in Eqs. (12) and (15). In Problems 17 and 18 we ask you to verify the entries in the last line of the following table.

Air Resistance	Maximum Height (ft)	Time Aloft (s)	Ascent Time (s)	Descent Time (s)	Impact Speed (ft/s)
0.0	122.5	10	5	5	49
(0.04)v	108.28	9.41	4.56	4.85	43.23
$(0.0011)v^2$	108.47	9.41	4.61	4.80	43.49



FIGURE 2.3.2. The height functions in Example 1 (without air resistance), Example 2 (with linear air resistance), and Example 3 (with air resistance proportional to the square of the velocity) are all plotted. The graphs of the latter two are visually indistinguishable.

>

Comparison of the last two lines of data here indicates little difference—for the motion of our crossbow bolt—between linear air resistance and air resistance proportional to the square of the velocity. And in Fig. 2.3.2, where the corresponding height functions are graphed, the difference is hardly visible. However, the difference between linear and nonlinear resistance can be significant in more complex situations—such as, for instance, the atmospheric reentry and descent of a space vehicle.

Variable Gravitational Acceleration

Unless a projectile in vertical motion remains in the immediate vicinity of the earth's surface, the gravitational acceleration acting on it is not constant. According to Newton's law of gravitation, the gravitational force of attraction between two point masses M and m located at a distance r apart is given by

$$F = \frac{GMm}{r^2},\tag{19}$$

where G is a certain empirical constant ($G \approx 6.6726 \times 10^{-11} \text{ N} \cdot (\text{m/kg})^2$ in mks units). The formula is also valid if either or both of the two masses are homogeneous spheres; in this case, the distance r is measured between the centers of the spheres.

The following example is similar to Example 2 in Section 1.2, but now we take account of lunar gravity.

Example 4 Lunar lander A lunar lander is free-falling toward the moon, and at an altitude of 53 kilometers above the lunar surface its downward velocity is measured at 1477 km/h. Its retrorockets, when fired in free space, provide a deceleration of $T = 4 \text{ m/s}^2$. At what height above the lunar surface should the retrorockets be activated to ensure a "soft touchdown" (v = 0 at impact)?

Solution Let r(t) denote the lander's distance from the center of the moon at time t (Fig. 2.3.3). When we combine the (constant) thrust acceleration T and the (negative) lunar acceleration $F/m = GM/r^2$ of Eq. (19), we get the (acceleration) differential equation

$$\frac{d^2r}{dt^2} = T - \frac{GM}{r^2},\tag{20}$$

where $M = 7.35 \times 10^{22}$ (kg) is the mass of the moon, which has a radius of $R = 1.74 \times 10^6$ meters (or 1740 km, a little over a quarter of the earth's radius). Noting that this second-order differential equation does not involve the independent variable *t*, we substitute

$$v = \frac{dr}{dt}, \quad \frac{d^2r}{dt^2} = \frac{dv}{dt} = \frac{dv}{dr} \cdot \frac{dr}{dt} = v\frac{dv}{dr}$$

(as in Eq. (36) of Section 1.6) and obtain the first-order equation

$$v\frac{dv}{dr} = T - \frac{GM}{r^2}$$

with the new independent variable r. Integration with respect to r now yields the equation

$$\frac{1}{2}v^2 = Tr + \frac{GM}{r} + C \tag{21}$$

that we can apply both before ignition (T = 0) and after ignition (T = 4).

Before ignition: Substitution of T = 0 in (21) gives the equation

$$\frac{1}{2}v^2 = \frac{GM}{r} + C_1$$
 (21a)

where the constant is given by $C_1 = v_0^2/2 - GM/r_0$ with

$$v_0 = -1477 \, \frac{\mathrm{km}}{\mathrm{h}} \times 1000 \, \frac{\mathrm{m}}{\mathrm{km}} \times \frac{1 \, \mathrm{h}}{3600 \, \mathrm{s}} = -\frac{14770}{36} \, \frac{\mathrm{m}}{\mathrm{s}}$$



FIGURE 2.3.3. The lunar lander descending to the surface of the moon.

and $r_0 = (1.74 \times 10^6) + 53,000 = 1.793 \times 10^6$ m (from the initial velocity–position measurement).

After ignition: Substitution of T = 4 and v = 0, r = R (at touchdown) into (21) gives

$$\frac{1}{2}v^2 = 4r + \frac{GM}{r} + C_2 \tag{21b}$$

where the constant $C_2 = -4R - GM/R$ is obtained by substituting the values v = 0, r = R at touchdown.

At the instant of ignition the lunar lander's position and velocity satisfy both (21a) and (21b). Therefore we can find its desired height *h* above the lunar surface at ignition by equating the right-hand sides in (21a) and (21b). This gives $r = \frac{1}{4}(C_1 - C_2) = 1.78187 \times 10^6$ and finally h = r - R = 41,870 meters (that is, 41.87 kilometers—just over 26 miles). Moreover, substitution of this value of *r* in (21a) gives the velocity v = -450 m/s at the instant of ignition.

Escape Velocity

In his novel *From the Earth to the Moon* (1865), Jules Verne raised the question of the initial velocity necessary for a projectile fired from the surface of the earth to reach the moon. Similarly, we can ask what initial velocity v_0 is necessary for the projectile to escape from the earth altogether. This will be so if its velocity v = dr/dt remains *positive* for all t > 0, so it continues forever to move away from the earth. With r(t) denoting the projectile's distance from the earth's center at time t (Fig. 2.3.4), we have the equation

$$\frac{dv}{dt} = \frac{d^2r}{dt^2} = -\frac{GM}{r^2},\tag{22}$$

similar to Eq. (20), but with T = 0 (no thrust) and with $M = 5.975 \times 10^{24}$ (kg) denoting the mass of the earth, which has an equatorial radius of $R = 6.378 \times 10^{6}$ (m). Substitution of the chain rule expression dv/dt = v(dv/dr) as in Example 4 gives

$$v\frac{dv}{dr} = -\frac{GM}{r^2}$$

Then integration of both sides with respect to r yields

$$\frac{1}{2}v^2 = \frac{GM}{r} + C.$$

Now $v = v_0$ and r = R when t = 0, so $C = \frac{1}{2}v_0^2 - GM/R$, and hence solution for v^2 gives

$$v^{2} = v_{0}^{2} + 2GM\left(\frac{1}{r} - \frac{1}{R}\right).$$
(23)

This implicit solution of Eq. (22) determines the projectile's velocity v as a function of its distance r from the earth's center. In particular,

$$v^2 > v_0^2 - \frac{2GM}{R}$$

so v will remain positive provided that $v_0^2 \ge 2GM/R$. Therefore, the escape velocity from the earth is given by

$$v_0 = \sqrt{\frac{2GM}{R}}.$$
(24)



FIGURE 2.3.4. A mass *m* at a great distance from the earth.

In Problem 27 we ask you to show that, if the projectile's initial velocity exceeds $\sqrt{2GM/R}$, then $r(t) \to \infty$ as $t \to \infty$, so it does, indeed, "escape" from the earth. With the given values of *G* and the earth's mass *M* and radius *R*, this gives $v_0 \approx 11,180 \text{ (m/s)}$ (about 36,680 ft/s, about 6.95 mi/s, about 25,000 mi/h).

Remark Equation (24) gives the escape velocity for any other (spherical) planetary body when we use *its* mass and radius. For instance, when we use the mass M and radius R for the moon given in Example 4, we find that escape velocity from the lunar surface is $v_0 \approx 2375$ m/s. This is just over one-fifth of the escape velocity from the earth's surface, a fact that greatly facilitates the return trip ("From the Moon to the Earth").

2.3 Problems

1. The acceleration of a Maserati is proportional to the difference between 250 km/h and the velocity of this sports car. If this machine can accelerate from rest to 100 km/h in 10 s, how long will it take for the car to accelerate from rest to 200 km/h?

Problems 2 through 8 explore the effects of resistance proportional to a power of the velocity.

Suppose that a body moves through a resisting medium with resistance proportional to its velocity v, so that dv/dt = -kv. (a) Show that its velocity and position at time t are given by

$$v(t) = v_0 e^{-kt}$$

and

$$x(t) = x_0 + \left(\frac{v_0}{k}\right)(1 - e^{-kt}).$$

(b) Conclude that the body travels only a finite distance, and find that distance.

- **3.** Suppose that a motorboat is moving at 40 ft/s when its motor suddenly quits, and that 10 s later the boat has slowed to 20 ft/s. Assume, as in Problem 2, that the resistance it encounters while coasting is proportional to its velocity. How far will the boat coast in all?
- 4. Consider a body that moves horizontally through a medium whose resistance is proportional to the *square* of the velocity v, so that $dv/dt = -kv^2$. Show that

$$v(t) = \frac{v_0}{1 + v_0 k t}$$

and that

$$x(t) = x_0 + \frac{1}{k}\ln(1 + v_0kt).$$

Note that, in contrast with the result of Problem 2, $x(t) \rightarrow +\infty$ as $t \rightarrow +\infty$. Which offers less resistance when the body is moving fairly slowly—the medium in this problem or the one in Problem 2? Does your answer seem consistent with the observed behaviors of x(t) as $t \rightarrow \infty$?

5. Assuming resistance proportional to the square of the velocity (as in Problem 4), how far does the motorboat of Problem 3 coast in the first minute after its motor quits?

6. Assume that a body moving with velocity v encounters resistance of the form $dv/dt = -kv^{3/2}$. Show that

$$v(t) = \frac{4v_0}{\left(kt\sqrt{v_0} + 2\right)^2}$$

and that

$$x(t) = x_0 + \frac{2}{k}\sqrt{v_0}\left(1 - \frac{2}{kt\sqrt{v_0} + 2}\right).$$

Conclude that under a $\frac{3}{2}$ -power resistance a body coasts only a finite distance before coming to a stop.

- 7. Suppose that a car starts from rest, its engine providing an acceleration of 10 ft/s^2 , while air resistance provides 0.1 ft/s^2 of deceleration for each foot per second of the car's velocity. (a) Find the car's maximum possible (limiting) velocity. (b) Find how long it takes the car to attain 90% of its limiting velocity, and how far it travels while doing so.
- 8. Rework both parts of Problem 7, with the sole difference that the deceleration due to air resistance now is $(0.001)v^2$ ft/s² when the car's velocity is *v* feet per second.

Problems 9 through 12 illustrate resistance proportional to the velocity.

9. A motorboat weighs 32,000 lb and its motor provides a thrust of 5000 lb. Assume that the water resistance is 100 pounds for each foot per second of the speed v of the boat. Then

$$1000\frac{dv}{dt} = 5000 - 100v.$$

If the boat starts from rest, what is the maximum velocity that it can attain?

- **10.** Falling parachutist A woman bails out of an airplane at an altitude of 10,000 ft, falls freely for 20 s, then opens her parachute. How long will it take her to reach the ground? Assume linear air resistance ρv ft/s², taking $\rho = 0.15$ without the parachute and $\rho = 1.5$ with the parachute. (*Suggestion:* First determine her height above the ground and velocity when the parachute opens.)
- **11. Falling paratrooper** According to a newspaper account, a paratrooper survived a training jump from 1200

ft when his parachute failed to open but provided some resistance by flapping unopened in the wind. Allegedly he hit the ground at 100 mi/h after falling for 8 s. Test the accuracy of this account. (*Suggestion:* Find ρ in Eq. (4) by assuming a terminal velocity of 100 mi/h. Then calculate the time required to fall 1200 ft.)

12. Nuclear waste disposal It is proposed to dispose of nuclear wastes—in drums with weight W = 640 lb and volume 8 ft³—by dropping them into the ocean ($v_0 = 0$). The force equation for a drum falling through water is

$$m\frac{dv}{dt} = -W + B + F_R,$$

where the buoyant force *B* is equal to the weight (at 62.5 lb/ft^3) of the volume of water displaced by the drum (Archimedes' principle) and *F_R* is the force of water resistance, found empirically to be 1 lb for each foot per second of the velocity of a drum. If the drums are likely to burst upon an impact of more than 75 ft/s, what is the maximum depth to which they can be dropped in the ocean without likelihood of bursting?

- 13. Separate variables in Eq. (12) and substitute $u = v \sqrt{\rho/g}$ to obtain the upward-motion velocity function given in Eq. (13) with initial condition $v(0) = v_0$.
- 14. Integrate the velocity function in Eq. (13) to obtain the upward-motion position function given in Eq. (14) with initial condition $y(0) = y_0$.
- **15.** Separate variables in Eq. (15) and substitute $u = v \sqrt{\rho/g}$ to obtain the downward-motion velocity function given in Eq. (16) with initial condition $v(0) = v_0$.
- 16. Integrate the velocity function in Eq. (16) to obtain the downward-motion position function given in Eq. (17) with initial condition $y(0) = y_0$.

Problems 17 and 18 apply Eqs. (12)–(17) to the motion of a crossbow bolt.

- 17. Consider the crossbow bolt of Example 3, shot straight upward from the ground (y = 0) at time t = 0 with initial velocity $v_0 = 49$ m/s. Take g = 9.8 m/s² and $\rho = 0.0011$ in Eq. (12). Then use Eqs. (13) and (14) to show that the bolt reaches its maximum height of about 108.47 m in about 4.61 s.
- 18. Continuing Problem 17, suppose that the bolt is now dropped ($v_0 = 0$) from a height of $y_0 = 108.47$ m. Then use Eqs. (16) and (17) to show that it hits the ground about 4.80 s later with an impact speed of about 43.49 m/s.

Problems 19 through 23 illustrate resistance proportional to the square of the velocity.

19. A motorboat starts from rest (initial velocity v(0) = v₀ = 0). Its motor provides a constant acceleration of 4 ft/s², but water resistance causes a deceleration of v²/400 ft/s². Find v when t = 10 s, and also find the *limiting velocity* as t → +∞ (that is, the maximum possible speed of the boat).

- **20.** An arrow is shot straight upward from the ground with an initial velocity of 160 ft/s. It experiences both the deceleration of gravity and deceleration $v^2/800$ due to air resistance. How high in the air does it go?
- **21.** If a ball is projected upward from the ground with initial velocity v_0 and resistance proportional to v^2 , deduce from Eq. (14) that the maximum height it attains is

$$y_{\max} = \frac{1}{2\rho} \ln \left(1 + \frac{\rho v_0^2}{g} \right).$$

- **22.** Suppose that $\rho = 0.075$ (in fps units, with g = 32 ft/s²) in Eq. (15) for a paratrooper falling with parachute open. If he jumps from an altitude of 10,000 ft and opens his parachute immediately, what will be his terminal speed? How long will it take him to reach the ground?
- **23.** Suppose that the paratrooper of Problem 22 falls freely for 30 s with $\rho = 0.00075$ before opening his parachute. How long will it now take him to reach the ground?

Problems 24 through 30 explore gravitational acceleration and escape velocity.

- 24. The mass of the sun is 329,320 times that of the earth and its radius is 109 times the radius of the earth. (a) To what radius (in meters) would the earth have to be compressed in order for it to become a *black hole*—the escape velocity from its surface equal to the velocity $c = 3 \times 10^8$ m/s of light? (b) Repeat part (a) with the sun in place of the earth.
- **25.** (a) Show that if a projectile is launched straight upward from the surface of the earth with initial velocity v_0 less than escape velocity $\sqrt{2GM/R}$, then the maximum distance from the center of the earth attained by the projectile is

$$r_{\rm max} = \frac{2GMR}{2GM - Rv_0^2}$$

where *M* and *R* are the mass and radius of the earth, respectively. (b) With what initial velocity v_0 must such a projectile be launched to yield a maximum altitude of 100 kilometers above the surface of the earth? (c) Find the maximum distance from the center of the earth, expressed in terms of earth radii, attained by a projectile launched from the surface of the earth with 90% of escape velocity.

- **26.** Suppose that you are stranded—your rocket engine has failed—on an asteroid of diameter 3 miles, with density equal to that of the earth with radius 3960 miles. If you have enough spring in your legs to jump 4 feet straight up on earth while wearing your space suit, can you blast off from this asteroid using leg power alone?
- 27. (a) Suppose a projectile is launched vertically from the surface r = R of the earth with initial velocity $v_0 = \sqrt{2GM/R}$, so $v_0^2 = k^2/R$ where $k^2 = 2GM$. Solve the differential equation $dr/dt = k/\sqrt{r}$ (from Eq. (23) in this section) explicitly to deduce that $r(t) \to \infty$ as $t \to \infty$.

(b) If the projectile is launched vertically with initial velocity $v_0 > \sqrt{2GM/R}$, deduce that

$$\frac{dr}{dt} = \sqrt{\frac{k^2}{r} + \alpha} > \frac{k}{\sqrt{r}}.$$

Why does it again follow that $r(t) \to \infty$ as $t \to \infty$?

28. (a) Suppose that a body is dropped ($v_0 = 0$) from a distance $r_0 > R$ from the earth's center, so its acceleration is $dv/dt = -GM/r^2$. Ignoring air resistance, show that it reaches the height $r < r_0$ at time

$$t = \sqrt{\frac{r_0}{2GM}} \left(\sqrt{rr_0 - r^2} + r_0 \cos^{-1} \sqrt{\frac{r}{r_0}} \right).$$

(*Suggestion:* Substitute $r = r_0 \cos^2 \theta$ to evaluate $\int \sqrt{r/(r_0 - r)} dr$.) (b) If a body is dropped from a height of 1000 km above the earth's surface and air resistance is neglected, how long does it take to fall and with what speed will it strike the earth's surface?

29. Suppose that a projectile is fired straight upward from the surface of the earth with initial velocity $v_0 < \sqrt{2GM/R}$. Then its height y(t) above the surface satisfies the initial value problem

$$\frac{d^2y}{dt^2} = -\frac{GM}{(y+R)^2}; \quad y(0) = 0, \quad y'(0) = v_0$$

Substitute dv/dt = v(dv/dy) and then integrate to obtain

$$v^2 = v_0^2 - \frac{2GMy}{R(R+y)}$$

for the velocity v of the projectile at height y. What maximum altitude does it reach if its initial velocity is 1 km/s?

30. In Jules Verne's original problem, the projectile launched from the surface of the earth is attracted by both the earth and the moon, so its distance r(t) from the center of the earth satisfies the initial value problem

$$\frac{d^2r}{dt^2} = -\frac{GM_e}{r^2} + \frac{GM_m}{(S-r)^2}; \quad r(0) = R, \quad r'(0) = v_0$$

where M_e and M_m denote the masses of the earth and the moon, respectively; R is the radius of the earth and S = 384,400 km is the distance between the centers of the earth and the moon. To reach the moon, the projectile must only just pass the point between the moon and earth where its net acceleration vanishes. Thereafter it is "under the control" of the moon, and falls from there to the lunar surface. Find the *minimal* launch velocity v_0 that suffices for the projectile to make it "From the Earth to the Moon."

2.3 Application Rocket Propulsion

Suppose that the rocket of Fig. 2.3.5 blasts off straight upward from the surface of the earth at time t = 0. We want to calculate its height y and velocity v = dy/dt at time t. The rocket is propelled by exhaust gases that exit (rearward) with constant speed c (relative to the rocket). Because of the combustion of its fuel, the mass m = m(t) of the rocket is variable.

To derive the equation of motion of the rocket, we use Newton's second law in the form

$$\frac{dP}{dt} = F,\tag{1}$$

where *P* is momentum (the product of mass and velocity) and *F* denotes net external force (gravity, air resistance, etc.). If the mass *m* of the rocket is constant so $m'(t) \equiv 0$ —when its rockets are turned off or burned out, for instance—then Eq. (1) gives

$$F = \frac{d(mv)}{dt} = m\frac{dv}{dt} + \frac{dm}{dt}v = m\frac{dv}{dt},$$

which (with dv/dt = a) is the more familiar form F = ma of Newton's second law.

But here *m* is not constant. Suppose *m* changes to $m + \Delta m$ and *v* to $v + \Delta v$ during the short time interval from *t* to $t + \Delta t$. Then the change in the momentum *of the rocket itself* is

$$\Delta P \approx (m + \Delta m)(v + \Delta v) - mv = m \,\Delta v + v \,\Delta m + \Delta m \,\Delta v.$$

But the system also includes the exhaust gases expelled during this time interval, with mass $-\Delta m$ and approximate velocity v - c. Hence the total change in momen-





tum during the time interval Δt is

$$\Delta P \approx (m \Delta v + v \Delta m + \Delta m \Delta v) + (-\Delta m)(v - c)$$

= $m \Delta v + c \Delta m + \Delta m \Delta v.$

Now we divide by Δt and take the limit as $\Delta t \rightarrow 0$, so $\Delta m \rightarrow 0$, assuming continuity of m(t). The substitution of the resulting expression for dP/dt in (1) yields the **rocket propulsion equation**

$$m\frac{dv}{dt} + c\frac{dm}{dt} = F.$$
 (2)

If $F = F_G + F_R$, where $F_G = -mg$ is a constant force of gravity and $F_R = -kv$ is a force of air resistance proportional to velocity, then Eq. (2) finally gives

$$m\frac{dv}{dt} + c\frac{dm}{dt} = -mg - kv.$$
(3)

Constant Thrust

Now suppose that the rocket fuel is consumed at the constant "burn rate" β during the time interval $[0, t_1]$, during which time the mass of the rocket decreases from m_0 to m_1 . Thus

$$m(0) = m_0, m(t_1) = m_1, (4)$$

$$m(t) = m_0 - \beta t, \frac{dm}{dt} = -\beta ext{ for } t \le t_1,$$

with burnout occurring at time $t = t_1$.

PROBLEM 1 Substitute the expressions in (4) into Eq. (3) to obtain the differential equation

$$(m_0 - \beta t)\frac{dv}{dt} + kv = \beta c - (m_0 - \beta t)g.$$
(5)

Solve this linear equation for

$$v(t) = v_0 M^{k/\beta} - \frac{g\beta t}{\beta - k} + \left(\frac{\beta c}{k} + \frac{gm_0}{\beta - k}\right) (1 - M^{k/\beta}),\tag{6}$$

where $v_0 = v(0)$ and

$$M = \frac{m(t)}{m_0} = \frac{m_0 - \beta t}{m_0}$$

denotes the rocket's **fractional mass** at time t.

No Resistance

PROBLEM 2 For the case of no air resistance, set k = 0 in Eq. (5) and integrate to obtain

$$v(t) = v_0 - gt + c \ln \frac{m_0}{m_0 - \beta t}.$$
(7)

Because $m_0 - \beta t_1 = m_1$, it follows that the velocity of the rocket at burnout $(t = t_1)$ is

$$v_1 = v(t_1) = v_0 - gt_1 + c \ln \frac{m_0}{m_1}.$$
(8)

PROBLEM 3 Start with Eq. (7) and integrate to obtain

$$v(t) = (v_0 + c)t - \frac{1}{2}gt^2 - \frac{c}{\beta}(m_0 - \beta t)\ln\frac{m_0}{m_0 - \beta t}.$$
(9)

It follows that the rocket's altitude at burnout is

$$y_1 = y(t_1) = (v_0 + c)t_1 - \frac{1}{2}gt_1^2 - \frac{cm_1}{\beta}\ln\frac{m_0}{m_1}.$$
 (10)

PROBLEM 4 The V-2 rocket that was used to attack London in World War II had an initial mass of 12,850 kg, of which 68.5% was fuel. This fuel burned uniformly for 70 seconds with an exhaust velocity of 2 km/s. Assume it encounters air resistance of 1.45 N per m/s of velocity. Then find the velocity and altitude of the V-2 at burnout under the assumption that it is launched vertically upward from rest on the ground.

PROBLEM 5 Actually, our basic differential equation in (3) applies without qualification only when the rocket is already in motion. However, when a rocket is sitting on its launch pad stand and its engines are turned on initially, it is observed that a certain time interval passes before the rocket actually "blasts off" and begins to ascend. The reason is that if v = 0 in (3), then the resulting initial acceleration

$$\frac{dv}{dt} = \frac{c}{m}\frac{dm}{dt} - g$$

of the rocket may be *negative*. But the rocket does not descend into the ground; it just "sits there" while (because *m* is decreasing) this calculated acceleration increases until it reaches 0 and (thereafter) positive values so the rocket can begin to ascend. With the notation introduced to describe the constant-thrust case, show that the rocket initially just "sits there" if the exhaust velocity *c* is less than m_0g/β , and that the time t_B which then elapses before actual blastoff is given by

$$t_B = \frac{m_0 g - \beta c}{\beta g}.$$

Free Space

Suppose finally that the rocket is accelerating in free space, where there is neither gravity nor resistance, so g = k = 0. With g = 0 in Eq. (8) we see that, as the mass of the rocket decreases from m_0 to m_1 , its increase in velocity is

$$\Delta v = v_1 - v_0 = c \ln \frac{m_0}{m_1}.$$
(11)

Note that Δv depends only on the exhaust gas speed *c* and the initial-to-final mass ratio m_0/m_1 , but does not depend on the burn rate β . For example, if the rocket blasts off from rest ($v_0 = 0$) and c = 5 km/s and $m_0/m_1 = 20$, then its velocity at burnout is $v_1 = 5 \ln 20 \approx 15$ km/s. Thus if a rocket initially consists predominantly of fuel, then it can attain velocities significantly greater than the (relative) velocity of its exhaust gases.

2.4 Numerical Approximation: Euler's Method

It is the exception rather than the rule when a differential equation of the general form

$$\frac{dy}{dx} = f(x, y)$$

can be solved exactly and explicitly by elementary methods like those discussed in Chapter 1. For example, consider the simple equation

$$\frac{dy}{dx} = e^{-x^2}.$$
(1)

A solution of Eq. (1) is simply an antiderivative of e^{-x^2} . But it is known that every antiderivative of $f(x) = e^{-x^2}$ is a **nonelementary** function—one that cannot be expressed as a finite combination of the familiar functions of elementary calculus. Hence no particular solution of Eq. (1) is finitely expressible in terms of elementary functions. Any attempt to use the symbolic techniques of Chapter 1 to find a simple explicit formula for a solution of (1) is therefore doomed to failure.

As a possible alternative, an old-fashioned computer plotter—one that uses an ink pen to draw curves mechanically—can be programmed to draw a solution curve that starts at the initial point (x_0, y_0) and attempts to thread its way through the slope field of a given differential equation y' = f(x, y). The procedure the plotter carries out can be described as follows.

- The plotter pen starts at the initial point (x_0, y_0) and moves a tiny distance along the slope segment though (x_0, y_0) . This takes it to the point (x_1, y_1) .
- At (x_1, y_1) the pen changes direction, and now moves a tiny distance along the slope segment through this new starting point (x_1, y_1) . This takes it to the next starting point (x_2, y_2) .
- At (x_2, y_2) the pen changes direction again, and now moves a tiny distance along the slope segment through (x_2, y_2) . This takes it to the next starting point (x_3, y_3) .

Figure 2.4.1 illustrates the result of continuing in this fashion—by a sequence of discrete straight-line steps from one starting point to the next. In this figure we see a polygonal curve consisting of line segments that connect the successive points $(x_0, y_0), (x_1, y_1), (x_2, y_2), (x_3, y_3), \ldots$ However, suppose that each "tiny distance" the pen travels along a slope segment—before the midcourse correction that sends it along a fresh new slope segment—is so very small that the naked eye cannot distinguish the individual line segments constituting the polygonal curve. Then the resulting polygonal curve looks like a smooth, continuously turning solution curve of the differential equation. Indeed, this is (in essence) how most of the solution curves shown in the figures of Chapter 1 were computer generated.

Leonhard Euler—the great 18th-century mathematician for whom so many mathematical concepts, formulas, methods, and results are named—did not have a computer plotter, and his idea was to do all this numerically rather than graphically. In order to approximate the solution of the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0,$$
 (2)

we first choose a fixed (horizontal) **step size** *h* to use in making each step from one point to the next. Suppose we've started at the initial point (x_0, y_0) and after *n* steps have reached the point (x_n, y_n) . Then the step from (x_n, y_n) to the next point (x_{n+1}, y_{n+1}) is illustrated in Fig. 2.4.2. The slope of the direction segment



FIGURE 2.4.1. The first few steps in approximating a solution curve.



FIGURE 2.4.2. The step from (x_n, y_n) to (x_{n+1}, y_{n+1}) .

through (x_n, y_n) is $m = f(x_n, y_n)$. Hence a horizontal change of h from x_n to x_{n+1} corresponds to a vertical change of $m \cdot h = h \cdot f(x_n, y_n)$ from y_n to y_{n+1} . Therefore the coordinates of the new point (x_{n+1}, y_{n+1}) are given in terms of the old coordinates by

$$x_{n+1} = x_n + h$$
, $y_{n+1} = y_n + h \cdot f(x_n, y_n)$.

Given the initial value problem in (2), **Euler's method** with step size h consists of starting with the initial point (x_0 , y_0) and applying the formulas

$x_1 = x_0 + h$	$y_1 = y_0 + h \cdot f(x_0, y_0)$
$x_2 = x_1 + h$	$y_2 = y_1 + h \cdot f(x_1, y_1)$
$x_3 = x_2 + h$	$y_3 = y_2 + h \cdot f(x_2, y_2)$
:	:
:	:
:	:

to calculate successive points (x_1, y_1) , (x_2, y_2) , (x_3, y_3) , ... on an approximate solution curve.

However, we ordinarily do not sketch the corresponding polygonal approximation. Instead, the numerical result of applying Euler's method is the sequence of *approximations*

$$y_1, y_2, y_3, \ldots, y_n, \ldots$$

to the true values

$$y(x_1), y(x_2), y(x_3), \dots, y(x_n), \dots$$

at the points $x_1, x_2, x_3, \ldots, x_n, \ldots$ of the *exact* (though unknown) solution y(x) of the initial value problem. These results typically are presented in the form of a table of approximate values of the desired solution.

ALGORITHM The Euler Method

Given the initial value problem

 $\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0,$ (2)

Euler's method with step size h consists of applying the iterative formula

$$y_{n+1} = y_n + h \cdot f(x_n, y_n) \quad (n \ge 0)$$
 (3)

to calculate successive approximations y_1, y_2, y_3, \ldots to the [true] values $y(x_1)$, $y(x_2), y(x_3), \ldots$ of the [exact] solution y = y(x) at the points x_1, x_2, x_3, \ldots , respectively.

The iterative formula in (3) tells us how to make the typical step from y_n to y_{n+1} and is the heart of Euler's method. Although the most important applications of Euler's method are to nonlinear equations, we first illustrate the method with a simple initial value problem whose exact solution is available, just for the purpose of comparison of approximate and actual solutions.

Example 1

Apply Euler's method to approximate the solution of the initial value problem

$$\frac{dy}{dx} = x + \frac{1}{5}y, \quad y(0) = -3,$$
(4)

- (a) first with step size h = 1 on the interval [0, 5],
- (b) then with step size h = 0.2 on the interval [0, 1].

Solution

(a) With $x_0 = 0$, $y_0 = -3$, $f(x, y) = x + \frac{1}{5}y$, and h = 1 the iterative formula in (3) yields the approximate values

$$y_{1} = y_{0} + h \cdot [x_{0} + \frac{1}{5}y_{0}] = (-3) + (1)[0 + \frac{1}{5}(-3)] = -3.6,$$

$$y_{2} = y_{1} + h \cdot [x_{1} + \frac{1}{5}y_{1}] = (-3.6) + (1)[1 + \frac{1}{5}(-3.6)] = -3.32,$$

$$y_{3} = y_{2} + h \cdot [x_{2} + \frac{1}{5}y_{2}] = (-3.32) + (1)[2 + \frac{1}{5}(-3.32)] = -1.984,$$

$$y_{4} = y_{3} + h \cdot [x_{3} + \frac{1}{5}y_{3}] = (-1.984) + (1)[3 + \frac{1}{5}(-1.984)] = 0.6192,$$
 and

$$y_{5} = y_{4} + h \cdot [x_{4} + \frac{1}{5}y_{4}] = (0.6912) + (1)[4 + \frac{1}{5}(0.6912)] \approx 4.7430$$

at the points $x_1 = 1$, $x_2 = 2$, $x_3 = 3$, $x_4 = 4$, and $x_5 = 5$. Note how the result of each calculation feeds into the next one. The resulting table of approximate values is

x	0	1	2	3	4	5
Approx. y	-3	-3.6	-3.32	-1.984	0.6912	4.7430

Figure 2.4.3 shows the graph of this approximation, together with the graphs of the Euler approximations obtained with step sizes h = 0.2 and 0.05, as well as the graph of the exact solution

$$y(x) = 22e^{x/5} - 5x - 25$$

that is readily found using the linear-equation technique of Section 1.5. We see that decreasing the step size increases the accuracy, but with any single approximation, the accuracy decreases with distance from the initial point.



FIGURE 2.4.3. Graphs of Euler approximations with step sizes h = 1, h = 0.2, and h = 0.05.

(b) Starting afresh with $x_0 = 0$, $y_0 = -3$, $f(x, y) = x + \frac{1}{5}y$, and h = 0.2, we get the approximate values

$$y_{1} = y_{0} + h \cdot [x_{0} + \frac{1}{5}y_{0}] = (-3) + (0.2)[0 + \frac{1}{5}(-3)] = -3.12,$$

$$y_{2} = y_{1} + h \cdot [x_{1} + \frac{1}{5}y_{1}] = (-3.12) + (0.2)[0.2 + \frac{1}{5}(-3.12)] \approx -3.205,$$

$$y_{3} = y_{2} + h \cdot [x_{2} + \frac{1}{5}y_{2}] \approx (-3.205) + (0.2)[0.4 + \frac{1}{5}(-3.205)] \approx -3.253,$$

$$y_{4} = y_{3} + h \cdot [x_{3} + \frac{1}{5}y_{3}] \approx (-3.253) + (0.2)[0.6 + \frac{1}{5}(-3.253)] \approx -3.263,$$

$$y_{5} = y_{4} + h \cdot [x_{4} + \frac{1}{5}y_{4}] \approx (-3.263) + (0.2)[0.8 + \frac{1}{5}(-3.263)] \approx -3.234$$

at the points $x_1 = 0.2$, $x_2 = 0.4$, $x_3 = 0.6$, $x_4 = 0.8$, and $x_5 = 1$. The resulting table of approximate values is

x	0	0.2	0.4	0.6	0.8	1
Approx. y	-3	-3.12	-3.205	-3.253	-3.263	-3.234

High accuracy with Euler's method usually requires a very small step size and hence a larger number of steps than can reasonably be carried out by hand. The application material for this section contains calculator and computer programs for automating Euler's method. One of these programs was used to calculate the table entries shown in Fig. 2.4.4. We see that 500 Euler steps (with step size h = 0.002) from x = 0 to x = 1 yield values that are accurate to within 0.001.

x	Approx y with $h = 0.2$	Approx y with $h = 0.02$	Approx y with $h = 0.002$	Actual value of y
0	-3.000	-3.000	-3.000	-3.000
0.2	-3.120	-3.104	-3.102	-3.102
0.4	-3.205	-3.172	-3.168	-3.168
0.6	-3.253	-3.201	-3.196	-3.195
0.8	-3.263	-3.191	-3.184	-3.183
1	-3.234	-3.140	-3.130	-3.129

FIGURE 2.4.4. Euler approximations with step sizes h = 0.2, h = 0.02, and h = 0.002.

Example 2 Falling baseball Suppose the baseball of Example 3 in Section 1.3 is simply dropped (instead of being thrown downward) from the helicopter. Then its velocity v(t) after t seconds satisfies the initial value problem

$$\frac{dv}{dt} = 32 - 0.16v, \quad v(0) = 0.$$
(5)

We use Euler's method with h = 1 to track the ball's increasing velocity at 1-second intervals for the first 10 seconds of fall. With $t_0 = 0$, $v_0 = 0$, F(t, v) = 32 - 0.16v, and h = 1 the iterative formula in (3) yields the approximate values

$$v_1 = v_0 + h \cdot [32 - 0.16v_0] = (0) + (1)[32 - 0.16(0)] = 32,$$

$$v_2 = v_1 + h \cdot [32 - 0.16v_1] = (32) + (1)[32 - 0.16(32)] = 58.88,$$

$$v_3 = v_2 + h \cdot [32 - 0.16v_2] = (58.88) + (1)[32 - 0.16(58.88)] \approx 81.46,$$

$$v_4 = v_3 + h \cdot [32 - 0.16v_3] = (81.46) + (1)[32 - 0.16(81.46)] \approx 100.43,$$
 and

$$v_5 = v_4 + h \cdot [32 - 0.16v_4] = (100.43) + (1)[32 - 0.16(100.43)] \approx 116.36.$$

Continuing in this fashion, we complete the h = 1 column of v-values shown in the table of Fig. 2.4.5—where we have rounded off velocity entries to the nearest foot per second. The values corresponding to h = 0.1 were calculated using a computer, and we see that they are accurate to within about 1 ft/s. Note also that after 10 seconds the falling ball has attained about 80% of its limiting velocity of 200 ft/s.

Local and Cumulative Errors

There are several sources of error in Euler's method that may make the approximation y_n to $y(x_n)$ unreliable for large values of n, those for which x_n is not sufficiently close to x_0 . The error in the linear approximation formula

$$y(x_{n+1}) \approx y_n + h \cdot f(x_n, y_n) = y_{n+1} \tag{6}$$

t	$\begin{array}{l} \text{Approx } v \\ \text{with } h = 1 \end{array}$	Approx v with $h = 0.1$	Actual value of v
1	32	30	30
2	59	55	55
3	81	77	76
4	100	95	95
5	116	111	110
6	130	124	123
7	141	135	135
8	150	145	144
9	158	153	153
10	165	160	160

FIGURE 2.4.5. Euler approximations in Example 2 with step sizes h = 1 and h = 0.1.

is the amount by which the tangent line at (x_n, y_n) departs from the solution curve through (x_n, y_n) , as illustrated in Fig. 2.4.6. This error, introduced at each step in the process, is called the **local error** in Euler's method.

The local error indicated in Fig. 2.4.6 would be the total error in y_{n+1} if the starting point y_n in (6) were an exact value, rather than merely an approximation to the actual value $y(x_n)$. But y_n itself suffers from the accumulated effects of all the local errors introduced at the previous steps. Thus the tangent line in Fig. 2.4.6 is tangent to the "wrong" solution curve—the one through (x_n, y_n) rather than the actual solution curve through the initial point (x_0, y_0) . Figure 2.4.7 illustrates this **cumulative error** in Euler's method; it is the amount by which the polygonal stepwise path from (x_0, y_0) departs from the actual solution curve through (x_0, y_0) .



FIGURE 2.4.7. The cumulative error in Euler's method.



FIGURE 2.4.6. The local error in Euler's method.

	y with	y with	y with	y with	Actual
x	h = 0.1	h = 0.02	h = 0.005	h = 0.001	у
0.1	1.1000	1.1082	1.1098	1.1102	1.1103
0.2	1.2200	1.2380	1.2416	1.2426	1.2428
0.3	1.3620	1.3917	1.3977	1.3993	1.3997
0.4	1.5282	1.5719	1.5807	1.5831	1.5836
0.5	1.7210	1.7812	1.7933	1.7966	1.7974
0.6	1.9431	2.0227	2.0388	2.0431	2.0442
0.7	2.1974	2.2998	2.3205	2.3261	2.3275
0.8	2.4872	2.6161	2.6422	2.6493	2.6511
0.9	2.8159	2.9757	3.0082	3.0170	3.0192
1.0	3.1875	3.3832	3.4230	3.4338	3.4366

FIGURE 2.4.8. Approximating the solution of dy/dx = x + y, y(0) = 1 with successively smaller step sizes.

The usual way of attempting to reduce the cumulative error in Euler's method is to decrease the step size *h*. The table in Fig. 2.4.8 shows the results obtained in approximating the exact solution $y(x) = 2e^x - x - 1$ of the initial value problem

$$\frac{dy}{dx} = x + y, \quad y(0) = 1,$$

using the successively smaller step sizes h = 0.1, h = 0.02, h = 0.005, and h = 0.001. We show computed values only at intervals of $\Delta x = 0.1$. For instance, with h = 0.001, the computation required 1000 Euler steps, but the value y_n is shown only when n is a multiple of 100, so that x_n is an integral multiple of 0.1.

By scanning the columns in Fig. 2.4.8 we observe that, for each fixed step size *h*, the error $y_{actual} - y_{approx}$ increases as *x* gets farther from the starting point $x_0 = 0$. But by scanning the rows of the table we see that for each fixed *x*, the error decreases as the step size *h* is reduced. The percentage errors at the final point x = 1 range from 7.25% with h = 0.1 down to only 0.08% with h = 0.001. Thus the smaller the step size, the more slowly does the error grow with increasing distance from the starting point.

The column of data for h = 0.1 in Fig. 2.4.8 requires only 10 steps, so Euler's method can be carried out with a hand-held calculator. But 50 steps are required to reach x = 1 with h = 0.02, 200 steps with h = 0.005, and 1000 steps with h = 0.001. A computer is almost always used to implement Euler's method when more than 10 or 20 steps are required. Once an appropriate computer program has been written, one step size is—in principle—just as convenient as another; after all, the computer hardly cares how many steps it is asked to carry out.

Why, then, do we not simply choose an exceedingly small step size (such as $h = 10^{-12}$), with the expectation that very great accuracy will result? There are two reasons for not doing so. The first is obvious: the time required for the computation. For example, the data in Fig. 2.4.8 were obtained using a hand-held calculator that carried out nine Euler steps per second. Thus it required slightly over one second to approximate y(1) with h = 0.1 and about 1 min 50 s with h = 0.001. But with $h = 10^{-12}$ it would require over 3000 years!

The second reason is more subtle. In addition to the local and cumulative errors discussed previously, the computer itself will contribute **roundoff error** at

each stage because only finitely many significant digits can be used in each calculation. An Euler's method computation with h = 0.0001 will introduce roundoff errors 1000 times as often as one with h = 0.1. Hence with certain differential equations, h = 0.1 might actually produce more accurate results than those obtained with h = 0.0001, because the cumulative effect of roundoff error in the latter case might exceed combined cumulative and roundoff error in the case h = 0.1.

The "best" choice of h is difficult to determine in practice as well as in theory. It depends on the nature of the function f(x, y) in the initial value problem in (2), on the exact code in which the program is written, and on the specific computer used. With a step size that is too large, the approximations inherent in Euler's method may not be sufficiently accurate, whereas if h is too small, then roundoff errors may accumulate to an unacceptable degree or the program may require too much time to be practical. The subject of *error propagation* in numerical algorithms is treated in numerical analysis courses and textbooks.

The computations in Fig. 2.4.8 illustrate the common strategy of applying a numerical algorithm, such as Euler's method, several times in succession, beginning with a selected number n of subintervals for the first application, then doubling n for each succeeding application of the method. Visual comparison of successive results often can provide an "intuitive feel" for their accuracy. In the next two examples we present graphically the results of successive applications of Euler's method.

Example 3 Approximate logistic solution The exact solution of the logistic initial value problem

$$\frac{dy}{dx} = \frac{1}{3}y(8-y), \quad y(0) = 1$$

is $y(x) = 8/(1 + 7e^{-8x/3})$. Figure 2.4.9 shows both the exact solution curve and approximate solution curves obtained by applying Euler's method on the interval $0 \le x \le 5$ with n = 5, n = 10, and n = 20 subintervals. Each of these "curves" actually consists of line segments joining successive points (x_n, y_n) and (x_{n+1}, y_{n+1}) . The Euler approximation with 5 subintervals is poor, and the approximation with 10 subintervals also overshoots the limiting value y = 8 of the solution before leveling off, but with 20 subintervals we obtain fairly good qualitative agreement with the actual behavior of the solution.





FIGURE 2.4.9. Approximating a logistic solution using Euler's method with n = 5, n = 10, and n = 20 subintervals.

FIGURE 2.4.10. Approximating the exact solution $y = e^{\sin x}$ using Euler's method with 50, 100, 200, and 400 subintervals.

Example 4

The exact solution of the initial value problem

$$\frac{dy}{dx} = y\cos x, \quad y(0) = 1$$

is the periodic function $y(x) = e^{\sin x}$. Figure 2.4.10 shows both the exact solution curve and approximate solution curves obtained by applying Euler's method on the interval $0 \le x \le 6\pi$

with n = 50, n = 100, n = 200, and n = 400 subintervals. Even with this many subintervals, Euler's method evidently has considerable difficulty keeping up with the oscillations in the actual solution. Consequently, the more accurate methods discussed in succeeding sections are needed for serious numerical investigations.

A Word of Caution

The data shown in Fig. 2.4.8 indicate that Euler's method works well in approximating the solution of dy/dx = x + y, y(0) = 1 on the interval [0, 1]. That is, for each fixed x it appears that the approximate values approach the actual value of y(x) as the step size h is decreased. For instance, the approximate values in the rows corresponding to x = 0.3 and x = 0.5 suggest that $y(0.3) \approx 1.40$ and $y(0.5) \approx 1.80$, in accord with the actual values shown in the final column of the table.

Example 5, in contrast, shows that some initial value problems are not so well behaved.

Example 5 Cautionary example Use Euler's method to approximate the solution of the initial value problem

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0) = 1 \tag{7}$$

on the interval [0, 1].

Solution Here $f(x, y) = x^2 + y^2$, so the iterative formula of Euler's method is

$$y_{n+1} = y_n + h \cdot (x_n^2 + y_n^2).$$
(8)

With step size h = 0.1 we obtain

$$y_1 = 1 + (0.1) \cdot [(0)^2 + (1)^2] = 1.1,$$

$$y_2 = 1.1 + (0.1) \cdot [(0.1)^2 + (1.1)^2] = 1.222,$$

$$y_3 = 1.222 + (0.1) \cdot [(0.2)^2 + (1.222)^2] \approx 1.3753,$$

and so forth. Rounded to four decimal places, the first ten values obtained in this manner are

$y_1 = 1.1000$	$y_6 = 2.1995$
$y_2 = 1.2220$	$y_7 = 2.7193$
$y_3 = 1.3753$	$y_8 = 3.5078$
$y_4 = 1.5735$	$y_9 = 4.8023$
$y_5 = 1.8371$	$y_{10} = 7.1895$

But instead of naively accepting these results as accurate approximations, we decided to use a computer to repeat the computations with smaller values of h. The table in Fig. 2.4.11 shows the results obtained with step sizes h = 0.1, h = 0.02, and h = 0.005. Observe that now the "stability" of the procedure in Example 1 is missing. Indeed, it seems obvious that something is going wrong near x = 1.

Figure 2.4.12 provides a graphical clue to the difficulty. It shows a slope field for $dy/dx = x^2 + y^2$, together with a solution curve through (0, 1) plotted using one of the more accurate approximation methods of the following two sections. It appears that this solution curve may have a vertical asymptote near x = 0.97. Indeed, an exact solution using Bessel functions (see Problem 34 in Section 11.4) can be used to show that $y(x) \to +\infty$ as $x \to 0.969811$ (approximately). Although Euler's method gives values (albeit spurious ones) at x = 1, the actual solution does not exist on the entire interval [0, 1]. Moreover, Euler's method is unable to "keep up" with the rapid changes in y(x) that occur as x approaches the infinite discontinuity near 0.969811.

The moral of Example 5 is that there are pitfalls in the numerical solution of certain initial value problems. Certainly it's pointless to attempt to approximate a solution on an interval where it doesn't even exist (or where it is not unique, in which

	y with	y with	y with
x	h = 0.1	h = 0.02	h = 0.005
0.1	1.1000	1.1088	1.1108
0.2	1.2220	1.2458	1.2512
0.3	1.3753	1.4243	1.4357
0.4	1.5735	1.6658	1.6882
0.5	1.8371	2.0074	2.0512
0.6	2.1995	2.5201	2.6104
0.7	2.7193	3.3612	3.5706
0.8	3.5078	4.9601	5.5763
0.9	4.8023	9.0000	12.2061
1.0	7.1895	30.9167	1502.2090



FIGURE 2.4.12. Solution of $dy/dx = x^2 + y^2$, y(0) = 1.

FIGURE 2.4.11. Attempting to approximate the solution of $dy/dx = x^2 + y^2$, y(0) = 1.

case there's no general way to predict which way the numerical approximations will branch at a point of nonuniqueness). One should never accept as accurate the results of applying Euler's method with a single fixed step size h. A second "run" with smaller step size (h/2, say, or h/5, or h/10) may give seemingly consistent results, thereby suggesting their accuracy, or it may—as in Example 5—reveal the presence of some hidden difficulty in the problem. Many problems simply require the more accurate and powerful methods that are discussed in the final two sections of this chapter.

2.4 Problems

In Problems 1 through 10, an initial value problem and its exact solution y(x) are given. Apply Euler's method twice to approximate to this solution on the interval $[0, \frac{1}{2}]$, first with step size h = 0.25, then with step size h = 0.1. Compare the three-decimal-place values of the two approximations at $x = \frac{1}{2}$ with the value $y(\frac{1}{2})$ of the actual solution.

1.
$$y' = -y$$
, $y(0) = 2$; $y(x) = 2e^{-x}$
2. $y' = 2y$, $y(0) = \frac{1}{2}$; $y(x) = \frac{1}{2}e^{2x}$
3. $y' = y + 1$, $y(0) = 1$; $y(x) = 2e^{x} - 1$
4. $y' = x - y$, $y(0) = 1$; $y(x) = 2e^{-x} + x - 1$
5. $y' = y - x - 1$, $y(0) = 1$; $y(x) = 2 + x - e^{x}$
6. $y' = -2xy$, $y(0) = 2$; $y(x) = 2e^{-x^{2}}$
7. $y' = -3x^{2}y$, $y(0) = 3$; $y(x) = 3e^{-x^{3}}$
8. $y' = e^{-y}$, $y(0) = 0$; $y(x) = \ln(x + 1)$
9. $y' = \frac{1}{4}(1 + y^{2})$, $y(0) = 1$; $y(x) = \tan \frac{1}{4}(x + \pi)$
10. $y' = 2xy^{2}$, $y(0) = 1$; $y(x) = \frac{1}{1 - x^{2}}$

Note: The application following this problem set lists illustrative calculator/computer programs that can be used in the remaining problems.

A programmable calculator or a computer will be useful for Problems 11 through 16. In each problem find the exact solution of the given initial value problem. Then apply Euler's method twice to approximate (to four decimal places) this solution on the given interval, first with step size h = 0.01, then with step size h = 0.005. Make a table showing the approximate values and the actual value, together with the percentage error in the more accurate approximation, for x an integral multiple of 0.2. Throughout, primes denote derivatives with respect to x.

11.
$$y' = y - 2$$
, $y(0) = 1$; $0 \le x \le 1$
12. $y' = \frac{1}{2}(y - 1)^2$, $y(0) = 2$; $0 \le x \le 1$
13. $yy' = 2x^3$, $y(1) = 3$; $1 \le x \le 2$
14. $xy' = y^2$, $y(1) = 1$; $1 \le x \le 2$
15. $xy' = 3x - 2y$, $y(2) = 3$; $2 \le x \le 3$
16. $y^2y' = 2x^5$, $y(2) = 3$; $2 \le x \le 3$

A computer with a printer is required for Problems 17 through 24. In these initial value problems, use Euler's method with step sizes h = 0.1, 0.02, 0.004, and 0.0008 to approximate to four decimal places the values of the solution at ten equally spaced points of the given interval. Print the results in tabular form with appropriate headings to make it easy to gauge the effect of varying the step size h. Throughout, primes denote derivatives with respect to x.

17.
$$y' = x^2 + y^2$$
, $y(0) = 0$; $0 \le x \le 1$
18. $y' = x^2 - y^2$, $y(0) = 1$; $0 \le x \le 2$

- **19.** $y' = x + \sqrt{y}, y(0) = 1; 0 \le x \le 2$
- **20.** $y' = x + \sqrt[3]{y}, y(0) = -1; 0 \le x \le 2$
- **21.** $y' = \ln y$, y(1) = 2; $1 \le x \le 2$ **22.** $y' = x^{2/3} + y^{2/3}$, y(0) = 1; $0 \le x \le 2$

- **23.** $y' = \sin x + \cos y, y(0) = 0; 0 \le x \le 1$ **24.** $y' = \frac{x}{1+y^2}, y(-1) = 1; -1 \le x \le 1$
- 25. Falling parachutist You bail out of the helicopter of Example 2 and immediately pull the ripcord of your parachute. Now k = 1.6 in Eq. (5), so your downward velocity satisfies the initial value problem

$$\frac{dv}{dt} = 32 - 1.6v, \quad v(0) = 0$$

(with t in seconds and v in ft/sec). Use Euler's method with a programmable calculator or computer to approximate the solution for $0 \le t \le 2$, first with step size h = 0.01and then with h = 0.005, rounding off approximate vvalues to one decimal place. What percentage of the limiting velocity 20 ft/sec has been attained after 1 second? After 2 seconds?

26. Deer population Suppose the deer population P(t) in a small forest initially numbers 25 and satisfies the logistic equation

$$\frac{dP}{dt} = 0.0225P - 0.0003P^2$$

(with t in months). Use Euler's method with a programmable calculator or computer to approximate the solution for 10 years, first with step size h = 1 and then with h = 0.5, rounding off approximate *P*-values to integral numbers of deer. What percentage of the limiting population of 75 deer has been attained after 5 years? After 10 years?

Use Euler's method with a computer system to find the desired solution values in Problems 27 and 28. Start with step size h = 0.1, and then use successively smaller step sizes until successive approximate solution values at x = 2 agree rounded off to two decimal places.

27. $y' = x^2 + y^2 - 1$, y(0) = 0; y(2) = ?**28.** $y' = x + \frac{1}{2}y^2$, y(-2) = 0; y(2) = ?

Problems 29 through 31 illustrate the unreliability of Euler's method near a discontinuity of the solution.

29. Consider the initial value problem

$$7x\frac{dy}{dx} + y = 0, \quad y(-1) = 1.$$

(a) Solve this problem for the exact solution

$$y(x) = -\frac{1}{x^{1/7}}$$

which has an infinite discontinuity at x = 0. (b) Apply Euler's method with step size h = 0.15 to approximate this solution on the interval $-1 \leq x \leq 0.5$. Note that, from these data alone, you might not suspect any difficulty near x = 0. The reason is that the numerical approximation "jumps across the discontinuity" to another solution of 7xy' + y = 0 for x > 0. (c) Finally, apply Euler's method with step sizes h = 0.03 and h = 0.006, but still printing results only at the original points x = -1.00, -0.85, -0.70, ..., 1.20, 1.35. and 1.50. Would you now suspect a discontinuity in the exact solution?

30. Apply Euler's method with successively smaller step sizes on the interval [0, 2] to verify empirically that the solution of the initial value problem

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0) = 0$$

has a vertical asymptote near x = 2.003147. (Contrast this with Example 2, in which y(0) = 1.)

31. The general solution of the equation

$$\frac{dy}{dx} = (1+y^2)\cos x$$

is $y(x) = \tan(C + \sin x)$. With the initial condition y(0) =0 the solution $y(x) = \tan(\sin x)$ is well behaved. But with y(0) = 1 the solution $y(x) = \tan\left(\frac{1}{4}\pi + \sin x\right)$ has a vertical asymptote at $x = \sin^{-1}(\pi/4) \approx 0.90334$. Use Euler's method to verify this fact empirically.

2.4 Application Implementing Euler's Method

Go to goo.gl/Dd91hc to download this application's computing resources including Maple/Mathematica/MATLAB/ Python/TI-Nspire.

Construction of a calculator or computer program to implement a numerical algorithm can sharpen one's understanding of the algorithm. Figure 2.4.13 lists TI-84 Plus and Python programs implementing Euler's method to approximate the solution of the initial value problem

$$\frac{dy}{dx} = x + y, \quad y(0) = 1$$

considered in this section. The comments provided in the final column should make these programs intelligible even if you have little familiarity with the Python and TI calculator programming languages. Python—freely available on the Internet—is

TI-84 Plus	Python	Comment
PROGRAM: EULER	# Program EULER	Program title
:10 \rightarrow N	N = 10	Number of steps
:0→X	X = 0.0	Initial x
:1→Y	Y = 1.0	Initial y
:1→T	x1 = 1.0	Final x
: (T-X) /N \rightarrow H	H = (X1-X)/N	Step size
:For(I,1,N)	for I in range(N):	Begin loop
:X+Y→F	F = X + Y	Function value
:Y+H*F→Y	Y = Y + H * F	Euler iteration
:X+H→X	X = X + H	New x
:Disp X,Y	print (X,Y)	Display results
:End	# END	End loop

FIGURE 2.4.13. TI-84 Plus and Python Euler's method programs.

a general-purpose computing language widely used in education and industry. In particular, its many add-on packages make Python a powerful scientific computing platform, with numerical, graphic, and symbolic capabilities approaching those of commercial systems such as MATLAB, *Mathematica*, and *Maple*. Further, Python code was designed with an emphasis on readability, making it well-suited to expressing basic mathematical algorithms.

To increase the number of steps (and thereby decrease the step size) you need only change the value of **N** specified in the first line of the program. To apply Euler's method to a different equation dy/dx = f(x, y), you need change only the single line that calculates the function value **F**.

Any other procedural programming language (such as FORTRAN or C^{++}) would follow the pattern illustrated by the parallel lines of TI-84 Plus and Python code in Fig. 2.4.13. Some of the modern functional programming languages mirror standard mathematical notation even more closely. Figure 2.4.14 shows a MATLAB implementation of Euler's method. The **euler** function takes as input the initial value **x**, the initial value **y**, the final value **x1** of x, and the desired number **n** of subintervals.

```
function yp = f(x, y)
yp = x + y;
                           % yp = y'
function [X,Y] = euler(x,y,x1,n)
h = (x1 - x)/n;
                           % step size
X = x;
                           % initial x
Y = Y;
                           % initial y
for i = 1:n
                           % begin loop
                           % Euler iteration
    y = y + h*f(x,y);
                           % new x
    x = x + h;
    X = [X; x];
                           % update x-column
                           % update y-column
    Y = [Y; y];
                           % end loop
    end
```

For instance, the MATLAB command

[X, Y] = euler(0, 1, 1, 10)

then generates the x_n - and y_n -data shown in the first two columns of the table of Fig. 2.4.8.

You should begin this project by implementing Euler's method with your own calculator or computer system. Test your program by first applying it to the initial value problem in Example 1, then to some of the problems for this section.

Famous Numbers Investigation

The following problems describe the numbers $e \approx 2.71828$, $\ln 2 \approx 0.69315$, and $\pi \approx 3.14159$ as specific values of solutions of certain initial value problems. In each case, apply Euler's method with $n = 50, 100, 200, \ldots$ subintervals (doubling *n* each time). How many subintervals are needed to obtain—twice in succession—the correct value of the target number rounded to three decimal places?

- 1. The number e = y(1), where y(x) is the solution of the initial value problem dy/dx = y, y(0) = 1.
- 2. The number $\ln 2 = y(2)$, where y(x) is the solution of the initial value problem dy/dx = 1/x, y(1) = 0.
- 3. The number $\pi = y(1)$, where y(x) is the solution of the initial value problem $dy/dx = 4/(1 + x^2)$, y(0) = 0.

Also explain in each problem what the point is—why the indicated famous number is the expected numerical result.

2.5 A Closer Look at the Euler Method

The Euler method as presented in Section 2.4 is not often used in practice, mainly because more accurate methods are available. But Euler's method has the advantage of simplicity, and a careful study of this method yields insights into the workings of more accurate methods, because many of the latter are extensions or refinements of the Euler method.

To compare two different methods of numerical approximation, we need some way to measure the accuracy of each. Theorem 1 tells what degree of accuracy we can expect when we use Euler's method.

THEOREM 1 The Error in the Euler Method

Suppose that the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0$$
 (1)

has a unique solution y(x) on the closed interval [a, b] with $a = x_0$, and assume that y(x) has a continuous second derivative on [a, b]. (This would follow from the assumption that f, f_x , and f_y are all continuous for $a \le x \le b$ and $c \le y \le d$, where $c \le y(x) \le d$ for all x in [a, b].) Then there exists a constant C such that the following is true: If the approximations $y_1, y_2, y_3, \ldots, y_k$ to the actual values $y(x_1), y(x_2), y(x_3), \ldots, y(x_k)$ at points of [a, b] are computed using Euler's method with step size h > 0, then

$$|y_n - y(x_n)| \le Ch \tag{2}$$

for each n = 1, 2, 3, ..., k.

Remark The error

 $y_{\text{actual}} - y_{\text{approx}} = y(x_n) - y_n$

in (2) denotes the [cumulative] error in Euler's method after n steps in the approximation, *exclusive* of roundoff error (as though we were using a perfect machine that made no roundoff errors). The theorem can be summarized by saying that *the error in Euler's method is of order* h; that is, the error is bounded by a [predetermined] constant C multiplied by the step size h. It follows, for instance, that (on a given closed interval) halving the step size cuts the maximum error in half; similarly, with step size h/10 we get 10 times the accuracy (that is, 1/10 the maximum error) as with step size h. Consequently, we can—in principle—get any degree of accuracy we want by choosing h sufficiently small.

We will omit the proof of this theorem, but one can be found in Chapter 7 of G. Birkhoff and G.-C. Rota, *Ordinary Differential Equations*, 4th ed. (New York: John Wiley, 1989). The constant C deserves some comment. Because C tends to increase as the maximum value of |y''(x)| on [a, b] increases, it follows that C must depend in a fairly complicated way on y, and actual computation of a value of C such that the inequality in (2) holds is usually impractical. In practice, the following type of procedure is commonly employed.

- 1. Apply Euler's method to the initial value problem in (1) with a reasonable value of h.
- 2. Repeat with h/2, h/4, and so forth, at each stage halving the step size for the next application of Euler's method.
- **3.** Continue until the results obtained at one stage agree—to an appropriate number of significant digits—with those obtained at the previous stage. Then the approximate values obtained at this stage are considered likely to be accurate to the indicated number of significant digits.

Example 1 Carry out this procedure with the initial value problem

$$\frac{dy}{dx} = -\frac{2xy}{1+x^2}, \quad y(0) = 1 \tag{3}$$

to approximate accurately the value y(1) of the solution at x = 1.

Solution

Using an Euler method program, perhaps one of those listed in Figs. 2.4.13 and 2.4.14, we begin with a step size h = 0.04 requiring n = 25 steps to reach x = 1. The table in Fig. 2.5.1 shows the approximate values of y(1) obtained with successively smaller values of h. The data suggest that the true value of y(1) is exactly 0.5. Indeed, the exact solution of the initial value problem in (3) is $y(x) = 1/(1 + x^2)$, so the true value of y(1) is exactly $\frac{1}{2}$.

h	Approximate y(1)	Actual y(1)	Error /h
0.04	0.50451	0.50000	0.11
0.02	0.50220	0.50000	0.11
0.01	0.50109	0.50000	0.11
0.005	0.50054	0.50000	0.11
0.0025	0.50027	0.50000	0.11
0.00125	0.50013	0.50000	0.10
0.000625	0.50007	0.50000	0.11
0.0003125	0.50003	0.50000	0.10

FIGURE 2.5.1. Table of values in Example 1.

The final column of the table in Fig. 2.5.1 displays the ratio of the magnitude of the error to *h*; that is, $|y_{actual} - y_{approx}|/h$. Observe how the data in this column



FIGURE 2.5.2. True and predicted values in Euler's method.

>

>

substantiate Theorem 1—in this computation, the error bound in (2) appears to hold with a value of C slightly larger than 0.1.

An Improvement in Euler's Method

As Fig. 2.5.2 shows, Euler's method is rather unsymmetrical. It uses the predicted slope $k = f(x_n, y_n)$ of the graph of the solution at the left-hand endpoint of the interval $[x_n, x_n + h]$ as if it were the actual slope of the solution over that entire interval. We now turn our attention to a way in which increased accuracy can easily be obtained; it is known as the *improved Euler method*.

Given the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0,$$
 (4)

suppose that after carrying out *n* steps with step size *h* we have computed the approximation y_n to the actual value $y(x_n)$ of the solution at $x_n = x_0 + nh$. We can use the Euler method to obtain a first estimate—which we now call u_{n+1} rather than y_{n+1} —of the value of the solution at $x_{n+1} = x_n + h$. Thus

$$u_{n+1} = y_n + h \cdot f(x_n, y_n) = y_n + h \cdot k_1.$$

Now that $u_{n+1} \approx y(x_{n+1})$ has been computed, we can take

$$k_2 = f(x_{n+1}, u_{n+1})$$

as a second estimate of the slope of the solution curve y = y(x) at $x = x_{n+1}$.

Of course, the approximate slope $k_1 = f(x_n, y_n)$ at $x = x_n$ has already been calculated. Why not *average* these two slopes to obtain a more accurate estimate of the average slope of the solution curve over the entire subinterval $[x_n, x_{n+1}]$? This idea is the essence of the *improved* Euler method. Figure 2.5.3 shows the geometry behind this method.

ALGORITHM The Improved Euler Method

Given the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0,$$

the **improved Euler method with step size** h consists in applying the iterative formulas

$$k_{1} = f(x_{n}, y_{n}),$$

$$u_{n+1} = y_{n} + h \cdot k_{1},$$

$$k_{2} = f(x_{n+1}, u_{n+1}),$$

$$y_{n+1} = y_{n} + h \cdot \frac{1}{2}(k_{1} + k_{2})$$
(5)

to compute successive approximations $y_1, y_2, y_3,...$ to the [true] values $y(x_1)$, $y(x_2), y(x_3),...$ of the [exact] solution y = y(x) at the points $x_1, x_2, x_3,...$, respectively.



FIGURE 2.5.3. The improved Euler method: Average the slopes of the tangent lines at (x_n, y_n) and (x_{n+1}, u_{n+1}) .

Remark The final formula in (5) takes the "Euler form"

$$y_{n+1} = y_n + h \cdot k$$

if we write

$$k = \frac{k_1 + k_2}{2}$$

for the approximate *average slope* on the interval $[x_n, x_{n+1}]$.

The improved Euler method is one of a class of numerical techniques known as **predictor-corrector** methods. First a predictor u_{n+1} of the next *y*-value is computed; then it is used to correct itself. Thus the **improved Euler method** with step size *h* consists of using the **predictor**

$$u_{n+1} = y_n + h \cdot f(x_n, y_n) \tag{6}$$

and the corrector

$$y_{n+1} = y_n + h \cdot \frac{1}{2} \left[f(x_n, y_n) + f(x_{n+1}, u_{n+1}) \right]$$
(7)

iteratively to compute successive approximations $y_1, y_2, y_2, ...$ to the values $y(x_1)$, $y(x_2), y(x_3), ...$ of the actual solution of the initial value problem in (4).

Remark Each improved Euler step requires two evaluations of the function f(x, y), as compared with the single function evaluation required for an ordinary Euler step. We naturally wonder whether this doubled computational labor is worth the trouble.

Answer Under the assumption that the exact solution y = y(x) of the initial value problem in (4) has a continuous third derivative, it can be proved—see Chapter 7 of Birkhoff and Rota—that the error in the improved Euler method is of order h^2 . This means that on a given bounded interval [a, b], each approximate value y_n satisfies the inequality

$$|y(x_n) - y_n| \le Ch^2,\tag{8}$$

where the constant C does not depend on h. Because h^2 is much smaller than h if h itself is small, this means that the improved Euler method is more accurate than Euler's method itself.

This advantage is offset by the fact that about twice as many computations are required. But the factor h^2 in (8) means that halving the step size results in 1/4 the maximum error, and with step size h/10 we get 100 times the accuracy (that is, 1/100 the maximum error) as with step size h.

Example 2 Figure 2.4.8 shows results of applying Euler's method to the initial value problem

$$\frac{dy}{dx} = x + y, \quad y(0) = 1 \tag{9}$$

with exact solution $y(x) = 2e^x - x - 1$. With f(x, y) = x + y in Eqs. (6) and (7), the predictor-corrector formulas for the improved Euler method are

$$u_{n+1} = y_n + h \cdot (x_n + y_n),$$

$$y_{n+1} = y_n + h \cdot \frac{1}{2} [(x_n + y_n) + (x_{n+1} + u_{n+1})].$$

With step size h = 0.1 we calculate

$$u_1 = 1 + (0.1) \cdot (0+1) = 1.1,$$

$$y_1 = 1 + (0.05) \cdot [(0+1) + (0.1+1.1)] = 1.11,$$

$$u_2 = 1.11 + (0.1) \cdot (0.1+1.11) = 1.231,$$

$$y_2 = 1.11 + (0.05) \cdot [(0.1+1.11) + (0.2+1.231)] = 1.24205$$

and so forth. The table in Fig. 2.5.4 compares the results obtained using the improved Euler method with those obtained previously using the "unimproved" Euler method. When the same step size h = 0.1 is used, the error in the Euler approximation to y(1) is 7.25%, but the error in the improved Euler approximation is only 0.24%.

	Euler Method, k = 0.1	Euler Method, k = 0.005	Improved Euler, h = 0.1	Actual
x	Values of y	Values of y	Values of y	y
0.1	1.1000	1.1098	1.1100	1.1103
0.2	1.2200	1.2416	1.2421	1.2428
0.3	1.3620	1.3977	1.3985	1.3997
0.4	1.5282	1.5807	1.5818	1.5836
0.5	1.7210	1.7933	1.7949	1.7974
0.6	1.9431	2.0388	2.0409	2.0442
0.7	2.1974	2.3205	2.3231	2.3275
0.8	2.4872	2.6422	2.6456	2.6511
0.9	2.8159	3.0082	3.0124	3.0192
1.0	3.1875	3.4230	3.4282	3.4366

x	Improved Euler, Approximate y	Actual y	
0.0	1.00000	1.00000	
0.1	1.11034	1.11034	
0.2	1.24280	1.24281	
0.3	1.39971	1.39972	
0.4	1.58364	1.58365	
0.5	1.79744	1.79744	
0.6	2.04423	2.04424	
0.7	2.32749	2.32751	
0.8	2.65107	2.65108	
0.9	3.01919	3.01921	
1.0	3 4 3 6 5 4	3 43656	

FIGURE 2.5.5. Improved Euler approximation to the solution of Eq. (9) with step size h = 0.005.

FIGURE 2.5.4. Euler and improved Euler approximations to the solution of dy/dx = x + y, y(0) = 1.

Indeed, the improved Euler method with h = 0.1 is more accurate (in this example) than the original Euler method with h = 0.005. The latter requires 200 evaluations of the function f(x, y), but the former requires only 20 such evaluations, so in this case the improved Euler method yields greater accuracy with only about one-tenth the work.

Figure 2.5.5 shows the results obtained when the improved Euler method is applied to the initial value problem in (9) using step size h = 0.005. Accuracy of five significant figures is apparent in the table. This suggests that, in contrast with the original Euler method, the improved Euler method is sufficiently accurate for certain practical applications—such as plotting solution curves.

	Improved Euler Approximation		
h	to y(1)	Error	$ \text{Error} /h^2$
0.04	0.500195903	-0.000195903	0.12
0.02	0.500049494	-0.000049494	0.12
0.01	0.500012437	-0.000012437	0.12
0.005	0.500003117	-0.000003117	0.12
0.0025	0.500000780	-0.00000780	0.12
0.00125	0.500000195	-0.000000195	0.12
0.000625	0.500000049	-0.00000049	0.12
0.0003125	0.500000012	-0.000000012	0.12

FIGURE 2.5.6. Improved Euler approximation to y(1) for $dy/dx = -2xy/(1 + x^2)$, y(0) = 1.

An improved Euler program (similar to the ones listed in the project material for this section) was used to compute approximations to the exact value y(1) = 0.5 of the solution $y(x) = 1/(1 + x^2)$ of the initial value problem

$$\frac{dy}{dx} = -\frac{2xy}{1+x^2}, \quad y(0) = 1 \tag{3}$$

of Example 1. The results obtained by successively halving the step size appear in the table in Fig. 2.5.6. Note that the final column of this table impressively corroborates the form of the error bound in (8), and that each halving of the step size reduces the error by a factor of almost exactly 4, as should happen if the error is proportional to h^2 .

In the following two examples we exhibit graphical results obtained by employing this strategy of successively halving the step size, and thus doubling the number of subintervals of a fixed interval on which we are approximating a solution.

Example 3

$$\frac{dy}{dx} = \frac{1}{3}y(8-y), \quad y(0) = 1.$$

In Example 3 of Section 2.4 we applied Euler's method to the logistic initial value problem

Figure 2.4.9 shows an obvious difference between the exact solution $y(x) = 8/(1 + 7e^{-8x/3})$ and the Euler approximation on $0 \le x \le 5$ using n = 20 subintervals. Figure 2.5.7 shows approximate solution curves plotted using the improved Euler's method.

The approximation with five subintervals is still bad—perhaps worse! It appears to level off considerably short of the actual limiting population M = 8. You should carry out at least the first two improved Euler steps manually to see for yourself how it happens that, after increasing appropriately during the first step, the approximate solution *decreases* in the second step rather than continuing to increase (as it should). In the project for this section we ask you to show empirically that the improved Euler approximate solution with step size h = 1 levels off at $y \approx 4.3542$.

In contrast, the approximate solution curve with n = 20 subintervals tracks the exact solution curve rather closely, and with n = 40 subintervals the exact and approximate solution curves are indistinguishable in Fig. 2.5.7. The table in Fig. 2.5.8 indicates that the improved Euler approximation with n = 200 subintervals is accurate rounded to three decimal places (that is, four significant digits) on the interval $0 \le x \le 5$. Because discrepancies in the fourth significant digit are not visually apparent at the resolution of an ordinary computer screen, the improved Euler method (using several hundred subintervals) is considered adequate for many graphical purposes.



FIGURE 2.5.7. Approximating a logistic solution using the improved Euler method with n = 5, n = 10, n = 20, and n = 40 subintervals.

x	Actual $y(x)$	Improved Euler with $n = 200$
0	1.0000	1.0000
1	5.3822	5.3809
2	7.7385	7.7379
3	7.9813	7.9812
4	7.9987	7.9987
5	7.9999	7.9999

FIGURE 2.5.8. Using the improved Euler method to approximate the actual solution of the initial value problem in Example 3.

Example 4

In Example 4 of Section 2.4 we applied Euler's method to the initial value problem

$$\frac{dy}{dx} = y\cos x, \quad y(0) = 1.$$

Figure 2.4.10 shows obvious visual differences between the periodic exact solution $y(x) = e^{\sin x}$ and the Euler approximations on $0 \le x \le 6\pi$ with as many as n = 400 subintervals.

Figure 2.5.9 shows the exact solution curve and approximate solution curves plotted using the improved Euler method with n = 50, n = 100, and n = 200 subintervals. The approximation obtained with n = 200 is indistinguishable from the exact solution curve, and the approximation with n = 100 is only barely distinguishable from it.

Although Figs. 2.5.7 and 2.5.9 indicate that the improved Euler method can provide accuracy that suffices for many graphical purposes, it does not provide the higher-precision numerical accuracy that sometimes is needed for more careful investigations. For instance, consider again the initial value problem

$$\frac{dy}{dx} = -\frac{2xy}{1+x^2}, \quad y(0) = 1$$

of Example 1. The final column of the table in Fig. 2.5.6 suggests that, if the improved Euler method is used on the interval $0 \le x \le 1$ with *n* subintervals and step size h = 1/n, then the resulting error *E* in the final approximation $y_n \approx y(1)$ is given by

$$E = |y(1) - y_n| \approx (0.12)h^2 = \frac{0.12}{n^2}.$$

If so, then 12-place accuracy (for instance) in the value y(1) would require that $(0.12)n^{-2} < 5 \times 10^{-13}$, which means that $n \ge 489,898$. Thus, roughly half a million steps of length $h \approx 0.000002$ would be required. Aside from the possible impracticality of this many steps (using available computational resources), the roundoff error resulting from so many successive steps might well overwhelm the cumulative error predicted by theory (which assumes exact computations in each separate step). Consequently, still more accurate methods than the improved Euler method are needed for such high-precision computations. Such a method is presented in Section 2.6.



FIGURE 2.5.9. Approximating the exact solution $y = e^{\sin x}$ using the improved Euler method with n = 50, 100, and 200 subintervals.

2.5 Problems

A hand-held calculator will suffice for Problems 1 through 10, where an initial value problem and its exact solution are given. Apply the improved Euler method to approximate this solution on the interval [0, 0.5] with step size h = 0.1. Construct a table showing four-decimal-place values of the approximate solution and actual solution at the points x = 0.1, 0.2, 0.3, 0.4,0.5.

1.
$$y' = -y$$
, $y(0) = 2$; $y(x) = 2e^{-x}$
2. $y' = 2y$, $y(0) = \frac{1}{2}$; $y(x) = \frac{1}{2}e^{2x}$
3. $y' = y + 1$, $y(0) = 1$; $y(x) = 2e^{x} - 1$
4. $y' = x - y$, $y(0) = 1$; $y(x) = 2e^{-x} + x - 1$
5. $y' = y - x - 1$, $y(0) = 1$; $y(x) = 2 + x - e^{x}$
6. $y' = -2xy$, $y(0) = 2$; $y(x) = 2e^{-x^{2}}$
7. $y' = -3x^{2}y$, $y(0) = 3$; $y(x) = 3e^{-x^{3}}$
8. $y' = e^{-y}$, $y(0) = 0$; $y(x) = \ln(x + 1)$
9. $y' = \frac{1}{4}(1 + y^{2})$, $y(0) = 1$; $y(x) = \tan \frac{1}{4}(x + \pi)$
10. $y' = 2xy^{2}$, $y(0) = 1$; $y(x) = \frac{1}{1 - x^{2}}$

Note: The application following this problem set lists illustrative calculator/computer programs that can be used in Problems 11 through 24.

A programmable calculator or a computer will be useful for Problems 11 through 16. In each problem find the exact solution of the given initial value problem. Then apply the improved Euler method twice to approximate (to five decimal places) this solution on the given interval, first with step size h = 0.01, then with step size h = 0.005. Make a table showing the approximate values and the actual value, together with the percentage error in the more accurate approximations, for x an integral multiple of 0.2. Throughout, primes denote derivatives with respect to x.

11.
$$y' = y - 2$$
, $y(0) = 1$; $0 \le x \le 1$
12. $y' = \frac{1}{2}(y - 1)^2$, $y(0) = 2$; $0 \le x \le 1$
13. $yy' = 2x^3$, $y(1) = 3$; $1 \le x \le 2$
14. $xy' = y^2$, $y(1) = 1$; $1 \le x \le 2$
15. $xy' = 3x - 2y$, $y(2) = 3$; $2 \le x \le 3$
16. $y^2y' = 2x^5$, $y(2) = 3$; $2 \le x \le 3$

A computer with a printer is required for Problems 17 through 24. In these initial value problems, use the improved Euler method with step sizes h = 0.1, 0.02, 0.004, and 0.0008 to approximate to five decimal places the values of the solution at ten equally spaced points of the given interval. Print the results in tabular form with appropriate headings to make it easy to gauge the effect of varying the step size h. Throughout, primes denote derivatives with respect to x.

17.
$$y' = x^2 + y^2$$
, $y(0) = 0$; $0 \le x \le 1$
18. $y' = x^2 - y^2$, $y(0) = 1$; $0 \le x \le 2$
19. $y' = x + \sqrt{y}$, $y(0) = 1$; $0 \le x \le 2$
20. $y' = x + \sqrt[3]{y}$, $y(0) = -1$; $0 \le x \le 2$
21. $y' = \ln y$, $y(1) = 2$; $1 \le x \le 2$
22. $y' = x^{2/3} + y^{2/3}$, $y(0) = 1$; $0 \le x \le 2$

.

23. $y' = \sin x + \cos y$, y(0) = 0; $0 \le x \le 1$ **24.** $v' = \frac{x}{x}$, $v(-1) = 1; -1 \le x \le 1$

$$1 + y^2, y(1) = 1, 1 \equiv x \equiv 1$$

25. Falling parachutist As in Problem 25 of Section 2.4, you bail out of a helicopter and immediately open your parachute, so your downward velocity satisfies the initial value problem

$$\frac{dv}{dt} = 32 - 1.6v, \quad v(0) = 0$$

(with *t* in seconds and *v* in ft/s). Use the improved Euler method with a programmable calculator or computer to approximate the solution for $0 \le t \le 2$, first with step size h = 0.01 and then with h = 0.005, rounding off approximate *v*-values to three decimal places. What percentage of the limiting velocity 20 ft/s has been attained after 1 second? After 2 seconds?

26. Deer population As in Problem 26 of Section 2.4, suppose the deer population P(t) in a small forest initially numbers 25 and satisfies the logistic equation

$$\frac{dP}{dt} = 0.0225P - 0.0003P^2$$

(with *t* in months). Use the improved Euler method with a programmable calculator or computer to approximate the solution for 10 years, first with step size h = 1 and then with h = 0.5, rounding off approximate *P*-values to three decimal places. What percentage of the limiting population of 75 deer has been attained after 5 years? After 10 years?

Decreasing Step Size

Use the improved Euler method with a computer system to find the desired solution values in Problems 27 and 28. Start with step size h = 0.1, and then use successively smaller step sizes until successive approximate solution values at x = 2 agree rounded off to four decimal places.

27.
$$y' = x^2 + y^2 - 1$$
, $y(0) = 0$; $y(2) = ?$

28.
$$y' = x + \frac{1}{2}y^2$$
, $y(-2) = 0$; $y(2) = ?$

29. Velocity-proportional resistance Consider the crossbow bolt of Example 2 in Section 2.3, shot straight upward from the ground with an initial velocity of 49 m/s. Because of linear air resistance, its velocity function v(t)satisfies the initial value problem

$$\frac{dv}{dt} = -(0.04)v - 9.8, \quad v(0) = 49$$

with exact solution $v(t) = 294e^{-t/25} - 245$. Use a calculator or computer implementation of the improved Euler method to approximate v(t) for $0 \le t \le 10$ using both n = 50 and n = 100 subintervals. Display the results at intervals of 1 second. Do the two approximations—each rounded to two decimal places—agree both with each other and with the exact solution? If the exact solution

were unavailable, explain how you could use the improved Euler method to approximate closely (a) the bolt's time of ascent to its apex (given in Section 2.3 as 4.56 s) and (b) its impact velocity after 9.41 s in the air.

30. Square-proportional resistance Consider now the crossbow bolt of Example 3 in Section 2.3. It still is shot straight upward from the ground with an initial velocity of 49 m/s, but because of air resistance proportional to the square of its velocity, its velocity function v(t) satisfies the initial value problem

$$\frac{dv}{dt} = -(0.0011)v|v| - 9.8, \quad v(0) = 49.$$

The symbolic solution discussed in Section 2.3 required

separate investigations of the bolt's ascent and its descent, with v(t) given by a tangent function during ascent and by a hyperbolic tangent function during descent. But the improved Euler method requires no such distinction. Use a calculator or computer implementation of the improved Euler method to approximate v(t) for $0 \le t \le 10$ using both n = 100 and n = 200 subintervals. Display the results at intervals of 1 second. Do the two approximations—each rounded to two decimal places—agree with each other? If an exact solution were unavailable, explain how you could use the improved Euler method to approximate closely (**a**) the bolt's time of ascent to its apex (given in Section 2.3 as 4.61 s) and (**b**) its impact velocity after 9.41 s in the air.

2.5 Application Improved Euler Implementation



Python/TI-Nspire.

Figure 2.5.10 lists TI-84 Plus and Python programs implementing the improved Euler method to approximate the solution of the initial value problem

$$\frac{dy}{dx} = x + y, \quad y(0) = 1$$

considered in Example 2 of this section. The comments provided in the final column should make these programs intelligible even if you have little familiarity with the Python and TI programming languages.

To apply the improved Euler method to a differential equation dy/dx = f(x, y), one need only replace **X+Y** throughout with the desired expression. To

TI-84 Plus	Python	Comment
PROGRAM: IMPEULER	# Program IMPEULER	Program title
	<pre>def F(X,Y): return X + Y</pre>	Define function f
:10→N	N = 10	No. of steps
:0→X	X = 0.0	Initial x
:1→Y	Y = 1.0	Initial y
:1→T	X1 = 1.0	Final x
: (T-X) /N \rightarrow H	H = (X1 - X) / N	Step size
:For(I,1,N)	for I in range(N):	Begin loop
:Y→Z	Y0 = Y	Save previous y
:X+Y→K	K1 = F(X, Y)	First slope
:Z+H*K→Y	Y = Y0 + H*K1	Predictor
:X+H→X	X = X + H	New x
:X+Y→L	K2 = F(X, Y)	Second slope
:(K+L)/2 \rightarrow K	K = (K1 + K2)/2	Average slope
:Z+H*K→Y	X = X0 + H * K	Corrector
:Disp X,Y	print (X,Y)	Display results
:End	# END	End loop

FIGURE 2.5.10. TI-84 Plus and Python improved Euler programs.

increase the number of steps (and thereby decrease the step size) one need only change the value of \mathbf{N} specified in the second line of the program.

Figure 2.5.11 exhibits one MATLAB implementation of the improved Euler method. The impeuler function takes as input the initial value x, the initial value y, the final value x1 of x, and the desired number n of subintervals. As output it produces the resulting column vectors x and y of x- and y-values. For instance, the MATLAB command

[X, Y] = impeuler(0, 1, 1, 10)

then generates the first and fourth columns of data shown in Fig. 2.5.4.

```
function yp = f(x, y)
yp = x + y;
                               % yp = y'
function [X,Y] = impeuler(x,y,x1,n)
h = (x1 - x)/n;
                               % step size
X = x;
                               % initial x
Y = Y;
                               % initial y
for i = 1:n;
                               % begin loop
    k1 = f(x,y);
                               % first slope
                             % second slope
    k2 = f(x+h,y+h*k1);
    k = (k1 + k2)/2;;
                              % average slope
    x = x + h;
                               % new x
                               % new y
    y = y + h*k;
    X = [X;x];
                               % update x-column
    Y = [Y;y];
                               % update y-column
                               % end loop
    end
```

FIGURE 2.5.11. MATLAB implementation of improved Euler method.

You should begin this project by implementing the improved Euler method with your own calculator or computer system. Test your program by applying it first to the initial value problem of Example 1, then to some of the problems for this section.

Famous Numbers Revisited

The following problems describe the numbers $e \approx 2.7182818$, $\ln 2 \approx 0.6931472$, and $\pi \approx 3.1415927$ as specific values of certain initial value problems. In each case, apply the improved Euler method with $n = 10, 20, 40, \ldots$ subintervals (doubling *n* each time). How many subintervals are needed to obtain—twice in succession—the correct value of the target number rounded to five decimal places?

- 1. The number e = y(1), where y(x) is the solution of the initial value problem dy/dx = y, y(0) = 1.
- 2. The number $\ln 2 = y(2)$, where y(x) is the solution of the initial value problem dy/dx = 1/x, y(1) = 0.
- 3. The number $\pi = y(1)$, where y(x) is the solution of the initial value problem $dy/dx = 4/(1 + x^2)$, y(0) = 0.

Logistic Population Investigation

Apply your improved Euler program to the initial value problem $dy/dx = \frac{1}{3}y(8-y)$, y(0) = 1 of Example 3. In particular, verify (as claimed) that the approximate

solution with step size h = 1 levels off at $y \approx 4.3542$ rather than at the limiting value y = 8 of the exact solution. Perhaps a table of values for $0 \le x \le 100$ will make this apparent.

For your own logistic population to investigate, consider the initial value problem

$$\frac{dy}{dx} = \frac{1}{n}y(m-y), \quad y(0) = 1$$

where *m* and *n* are (for instance) the largest and smallest nonzero digits in your student ID number. Does the improved Euler approximation with step size h = 1 level off at the "correct" limiting value of the exact solution? If not, find a smaller value of *h* so that it does.

Periodic Harvesting and Restocking

The differential equation

$$\frac{dy}{dt} = ky(M - y) - h\sin\left(\frac{2\pi t}{P}\right)$$

models a logistic population that is periodically harvested and restocked with period P and maximal harvesting/restocking rate h. A numerical approximation program was used to plot the typical solution curves for the case k = M = h = P = 1 that are shown in Fig. 2.5.12. This figure suggests—although it does not prove—the existence of a threshold initial population such that

- Beginning with an initial population above this threshold, the population oscillates (perhaps with period P?) around the (unharvested) stable limiting population $y(t) \equiv M$, whereas
- The population dies out if it begins with an initial population below this threshold.

Use an appropriate plotting utility to investigate your own logistic population with periodic harvesting and restocking (selecting typical values of the parameters k, M, h, and P). Do the observations indicated here appear to hold for your population?

2.6 The Runge–Kutta Method

We now discuss a method for approximating the solution y = y(x) of the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0$$
 (1)

that is considerably more accurate than the improved Euler method and is more widely used in practice than any of the numerical methods discussed in Sections 2.4 and 2.5. It is called the *Runge–Kutta method*, after the German mathematicians who developed it, Carl Runge (1856–1927) and Wilhelm Kutta (1867–1944).

With the usual notation, suppose that we have computed the approximations $y_1, y_2, y_3, \ldots, y_n$ to the actual values $y(x_1), y(x_2), y(x_3), \ldots, y(x_n)$ and now want to compute $y_{n+1} \approx y(x_{n+1})$. Then

$$y(x_{n+1}) - y(x_n) = \int_{x_n}^{x_{n+1}} y'(x) \, dx = \int_{x_n}^{x_n+h} y'(x) \, dx \tag{2}$$



FIGURE 2.5.12. Solution curves of $dy/dt = y(1 - y) - \sin 2\pi t$.

by the fundamental theorem of calculus. Next, Simpson's rule for numerical integration yields

$$y(x_{n+1}) - y(x_n) \approx \frac{h}{6} \left[y'(x_n) + 4y'\left(x_n + \frac{h}{2}\right) + y'(x_{n+1}) \right].$$
 (3)

Hence we want to define y_{n+1} so that

$$y_{n+1} \approx y_n + \frac{h}{6} \left[y'(x_n) + 2y'\left(x_n + \frac{h}{2}\right) + 2y'\left(x_n + \frac{h}{2}\right) + y'(x_{n+1}) \right];$$
 (4)

we have split $4y'(x_n + \frac{1}{2}h)$ into a sum of two terms because we intend to approximate the slope $y'(x_n + \frac{1}{2}h)$ at the midpoint $x_n + \frac{1}{2}h$ of the interval $[x_n, x_{n+1}]$ in two different ways.

On the right-hand side in (4), we replace the [true] slope values $y'(x_n)$, $y'(x_n + \frac{1}{2}h)$, $y'(x_n + \frac{1}{2}h)$, and $y'(x_{n+1})$, respectively, with the following estimates.

$$k_1 = f(x_n, y_n) \tag{5a}$$

• This is the Euler method slope at x_n .

$$k_2 = f\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}hk_1\right)$$
(5b)

• This is an estimate of the slope at the midpoint of the interval $[x_n, x_{n+1}]$ using the Euler method to predict the ordinate there.

$$k_3 = f\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}hk_2\right)$$
(5c)

• This is an improved Euler value for the slope at the midpoint.

≻

>

>

$$k_4 = f(x_{n+1}, y_n + hk_3)$$
(5d)

• This is the Euler method slope at x_{n+1} , using the improved slope k_3 at the midpoint to step to x_{n+1} .

When these substitutions are made in (4), the result is the iterative formula

$$y_{n+1} = y_n + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4).$$
(6)

The use of this formula to compute the approximations y_1, y_2, y_3, \ldots successively constitutes the **Runge–Kutta method**. Note that Eq. (6) takes the "Euler form"

$$y_{n+1} = y_n + h \cdot k$$

if we write

$$k = \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4) \tag{7}$$

for the approximate *average slope* on the interval $[x_n, x_{n+1}]$.

The Runge–Kutta method is a *fourth-order* method—it can be proved that the cumulative error on a bounded interval [a, b] with $a = x_0$ is of order h^4 . (Thus the iteration in (6) is sometimes called the *fourth-order* Runge–Kutta method because it is possible to develop Runge–Kutta methods of other orders.) That is,

$$|y(x_n) - y_n| \le Ch^4,\tag{8}$$

where the constant C depends on the function f(x, y) and the interval [a, b], but does not depend on the step size h. The following example illustrates this high accuracy in comparison with the lower-order accuracy of our previous numerical methods. We first apply the Runge-Kutta method to the illustrative initial value problem

$$\frac{dy}{dx} = x + y, \quad y(0) = 1$$
 (9)

that we considered in Fig. 2.4.8 of Section 2.4 and again in Example 2 of Section 2.5. The exact solution of this problem is $y(x) = 2e^x - x - 1$. To make a point we use h = 0.5, a larger step size than in any previous example, so only two steps are required to go from x = 0 to x = 1.

In the first step we use the formulas in (5) and (6) to calculate

 $k_1 = 0 + 1 = 1,$ $k_2 = (0 + 0.25) + (1 + (0.25) \cdot (1)) = 1.5,$ $k_3 = (0 + 0.25) + (1 + (0.25) \cdot (1.5)) = 1.625,$ $k_4 = (0.5) + (1 + (0.5) \cdot (1.625)) = 2.3125,$

and then

Example 1

$$y_1 = 1 + \frac{0.5}{6} [1 + 2 \cdot (1.5) + 2 \cdot (1.625) + 2.3125] \approx 1.7969.$$

Similarly, the second step yields $y_2 \approx 3.4347$.

Figure 2.6.1 presents these results together with the results (from Fig. 2.5.4) of applying the improved Euler method with step size h = 0.1. We see that even with the larger step size, the Runge–Kutta method gives (for this problem) four to five times the accuracy (in terms of relative percentage errors) of the improved Euler method.

	Improved Euler		Runge–Kutta		
x	y with $h = 0.1$	Percent Error	y with $h = 0.5$	Percent Error	Actual y
0.0	1.0000	0.00%	1.0000	0.00%	1.0000
0.5	1.7949	0.14%	1.7969	0.03%	1.7974
1.0	3.4282	0.24%	3.4347	0.05%	3.4366

FIGURE 2.6.1. Runge–Kutta and improved Euler results for the initial value problem dy/dx = x + y, y(0) = 1.

It is customary to measure the computational labor involved in solving dy/dx = f(x, y) numerically by counting the number of evaluations of the function f(x, y) that are required. In Example 1, the Runge–Kutta method required eight evaluations of f(x, y) = x + y (four at each step), whereas the improved Euler method required 20 such evaluations (two for each of 10 steps). Thus the Runge–Kutta method gave over four times the accuracy with only 40% of the labor.

Computer programs implementing the Runge–Kutta method are listed in the project material for this section. Figure 2.6.2 shows the results obtained by applying the improved Euler and Runge–Kutta methods to the problem dy/dx = x + y, y(0) = 1 with the same step size h = 0.1. The relative error in the improved Euler value at x = 1 is about 0.24%, but for the Runge–Kutta value it is 0.00012%. In this comparison the Runge–Kutta method is about 2000 times as accurate, but requires only twice as many function evaluations, as the improved Euler method.

The error bound

$$|y(x_n) - y_n| \le Ch^4 \tag{8}$$

for the Runge–Kutta method results in a rapid decrease in the magnitude of errors when the step size h is reduced (except for the possibility that very small step sizes may result in unacceptable roundoff errors). It follows from the inequality in (8) that (on a fixed bounded interval) halving the step size decreases the absolute error by
130 Chapter 2 Mathematical Models and Numerical Methods

x	Improved Euler y	Runge–Kutta y	Actual y
0.1	1.1100	1.110342	1.110342
0.2	1.2421	1.242805	1.242806
0.3	1.3985	1.399717	1.399718
0.4	1.5818	1.583648	1.583649
0.5	1.7949	1.797441	1.797443
0.6	2.0409	2.044236	2.044238
0.7	2.3231	2.327503	2.327505
0.8	2.6456	2.651079	2.651082
0.9	3.0124	3.019203	3.019206
1.0	3.4282	3.436559	3.436564

FIGURE 2.6.2. Runge–Kutta and improved Euler results for the initial value problem dy/dx = x + y, y(0) = 1, with the same step size h = 0.1.

a factor of $(\frac{1}{2})^4 = \frac{1}{16}$. Consequently, the common practice of successively halving the step size until the computed results "stabilize" is particularly effective with the Runge–Kutta method.

Example 2

lem

Infinite discontinuity In Example 5 of Section 2.4 we saw that Euler's method is not adequate to approximate the solution y(x) of the initial value problem

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0) = 1 \tag{10}$$

as x approaches the infinite discontinuity near x = 0.969811 (see Fig. 2.6.3). Now we apply the Runge–Kutta method to this initial value problem.

Figure 2.6.4 shows Runge–Kutta results on the interval [0.0, 0.9], computed with step sizes h = 0.1, h = 0.05, and h = 0.025. There is still some difficulty near x = 0.9, but it seems safe to conclude from these data that $y(0.5) \approx 2.0670$.

x	y with $h = 0.1$	<i>y</i> with $h = 0.05$	<i>y</i> with $h = 0.025$
0.1	1.1115	1.1115	1.1115
0.3	1.4397	1.4397	1.4397
0.5	2.0670	2.0670	2.0670
0.7	3.6522	3.6529	3.6529
0.9	14.0218	14.2712	14.3021



FIGURE 2.6.3. Solutions of $dy/dx = x^2 + y^2$, y(0) = 1.

FIGURE 2.6.4. Approximating the solution of the initial value problem in Eq. (10).

We therefore begin anew and apply the Runge-Kutta method to the initial value prob-

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0.5) = 2.0670.$$
 (11)

Figure 2.6.5 shows results on the interval [0.5, 0.9], obtained with step sizes h = 0.01, h = 0.005, and h = 0.0025. We now conclude that $y(0.9) \approx 14.3049$.

Finally, Fig. 2.6.6 shows results on the interval [0.90, 0.95] for the initial value problem

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0.9) = 14.3049,$$
 (12)

obtained using step sizes h = 0.002, h = 0.001, and h = 0.0005. Our final approximate result is $y(0.95) \approx 50.4723$. The actual value of the solution at x = 0.95 is $y(0.95) \approx 50.471867$.

x	<i>y</i> with $h = 0.01$	<i>y</i> with $h = 0.005$	<i>y</i> with $h = 0.0025$
0.5	2.0670	2.0670	2.0670
0.6	2.6440	2.6440	2.6440
0.7	3.6529	3.6529	3.6529
0.8	5.8486	5.8486	5.8486
0.9	14.3048	14.3049	14.3049

FIGURE 2.6.5. Approximating the solution of the initial value problem in Eq. (11).

x	<i>y</i> with $h = 0.002$	<i>y</i> with $h = 0.001$	<i>y</i> with $h = 0.0005$
0.90	14.3049	14.3049	14.3049
0.91	16.7024	16.7024	16.7024
0.92	20.0617	20.0617	20.0617
0.93	25.1073	25.1073	25.1073
0.94	33.5363	33.5363	33.5363
0.95	50.4722	50.4723	50.4723

FIGURE 2.6.6. Approximating the solution of the initial value problem in Eq. (12).

Our slight overestimate results mainly from the fact that the four-place initial value in (12) is (in effect) the result of rounding up the actual value $y(0.9) \approx 14.304864$; such errors are magnified considerably as we approach the vertical asymptote.

Example 3 Skydiver A skydiver with a mass of 60 kg jumps from a helicopter hovering at an initial altitude of 5 kilometers. Assume that she falls vertically with initial velocity zero and experiences an upward force F_R of air resistance given in terms of her velocity v (in meters per second) by

$$F_R = (0.0096)(100v + 10v^2 + v^3)$$

(in newtons, and with the coordinate axis directed downward so that v > 0 during her descent to the ground). If she does not open her parachute, what will be her terminal velocity? How fast will she be falling after 5 s have elapsed? After 10 s? After 20 s?

Solution Newton's law F = ma gives

$$m\frac{dv}{dt} = mg - F_R;$$

that is,

$$60\frac{dv}{dt} = (60)(9.8) - (0.0096)(100v + 10v^2 + v^3)$$
(13)

because m = 60 and g = 9.8. Thus the velocity function v(t) satisfies the initial value problem

$$\frac{dv}{dt} = f(v), \quad v(0) = 0,$$
 (14)

where

$$f(v) = 9.8 - (0.00016)(100v + 10v^2 + v^3).$$
⁽¹⁵⁾

The skydiver reaches her terminal velocity when the forces of gravity and air resistance balance, so f(v) = 0. We can therefore calculate her terminal velocity immediately by solving the equation

$$f(v) = 9.8 - (0.00016)(100v + 10v^2 + v^3) = 0.$$
 (16)

Figure 2.6.7 shows the graph of the function f(v) and exhibits the single real solution $v \approx$ 35.5780 (found graphically or by using a calculator or computer **Solve** procedure). Thus the skydiver's terminal speed is approximately 35.578 m/s, about 128 km/h (almost 80 mi/h).

t (s)	v (m/s)	<i>t</i> (s)	v (m/s)
0	0	11	35.541
1	9.636	12	35.560
2	18.386	13	35.569
3	25.299	14	35.574
4	29.949	15	35.576
5	32.678	16	35.577
6	34.137	17	35.578
7	34.875	18	35.578
8	35.239	19	35.578
9	35.415	20	35.578
10	35.500		

60 40 20

 $\widehat{\underbrace{a}}_{-20}^{0} 0$ -40-60

FIGURE 2.6.7. Graph of f(v) =9.8 - (0.00016)(100v + 10v² + v³).

0

20

40 60 80 100

-60-40-20

FIGURE 2.6.8. The skydiver's velocity data.

Figure 2.6.8 shows the results of Runge–Kutta approximations to the solution of the initial value problem in (14); the step sizes h = 0.2 and h = 0.1 yield the same results (to three decimal places). Observe that the terminal velocity is effectively attained in only 15 s. But the skydiver's velocity is 91.85% of her terminal velocity after only 5 s, and 99.78% after 10 s.

The final example of this section contains a *warning*: For certain types of initial value problems, the numerical methods we have discussed are not nearly so successful as in the previous examples.

4 Consider the seemingly innocuous initial value problem

$$\frac{dy}{dx} = 5y - 6e^{-x}, \quad y(0) = 1 \tag{17}$$

whose exact solution is $y(x) = e^{-x}$. The table in Fig. 2.6.9 shows the results obtained by applying the Runge–Kutta method on the interval [0, 4] with step sizes h = 0.2, h = 0.1, and h = 0.05. Obviously these attempts are spectacularly unsuccessful. Although $y(x) = e^{-x} \rightarrow 0$ as $x \rightarrow +\infty$, it appears that our numerical approximations are headed toward $-\infty$ rather than zero.

	Runge–Kutta y	Runge–Kutta y	Runge–Kutta y	
x	with $h = 0.2$	with $h = 0.1$	with $h = 0.05$	Actual y
0.4	0.66880	0.67020	0.67031	0.67032
0.8	0.43713	0.44833	0.44926	0.44933
1.2	0.21099	0.29376	0.30067	0.30119
1.6	-0.46019	0.14697	0.19802	0.20190
2.0	-4.72142	-0.27026	0.10668	0.13534
2.4	-35.53415	-2.90419	-0.12102	0.09072
2.8	-261.25023	-22.05352	-1.50367	0.06081
3.2	-1,916.69395	-163.25077	-11.51868	0.04076
3.6	-14059.35494	-1205.71249	-85.38156	0.02732
4.0	-103,126.5270	-8903.12866	-631.03934	0.01832

FIGURE 2.6.9. Runge–Kutta attempts to solve numerically the initial value problem in Eq. (17).



FIGURE 2.6.10. Direction field and solution curves for $dy/dx = 5y - 6e^{-x}$.

The explanation lies in the fact that the general solution of the equation $dy/dx = 5y - 6e^{-x}$ is

y

$$(x) = e^{-x} + Ce^{5x}.$$
 (18)

The particular solution of (17) satisfying the initial condition y(0) = 1 is obtained with C = 0. But any departure, however small, from the exact solution $y(x) = e^{-x}$ —even if due only to roundoff error—introduces [in effect] a nonzero value of *C* in Eq. (18). And as indicated in Fig. 2.6.10, all solution curves of the form in (18) with $C \neq 0$ diverge rapidly away from the one with C = 0, even if their initial values are close to 1.

Difficulties of the sort illustrated by Example 4 sometimes are unavoidable, but one can at least hope to recognize such a problem when it appears. Approximate values whose order of magnitude varies with changing step size are a common indicator of such instability. These difficulties are discussed in numerical analysis textbooks and are the subject of current research in the field.

2.6 **Problems**

A hand-held calculator will suffice for Problems 1 through 10, where an initial value problem and its exact solution are given. Apply the Runge–Kutta method to approximate this solution on the interval [0, 0.5] with step size h = 0.25. Construct a table showing five-decimal-place values of the approximate solution and actual solution at the points x = 0.25 and 0.5.

1.
$$y' = -y$$
, $y(0) = 2$; $y(x) = 2e^{-x}$
2. $y' = 2y$, $y(0) = \frac{1}{2}$; $y(x) = \frac{1}{2}e^{2x}$
3. $y' = y + 1$, $y(0) = 1$; $y(x) = 2e^{x} - 1$
4. $y' = x - y$, $y(0) = 1$; $y(x) = 2e^{-x} + x - 1$
5. $y' = y - x - 1$, $y(0) = 1$; $y(x) = 2 + x - e^{x}$
6. $y' = -2xy$, $y(0) = 2$; $y(x) = 2e^{-x^{2}}$
7. $y' = -3x^{2}y$, $y(0) = 3$; $y(x) = 3e^{-x^{3}}$
8. $y' = e^{-y}$, $y(0) = 0$; $y(x) = \ln(x + 1)$
9. $y' = \frac{1}{4}(1 + y^{2})$, $y(0) = 1$; $y(x) = \frac{1}{1 - x^{2}}$

Note: The application following this problem set lists illustrative calculator/computer programs that can be used in the remaining problems.

A programmable calculator or a computer will be useful for Problems 11 through 16. In each problem find the exact solution of the given initial value problem. Then apply the Runge– Kutta method twice to approximate (to five decimal places) this solution on the given interval, first with step size h = 0.2, then with step size h = 0.1. Make a table showing the approximate values and the actual value, together with the percentage error in the more accurate approximation, for x an integral multiple of 0.2. Throughout, primes denote derivatives with respect to x.

11.
$$y' = y - 2$$
, $y(0) = 1$; $0 \le x \le 1$
12. $y' = \frac{1}{2}(y - 1)^2$, $y(0) = 2$; $0 \le x \le 1$
13. $yy' = 2x^3$, $y(1) = 3$; $1 \le x \le 2$
14. $xy' = y^2$, $y(1) = 1$; $1 \le x \le 2$

- **15.** $xy' = 3x 2y, y(2) = 3; 2 \le x \le 3$
- **16.** $y^2 y' = 2x^5$, y(2) = 3; $2 \le x \le 3$

A computer with a printer is required for Problems 17 through 24. In these initial value problems, use the Runge–Kutta method with step sizes h = 0.2, 0.1, 0.05, and 0.025 to approximate to six decimal places the values of the solution at five equally spaced points of the given interval. Print the results in tabular form with appropriate headings to make it easy to gauge the effect of varying the step size h. Throughout, primes denote derivatives with respect to x.

- **17.** $y' = x^2 + y^2$, y(0) = 0; $0 \le x \le 1$ **18.** $y' = x^2 - y^2$, y(0) = 1; $0 \le x \le 2$
- **19.** $y' = x + \sqrt{y}, y(0) = 1; 0 \le x \le 2$
- **20.** $y' = x + \sqrt[3]{y}, y(0) = -1; 0 \le x \le 2$
- **21.** $y' = \ln y, y(1) = 2; 1 \le x \le 2$
- **22.** $y' = x^{2/3} + y^{2/3}$, y(0) = 1; $0 \le x \le 2$
- **23.** $y' = \sin x + \cos y$, y(0) = 0; $0 \le x \le 1$

24.
$$y' = \frac{x}{1+y^2}, y(-1) = 1; -1 \le x \le 1$$

25. Falling parachutist As in Problem 25 of Section 2.5, you bail out of a helicopter and immediately open your parachute, so your downward velocity satisfies the initial value problem

$$\frac{dv}{dt} = 32 - 1.6v, \quad v(0) = 0$$

(with *t* in seconds and *v* in ft/s). Use the Runge–Kutta method with a programmable calculator or computer to approximate the solution for $0 \le t \le 2$, first with step size h = 0.1 and then with h = 0.05, rounding off approximate *v*-values to three decimal places. What percentage of the limiting velocity 20 ft/s has been attained after 1 second? After 2 seconds?

26. Deer population As in Problem 26 of Section 2.5, suppose the deer population P(t) in a small forest initially numbers 25 and satisfies the logistic equation

$$\frac{dP}{dt} = 0.0225P - 0.0003P^2$$

(with *t* in months). Use the Runge–Kutta method with a programmable calculator or computer to approximate the solution for 10 years, first with step size h = 6 and then with h = 3, rounding off approximate *P*-values to four decimal places. What percentage of the limiting population of 75 deer has been attained after 5 years? After 10 years?

Use the Runge–Kutta method with a computer system to find the desired solution values in Problems 27 and 28. Start with step size h = 1, and then use successively smaller step sizes until successive approximate solution values at x = 2 agree rounded off to five decimal places.

27.
$$y' = x^2 + y^2 - 1$$
, $y(0) = 0$; $y(2) = ?$
28. $y' = x + \frac{1}{2}y^2$, $y(-2) = 0$; $y(2) = ?$

Velocity-Acceleration Problems

In Problems 29 and 30, the linear acceleration a = dv/dt of a moving particle is given by a formula dv/dt = f(t, v), where the velocity v = dy/dt is the derivative of the function y = y(t) giving the position of the particle at time t. Suppose that the velocity v(t) is approximated using the Runge–Kutta method to solve numerically the initial value problem

$$\frac{dv}{dt} = f(t, v), \quad v(0) = v_0.$$
 (19)

That is, starting with $t_0 = 0$ and v_0 , the formulas in Eqs. (5) and (6) are applied—with t and v in place of x and y—to calculate the successive approximate velocity values v_1 , v_2 , v_3 , ..., v_m at the successive times t_1 , t_2 , t_3 , ..., t_m (with $t_{n+1} = t_n + h$). Now suppose that we also want to approximate the distance y(t) traveled by the particle. We can do this by beginning with the initial position $y(0) = y_0$ and calculating

$$y_{n+1} = y_n + v_n h + \frac{1}{2}a_n h^2 \tag{20}$$

(n = 1, 2, 3, ...), where $a_n = f(t_n, v_n) \approx v'(t_n)$ is the particle's approximate acceleration at time t_n . The formula in (20) would give the correct increment (from y_n to y_{n+1}) if the acceleration a_n remained constant during the time interval $[t_n, t_{n+1}]$.

Thus, once a table of approximate velocities has been calculated, Eq. (20) provides a simple way to calculate a table of corresponding successive positions. This process is illustrated in the project for this section, by beginning with the velocity data in Fig. 2.6.8 (Example 3) and proceeding to follow the skydiver's position during her descent to the ground.

29. Consider again the crossbow bolt of Example 2 in Section 2.3, shot straight upward from the ground with an initial velocity of 49 m/s. Because of linear air resistance, its velocity function v = dy/dt satisfies the initial value problem

$$\frac{dv}{dt} = -(0.04)v - 9.8, \quad v(0) = 49$$

with exact solution $v(t) = 294e^{-t/25} - 245$. (a) Use a calculator or computer implementation of the Runge–Kutta method to approximate v(t) for $0 \le t \le 10$ using both n = 100 and n = 200 subintervals. Display the results at intervals of 1 second. Do the two approximations—each rounded to four decimal places—agree both with each other and with the exact solution? (b) Now use the velocity data from part (a) to approximate y(t) for $0 \le t \le 10$ using n = 200 subintervals. Display the results at intervals of 1 second. Do these approximate position values—each rounded to two decimal places—agree with the exact solution

$$y(t) = 7350 \left(1 - e^{-t/25} \right) - 245t?$$

(c) If the exact solution were unavailable, explain how you could use the Runge–Kutta method to approximate closely the bolt's times of ascent and descent and the maximum height it attains.

30. Now consider again the crossbow bolt of Example 3 in Section 2.3. It still is shot straight upward from the ground with an initial velocity of 49 m/s, but because of air resistance proportional to the square of its velocity, its velocity function v(t) satisfies the initial value problem

$$\frac{dv}{dt} = -(0.0011)v|v| - 9.8, \quad v(0) = 49.$$

Beginning with this initial value problem, repeat parts (a) through (c) of Problem 29 (except that you may need n = 200 subintervals to get four-place accuracy in part (a) and n = 400 subintervals for two-place accuracy in part (b)). According to the results of Problems 17 and 18 in Section 2.3, the bolt's velocity and position functions during ascent and descent are given by the following formulas.

Ascent: $v(t) = (94.388) \tan(0.478837 - [0.103827]t),$ y(t) = 108.465 $+ (909.091) \ln (\cos(0.478837 - [0.103827]t));$ Descent: $v(t) = -(94.388) \tanh(0.103827[t - 4.6119]),$ y(t) = 108.465

 $-(909.091) \ln (\cosh(0.103827[t-4.6119])).$

2.6 Application Runge–Kutta Implementation



computing resources including Maple/Mathematica/MATLAB/ Python/TI-Nspire. Figure 2.6.11 lists TI-Nspire CX CAS and Python programs implementing the Runge–Kutta method to approximate the solution of the initial value problem

$$\frac{dy}{dx} = x + y, \quad y(0) = 1$$

considered in Example 1 of this section. The comments provided in the final column should make these programs intelligible even if you have little familiarity with the Python and TI programming languages.

To apply the Runge-Kutta method to a different equation dy/dx = f(x, y), one need only change the initial line of the program, in which the function f is defined. To increase the number of steps (and thereby decrease the step size), one need only change the value of n specified in the second line of the program.

Figure 2.6.12 exhibits a MATLAB implementation of the Runge-Kutta method. Suppose that the function f describing the differential equation y' = f(x, y) has been defined. Then the **rk** function takes as input the initial value **x**, the initial value **y**, the final value **x1** of x, and the desired number **n** of subintervals. As output it produces the resulting column vectors **x** and **y** of x- and y-values. For instance, the MATLAB command

TI-Nspire CX CAS	Python	Comment
Define rk()=Prgm	# Program RK	Program title
f(x,y) := x+y	def F(X,Y): return X + Y	Define function f
n:=10	N = 10	No. of steps
x:=0.0	X = 0.0	Initial x
y:=1.0	Y = 1.0	Initial y
x1:=1.0	X1 = 1.0	Final x
h:=(x1-x)/n	H = (X1 - X) / N	Step size
For i,1,n	for I in range(N):	Begin loop
x0:=x	X0 = X	Save previous x
y0:=y	Y0 = Y	Save previous y
k1:=f(x,y)	K1 = F(X, Y)	First slope
x:=x0+h/2	X = X0 + H/2	Midpoint
y:=y0+(h*k1)/2	Y = Y0 + H*K1/2	Midpt predictor
k2:=f(x,y)	K2 = F(X, Y)	Second slope
y:=y0+(h*k2)/2	Y = Y0 + H*K2/2	Midpt predictor
k3:=f(x,y)	K3 = F(X, Y)	Third slope
x := x0+h	X = X0 + H	New x
y:=y0+h*k3	Y = Y0 + H*K3	Endpt predictor
k4:=f(x,y)	K4 = F(X, Y)	Fourth slope
k:=(k1+2*k2+2*k3+k4)/6	K = (K1+2*K2+2*K3 + K4)/6	Average slope
y:=y0+h*k	Y = Y0 + H * K	Corrector
Disp x,y	print (X,Y)	Display results
EndFor	# END	End loop
EndPrgm		

```
function yp = f(x, y)
yp = x + y;
                                    % yp = y'
function [X,Y] = rk(x,y,x1,n)
h = (x1 - x)/n;
                                    % step size
X = x;
                                    % initial x
Y = y;
                                    % initial y
for i = 1:n
                                    % begin loop
    k1 = f(x,y);
                                    % first slope
    k2 = f(x+h/2,y+h*k1/2);
                                    % second slope
    k3 = f(x+h/2,y+h*k2/2);
                                    % third slope
    k4 = f(x+h,y+h*k3);
                                    % fourth slope
    k = (k1+2*k2+2*k3+k4)/6;
                                    % average slope
    x = x + h;
                                    % new x
    y = y + h*k;
                                    % new y
    X = [X;x];
                                    % update x-column
    Y = [Y;y];
                                    % update y-column
                                    % end loop
    end
```

FIGURE 2.6.12. MATLAB implementation of the Runge–Kutta method.

[X, Y] = rk(0, 1, 1, 10)

then generates the first and third columns of data shown in the table in Fig. 2.6.2.

You should begin this project by implementing the Runge–Kutta method with your own calculator or computer system. Test your program by applying it first to the initial value problem in Example 1, then to some of the problems for this section.

Famous Numbers Revisited, One Last Time

The following problems describe the numbers

```
e \approx 2.71828182846, \ln 2 \approx 0.69314718056, and \pi \approx 3.14159265359
```

as specific values of certain initial value problems. In each case, apply the Runge– Kutta method with n = 10, 20, 40, ... subintervals (doubling *n* each time). How many subintervals are needed to obtain—twice in succession—the correct value of the target number rounded to nine decimal places?

- 1. The number e = y(1), where y(x) is the solution of the initial value problem dy/dx = y, y(0) = 1.
- 2. The number $\ln 2 = y(2)$, where y(x) is the solution of the initial value problem dy/dx = 1/x, y(1) = 0.
- 3. The number $\pi = y(1)$, where y(x) is the solution of the initial value problem $dy/dx = 4/(1 + x^2)$, y(0) = 0.

The Skydiver's Descent

The following MATLAB function describes the skydiver's acceleration function in Example 3.

```
function vp = f(t,v)
vp = 9.8 - 0.00016*(100*v + 10*v^2 + v^3);
```

Then the commands

```
k = 200 % 200 subintervals
[t,v] = rk(0, 0, 20, k); % Runge-Kutta approximation
[t(1:10:k+1); v(1:10:k+1)] % Display every 10th entry
```

produce the table of approximate velocities shown in Fig. 2.6.8. Finally, the commands

t (s)	v (m/s)	y (m)
0	0	0
2	18.386	18.984
4	29.949	68.825
6	34.137	133.763
8	35.239	203.392
10	35.500	274.192
12	35.560	345.266
14	35.574	416.403
16	35.577	487.555
18	35.578	558.710
20	35.578	629.866

FIGURE 2.6.13. The skydiver's velocity and position data.

y = zeros(k+1, 1):	% initialize y
h = 0.1;	% step size
for $n = 1:k$	% for n = 1 to k
a = f(t(n), v(n)):	% acceleration
$y(n+1) = y(n) + v(n) * h + 0.5 * a * h^2;$	% Equation (20)
end	% end loop
[t(1:20:k+1), v(1:20:k+1), v(1:20:k+1)]	% each 20th entry

carry out the position function calculations described in Eq. (20) in the instructions for Problems 29 and 30. The results of these calculations are shown in the table in Fig. 2.6.13. It appears that the skydiver falls 629.866 m during her first 20 s of descent, and then free falls the remaining 4370.134 meters to the ground at her terminal speed of 35.578 m/s. Hence her total time of descent is $20 + (4370.134/35.578) \approx 142.833 \text{ s}$, or about 2 min 23 s.

For an individual problem to solve after implementing these methods using an available computer system, analyze your own skydive (perhaps from a different height), using your own mass *m* and a plausible air-resistance force of the form $F_R = av + bv^2 + cv^3$.



Linear Systems and Matrices

3.1 Introduction to Linear Systems

The subject of **linear algebra** centers around the problem of solving systems of (algebraic) *linear equations*. In high school algebra the *method of elimination* is used to solve systems of two and three (simultaneous) linear equations, with attention ordinarily confined to systems that have one and only one solution, and with applications to "word problems" that have one and only one "answer." In this section we introduce some of the basic terminology of linear algebra by reviewing the elementary technique of elimination from a slightly more general viewpoint. In subsequent sections we apply this same technique to the solution of systems involving many linear equations in many unknowns. The applications of the method of elimination are diverse and important, because so many mathematical problems involve the solution of systems of linear equations.

Recall that if a, b, and c are constants with a and b not both zero, then the graph of the equation

$$ax + by = c \tag{1}$$

is a (straight) *line* in the *xy*-plane. For this reason, an equation of the form in (1) is called a **linear equation** in the **variables** x and y. Similarly, an equation that can be written in the form

$$ax + by + cz = d \tag{2}$$

is called **linear** in the three variables x, y, and z (even though its graph in xyz-space is a plane rather than a line). Thus the equations

$$3x - 2y = 5$$
 and $x - 7y + 5z = -11$

are linear, because they involve only the first powers of the variables. By contrast, the equations

$$x + y + xy = 5$$
 and $x^2 + y^3 + \sqrt{z} = 1$

are *not* linear, because they cannot be rewritten to eliminate the higher powers, roots, and products of the variables.

Two Equations in Two Unknowns

A system of linear equations (also called a **linear system**) is simply a finite collection of linear equations involving certain variables. Sometimes we refer to the variables as the "unknowns" in the system. Thus a system of two linear equations in two unknowns x and y may be written in the form

$$a_1x + b_1y = c_1 a_2x + b_2y = c_2.$$
(3)

By a **solution** of the system in (3) is meant a pair (x, y) of values—normally real numbers—that satisfy *both* equations *simultaneously*.

Example 1The values
$$x = 2, y = -1$$
 constitute a solution of the system $2x - y = 5$
 $x + 2y = 0$ (4)because both $2(2) - (-1) = 5$ and $(2) + 2(-1) = 0$. The values $x = 3, y = 1$ satisfy the first
equation in (4) but do not satisfy the second. Hence (3, 1) is not a solution of the system in
(4).Example 2The linear system

$$\begin{aligned} x + y &= 1\\ 2x + 2y &= 3 \end{aligned}$$
(5)

has no solution at all, because if x + y = 1, then 2x + 2y = 2, and so $2x + 2y \neq 3$. Thus, two numbers that satisfy the first equation in (5) cannot simultaneously satisfy the second.

A linear system is said to be **consistent** if it has at least one solution and **inconsistent** if it has none. Thus the system in Example 1 is consistent, whereas the system in Example 2 is inconsistent.

Three Possibilities

Given a system of linear equations, we ask: What is the set of all solutions of the system? In brief, what is the **solution set** of the system?

In the case of a linear system

$$a_1 x + b_1 y = c_1 a_2 x + b_2 y = c_2$$
(6)

of two equations in two unknowns, we can use our knowledge of elementary geometry to sort out the possibilities for its solution set. If neither equation in (6) has both left-hand coefficients zero, then their graphs in the *xy*-plane are two straight lines L_1 and L_2 . Then exactly one of the following situations must hold.

- The lines L_1 and L_2 intersect at a single point (as in Figure 3.1.1).
- The lines L_1 and L_2 are parallel nonintersecting lines (as in Figure 3.1.2).
- The lines L_1 and L_2 coincide—they actually are the same line (see Figure 3.1.3).

A pair (x, y) of real numbers constitutes a solution of the system in (6) if and only if the point (x, y) in the coordinate plane lies on *both* the two lines L_1 and L_2 . In the case shown in Figure 3.1.1, there is exactly one such point. In the case shown in Figure 3.1.2, there is no such point, and in the case in Figure 3.1.3, there are infinitely many such points—every point on the line $L_1 = L_2$ is such a point. We therefore see that there are just *three possibilities* for a linear system of two equations in two unknowns: It has either



FIGURE 3.1.1. Two intersecting lines: a unique solution.

FIGURE 3.1.2. Two parallel lines: no solution.



FIGURE 3.1.3. Two coincident lines: infinitely many solutions.

- exactly one solution;
- no solution; or
- infinitely many solutions.

It is a fundamental fact of linear algebra (which we will establish in Section 3.3) that, however many equations and variables may appear in a linear system, precisely these same three possibilities occur: A system of m linear equations in n variables either has a **unique** solution (that is, exactly one solution), or it has no solution, or it has infinitely many solutions. Thus it is impossible, for instance, for a linear system to have exactly 2 solutions or to have exactly 17 solutions.

The Method of Elimination

The next three examples illustrate how we can use the elementary method of elimination to solve a system of two equations in two unknowns. To **solve** a system means to determine what its solution set is. The basic idea of the method is this:

- First, we add an appropriate constant multiple of the first equation to the second equation. The idea is to choose the constant in such a way as to eliminate the variable *x* from the second equation.
- Next, the new second equation contains only the variable *y*, so we readily solve it for the value of *y*.
- Finally, we determine the value of x by "back substitution" of this value of y in the first equation.

Examples 3 through 5 illustrate this method in the three cases corresponding to Figures 3.1.1 through 3.1.3 (respectively).

Example 3 In order to solve the system

$$5x + 3y = 1 x - 2y = 8,$$
(7)

we first interchange the two equations:

$$\begin{aligned} x - 2y &= 8\\ 5x + 3y &= 1. \end{aligned}$$

Then we multiply the first equation by -5 and add the resulting terms to the second (without

3.1 INTRODUCTION TO LINEAR SYSTEMS 41

changing the first equation). The result is

$$\begin{array}{rcl} x - & 2y = & 8 \\ & 13y = -39. \end{array}$$
(8)

Now the second equation immediately yields the value y = -3, and *back substitution* of this value in the first equation yields

$$x = 2y + 8 = 2(-3) + 8 = 2$$

Taking it as obvious (for the moment) that the systems in (7) and (8) have the same solution set, we conclude that the original system in (7) has the *unique solution* x = 2, y = -3.

Example 4 To solve the

$$2x + 6y = 4 3x + 9y = 11,$$
(9)

we first multiply the first equation by $\frac{1}{2}$ and get

$$x + 3y = 2$$
$$3x + 9y = 11$$

We next multiply the first equation by -3 and add each term to the corresponding term in the second equation. The result is

$$x + 3y = 2
 0 = 5
 (10)$$

What, however, are we to make of the new second equation, 0 = 5? The system in (10) actually is

$$x + 3y = 2$$
$$0x + 0y = 5.$$

Because 0 is simply not equal to 5, there are *no* values of x and y that satisfy the second equation. Hence there certainly can be no values that satisfy both simultaneously. We conclude that the original system in (9) has *no solution*.

Example 5

If, instead of the system in (9), we had begun in Example 4 with the system

$$2x + 6y = 4 (11) 3x + 9y = 6$$

and performed the same operations, we would have obtained, instead of (10), the system

$$x + 3y = 2
 0 = 0
 (12)$$

Here, 0 = 0 is shorthand for the equation

$$0x + 0y = 0,$$

which is satisfied by *all* values of x and y. In terms of restrictions or conditions on x and y, one of our original two equations has in effect disappeared, leaving us with the single equation

$$x + 3y = 2.$$
 (13)

Of course this is hardly surprising, because each equation in (11) is a multiple of the one in (13); in some sense we really had only one equation to begin with. At any rate, we can substitute any value of y we please in (13) and then solve for x. Thus our system in (11) has *infinitely many solutions*. To describe them explicitly, let us write y = t, where the **parameter** t is a new independent variable that we will use to generate solution pairs (x, y). Then Equation (13) yields x = 2 - 3t, so our infinite solution set of the system in (11) may be described as follows:

$$x = 2 - 3t, \quad y = t$$

as the arbitrary parameter t ranges over the set of all real numbers. For instance, t = 2 yields the solution (-4, 2), and t = -3 yields the solution (11, -3).

Remark There is some latitude in how we parameterize the solutions of a system having infinitely many solutions, including the names of any parameters we introduce for this purpose. For example, we could parameterize the solutions of the system in (11) alternatively by writing x = s in (13), thereby getting the *different* parameterization

$$x = s$$
, $y = \frac{1}{3}(2-s)$

that yields the *same* solutions. For instance, the parameter values s = -4 and s = 11 here give the previously noted solutions (-4, 2) and (11, -3) of the original system in (13).

Comment These three examples illustrate the basic features of the method of elimination, which involves "transforming" a given linear system by means of a sequence of successive steps that do not change the solutions of the system. Each of these steps consists of performing one of the following three **elementary operations**:

- **1.** Multiply one equation by a nonzero constant.
- 2. Interchange two equations.
- **3.** Add a constant multiple of (the terms of) one equation to (corresponding terms of) another equation.

In subsequent sections of this chapter we discuss the systematic use of these elementary operations to eliminate variables successively in any linear system, whatever the number of equations and variables. In this way we will see that every linear system corresponds to precisely one of the three situations illustrated in Examples 3–5. That is, either

- we discover (as in Example 3) a unique solution of the system; or
- we eventually arrive at an inconsistent equation (as in Example 4), so that the system has no solution; or
- we find that the original system has infinitely many solutions.

Three Equations in Three Unknowns

In the case of a system of three linear equations in three variables x, y, and z, we may proceed as follows: Assuming that the first equation involves x, use the first equation to eliminate x from the second and third equations by adding appropriate multiples of the first equation. Then, assuming that the new second equation involves y, use the new second equation to eliminate y from the new third equation. Solve the new third equation for z, back-substitute in the second equation to determine y, and, finally, back-substitute y and z in the first equation to find x. The following two examples illustrate this procedure.

Example 6 Solve the linear system

$$x + 2y + z = 4$$

$$3x + 8y + 7z = 20$$

$$2x + 7y + 9z = 23.$$

(14)

Solution First, we add -3 times the first equation to the second equation; the result is

$$x + 2y + z = 4$$
$$2y + 4z = 8$$
$$2x + 7y + 9z = 23$$

Then addition of -2 times the first equation to the third equation yields

х

$$+2y + z = 4$$

 $2y + 4z = 8$
 $3y + 7z = 15.$

3.1 INTRODUCTION TO LINEAR SYSTEMS 43

We have now eliminated x from the second and third equations. To simplify the process of eliminating y from the third equation, we multiply the second equation by $\frac{1}{2}$, to obtain

$$x + 2y + z = 4$$
$$y + 2z = 4$$
$$3y + 7z = 15.$$

Finally, addition of -3 times the second equation to the third equation gives

$$\begin{aligned}
 x + 2y + z &= 4 \\
 y + 2z &= 4 \\
 z &= 3.
 \end{aligned}$$
(15)

This system has a *triangular* form that makes its solution easy. By back substitution of z = 3 in the second equation in (15), we find that

$$y = 4 - 2z = 4 - 2(3) = -2;$$

then the first equation yields

$$x = 4 - 2y - z$$

= 4 - 2(-2) - (3) = 5.

Thus our original system in (14) has the unique solution x = 5, y = -2, z = 3.

Comment The steps by which we transformed (14) into (15) show that every solution of the system in (14) is a solution of the system in (15). But these steps can be reversed to show similarly that every solution of the system in (15) is also a solution of the system in (14). Thus the two systems are equivalent in the sense that they have the same solution set. The computation at the end of Example 6 shows that (15) has the unique solution (x, y, z) = (5, -2, 3), and it follows that this also is the unique solution of the original system in (14).

	Solve the system	Example 7
	3x - 8y + 10z = 22	
(16)	x - 3y + 2z = 5	
	2x - 9y - 8z = -11.	
interchange the first two equations	In order to avoid fractions in the elimination of x , we first i to get	Solution
	x - 3y + 2z = 5	
	3x - 8y + 10z = 22	
	2x - 9y - 8z = -11.	
	Addition of -3 times the first equation to the second gives	

$$x - 3y + 2z = 5$$
$$y + 4z = 7$$
$$2x - 9y - 8z = -11$$

and then addition of -2 times the first equation to the third equation gives

$$x - 3y + 2z = 5$$
$$y + 4z = 7$$
$$-3y - 12z = -21$$

Finally, addition of 3 times the second equation to the third equation yields



FIGURE 3.1.4. The three planes of Example 7.

Because the third equation has disappeared, we can choose z = t arbitrarily and then solve for x and y:

$$y = 7 - 4z = 7 - 4t,$$

$$x = 5 + 3y - 2z$$

$$= 5 + 3(7 - 4t) - 2t = 26 - 14t.$$

Thus our original system in (16) has infinitely many solutions. Moreover, one convenient way of describing them is this:

$$x = 26 - 14t, \quad y = 7 - 4t, \quad z = t.$$
(18)

The arbitrary parameter t can take on all real number values, and in so doing generates all the (infinitely many) solutions of the original system in (16). Figure 3.1.4 shows the three planes corresponding to the three equations in (16). These planes intersect in a straight line that is parameterized by the equations in (18).

A Differential Equations Application

In Chapter 1 we saw that a general solution of a first-order differential equation involves an arbitrary constant that must be determined in order to satisfy a given initial condition. The next example illustrates the fact that a general solution of a second-order differential equation typically involves *two* arbitrary constants. An initial value problem for such a differential equation ordinarily involves *two* initial conditions, whose imposition leads to a system of two equations with the two arbitrary constants as unknowns.

Example 8

$$y(x) = Ae^{3x} + Be^{-3x}, (19)$$

then differentiation yields

If A and B are constants and

$$y'(x) = 3Ae^{3x} - 3Be^{-3x}$$
(20)

and

$$y''(x) = 9Ae^{3x} + 9Be^{-3x} = 9y(x).$$

Thus the function y(x) defined by (19) satisfies the second-order differential equation

$$y'' - 9y = 0. (21)$$

Now, suppose that we want to solve the initial value problem consisting of this equation and the two initial conditions

$$y(0) = 7, \quad y'(0) = 9.$$
 (22)

Then substitution of x = 0 in (19) and (20) yields the linear system

$$\begin{array}{rr} A + & B = 7 \\ 3A - 3B = 9 \end{array}$$

that we readily solve for A = 5, B = 2. It follows that the particular solution

$$y(x) = 5e^{3x} + 2e^{-3x}$$

satisfies both the differential equation (21) and the initial conditions in (22).

In Chapter 5 we will study higher-order differential equations and initial value problems in detail, after developing the necessary methods of linear algebra in the intervening chapters.

3.1 Problems

In each of Problems 1-22, use the method of elimination to determine whether the given linear system is consistent or inconsistent. For each consistent system, find the solution if it is unique; otherwise, describe the infinite solution set in terms of an arbitrary parameter t (as in Examples 5 and 7).

2. 3x + 2y = 91. x + 3y = 92x + y = 8x - y = 83. 2x + 3y = 14. 5x - 6y = 13x + 5y = 36x - 5y = 106. 4x - 2y = 45. x + 2y = 42x + 4y = 96x - 3v = 7x - 4y = -108. 3x - 6y = 127. -2x + 8y = 202x - 4v = 89. x + 5y + z = 210. x + 3y + 2z = 22x + y - 2z = 12x + 7y + 7z = -1x + 7y + 2z = 32x + 5y + 2z = 7**11.** 2x + 7y + 3z = 11**12.** 3x + 5y - z = 13x + 3y + 2z = 22x + 7y + z = 283x + 7y + 9z = -12x + 7y + 2z = 32**13.** 3x + 9y + 7z = 0**14.** 4x + 9y + 12z = -12x + 7y + 4z = 03x + y + 16z = -462x + 6y + 5z = 02x + 7y + 3z = 1915. x + 3y + 2z = 516. x - 3y + 2z = 6x - y + 3z = 3x + 4y - z = 45x + 6y + z = 203x + y + 8z = 10**17.** 2x - y + 4z = 7**18.** x + 5y + 6z = 33x + 2y - 2z = 35x + 2y - 10z = 15x + y + 2z = 158x + 17y + 8z = 5**20.** 2x + 3y + 7z = 15**19.** x - 2y + z = 22x - y - 4z = 13x + 4y + z = 20x - y - z = 5x + 2y + 3z = 10**21.** x + y - z = 5**22.** 4x - 2y + 6z = 03x + y + 3z = 11x - y - z = 04x + y + 5z = 142x - y + 3z = 0

In each of Problems 23–30, a second-order differential equation and its general solution y(x) are given. Determine the constants A and B so as to find a solution of the differential equation that satisfies the given initial conditions involving y(0) and y'(0).

- **23.** y'' + 4y = 0, $y(x) = A \cos 2x + B \sin 2x$, y(0) = 3, y'(0) = 8
- **24.** y'' 9y = 0, $y(x) = A \cosh 3x + B \sinh 3x$, y(0) = 5, y'(0) = 12
- **25.** y'' 25y = 0, $y(x) = Ae^{5x} + Be^{-5x}$, y(0) = 10, y'(0) = 20
- **26.** y'' 121y = 0, $y(x) = Ae^{11x} + Be^{-11x}$, y(0) = 44, y'(0) = 22
- **27.** y'' + 2y' 15y = 0, $y(x) = Ae^{3x} + Be^{-5x}$, y(0) = 40, y'(0) = -16
- **28.** y'' 10y' + 21y = 0, $y(x) = Ae^{3x} + Be^{7x}$, y(0) = 15, y'(0) = 13

- **29.** 6y'' 5y' + y = 0, $y(x) = Ae^{x/2} + Be^{x/3}$, y(0) = 7, y'(0) = 11
- **30.** 15y'' + y' 28y = 0, $y(x) = Ae^{4x/3} + Be^{-7x/5}$, y(0) = 41, y'(0) = 164
- **31.** A system of the form

$$a_1x + b_1y = 0$$
$$a_2x + b_2y = 0,$$

in which the constants on the right-hand side are all zero, is said to be **homogeneous**. Explain by geometric reasoning why such a system has either a unique solution or infinitely many solutions. In the former case, what is the unique solution?

32. Consider the system

$$a_1x + b_1y + c_1z = d_1$$
$$a_2x + b_2y + c_2z = d_2$$

of two equations in three unknowns.

- (a) Use the fact that the graph of each such equation is a plane in *xyz*-space to explain why such a system always has either no solution or infinitely many solutions.
- (b) Explain why the system must have infinitely many solutions if $d_1 = 0 = d_2$.
- 33. The linear system

$$a_1x + b_1y = c_1$$
$$a_2x + b_2y = c_2$$
$$a_3x + b_3y = c_3$$

of three equations in two unknowns represents three lines L_1 , L_2 , and L_3 in the *xy*-plane. Figure 3.1.5 shows six possible configurations of these three lines. In each case describe the solution set of the system.

34. Consider the linear system

 $a_1x + b_1y + c_1z = d_1$ $a_2x + b_2y + c_2z = d_2$ $a_3x + b_3y + c_3z = d_3$

of three equations in three unknowns to represent three planes P_1 , P_2 , and P_3 in *xyz*-space. Describe the solution set of the system in each of the following cases.

- (a) The three planes are parallel and distinct.
- (b) The three planes coincide— $P_1 = P_2 = P_3$.
- (c) P_1 and P_2 coincide and are parallel to P_3 .
- (d) P_1 and P_2 intersect in a line L that is parallel to P_3 .
- (e) P_1 and P_2 intersect in a line L that lies in P_3 .
- (f) P_1 and P_2 intersect in a line L that intersects P_3 in a single point.



FIGURE 3.1.5. Three lines in the plane (Problem 33).

3.2 Matrices and Gaussian Elimination

In Example 6 of Section 3.1 we applied the method of elimination to solve the linear system

$$1x + 2y + 1z = 4$$

$$3x + 8y + 7z = 20$$

$$2x + 7y + 9z = 23.$$
(1)

There we employed elementary operations to transform this system into the equivalent system

$$1x + 2y + 1z = 4$$

$$0x + 1y + 2z = 4$$

$$0x + 0y + 1z = 3,$$

(2)

which we found easy to solve by back substitution. Here we have printed in color the coefficients and constants (including the 0s and 1s that would normally be omitted) because everything else—the symbols x, y, and z for the variables and the + and = signs—is excess baggage that means only extra writing, for we can keep track of these symbols mentally. In effect, in Example 6 we used an appropriate sequence of operations to transform the array

$$\begin{bmatrix} 1 & 2 & 1 & 4 \\ 3 & 8 & 7 & 20 \\ 2 & 7 & 9 & 23 \end{bmatrix}$$
(3)

of coefficients and constants in (1) into the array

$$\begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$
(4)

of constants and coefficients in (2).

3.2 MATRICES AND GAUSSIAN ELIMINATION 147

Rectangular arrays of numbers like those in (3) and (4) are called *matrices*. Thus, a **matrix** is simply a rectangular array of numbers, which are called the **entries** or **elements** of the matrix. The **size**, or **shape**, of a matrix is specified by telling how many horizontal **rows** and vertical **columns** it has. Each matrix in (3) and (4) has three rows and four columns; therefore, each is a 3×4 (read "three by four") matrix. We *always* specify *first* the number of rows and *then* the number of columns.

Example 1 The matrices

$$\begin{bmatrix} 3 & -7 \\ -2 & 5 \end{bmatrix} \begin{bmatrix} 3 & 0 & -1 & 5 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ -3 \end{bmatrix}$$

have sizes 2×2 , 1×4 , and 3×1 , respectively.

Coefficient Matrices

A general system of *m* linear equations in the *n* variables $x_1, x_2, ..., x_n$ may be written in the form

$$a_{11}x_{1} + a_{12}x_{2} + a_{13}x_{3} + \dots + a_{1n}x_{n} = b_{1}$$

$$a_{21}x_{1} + a_{22}x_{2} + a_{23}x_{3} + \dots + a_{2n}x_{n} = b_{2}$$

$$\vdots$$

$$a_{m1}x_{1} + a_{m2}x_{2} + a_{m3}x_{3} + \dots + a_{mn}x_{n} = b_{m}.$$
(5)

Observe that a_{ij} denotes the (constant) coefficient in the *i*th equation of the *j*th variable x_j and that b_i denotes the constant on the right-hand side in the *i*th equation. Thus the first subscript *i* refers to the equation and the second subscript *j* to the variable. For instance, a_{32} denotes the coefficient of x_2 in the 3rd equation. This scheme of systematic double subscripts enables us to specify readily the location of each coefficient in a system with many equations and variables. It also helps us get rid of the excess baggage in (5) by focusing our attention on the coefficients.

The **coefficient matrix** of the linear system in (5) is the $m \times n$ matrix

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}.$$
 (6)

We will use **boldface** capital letters to denote matrices and lowercase letters to denote numbers. When we want to refer more briefly to the matrix **A** and its entries, we can write

$$\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix} = \begin{bmatrix} \vdots \\ \cdots & a_{ij} & \cdots \\ \vdots \\ (j \text{ th column}) \end{bmatrix} (i \text{ th row})$$
(7)

The first subscript *i* specifies the row and the second subscript *j* the column of **A** in which the element a_{ij} appears:

First subscript	Second subscript	
Row	Column	

Example 2

Although double subscripts may seem tedious when first encountered, their usefulness should not be underestimated. For instance, they are consistent with the notation used in most programming languages. Many practical applications lead to linear systems with hundreds or even thousands of variables and equations. In a typical computer system, the coefficient matrix **A** of such a system would be represented by a two-dimensional array in which A(i, j) denotes a_{ij} . With this approach, the computer can store a 100×100 matrix in the same way that it stores a 3×3 matrix.

The matrices in (3) and (4) include not only the coefficients but also the constants on the right-hand sides in the corresponding linear systems in (1) and (2). Let us write

$$\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} \tag{8}$$

for the column of constants in the general system in (5). An $m \times 1$ matrix—that is, one with a single column—is often called a (**column**) vector and is denoted by a boldface letter. When we adjoin the constant vector **b** to the coefficient matrix **A** (as a final column), we get the matrix

$$\begin{bmatrix} \mathbf{A} & \mathbf{b} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix}.$$
 (9)

This $m \times (n + 1)$ matrix is called the **augmented coefficient matrix**, or simply the augmented matrix, of the $m \times n$ system in (5).

Although it takes a good deal of notation to describe the augmented matrix of a *general* linear system, it is a very simple matter to write the augmented matrix of a *particular* system. If the system is written in chalk on a blackboard with the variables in the same order in each equation, we merely erase all the x_j 's and the plus and equal signs (retaining a minus sign for each negative coefficient) and insert a zero in each spot where a variable is missing in an equation.

The augmented coefficient matrix of the system

$$2x_1 + 3x_2 - 7x_3 + 4x_4 = 6$$

$$x_2 + 3x_3 - 5x_4 = 0$$

$$-x_1 + 2x_2 - 9x_4 = 17$$

of three equations in four variables is the 3×5 matrix

Γ	2	3	-7	4	6]	
	0	1	3	-5	0	
L	-1	2	0	-9	17	

Elementary Row Operations

In Section 3.1 we described the three elementary operations that are used in the method of elimination. To each of these corresponds an *elementary row operation* on the augmented matrix of the system. For instance, when we interchange two equations in the system, we interchange the corresponding two rows of its augmented coefficient matrix.

DEFINITION Elementary Row Operations

The following are the three types of **elementary row operations** on the matrix **A**:

- 1. Multiply any (single) row of A by a nonzero constant.
- 2. Interchange two rows of A.
- 3. Add a constant multiple of one row of A to another row.

Figure 3.2.1 shows notation that we use to describe elementary row operations briefly. For instance,

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \xrightarrow{(3)R_1 + R_2} \begin{bmatrix} 1 & 2 \\ 6 & 10 \end{bmatrix}$$
(10)

shows the result of adding 3 times row 1 to row 2. Note that, when we perform the operation $(c)R_p + R_q$ on a matrix—that is, when we add c times row p to row q—row p itself remains unchanged.

Туре	Row Operation	Notation		
1	Multiply row p by c	cR_p		
2	Interchange row p and row q	$SWAP(R_p, R_q)$		
3	Add c times row p to row q	$(c)R_p + R_q$		

FIGURE 3.2.1. Notation for elementary row operations.

Example 3 To solve the system

$$x_1 + 2x_2 + x_3 = 4$$

$$3x_1 + 8x_2 + 7x_3 = 20$$

$$2x_1 + 7x_2 + 9x_3 = 23,$$

(11)

whose augmented coefficient matrix is exhibited in (3), we carry out the following sequence of elementary row operations, corresponding to the steps in the solution of Example 6 in Section 3.1:

$$\begin{bmatrix} 1 & 2 & 1 & 4 \\ 3 & 8 & 7 & 20 \\ 2 & 7 & 9 & 23 \end{bmatrix}$$
(12)

$$\xrightarrow{(-3)R_1 + R_2} \qquad \begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 2 & 4 & 8 \\ 2 & 7 & 9 & 23 \end{bmatrix}$$

$$\xrightarrow{(-2)R_1 + R_3} \qquad \begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 2 & 4 & 8 \\ 0 & 3 & 7 & 15 \end{bmatrix}$$

$$\xrightarrow{\left(\frac{1}{2}\right)R_2} \qquad \begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 1 & 2 & 4 \\ 0 & 3 & 7 & 15 \end{bmatrix}$$

$$\xrightarrow{(-3)R_2 + R_3} \qquad \begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 3 \end{bmatrix}.$$
(13)

The final matrix here is the augmented coefficient matrix of the system

$$x_1 + 2x_2 + x_3 = 4 x_2 + 2x_3 = 4. x_3 = 3$$
(14)

whose unique solution (readily found by back substitution) is $x_1 = 5$, $x_2 = -2$, $x_3 = 3$.

It is not quite self-evident that a sequence of elementary row operations produces a linear system having the same solution set as the original system. To state the pertinent result concisely, we need the following definition.

DEFINITION Row-Equivalent Matrices

Two matrices are called **row equivalent** if one can be obtained from the other by a (finite) sequence of elementary row operations.

Thus the two matrices in (10) are row equivalent, as are the two matrices in (12) and (13). In Problem 29 we ask you to show that if **B** can be obtained from **A** by elementary row operations, then **A** can be obtained from **B** by elementary row operations. This follows from the observation that row operations are "reversible." In Problem 30 we suggest a proof of the following theorem.

THEOREM 1 Equivalent Systems and Equivalent Matrices

If the augmented coefficient matrices of two linear systems are row equivalent, then the two systems have the same solution set.

Thus, the linear systems in (11) and (14) have the same solution set, because their augmented matrices in (12) and (13) are row equivalent.

Echelon Matrices

Up to this point we have been somewhat informal in our description of the method of elimination. Its objective is to transform a given linear system, by elementary row operations, into one for which back substitution leads easily and routinely to a solution. The following definition tells what should be the appearance of the augmented coefficient matrix of the transformed system.

DEFINITION Echelon Matrix

The matrix \mathbf{E} is called an **echelon matrix** provided it has the following two properties:

- **1.** Every row of **E** that consists entirely of zeros (if any) lies *beneath* every row that contains a nonzero element.
- 2. In each row of **E** that contains a nonzero element, the *first* nonzero element lies strictly to the *right* of the first nonzero element in the preceding row (if there is a preceding row).

Echelon matrices are sometimes called *row-echelon matrices*. Property 1 says that if **E** has any all-zero rows, then they are grouped together at the bottom of the matrix. The first (from the left) *nonzero* element in each of the other rows is called its **leading entry**. Property 2 says that the leading entries form a "descending

staircase" pattern from upper left to lower right, as in the following echelon matrix.

$$\mathbf{E} = \begin{bmatrix} \mathbf{2} & -1 & 0 & 4 & 7\\ 0 & \mathbf{1} & \mathbf{2} & 0 & -5\\ 0 & 0 & 0 & \mathbf{3} & 0\\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Here we have highlighted the leading entries and indicated the descending staircase pattern.

The following list exhibits *all* possibilities for the form of a 3×3 echelon matrix, with asterisks denoting the (nonzero) leading entries; a # denotes an element that may be either zero or nonzero.

$\begin{bmatrix} * \\ 0 \\ 0 \end{bmatrix}$	# * 0	# # *	$\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$			
$\begin{bmatrix} * \\ 0 \\ 0 \end{bmatrix}$	# * 0	# # 0	$\begin{bmatrix} * & \# \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$	# * 0	$\begin{bmatrix} 0\\0\\0 \end{bmatrix}$	* 0 0	# * 0
$\begin{bmatrix} \ast \\ 0 \\ 0 \end{bmatrix}$	# 0 0	$\begin{bmatrix} \# \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \ast \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$	# 0 0	$\left[\begin{array}{c} 0\\ 0\\ 0\\ 0\end{array}\right]$	0 0 0	* 0 0

It follows from Properties 1 and 2 that the elements *beneath* any leading entry in the same column are all zero. The matrices

	[1]	3	-2			0	1	5
$\mathbf{A} =$	0	0	0	and	B =	1	3	-2
	0	1	5			0	0	0_

are not echelon matrices because **A** does not have Property 1 and **B** does not have Property 2.

Suppose that a linear system is in **echelon form**—its augmented matrix is an echelon matrix. Then those variables that correspond to *columns* containing leading entries are called **leading variables**; all the other variables are called **free variables**. The following algorithm describes the process of **back substitution** to solve such a system.

ALGORITHM Back Substitution

To solve a consistent linear system in echelon form by back substitution, carry out the following steps.

- **1.** Set each free variable equal to an arbitrary parameter (a different parameter for each free variable).
- 2. Solve the *final* (nonzero) equation for its leading variable.
- **3.** Substitute the result in the next-to-last equation and then solve for its leading variable.
- **4.** Continuing in this fashion, work upward through the system of equations until all variables have been determined.

Example 4 The augmented coefficient matrix of the system

$$x_{1} - 2x_{2} + 3x_{3} + 2x_{4} + x_{5} = 10$$

$$x_{3} + 2x_{5} = -3$$

$$x_{4} - 4x_{5} = 7$$
(15)

is the echelon matrix

1	-2	3	2	1	10		
0	0	1	0	2	-3	. (1	6)
0	0	0	1	-4	7		

The leading entries are in the first, third, and fourth columns. Hence x_1 , x_3 , and x_4 are the leading variables and x_2 and x_5 are the free variables. To solve the system by back substitution, we first write

$$x_2 = s, \quad x_5 = t,$$
 (17a)

where s and t are arbitrary parameters. Then the third equation in (15) gives

$$x_4 = 7 + 4x_5 = 7 + 4t; \tag{17b}$$

the second equation gives

$$x_3 = -3 - 2x_5 = -3 - 2t; \tag{17c}$$

finally, the first equation in (15) yields

$$x_1 = 10 + 2x_2 - 3x_3 - 2x_4 - x_5$$

= 10 + 2s - 3(-3 - 2t) - 2(7 + 4t) - t.

Therefore,

$$x_1 = 5 + 2s - 3t. \tag{17d}$$

Thus the system in (15) has an infinite solution set consisting of all $(x_1, x_2, x_3, x_4, x_5)$ given in terms of the two parameters *s* and *t* as follows:

$$x_1 = 5 + 2s - 3t$$

$$x_2 = s$$

$$x_3 = -3 - 2t$$

$$x_4 = 7 + 4t$$

$$x_5 = t.$$

For instance, with s = 2 and t = 1 we get the solution $x_1 = 6$, $x_2 = 2$, $x_3 = -5$, $x_4 = 11$, and $x_5 = 1$.

Gaussian Elimination

Because we can use back substitution to solve any linear system already in echelon form, it remains only to establish that we can transform any matrix (using elementary row operations) into an echelon matrix. The procedure that we have already illustrated in several examples is known as **Gaussian elimination**. The following systematic description of the procedure makes it clear that it succeeds with *any* given matrix **A**.

ALGORITHM Gaussian Elimination

- 1. Locate the first column of A that contains a nonzero element.
- 2. If the first (top) entry in this column is zero, interchange the first row of A with a row in which the corresponding entry is nonzero.
- **3.** Now the first entry in our column is nonzero. Replace the entries below it in the same column with zeros by adding appropriate multiples of the first row of **A** to lower rows.
- **4.** After Steps 1–3, the matrix looks like the matrix below, although there may be several initial columns of zeros or even none at all. Perform Steps 1–3 on the indicated lower right matrix A_1 .
- 5. Repeat this cycle of steps until an echelon matrix is obtained.

$$\mathbf{A} = \begin{bmatrix} 0 & \ast & \# & \# & \cdots & \# \\ 0 & 0 & & & \\ 0 & 0 & & & \mathbf{A}_1 \\ \vdots & \vdots & & & \\ 0 & 0 & & & & \end{bmatrix}$$
(17)

In brief, we work on the matrix **A** one column at a time, from left to right. In each column containing a leading entry (perhaps after a row interchange), we "clear out" the nonzero elements below it and then move on to the next column.

Example 5	To solve the system	
	$x_1 - 2x_2 + 3x_3 + 2x_4 + x_5 = 10$	
	$2x_1 - 4x_2 + 8x_3 + 3x_4 + 10x_5 = 7$	(18)
	$3x_1 - 6x_2 + 10x_3 + 6x_4 + 5x_5 = 27,$	
	we reduce its augmented coefficient matrix to echelon form as follows.	
	$\begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10 \\ 2 & -4 & 8 & 3 & 10 & 7 \\ 3 & -6 & 10 & 6 & 5 & 27 \end{bmatrix}$	
	$\xrightarrow{(-2)R_1 + R_2} \begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10\\ 0 & 0 & 2 & -1 & 8 & -13\\ 3 & -6 & 10 & 6 & 5 & 27 \end{bmatrix}$	
	$\xrightarrow{(-3)R_1 + R_3} \begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10\\ 0 & 0 & 2 & -1 & 8 & -13\\ 0 & 0 & 1 & 0 & 2 & -3 \end{bmatrix}$	
	$\xrightarrow{\text{SWAP}(R_2, R_3)} \begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10 \\ 0 & 0 & 1 & 0 & 2 & -3 \\ 0 & 0 & 2 & -1 & 8 & -13 \end{bmatrix}$	
	$\xrightarrow{(-2)R_2 + R_3} \begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10\\ 0 & 0 & 1 & 0 & 2 & -3\\ 0 & 0 & 0 & -1 & 4 & -7 \end{bmatrix}$	
	$\xrightarrow{(-1)R_3} \begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10\\ 0 & 0 & 1 & 0 & 2 & -3\\ 0 & 0 & 0 & 1 & -4 & 7 \end{bmatrix}$	

Our final result is the echelon matrix in (16), so by Eqs. (17a)-(17d) in Example 4, the infinite solution set of the system in (18) is described in terms of arbitrary parameters *s* and *t*

as follows:

$$x_{1} = 5 + 2s - 3t$$

$$x_{2} = s$$

$$x_{3} = -3 - 2t$$

$$x_{4} = 7 + 4t$$

$$x_{5} = t.$$
(19)

Thus, the substitution of any two specific values for *s* and *t* in (19) yields a particular solution $(x_1, x_2, x_3, x_4, x_5)$ of the system, and each of the system's infinitely many different solutions is the result of some such substitution.

Examples 3 and 5 illustrate the ways in which Gaussian elimination can result in either a unique solution or infinitely many solutions. On the other hand, if the reduction of the augmented matrix to echelon form leads to a row of the form

$$0 \ 0 \ \cdots \ 0 \ 0 \ *$$

where the asterisk denotes a nonzero entry in the last column, then we have an inconsistent equation,

$$0x_1 + 0x_2 + \dots + 0x_n = *,$$

and therefore the system has no solution.

10

Remark We use algorithms such as the back substitution and Gaussian elimination algorithms of this section to outline the basic computational procedures of linear algebra. In modern numerical work, these procedures often are implemented on a computer. For instance, linear systems of more than four equations are usually solved in practice by using a computer to carry out the process of Gaussian elimination.

3.2 Problems

The linear systems in Problems 1–10 are in echelon form. Solve each by back substitution.

1. $x_1 + x_2 + 2x_3 = 5$ $x_2 + 3x_3 = 6$ **2.** $2x_1 - 5x_2 + x_3 = 2$ $3x_2 - 2x_3 = 9$ $x_3 = 2$ $x_3 = -3$ **3.** $x_1 - 3x_2 + 4x_3 = 7$ 4. $x_1 - 5x_2 + 2x_3 = 10$ $x_2 - 5x_3 = 2$ $x_2 - 7x_3 = 5$ 5. $x_1 + x_2 - 2x_3 + x_4 = 9$ $x_2 - x_3 + 2x_4 = 1$ $x_3 - 3x_4 = 5$ 6. $x_1 - 2x_2 + 5x_3 - 3x_4 = 7$ $x_2 - 3x_3 + 2x_4 = 3$ $x_4 = -4$ 7. $x_1 + 2x_2 + 4x_3 - 5x_4 = 17$ $x_2 - 2x_3 + 7x_4 = 7$ 8. $x_1 - 10x_2 + 3x_3 - 13x_4 = 5$ $x_3 + 3x_4 = 10$ **9.** $2x_1 + x_2 + x_3 + x_4 = 6$ $3x_2 - x_3 - 2x_4 = 2$ $3x_3 + 4x_4 = 9$ $x_4 = 6$

In Problems 11–22, use elementary row operations to transform each augmented coefficient matrix to echelon form. Then solve the system by back substitution.

11. $2x_1 + 8x_2 + 3x_3 = 2$ 12. $3x_1 + x_2 - 3x_3 = 6$ $x_1 + 3x_2 + 2x_3 = 5$ $2x_1 + 7x_2 + x_3 = -9$ $2x_1 + 7x_2 + 4x_3 = 8$ $2x_1 + 5x_2 = -5$ 14. $3x_1 - 6x_2 - 2x_3 = 1$ 13. $x_1 + 3x_2 + 3x_3 = 13$ $2x_1 + 5x_2 + 4x_3 = 23$ $2x_1 - 4x_2 + x_3 = 17$ $2x_1 + 7x_2 + 8x_3 = 29$ $x_1 - 2x_2 - 2x_3 = -9$ **15.** $3x_1 + x_2 - 3x_3 = -4$ **16.** $2x_1 + 5x_2 + 12x_3 = 6$ $x_1 + x_2 + x_3 = 1$ $3x_1 + x_2 + 5x_3 = 12$ $5x_1 + 6x_2 + 8x_3 = 8$ $5x_1 + 8x_2 + 21x_3 = 17$ 17. $x_1 - 4x_2 - 3x_3 - 3x_4 = 4$ $2x_1 - 6x_2 - 5x_3 - 5x_4 =$ 5 $3x_1 - x_2 - 4x_3 - 5x_4 = -7$ **18.** $3x_1 - 6x_2 + x_3 + 13x_4 = 15$ $3x_1 - 6x_2 + 3x_3 + 21x_4 = 21$

$$2x_1 - 4x_2 + 5x_3 + 26x_4 = 23$$

3.2 MATRICES AND GAUSSIAN ELIMINATION 155

19. $3x_1 + x_2 + x_3 + 6x_4 = 14$ $x_1 - 2x_2 + 5x_3 - 5x_4 = -7$ $4x_1 + x_2 + 2x_3 + 7x_4 = 17$ **20.** $2x_1 + 4x_2 - x_3 - 2x_4 + 2x_5 = 6$ $x_1 + 3x_2 + 2x_3 - 7x_4 + 3x_5 = 9$ $5x_1 + 8x_2 - 7x_3 + 6x_4 + x_5 = 4$ = 6 **21.** $x_1 + x_2 + x_3$ $2x_1 - 2x_2 - 5x_3$ = -13 $3x_1 + x_3 + x_4 = 13$ $4x_1 - 2x_2 - 3x_3 + x_4 = 1$ **22.** $4x_1 - 2x_2 - 3x_3 + x_4 =$ 3 $2x_1 - 2x_2 - 5x_3 = -10$ $4x_1 + x_2 + 2x_3 + x_4 = 17$ $3x_1$ $+ x_3 + x_4 = 12$

In Problems 23–27, determine for what values of k each system has (a) a unique solution; (b) no solution; (c) infinitely many solutions.

- **23.** 3x + 2y = 1**24.** 3x + 2y = 06x + ky = 06x + 4y = k
- **25.** 3x + 2y = 11**26.** 3x + 2y = 17x + 5y = k
- 6x + ky = 21**27.** x + 2y + z = 32x - y - 3z = 54x + 3y - z = k
- **28.** Under what condition on the constants a, b, and c does the system

$$2x - y + 3z = a$$
$$x + 2y + z = b$$
$$7x + 4y + 9z = c$$

have a unique solution? No solution? Infinitely many solutions?

- **29.** This problem deals with the reversibility of elementary row operations.
 - (a) If the elementary row operation cR_p changes the matrix **A** to the matrix **B**, show that $(1/c)R_p$ changes **B** to A.
 - (b) If SWAP (R_p, R_q) changes A to B, show that $SWAP(R_p, R_q)$ also changes **B** to **A**.
 - (c) If $cR_p + R_q$ changes A to B, show that $(-c)R_p + R_q$ changes **B** to **A**.
 - (d) Conclude that if A can be transformed into B by a finite sequence of elementary row operations, then **B** can similarly be transformed into A.
- **30.** This problem outlines a proof that two linear systems LS_1 and LS_2 are equivalent (that is, have the same solution set) if their augmented coefficient matrices A_1 and A_2 are row equivalent.
 - (a) If a *single* elementary row operation transforms A_1 to A_2 , show directly—considering separately the three cases—that every solution of LS_1 is also a solution of LS_2 .
 - (b) Explain why it now follows from Problem 29 that every solution of either system is also a solution of the other system; thus the two systems have the same solution set.

3.2 Application Automated Row Reduction

Go to goo.gl/9szcxW to download this application's computing resources including Maple/Mathematica/MATLAB.

Computer algebra systems are often used to ease the labor of matrix computations, including elementary row operations. The 3×4 augmented coefficient matrix of Example 3 can be entered with the *Maple* command

```
with(linalg):
A := array([[1, 2, 1, 4]])
             [3, 8, 7, 20],
             [2, 7, 9, 23]]);
```

or the Mathematica command

 $A = \{\{1, 2, 1, 4\},\$ $\{3, 8, 7, 20\},\$ $\{2, 7, 9, 23\}\}$

or the MATLAB command

A = [1 2 1 4]3 8 7 20 2 7 9 231

The *Maple* linalg package has built-in elementary row operations that can be used to carry out the reduction of **A** exhibited in Example 3, as follows:



Α	:=	addrow(A,1,2,-3);	#	(-3)R1 + R2
Α	:=	addrow(A,1,3,-2);	#	(-2)R1 + R3
Α	:=	mulrow(A,2,1/2);	#	(1/2)R2
A	:=	addrow(A,2,3,-3);	#	(-3)R2 + R3

The inserted remarks, which follow the notation of Fig. 3.2.1, should make clear the structure of these row operations in *Maple*.

In *Mathematica* the *i*th row of the matrix **A** is denoted by **A**[[i]], and we can operate directly on rows to carry out the reduction of Example 3, as follows:

A[[2]] =	(-3)A[[1]] + A[[2]];	(*	(-3)R1 + R2	*)
A[[3]] =	(-2)A[[1]] + A[[3]];	(*	(-2)R1 + R3	*)
A[[2]] =	(1/2)A[[2]];	(*	(1/2)R2	*)
A[[3]] =	(-3)A[[2]] + A[[3]];	(*	(-3)R2 + R3	*)

In MATLAB, the *i*th row of the matrix **A** is denoted by **A(i,:)**, and we can operate similarly on rows to carry out the reduction to echelon form, as follows:

A(2,:)	= (-3)*A(1,:)	+ A(2,:)	%	(-3)R1 + R2
A(3,:)	= (-2)*A(1,:)	+ A(3,:)	%	(-2)R1 + R3
A(2,:)	= (1/2) * A(2,:)		%	(1/2)R2
A(3,:)	= (-3)*A(2,:)	+ A(3,:)	%	(-3)R2 + R3

You should verify (by using the computer algebra system of your choice) that these operations yield the echelon matrix

[1]	2	1	4	
0	1	2	4	.
0	0	1	3	

If you wanted to interchange the 1st and 3rd rows—and proceed to solve the corresponding system by "forward substitution" rather than back substitution—this could be done as follows:

<pre>A := swaprow(A,1,3);</pre>	(Maple)
$A[[{1,3}]] = A[[{3,1}]];$	(Mathematica)
A([1 3],:) = A([3 1],:)	(MATLAB)

These "automated" elementary row operations can be used in Problems 11–22 of this section.

3.3 Reduced Row-Echelon Matrices

The result of the process of Gaussian elimination described in Section 3.2 is not uniquely determined. That is, two different sequences of elementary row operations, both starting with the same matrix \mathbf{A} , may yield two different echelon matrices, each of course still row equivalent to \mathbf{A} . For instance, the two echelon matrices

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix} \text{ and } \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}$$
(1)

are readily seen (as in Problem 31) to be row equivalent. Hence it is possible to begin with an appropriate 3×3 matrix and derive by Gaussian elimination either of the two different echelon matrices in (1).

3.3 REDUCED ROW-ECHELON MATRICES 157

A full understanding of the structure of systems of linear equations depends on the definition of a special class of echelon matrices, a class having the property that every matrix in it is row equivalent to one and only one of these special echelon matrices. Recall that an echelon matrix \mathbf{E} is one that has the following two properties:

- **1.** Every all-zero row of **E** lies beneath every row that contains a nonzero element.
- **2.** The leading nonzero entry in each row lies to the right of the leading nonzero entry in the preceding row.

A *reduced* echelon matrix (sometimes called a reduced row-echelon matrix) has two additional properties.

DEFINITION Reduced Echelon Matrix

A **reduced echelon matrix E** is an echelon matrix that has—in addition to Properties 1 and 2—the following properties:

- **3.** Each leading entry of **E** is 1.
- 4. Each leading entry of E is the only nonzero element in its column.

A matrix is said to be in **reduced echelon form** if it is a reduced echelon matrix. Similarly, a linear system is in **reduced echelon form** if its augmented coefficient matrix is a reduced echelon matrix.

Example I	The following matrices are reduced echelon matrices.								
		$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} 1\\0\\0 \end{bmatrix}$	$\begin{array}{ccc} 0 & -3 \\ 1 & 4 \\ 0 & 0 \end{array}$	$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$				
		$\begin{bmatrix} 1 & -2 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$	$\begin{bmatrix} -7\\5\\0 \end{bmatrix}$				
	The echelon matrices								

	1	0	0]			[1]	0	2]
$\mathbf{A} =$	0	2	0	and	B =	0	1	0
	0	0	0			0	0	1

are not in *reduced* echelon form, because **A** does not have Property 3 and **B** does not have Property 4.

The process of transforming a matrix **A** into reduced echelon form is called **Gauss-Jordan elimination**.

ALGORITHM Gauss-Jordan Elimination

- 1. First transform A into echelon form by Gaussian elimination.
- **2.** Then divide each element of each nonzero *row* by its leading entry (to satisfy Property 3).
- **3.** Finally, use each leading 1 to "clear out" any remaining nonzero elements in its *column* (to satisfy Property 4).

The reduced echelon form of a matrix *is* unique. The special class of matrices that we mentioned at the beginning of this discussion is simply the class of all reduced echelon matrices. A proof of the following theorem may be found in Section

4.2 of B. Noble and J. W. Daniel, *Applied Linear Algebra*, 3rd ed., Hoboken, NJ: Pearson, 1988.

THEOREM 1 Unique Reduced Echelon Form

Every matrix is row equivalent to one and only one reduced echelon matrix.

Example 2 Find the reduced echelon form of the matrix

 $\mathbf{A} = \begin{bmatrix} 1 & 2 & 1 & 4 \\ 3 & 8 & 7 & 20 \\ 2 & 7 & 9 & 23 \end{bmatrix}.$

Solution In Example 3 of Section 3.2 we found the echelon form

[1]	2	1	4]
0	1	2	4	
0	0	1	3_	

which already satisfies Property 3. To clear out columns 2 and 3 (in order to satisfy Property 4), we continue the reduction as follows.

$$\begin{bmatrix} 1 & 2 & 1 & 4 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 3 \end{bmatrix} \xrightarrow{(-2)R_2 + R_1} \begin{bmatrix} 1 & 0 & -3 & -4 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$
$$\xrightarrow{(-2)R_3 + R_2} \begin{bmatrix} 1 & 0 & -3 & -4 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$
$$\xrightarrow{(3)R_3 + R_1} \begin{bmatrix} 1 & 0 & 0 & 5 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{bmatrix}$$

For instance, we see immediately from this reduced echelon form that the linear system

$$x + 2y + z = 4$$

 $3x + 8y + 7z = 20$
 $2x + 7y + 9z = 23$

with augmented coefficient matrix **A** has the unique solution x = 5, y = -2, z = 3.

Example 3 To use Gauss-Jordan elimination to solve the linear system

$$x_1 + x_2 + x_3 + x_4 = 12$$

$$x_1 + 2x_2 + 5x_4 = 17$$

$$3x_1 + 2x_2 + 4x_3 - x_4 = 31,$$
(2)

we transform its augmented coefficient matrix into reduced echelon form, as follows:

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 12\\ 1 & 2 & 0 & 5 & 17\\ 3 & 2 & 4 & -1 & 31 \end{bmatrix} \xrightarrow{(-1)R_1 + R_2} \begin{bmatrix} 1 & 1 & 1 & 1 & 12\\ 0 & 1 & -1 & 4 & 5\\ 3 & 2 & 4 & -1 & 31 \end{bmatrix} \xrightarrow{(-3)R_1 + R_3} \begin{bmatrix} 1 & 1 & 1 & 1 & 12\\ 0 & 1 & -1 & 4 & 5\\ 0 & -1 & 1 & -4 & -5 \end{bmatrix}$$

3.3 REDUCED ROW-ECHELON MATRICES159

$$\begin{array}{cccc} \underbrace{(1)R_2 + R_3} \\ \hline \end{array} & \begin{bmatrix} 1 & 1 & 1 & 1 & 12 \\ 0 & 1 & -1 & 4 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \\ \\ \underbrace{(-1)R_2 + R_1} \\ \hline \end{array} & \begin{bmatrix} 1 & 0 & 2 & -3 & 7 \\ 0 & 1 & -1 & 4 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Thus the reduced echelon form of the system in (2) is

$$\begin{aligned} x_1 &+ 2x_3 - 3x_4 &= 7 \\ x_2 - x_3 + 4x_4 &= 5 \\ 0 &= 0. \end{aligned}$$
 (3)

The leading variables are x_1 and x_2 ; the free variables are x_3 and x_4 . If we set

$$x_3 = s$$
 and $x_4 = t$,

then (3) immediately yields

$$x_1 = 7 - 2s + 3t, x_2 = 5 + s - 4t.$$

As a practical matter, Gauss-Jordan elimination generally offers no significant computational advantage over Gaussian elimination (transformation to nonreduced echelon form) followed by back substitution. Therefore, Gaussian elimination is commonly employed in practical procedures and in computer programs used to solve linear systems numerically.

The Three Possibilities

 x_{j_1}

The chief importance of Gauss-Jordan elimination stems from the fact that the reduced echelon form of a general linear system

$$a_{11}x_{1} + a_{12}x_{2} + a_{13}x_{3} + \dots + a_{1n}x_{n} = b_{1}$$

$$a_{21}x_{1} + a_{22}x_{2} + a_{23}x_{3} + \dots + a_{2n}x_{n} = b_{2}$$

$$\vdots$$

$$a_{m1}x_{1} + a_{m2}x_{2} + a_{m3}x_{3} + \dots + a_{mn}x_{n} = b_{m}$$
(4)

n

most clearly exhibits its underlying structure and enables us most readily to answer questions about the number and type of its solutions. If the leading variables are $x_{j_1}, x_{j_2}, \ldots, x_{j_r}$, then the reduced echelon form of the system in (4) looks like

$$+\sum_{k=r+1}^{n} c_{1k} x_{j_{k}} = d_{1}$$

$$x_{j_{2}} + \sum_{k=r+1}^{n} c_{2k} x_{j_{k}} = d_{2}$$

$$\vdots$$

$$x_{j_{r}} + \sum_{k=r+1}^{n} c_{rk} x_{j_{k}} = d_{r}$$

$$0 = d_{r+1}$$

$$\vdots$$

$$0 = d_{m},$$
(5)

where the summations involve only the (non-leading) free variables $x_{j_{r+1}}, \ldots, x_{j_n}$.

Now if any of the constants $d_{r+1}, d_{r+2}, \ldots, d_m$ in (5) is nonzero, then clearly the system has *no* real solution. On the other hand, suppose that $d_{r+1} = d_{r+2} = \cdots = d_m = 0$. If r < n, then there are free variables that we can set equal to arbitrary parameters, and so the system has *infinitely many* solutions. If instead r = n, then the summations in (5) disappear and we have the *unique* solution $x_1 = d_1, x_2 = d_2, \ldots, x_n = d_n$. These observations establish the following result, to which we have already alluded.

THEOREM 2 The Three Possibilities

A linear system of equations has either

- a unique solution, or
- no solution, or
- infinitely many solutions.

Homogeneous Systems

The linear system in (4) is called **homogeneous** provided that the constants b_1, b_2, \ldots, b_m on the right-hand side are all zero. Thus a homogeneous system of *m* equations in *n* variables has the form

$$a_{11}x_{1} + a_{12}x_{2} + a_{13}x_{3} + \dots + a_{1n}x_{n} = 0$$

$$a_{21}x_{1} + a_{22}x_{2} + a_{23}x_{3} + \dots + a_{2n}x_{n} = 0$$

$$\vdots$$

$$a_{m1}x_{1} + a_{m2}x_{2} + a_{m3}x_{3} + \dots + a_{mn}x_{n} = 0.$$
(6)

Every homogeneous system obviously has at least the trivial solution

$$x_1 = 0, \quad x_2 = 0, \dots, x_n = 0.$$
 (7)

Thus (by Theorem 2) we know from the outset that *every homogeneous linear system either has only the trivial solution or has infinitely many solutions*. If it has a *nontrivial* solution—one with not every x_i equal to zero—then it must have infinitely many solutions.

An important special case, in which a nontrivial solution is guaranteed, is that of a homogeneous system with *more variables than equations*: m < n. To see why there must be a solution, consider the reduced echelon system in (5) with the constants on the right-hand side all zero (because the original system is homogeneous). The number r of leading variables is at most the number m of equations (because there is at most one leading variable per equation). If m < n, it then follows that r < n, so there is at least one free variable that can be set equal to an arbitrary parameter, thereby yielding infinitely many solutions. This argument establishes the following key result.

THEOREM 3 Homogeneous Systems with More Variables than Equations

Every homogeneous linear system with more variables than equations has infinitely many solutions. **Example 4** The homogeneous linear system

 $47x_1 - 73x_2 + 56x_3 + 21x_4 = 0$ $19x_1 + 81x_2 - 17x_3 - 99x_4 = 0$ $53x_1 + 62x_2 + 39x_3 + 25x_4 = 0$

of three equations in four unknowns necessarily has infinitely many solutions. The only question (which we could answer by reducing the system to echelon form) is whether the system has one, two, or three free variables.

The situation is different for a *nonhomogeneous* system with more variables than equations. The simple example

$$x_1 + x_2 + x_3 = 0$$
$$x_1 + x_2 + x_3 = 1$$

shows that such a system may be inconsistent. But if it is consistent—meaning that $d_{r+1} = \cdots = d_m = 0$ in the reduced echelon form in (5)—then the fact that m < n implies (just as in the proof of Theorem 3) that there is at least one free variable, and that the system therefore has infinitely many solutions. Hence *every* nonhomogeneous system with more variables than equations either has no solution or has infinitely many solutions.

Equal Numbers of Equations and Variables

 x_1

An especially important case in the theory of linear systems is that of a homogeneous system

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = 0$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = 0$$

$$\vdots$$

$$a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n = 0$$

(8)

with the *same* number *n* of variables and equations. The coefficient matrix $\mathbf{A} = [a_{ij}]$ then has the same number of rows and columns and thus is an $n \times n$ square matrix.

Here we are most interested in the situation when (8) has *only* the trivial solution $x_1 = x_2 = \cdots = x_n = 0$. This occurs if and only if the reduced echelon system contains *no* free variables. That is, all *n* of the variables x_1, x_2, \ldots, x_n must be leading variables. Because the system consists of exactly *n* equations, we conclude that the reduced echelon system is simply

$$= 0$$

$$x_2 = 0$$

$$x_3 = 0$$

$$\therefore \qquad \vdots$$

$$x_n = 0,$$

and, therefore, that the reduced echelon form of the coefficient matrix A is the matrix

$$\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}.$$
(9)

Such a (square) matrix, with ones on its **principal diagonal** (the one from upper left to lower right) and zeros elsewhere, is called an **identity matrix** (for reasons given in Section 3.4). For instance, the 2×2 and 3×3 identity matrices are

	[1	0	0]	
and	0	1	0	
	0	0	1	

The matrix in (9) is the $n \times n$ identity matrix. With this terminology, the preceding argument establishes the following theorem.

THEOREM 4 Homogeneous Systems with Unique Solutions

Let **A** be an $n \times n$ matrix. Then the homogeneous system with coefficient matrix **A** has only the trivial solution if and only if **A** is row equivalent to the $n \times n$ identity matrix.

Example 5

The computation in Example 2 (disregarding the fourth column in each matrix there) shows that the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 8 & 7 \\ 2 & 7 & 9 \end{bmatrix}$$

is row equivalent to the 3×3 identity matrix. Hence Theorem 4 implies that the homogeneous system

$$x_1 + 2x_2 + x_3 = 0$$

$$3x_1 + 8x_2 + 7x_3 = 0$$

$$2x_1 + 7x_2 + 9x_3 = 0$$

with coefficient matrix **A** has only the trivial solution $x_1 = x_2 = x_3 = 0$.

3.3 **Problems**

Find the reduced echelon form of each of the matrices given in Problems 1–20.

1.	$\left[\begin{array}{rrr}1&2\\3&7\end{array}\right]$	2. $\begin{bmatrix} 3 & 7 \\ 2 & 5 \end{bmatrix}$
3.	$\begin{bmatrix} 3 & 7 & 15 \\ 2 & 5 & 11 \end{bmatrix}$	4. $\begin{bmatrix} 3 & 7 & -1 \\ 5 & 2 & 8 \end{bmatrix}$
5.	$\begin{bmatrix} 1 & 2 & -11 \\ 2 & 3 & -19 \end{bmatrix}$	6. $\begin{bmatrix} 1 & -2 & 19 \\ 4 & -7 & 70 \end{bmatrix}$
7.	$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 1 \\ 2 & 1 & 9 \end{bmatrix}$	$8. \begin{bmatrix} 1 & -4 & -5 \\ 3 & -9 & 3 \\ 1 & -2 & 3 \end{bmatrix}$
9.	$\begin{bmatrix} 5 & 2 & 18 \\ 0 & 1 & 4 \\ 4 & 1 & 12 \end{bmatrix}$	10. $\begin{bmatrix} 5 & 2 & -5 \\ 9 & 4 & -7 \\ 4 & 1 & -7 \end{bmatrix}$
11.	$\begin{bmatrix} 3 & 9 & 1 \\ 2 & 6 & 7 \\ 1 & 3 & -6 \end{bmatrix}$	12. $\begin{bmatrix} 1 & -4 & -2 \\ 3 & -12 & 1 \\ 2 & -8 & 5 \end{bmatrix}$
13.	$\begin{bmatrix} 2 & 7 & 4 & 0 \\ 1 & 3 & 2 & 1 \\ 2 & 6 & 5 & 4 \end{bmatrix}$	1 3 2 5 2 5 2 3 2 7 7 22

15.	2 1 2	-1 7	4 -4 19	$\frac{2}{3}$ -3		
16.	$\begin{bmatrix} 1\\ 2\\ 2 \end{bmatrix}$	3 1 4 2 7 3	15 22 34 1	7 8 7	-	
17.	$\begin{bmatrix} 1\\1\\2 \end{bmatrix}$	$ \begin{array}{c} 1 \\ -2 \\ 3 \end{array} $	$1 \\ -2 \\ -1$	-1 8 3	$\begin{bmatrix} -4 \\ -1 \\ 11 \end{bmatrix}$	
18.	$\begin{bmatrix} 1\\ 2\\ 2 \end{bmatrix}$	$-2 \\ 3 \\ 5$	-5 18 26	-12 11 21	2 1 I 9 I 11	
19.	$\begin{bmatrix} 2\\1\\1 \end{bmatrix}$	7 3 0	$-10 \\ -4 \\ 2$	— 1 — —	$\begin{bmatrix} 19 & 13 \\ -8 & 6 \\ 1 & 3 \end{bmatrix}$	
20.	3 5 2	6 10 4	1 8 5	7 18 9	13 47 26	

21–30. Use the method of Gauss-Jordan elimination (transforming the augmented matrix into reduced echelon form) to solve Problems 11–20 in Section 3.2.

3.3 REDUCED ROW-ECHELON MATRICES163

- **31.** Show that the two matrices in (1) are both row equivalent to the 3×3 identity matrix (and hence, by Theorem 1, to each other).
- **32.** Show that the 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

is row equivalent to the 2 × 2 identity matrix provided that $ad - bc \neq 0$.

- **33.** List all possible reduced row-echelon forms of a 2×2 matrix, using asterisks to indicate elements that may be either zero or nonzero.
- **34.** List all possible reduced row-echelon forms of a 3×3 matrix, using asterisks to indicate elements that may be either zero or nonzero.
- 35. Consider the homogeneous system

$$ax + by = 0$$
$$cx + dy = 0.$$

- (a) If $x = x_0$ and $y = y_0$ is a solution and k is a real number, then show that $x = kx_0$ and $y = ky_0$ is also a solution.
- (b) If $x = x_1$, $y = y_1$ and $x = x_2$, $y = y_2$ are both solutions, then show that $x = x_1 + x_2$, $y = y_1 + y_2$ is a solution.

- **36.** Suppose that $ad bc \neq 0$ in the homogeneous system of Problem 35. Use Problem 32 to show that its only solution is the trivial solution.
- **37.** Show that the homogeneous system in Problem 35 has a nontrivial solution if and only if ad bc = 0.
- **38.** Use the result of Problem 37 to find all values of c for which the homogeneous system

$$(c+2)x + 3y = 0$$

 $2x + (c-3)y = 0$

has a nontrivial solution.

- **39.** Consider a *homogeneous* system of three equations in three unknowns. Suppose that the third equation is the sum of some multiple of the first equation and some multiple of the second equation. Show that the system has a nontrivial solution.
- 40. Let E be an echelon matrix that is row equivalent to the matrix A. Show that E has the same number of nonzero rows as does the reduced echelon form E* of A. Thus the number of nonzero rows in an echelon form of A is an "invariant" of the matrix A. *Suggestion*: Consider reducing E to E*.

3.3 Application Automated Row Reduction



Go to goo.gl/7ryusP to download this application's computing resources including *Maple/Mathematica*/MATLAB. Most computer algebra systems include commands for the immediate reduction of matrices to reduced echelon form. For instance, if the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 1 & 4 \\ 3 & 8 & 7 & 20 \\ 2 & 7 & 9 & 23 \end{bmatrix}$$

of Example 2 has been entered—as illustrated in the 3.2 Application—then the *Maple* command

with(linalg): R := rref(A);

or the Mathematica command

R = RowReduce[A] // MatrixForm

or the MATLAB command

R = rref(A)

or the Wolfram Alpha query

row reduce ((1, 2, 1, 4), (3, 8, 7, 20), (2, 7, 9, 23))

produces the reduced echelon matrix

$$\mathbf{R} = \begin{bmatrix} 1 & 0 & 0 & 5\\ 0 & 1 & 0 & -2\\ 0 & 0 & 1 & 3 \end{bmatrix}$$

2 1 4 20 8 7 20 2 7 .9 23 0 0 5 Γ1 0 1 0 -2 rref(a) RAD AUTO FUNC 2/30

F3 F4 F4 F5 F5 F6 F6 UP

FIGURE 3.3.1. Finding a reduced row-echelon matrix with a TI-89 calculator.

that exhibits the solution of the linear system having augmented coefficient matrix **A**. The same calculation is illustrated in the calculator screen of Fig. 3.3.1. Solve

similarly the systems in the following problems (which apparently would be somewhat tedious to solve by manual row reduction):

1. 17x + 42y - 36z = 213**2.** 32x + 57y - 41z = 71313x + 45y - 34z = 22623x + 43y - 37z = 13012x + 47y - 35z = 19742x - 61v + 39z = 221**3.** 231x + 157y - 241z = 420**4.** 837x + 667y - 729z = 1659323x + 181y - 375z = 412152x - 179y - 975z = 1630542x + 328y - 759z = 1645542x + 161y - 759z = 4195. 49w - 57x + 37y - 59z = 976. 64w - 57x + 97y - 67z = 48573w - 15x - 19y - 22z = 9992w + 77x - 34y - 37z = 48652w - 51x + 14y - 29z = 8944w - 34x + 53y - 34z = 46513w - 27x + 27y - 25z = 7327w + 57x - 69v + 29z = 464

3.4 Matrix Operations

As yet, we have used matrices only to simplify our record keeping in the solution of linear systems. But it turns out that matrices can be added and multiplied in ways similar to the ways in which numbers are added and multiplied and that these operations with matrices have far-reaching applications.

At the level of this text everyone "knows" that 2 + 3 = 5, and we do not dwell on the underlying meaning of this equation. But in the case of matrices we must begin with precise definitions of what the familiar language of algebra is to mean when it is applied to matrices rather than to numbers.

Two matrices **A** and **B** of the same size—the same number of rows and the same number of columns—are called **equal** provided that each element of **A** is equal to the corresponding element of **B**. Thus two matrices of the same size are equal provided they are *elementwise equal*, and we write $\mathbf{A} = \mathbf{B}$ to denote equality of the two matrices **A** and **B**.

Example 1

If

If

$$\mathbf{A} = \begin{bmatrix} 3 & 4 \\ 5 & 6 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}, \quad \text{and} \quad \mathbf{C} = \begin{bmatrix} 3 & 4 & 7 \\ 5 & 6 & 8 \end{bmatrix}$$

then $\mathbf{A} \neq \mathbf{B}$ because $a_{22} = 6$, whereas $b_{22} = 7$, and $\mathbf{A} \neq \mathbf{C}$ because the matrices \mathbf{A} and \mathbf{C} are not of the same size.

The next two definitions are further examples of "doing it elementwise."

DEFINITION Addition of Matrices

If $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} b_{ij} \end{bmatrix}$ are matrices of the same size, then their sum $\mathbf{A} + \mathbf{B}$ is the matrix obtained by adding corresponding elements of the matrices \mathbf{A} and \mathbf{B} . That is,

$$\mathbf{A} + \mathbf{B} = \begin{bmatrix} a_{ij} + b_{ij} \end{bmatrix},\tag{1}$$

where the notation on the right signifies that the element in the *i*th row and *j*th column of the matrix $\mathbf{A} + \mathbf{B}$ is $a_{ij} + b_{ij}$.

Example 2

 $\mathbf{A} = \begin{bmatrix} 3 & 0 & -1 \\ 2 & -7 & 5 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 4 & -3 & 6 \\ 9 & 0 & -2 \end{bmatrix}, \text{ and } \mathbf{C} = \begin{bmatrix} 3 & -2 \\ -1 & 6 \end{bmatrix},$

then

$$\mathbf{A} + \mathbf{B} = \begin{bmatrix} 7 & -3 & 5\\ 11 & -7 & 3 \end{bmatrix},$$

but the sum A + C is not defined because the matrices A and C are not of the same size.

DEFINITION Multiplication of a Matrix by a Number

If $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ is a matrix and *c* is a number, then *c* \mathbf{A} is the matrix obtained by multiplying each element of \mathbf{A} by *c*. That is,

$$c\mathbf{A} = \begin{bmatrix} ca_{ij} \end{bmatrix}. \tag{2}$$

Using multiplication of a matrix by a scalar, we define the **negative** $-\mathbf{A}$ of the matrix \mathbf{A} and the **difference** $\mathbf{A} - \mathbf{B}$ of the two matrices \mathbf{A} and \mathbf{B} by writing

$$-\mathbf{A} = (-1)\mathbf{A}$$
 and $\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$.

Example 3 If **A** and **B** are the 2 × 3 matrices of Example 2, then $3\mathbf{A} = \begin{bmatrix} 9 & 0 & -3 \\ 6 & -21 & 15 \end{bmatrix}, \quad -\mathbf{B} = \begin{bmatrix} -4 & 3 & -6 \\ -9 & 0 & 2 \end{bmatrix},$

and

$$3\mathbf{A} - \mathbf{B} = \begin{bmatrix} 5 & 3 & -9 \\ -3 & -21 & 17 \end{bmatrix}.$$

Vectors

Our first application of these matrix operations is to vectors. As mentioned in Section 3.2, a **column vector** (or simply **vector**) is merely an $n \times 1$ matrix, one having a single column. We normally use boldface lowercase letters, rather than lightface uppercase letters, to denote vectors. If

$$\mathbf{a} = \begin{bmatrix} 6\\-2\\5 \end{bmatrix}$$
 and $\mathbf{b} = \begin{bmatrix} -2\\3\\-4 \end{bmatrix}$,

then we can form such combinations as

$$3\mathbf{a} + 2\mathbf{b} = \begin{bmatrix} 18\\-6\\15 \end{bmatrix} + \begin{bmatrix} -4\\6\\-8 \end{bmatrix} = \begin{bmatrix} 14\\0\\7 \end{bmatrix}.$$

Largely for typographical reasons, we sometimes write

$$\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = (a_1, a_2, \dots, a_n). \tag{3}$$

That is, $(a_1, a_2, ..., a_n)$ is simply another notation for the *column vector* with elements $a_1, a_2, ..., a_n$. It should not be confused with the *row vector*

$$\begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix}. \tag{4}$$
A row vector is a $1 \times n$ (rather than $n \times 1$) matrix having a single row, and

$$(3,2,1) = \begin{bmatrix} 3\\2\\1 \end{bmatrix} \neq \begin{bmatrix} 3 & 2 & 1 \end{bmatrix}$$

because the two matrices here have different sizes (even though they have the same elements).

Now consider the linear system

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n = b_m$$
(5)

of *m* equations in *n* variables. We may regard a solution of this system as a *vector*

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = (x_1, x_2, x_3, \dots, x_n) \tag{6}$$

whose elements satisfy each of the equations in (5). If we want to refer explicitly to the number of elements, we may call \mathbf{x} an *n*-vector.

Example 4 Consider the homogeneous system

 $x_1 + 3x_2 - 15x_3 + 7x_4 = 0$ $x_1 + 4x_2 - 19x_3 + 10x_4 = 0$ $2x_1 + 5x_2 - 26x_3 + 11x_4 = 0.$ (7)

We find readily that the reduced echelon form of the augmented coefficient matrix of this system is

 $\begin{bmatrix} 1 & 0 & -3 & -2 & 0 \\ 0 & 1 & -4 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$

Hence x_1 and x_2 are leading variables and x_3 and x_4 are free variables. In the manner of Sections 3.2 and 3.3, we therefore see that the infinite solution set of the system in (7) is described by the equations

$$x_{4} = t, x_{3} = s, x_{2} = 4s - 3t, x_{1} = 3s + 2t$$
(8)

in terms of the arbitrary parameters s and t.

Now let us write the solution $\mathbf{x} = (x_1, x_2, x_3, x_4)$ in vector notation. The equations in (8) yield

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3s + 2t \\ 4s - 3t \\ s \\ t \end{bmatrix},$$

and "separating" the s and t parts gives

$$\mathbf{x} = \begin{bmatrix} 3s \\ 4s \\ s \\ 0 \end{bmatrix} + \begin{bmatrix} 2t \\ -3t \\ 0 \\ t \end{bmatrix} = s \begin{bmatrix} 3 \\ 4 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} 2 \\ -3 \\ 0 \\ 1 \end{bmatrix}$$

-that is,

$$\mathbf{x} = s(3, 4, 1, 0) + t(2, -3, 0, 1) = s\mathbf{x}_1 + t\mathbf{x}_2.$$
(9)

Equation (9) expresses in **vector form** the general solution of the linear system in (7). It says that the vector **x** is a solution if and only if **x** is a *linear combination*—a sum of multiples—of the particular solutions $\mathbf{x}_1 = (3, 4, 1, 0)$ and $\mathbf{x}_2 = (2, -3, 0, 1)$. The parameters *s* and *t* are simply the coefficients in this "sum of multiples."

In the same manner as that in which we derived Eq. (9) from the equations in (8), the general solution of every homogeneous linear system can be expressed as a linear combination of particular solution vectors. For this reason (as well as others), linear combinations of vectors will play a central role in succeeding chapters.

Matrix Multiplication

The first surprise is that matrices are *not* multiplied elementwise. The initial purpose of matrix multiplication is to simplify the notation for systems of linear equations. If we write

$$\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}, \quad (10)$$

then \mathbf{A} , \mathbf{x} , and \mathbf{b} are, respectively, the coefficient matrix, the unknown vector, and the constant vector for the linear system in (5). We want to define the matrix product $\mathbf{A}\mathbf{x}$ in such a way that the entire system of linear equations reduces to the single matrix equation

$$\mathbf{A}\mathbf{x} = \mathbf{b}.\tag{11}$$

The first step is to define the product of a row vector **a** and a column vector **b**,

$$\mathbf{a} = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix}$$
 and $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$,

each having *n* elements. In this case, the product **ab** is *defined* to be

$$\mathbf{ab} = a_1 b_1 + a_2 b_2 + \dots + a_n b_n. \tag{12}$$

Thus **ab** is the *sum of products* of corresponding elements of **a** and **b**. For instance,

$$\begin{bmatrix} 2 & -3 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = (2)(3) + (-3)(5) = -9$$

and

$$\begin{bmatrix} 3 & 0 & -1 & 7 \end{bmatrix} \begin{bmatrix} 5 \\ 2 \\ -3 \\ 4 \end{bmatrix} = 3 \cdot 5 + 0 \cdot 2 + (-1)(-3) + 7 \cdot 4 = 46.$$

Note that if

$$\mathbf{a} = \begin{bmatrix} a_1 & a_2 & \cdots & a_n \end{bmatrix}$$
 and $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$

then

$$\mathbf{a}\mathbf{x} = a_1x_1 + a_2x_2 + \dots + a_nx_n.$$

Hence the single equation

$$a_1 x_1 + a_2 x_2 + \dots + a_n x_n = b \tag{13}$$

reduces to the equation

$$\mathbf{a}\mathbf{x} = b,\tag{14}$$

which is a step toward the objective expressed in Eq. (11). This observation is the underlying motivation for the following definition.

DEFINITION Matrix Multiplication

Suppose that **A** is an $m \times p$ matrix and **B** is a $p \times n$ matrix. Then the **product AB** is the $m \times n$ matrix defined as follows: The element of **AB** in its *i*th row and *j* th column is the *sum of products* of corresponding elements in the *i*th row of **A** and the *j* th column of **B**.

That is, if the *i*th row of **A** is

$$\begin{bmatrix} a_{i1} & a_{i2} & a_{i3} & \cdots & a_{ip} \end{bmatrix}$$

and the *j* th column of **B** is

$$\begin{bmatrix} b_{1j} \\ b_{2j} \\ b_{3j} \\ \vdots \\ b_{pj} \end{bmatrix},$$

then the element in the *i*th row and *j*th column of the product **AB** is

$$a_{i1}b_{1j} + a_{i2}b_{2j} + a_{i3}b_{3j} + \dots + a_{ip}b_{pj}.$$

Example 5

$$\mathbf{A} = \begin{bmatrix} 2 & -1 \\ -4 & 3 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} 1 & 5 \\ 3 & 7 \end{bmatrix},$$

then m = p = n = 2, so **AB** will also be a 2 × 2 matrix. To find **AB**, we calculate sums of products as follows:

AB, row 1, column 1 :
$$(2)(1) + (-1)(3) = -1$$
;
AB, row 1, column 2 : $(2)(5) + (-1)(7) = 3$;
AB, row 2, column 1 : $(-4)(1) + (3)(3) = 5$;
AB, row 2, column 2 : $(-4)(5) + (3)(7) = 1$.

Thus

If

$$\mathbf{AB} = \begin{bmatrix} -1 & 3\\ 5 & 1 \end{bmatrix}.$$

For your first practice with matrix multiplication, you should compute

$$\mathbf{BA} = \begin{bmatrix} 1 & 5 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -4 & 3 \end{bmatrix} = \begin{bmatrix} -18 & 14 \\ -22 & 18 \end{bmatrix}.$$

Note that $AB \neq BA$. This shows that multiplication of matrices is *not* commutative! We must therefore be careful about the order in which we write the matrices in a matrix product.

The definition of the matrix product bears careful examination to see how it fits together. First, the fact that **A** is $m \times p$ and **B** is $p \times n$ implies that the number of *columns* of **A** is equal to the number of *rows* of **B**. If so, then the size of the product **AB** is obtained by a sort of cancellation of the "inside" dimensions:



If the inside dimensions are not equal, then the product **AB** is *not defined*.

Example 6

If **A** is a 3×2 matrix and **B** is a 2×3 matrix, then **AB** will be a 3×3 matrix, whereas **BA** will be a 2×2 matrix. If **C** is a 3×5 matrix and **D** is a 5×7 matrix, then **CD** will be a 3×7 matrix, but **DC** is undefined.

To emphasize the fact that the *ij*th element of AB is the product of the *i*th row of A and the *j*th column of B, we can write



where $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$ denote the *m* row vectors of **A** and $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n$ denote the *n* column vectors of **B**. More briefly, if

$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_m \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \cdots & \mathbf{b}_n \end{bmatrix}$$

in terms of the rows of A and the columns of B, then

$$\mathbf{AB} = \begin{bmatrix} \mathbf{a}_i \mathbf{b}_j \end{bmatrix}. \tag{15}$$

Therefore, as mentioned earlier, the *ij* th element $\mathbf{a}_i \mathbf{b}_j$ of **AB** is given in terms of elements of **A** and **B** by

$$\mathbf{a}_i \mathbf{b}_j = \begin{bmatrix} a_{i1} & a_{i2} & \cdots & a_{ip} \end{bmatrix} \begin{bmatrix} b_{1j} \\ b_{2j} \\ \vdots \\ b_{pj} \end{bmatrix} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj}.$$

That is,

$$\mathbf{a}_i \mathbf{b}_j = \sum_{k=1}^p a_{ik} b_{kj}.$$
 (16)

One can visualize "pouring the *i*th row of **A** down the *j*th column of **B**" until elements match in pairs, then forming the sum of the products of these pairs, to obtain the element c_{ij} of the matrix $\mathbf{C} = \mathbf{AB}$.

Suggestion The key to accuracy and confidence in computing matrix products lies in doing it systematically. Always perform your computations in the same order. *First* calculate the elements of the *first* row of **AB** by multiplying the *first* row of **A** by the successive columns of **B**; *second*, calculate the elements of the *second* row of **AB** by multiplying the *second* row of **A** by the successive columns of **B**; and so forth.

Computing systems often are used for the calculation of products of "large" matrices. If the matrices **A** and **B**, with appropriate sizes, have been entered—as illustrated in the 3.2 Application—then the *Maple* command

with(linalg) : C := multiply(A,B),

or the Mathematica command

C = A.B,

or the MATLAB command

C = A*B

immediately yield the product matrix $\mathbf{C} = \mathbf{A}\mathbf{B}$.

Matrix Equations

If $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ is an $m \times n$ coefficient matrix and $\mathbf{x} = (x_1, x_2, \dots, x_n)$ is an $n \times 1$ variable (column) matrix, then the product $\mathbf{A}\mathbf{x}$ is the $m \times 1$ matrix

$$\mathbf{A}\mathbf{x} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$
$$= \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix} \stackrel{(?)}{=} \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = \mathbf{b}.$$

We therefore see that

$$\mathbf{x} = \mathbf{b} \tag{17}$$

if and only if $\mathbf{x} = (x_1, x_2, ..., x_n)$ is a solution of the linear system in (5). Thus, matrix multiplication enables us to "boil down" a system of *m* scalar equations in *n* unknowns to the single matrix equation in (17), which is analogous in notation to the single scalar equation ax = b in a single variable *x*.

A

Example 7

The system

$$3x_1 - 4x_2 + x_3 + 7x_4 = 10$$

$$4x_1 - 5x_3 + 2x_4 = 0$$

$$x_1 + 9x_2 + 2x_3 - 6x_4 = 5$$

of three equations in four unknowns is equivalent to the single matrix equation

$$\begin{bmatrix} 3 & -4 & 1 & 7 \\ 4 & 0 & -5 & 2 \\ 1 & 9 & 2 & -6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 10 \\ 0 \\ 5 \end{bmatrix}.$$

Matrix Algebra

The definitions of matrix addition and multiplication can be used to establish the rules of matrix algebra listed in the following theorem.

THEOREM 1 Rules of Matrix Algebra

If **A**, **B**, and **C** are matrices of appropriate sizes to make the indicated operations possible, then the following identities hold.

Commutative law of addition:	$\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$
Associative law of addition:	$\mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C}$
Associative law of multiplication:	$\mathbf{A}(\mathbf{B}\mathbf{C}) = (\mathbf{A}\mathbf{B})\mathbf{C}$
Distributive laws:	$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{A}\mathbf{B} + \mathbf{A}\mathbf{C}$
and	
	$(\mathbf{A} + \mathbf{B})\mathbf{C} = \mathbf{A}\mathbf{C} + \mathbf{B}\mathbf{C}$

The only verification that is not entirely routine is that of the associative law of multiplication; see Problem 44 for an outline. Each of the others follows quickly from the corresponding law for the ordinary arithmetic of real numbers. As an illustration, we prove the first distributive law. Suppose that $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ is an $m \times p$ matrix and that $\mathbf{B} = \begin{bmatrix} b_{ij} \end{bmatrix}$ and $\mathbf{C} = \begin{bmatrix} c_{ij} \end{bmatrix}$ are $p \times n$ matrices. Then

$$\mathbf{B} + \mathbf{C} = \left[b_{ij} + c_{ij} \right],$$

so by (16) the *ij*th element of the $m \times n$ matrix A(B + C) is

$$\sum_{k=1}^{p} a_{ik} (b_{kj} + c_{kj}).$$
(18)

The *ij*th element of the $m \times n$ matrix **AB** + **AC** is

$$\sum_{k=1}^{p} a_{ik} b_{kj} + \sum_{k=1}^{p} a_{ik} c_{kj} = \sum_{k=1}^{p} (a_{ik} b_{kj} + a_{ik} c_{kj}).$$
(19)

But the distributive law for real numbers, a(b + c) = ab + ac, tells us that corresponding terms of the sums in (18) and (19) are equal. Hence, the *ij*th terms of the two $m \times n$ matrices A(B + C) and AB + AC are equal, and so these matrices are equal: A(B + C) = AB + AC.

If a and b are real numbers, then rules such as

$$(a+b)\mathbf{C} = a\mathbf{C} + b\mathbf{C}, \quad (ab)\mathbf{C} = a(b\mathbf{C}), \quad a(\mathbf{B}\mathbf{C}) = (a\mathbf{B})\mathbf{C}$$

are even easier to verify. What all these rules amount to is this: In matrix manipulations, pairs of parentheses can be inserted or deleted in the same ways as in the ordinary algebra of real numbers. But not all of the rules of "ordinary" algebra carry over to matrix algebra. In Example 5 we saw that multiplication of matrices is *not* commutative—in general, $AB \neq BA$. Other exceptions are associated with zero matrices. A zero matrix is one whose elements are *all* zero, such as

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

We ordinarily denote a zero matrix (whatever its size) by **0**. It should be clear that for any matrix **A**,

$$\mathbf{0} + \mathbf{A} = \mathbf{A} = \mathbf{A} + \mathbf{0}, \quad \mathbf{A}\mathbf{0} = \mathbf{0}, \quad \text{and} \quad \mathbf{0}\mathbf{A} = \mathbf{0},$$

where in each case $\mathbf{0}$ is a zero matrix of appropriate size. Thus zero matrices appear to play a role in the arithmetic of matrices similar to the role of the real number 0 in ordinary arithmetic.

For real numbers, the following two rules are familiar:

- If ab = ac and $a \neq 0$, then b = c
 - (the "cancellation law").
- If ad = 0, then either a = 0 or d = 0.

The following example shows that matrices do not obey either of these rules.

Example 8

$$\mathbf{A} = \begin{bmatrix} 4 & 1 & -2 & 7 \\ 3 & 1 & -1 & 5 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 1 & 5 \\ 3 & -1 \\ -2 & 4 \\ 2 & -3 \end{bmatrix}, \quad \text{and} \quad \mathbf{C} = \begin{bmatrix} 3 & 4 \\ 2 & 1 \\ -2 & 3 \\ 1 & -3 \end{bmatrix},$$

then $\mathbf{B} \neq \mathbf{C}$, but

$$\mathbf{AB} = \begin{bmatrix} 25 & -10\\ 18 & -5 \end{bmatrix} = \mathbf{AC}. \quad \text{(Check this!)}$$

Thus the cancellation law does not generally hold for matrices. If

$$\mathbf{D} = \mathbf{B} - \mathbf{C} = \begin{bmatrix} -2 & 1\\ 1 & -2\\ 0 & 1\\ 1 & 0 \end{bmatrix},$$

then

If

$$\mathbf{A}\mathbf{D} = \begin{bmatrix} 0 & 0\\ 0 & 0 \end{bmatrix} = \mathbf{0}$$

despite the fact that neither **A** nor **D** is a zero matrix. See Problems 31–38 for additional ways in which the algebra of matrices differs significantly from the familiar algebra of real numbers.

Recall that an *identity matrix* is a square matrix I that has ones on its principal diagonal and zeros elsewhere. Identity matrices play a role in matrix arithmetic which is strongly analogous to that of the real number 1, for which $a \cdot 1 = 1 \cdot a = a$ for all values of the real number a. For instance, you can check that

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

Similarly, if

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \text{ and } \mathbf{I} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

0 7

then AI = IA = A. For instance, the element in the second row and third column of AI is

$$(a_{21})(0) + (a_{22})(0) + (a_{23})(1) = a_{23}.$$

If a is a nonzero real number and $b = a^{-1}$, then ab = ba = 1. Given a nonzero square matrix **A**, the question as to whether there exists an *inverse matrix* **B**, one such that AB = BA = I, is more complicated and is investigated in Section 3.5.

3.4 Problems

In Problems 1–4, two matrices **A** and **B** and two numbers *c* and *d* are given. Compute the matrix $c\mathbf{A} + d\mathbf{B}$.

1.
$$\mathbf{A} = \begin{bmatrix} 3 & -5 \\ 2 & 7 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} -1 & 0 \\ 3 & -4 \end{bmatrix}, c = 3, d = 4$$

2. $\mathbf{A} = \begin{bmatrix} 2 & 0 & -3 \\ -1 & 5 & 6 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} -2 & 3 & 1 \\ 7 & 1 & 5 \end{bmatrix}, c = 5, d = -3$
3. $\mathbf{A} = \begin{bmatrix} 5 & 0 \\ 0 & 7 \\ 3 & -1 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} -4 & 5 \\ 3 & 2 \\ 7 & 4 \end{bmatrix}, c = -2, d = 4$
4. $\mathbf{A} = \begin{bmatrix} 2 & -1 & 0 \\ 4 & 0 & -3 \\ 5 & -2 & 7 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 6 & -3 & -4 \\ 5 & 2 & -1 \\ 0 & 7 & 9 \end{bmatrix}, c = 7, d = 5$

In Problems 5–12, two matrices **A** and **B** are given. Calculate whichever of the matrices **AB** and **BA** is defined.

5.
$$\mathbf{A} = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} -4 & 2 \\ 1 & 3 \end{bmatrix}$$

6. $\mathbf{A} = \begin{bmatrix} 1 & 0 & -3 \\ 3 & 2 & 4 \\ 2 & -3 & 5 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 7 & -4 & 3 \\ 1 & 5 & -2 \\ 0 & 3 & 9 \end{bmatrix}$
7. $\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$
8. $\mathbf{A} = \begin{bmatrix} 1 & 0 & 3 \\ 2 & -5 & 4 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 3 \\ -1 & 4 \\ 6 & 5 \end{bmatrix}$
9. $\mathbf{A} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 0 & -2 \\ 3 & 1 \\ -4 & 5 \end{bmatrix}$
10. $\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} -1 & 0 & 4 \\ 3 & -2 & 5 \end{bmatrix}$
11. $\mathbf{A} = \begin{bmatrix} 3 & -5 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 2 & 7 & 5 & 6 \\ -1 & 4 & 2 & 3 \end{bmatrix}$
12. $\mathbf{A} = \begin{bmatrix} 1 & 0 & 3 & -2 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 2 & -7 & 5 \\ 3 & 9 & 10 \end{bmatrix}$

In Problems 13–16, three matrices \mathbf{A} , \mathbf{B} , and \mathbf{C} are given. Verify by computation of both sides the associative law $\mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}$.

13.
$$\mathbf{A} = \begin{bmatrix} 3 & 1 \\ -1 & 4 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 2 & 5 \\ -3 & 1 \end{bmatrix}, \mathbf{C} = \begin{bmatrix} 0 & 1 \\ 2 & 3 \end{bmatrix}$$

14. $\mathbf{A} = \begin{bmatrix} 2 & -1 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 2 & 5 \\ -3 & 1 \end{bmatrix}, \mathbf{C} = \begin{bmatrix} 6 \\ -5 \end{bmatrix}$

15.
$$\mathbf{A} = \begin{bmatrix} 3\\2 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}, \mathbf{C} = \begin{bmatrix} 2 & 0\\0 & 3\\1 & 4 \end{bmatrix}$$

16. $\mathbf{A} = \begin{bmatrix} 2 & 0\\0 & 3\\1 & 4 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 1 & -1\\3 & -2 \end{bmatrix},$
 $\mathbf{C} = \begin{bmatrix} 1 & 0 & -1 & 2\\3 & 2 & 0 & 1 \end{bmatrix}$

In Problems 17–22, first write each given homogeneous system in the matrix form $\mathbf{A}\mathbf{x} = \mathbf{0}$. Then find the solution in vector form, as in Eq. (9).

17.
$$x_1 - 5x_3 + 4x_4 = 0$$

 $x_2 + 2x_3 - 7x_4 = 0$
18. $x_1 - 3x_2 + 6x_4 = 0$
 $x_3 + 9x_4 = 0$
19. $x_1 + 3x_4 - x_5 = 0$
 $x_2 - 2x_4 + 6x_5 = 0$
 $x_3 + x_4 - 8x_5 = 0$
20. $x_1 - 3x_2 + 7x_5 = 0$
 $x_3 - 2x_5 = 0$
 $x_4 - 10x_5 = 0$
21. $x_1 - x_3 + 2x_4 + 7x_5 = 0$
 $x_2 + 2x_3 - 3x_4 + 4x_5 = 0$
22. $x_1 - x_2 + 7x_4 + 3x_5 = 0$
 $x_3 - x_4 - 2x_5 = 0$

Problems 23 through 26 introduce the idea—developed more fully in the next section—of a multiplicative inverse of a square matrix.

23. Let

$$\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

 $\mathbf{I} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$

and

Find **B** so that AB = I = BA as follows: First equate entries on the two sides of the equation AB = I. Then solve the resulting four equations for *a*, *b*, *c*, and *d*. Finally verify that BA = I as well.

24. Repeat Problem 23, but with A replaced by the matrix

$$\mathbf{A} = \left[\begin{array}{cc} 3 & 4 \\ 5 & 7 \end{array} \right]$$

25. Repeat Problem 23, but with A replaced by the matrix

$$\mathbf{A} = \begin{bmatrix} 5 & 7 \\ 2 & 3 \end{bmatrix}$$

26. Use the technique of Problem 23 to show that if

$$\mathbf{A} = \begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix}.$$

then there does *not* exist a matrix **B** such that AB = I. Suggestion: Show that the system of four equations in *a*, *b*, *c*, and *d* is inconsistent.

27. A diagonal matrix is a square matrix of the form

$$\begin{bmatrix} a_1 & 0 & 0 & \cdots & 0 \\ 0 & a_2 & 0 & \cdots & 0 \\ 0 & 0 & a_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_n \end{bmatrix}$$

in which every element off the main diagonal is zero. Show that the product **AB** of two $n \times n$ diagonal matrices **A** and **B** is again a diagonal matrix. State a concise rule for quickly computing **AB**. Is it clear that **AB** = **BA**? Explain.

Problems 28 through 30 develop a method of computing powers of a square matrix.

28. The positive integral powers of a square matrix **A** are defined as follows:

$$\mathbf{A}^1 = \mathbf{A}, \quad \mathbf{A}^2 = \mathbf{A}\mathbf{A}, \quad \mathbf{A}^3 = \mathbf{A}\mathbf{A}^2, \\ \mathbf{A}^4 = \mathbf{A}\mathbf{A}^3, \dots, \quad \mathbf{A}^{n+1} = \mathbf{A}\mathbf{A}^n, \dots$$

Suppose that *r* and *s* are positive integers. Prove that $\mathbf{A}^{r}\mathbf{A}^{s} = \mathbf{A}^{r+s}$ and that $(\mathbf{A}^{r})^{s} = \mathbf{A}^{rs}$ (in close analogy with the laws of exponents for real numbers).

29. If
$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$
, then show that
$$\mathbf{A}^2 = (a+d)\mathbf{A} - (ad-bc)\mathbf{I},$$

where I denotes the 2×2 identity matrix. Thus every 2×2 matrix A satisfies the equation

$$\mathbf{A}^2 - (\text{trace } \mathbf{A})\mathbf{A} + (\det \mathbf{A})\mathbf{I} = \mathbf{0}$$

where det $\mathbf{A} = ad - bc$ denotes the determinant of the matrix \mathbf{A} , and trace \mathbf{A} denotes the sum of its diagonal elements. This result is the 2-dimensional case of the Cayley-Hamilton theorem of Section 6.3.

30. The formula in Problem 29 can be used to compute A^2 without an explicit matrix multiplication. It follows that

$$\mathbf{A}^3 = (a+d)\mathbf{A}^2 - (ad-bc)\mathbf{A}$$

without an explicit matrix multiplication,

$$\mathbf{A}^4 = (a+d)\mathbf{A}^3 - (ad-bc)\mathbf{A}^2,$$

and so on. Use this method to compute A^2 , A^3 , A^4 , and A^5 given

$$\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}.$$

Problems 31–38 illustrate ways in which the algebra of matrices is not analogous to the algebra of real numbers.

- **31.** (a) Suppose that A and B are the matrices of Example 5. Show that $(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B}) \neq \mathbf{A}^2 - \mathbf{B}^2$.
 - (b) Suppose that A and B are square matrices with the property that AB = BA. Show that $(A + B)(A B) = A^2 B^2$.
- 32. (a) Suppose that A and B are the matrices of Example 5. Show that $(A + B)^2 \neq A^2 + 2AB + B^2$.
 - (b) Suppose that A and B are square matrices such that AB = BA. Show that $(A + B)^2 = A^2 + 2AB + B^2$.
- **33.** Find four different 2×2 matrices **A**, with each main diagonal element either +1 or -1, such that $\mathbf{A}^2 = \mathbf{I}$.
- **34.** Find a 2×2 matrix **A** with each element +1 or -1 such that $\mathbf{A}^2 = \mathbf{0}$. The formula of Problem 29 may be helpful.
- **35.** Use the formula of Problem 29 to find a 2×2 matrix **A** such that $\mathbf{A} \neq \mathbf{0}$ and $\mathbf{A} \neq \mathbf{I}$ but such that $\mathbf{A}^2 = \mathbf{A}$.
- **36.** Find a 2×2 matrix **A** with each main diagonal element zero such that $\mathbf{A}^2 = \mathbf{I}$.
- **37.** Find a 2 × 2 matrix **A** with each main diagonal element zero such that $\mathbf{A}^2 = -\mathbf{I}$.
- **38.** This is a continuation of the previous two problems. Find two nonzero 2×2 matrices **A** and **B** such that $A^2 + B^2 = 0$.
- **39.** Use matrix multiplication to show that if \mathbf{x}_1 and \mathbf{x}_2 are two solutions of the homogeneous system $\mathbf{A}\mathbf{x} = \mathbf{0}$ and c_1 and c_2 are real numbers, then $c_1\mathbf{x}_1 + c_2\mathbf{x}_2$ is also a solution.
- **40.** (a) Use matrix multiplication to show that if \mathbf{x}_0 is a solution of the homogeneous system $A\mathbf{x} = \mathbf{0}$ and \mathbf{x}_1 is a solution of the nonhomogeneous system $A\mathbf{x} = \mathbf{b}$, then $\mathbf{x}_0 + \mathbf{x}_1$ is also a solution of the nonhomogeneous system.
 - (b) Suppose that \mathbf{x}_1 and \mathbf{x}_2 are solutions of the nonhomogeneous system of part (a). Show that $\mathbf{x}_1 \mathbf{x}_2$ is a solution of the homogeneous system $\mathbf{A}\mathbf{x} = \mathbf{0}$.
- 41. This is a continuation of Problem 32. Show that if A and B are square matrices such that AB = BA, then

$$(\mathbf{A} + \mathbf{B})^3 = \mathbf{A}^3 + 3\mathbf{A}^2\mathbf{B} + 3\mathbf{A}\mathbf{B}^2 + \mathbf{B}^3$$

and

$$(\mathbf{A} + \mathbf{B})^4 = \mathbf{A}^4 + 4\mathbf{A}^3\mathbf{B} + 6\mathbf{A}^2\mathbf{B}^2 + 4\mathbf{A}\mathbf{B}^3 + \mathbf{B}^4.$$

42. Let

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$$
$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix} = \mathbf{I} + \mathbf{N}.$$

(a) Show that $N^2 \neq 0$ but $N^3 = 0$.

(b) Use the binomial formulas of Problem 41 to compute

$$A^2 = (I + N)^2 = I + 2N + N^2,$$

 $A^3 = (I + N)^3 = I + 3N + 3N^2,$

and

$$A^4 = (I + N)^4 = I + 4N + 6N^2.$$

43. Consider the 3×3 matrix

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}.$$

First verify by direct computation that $A^2 = 3A$. Then conclude that $A^{n+1} = 3^n A$ for every positive integer *n*.

3.5 Inverses of Matrices

44. Let $\mathbf{A} = \begin{bmatrix} a_{hi} \end{bmatrix}$, $\mathbf{B} = \begin{bmatrix} b_{ij} \end{bmatrix}$, and $\mathbf{C} = \begin{bmatrix} c_{jk} \end{bmatrix}$ be matrices of sizes $m \times n$, $n \times p$, and $p \times q$, respectively. To establish the associative law $\mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}$, proceed as follows. By Equation (16) the *hj*th element of \mathbf{AB} is

$$\sum_{i=1}^{n} a_{hi} b_{ij}.$$

By another application of Equation (16), the *hk*th element of (AB)C is

$$\sum_{j=1}^{p} \left(\sum_{i=1}^{n} a_{hi} b_{ij} \right) c_{jk} = \sum_{i=1}^{n} \sum_{j=1}^{p} a_{hi} b_{ij} c_{jk}.$$

Show similarly that the double sum on the right is also equal to the *hk*th element of A(BC). Hence the $m \times q$ matrices (AB)C and A(BC) are equal.

Recall that the $n \times n$ identity matrix is the diagonal matrix

$$\mathbf{I} = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}$$
(1)

having ones on its main diagonal and zeros elsewhere. It is not difficult to deduce directly from the definition of the matrix product that I acts like an identity for matrix multiplication:

$$\mathbf{A}\mathbf{I} = \mathbf{A} \quad \text{and} \quad \mathbf{I}\mathbf{B} = \mathbf{B} \tag{2}$$

if the sizes of **A** and **B** are such that the products **AI** and **IB** are defined. It is, nevertheless, instructive to derive the identities in (2) formally from the two basic facts about matrix multiplication that we state below. First, recall that the notation

$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 & \cdots & \mathbf{a}_n \end{bmatrix}$$
(3)

expresses the $m \times n$ matrix **A** in terms of its column vectors $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \dots, \mathbf{a}_n$.

Fact 1 Ax in terms of columns of A

If
$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}$$
 and $\mathbf{x} = (x_1, x_2, \dots, x_n)$ is an *n*-vector, then

$$\mathbf{A}\mathbf{x} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \dots + x_n\mathbf{a}_n.$$
(4)

The reason is that when each row vector of **A** is multiplied by the column vector **x**, its *j* th element is multiplied by x_j .

Fact 2 AB in terms of columns of B

If **A** is an $m \times n$ matrix and **B** = $\begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \cdots & \mathbf{b}_p \end{bmatrix}$ is an $n \times p$ matrix, then

$$\mathbf{AB} = \begin{bmatrix} \mathbf{Ab}_1 & \mathbf{Ab}_2 & \cdots & \mathbf{Ab}_p \end{bmatrix}.$$
(5)

That is, the *j*th column of AB is the product of A and the *j*th column of B. The reason is that the elements of the *j*th column of AB are obtained by multiplying the individual rows of A by the *j*th column of B.

Example 1

The third column of the product **AB** of the matrices

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 0 \\ 4 & 0 & 3 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} 3 & 7 & 5 & -4 \\ -2 & 6 & 3 & 6 \\ 5 & 1 & -2 & -1 \end{bmatrix}$$

is

$$\mathbf{Ab}_3 = \begin{bmatrix} 2 & -1 & 0 \\ 4 & 0 & 3 \end{bmatrix} \begin{bmatrix} 5 \\ 3 \\ -2 \end{bmatrix} = \begin{bmatrix} 7 \\ 14 \end{bmatrix}.$$

To prove that $\mathbf{AI} = \mathbf{A}$, note first that

$$\mathbf{I} = \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \cdots & \mathbf{e}_n \end{bmatrix},\tag{6}$$

where the *j* th column vector of **I** is the *j* th **basic unit vector**

$$\mathbf{e}_{j} = \begin{bmatrix} 0\\ \vdots\\ 1\\ \vdots\\ 0 \end{bmatrix} \leftarrow j \text{ th entry.}$$
(7)

If $\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}$, then Fact 1 yields

$$\mathbf{A}\mathbf{e}_{j} = 0 \cdot \mathbf{a}_{1} + \dots + 1 \cdot \mathbf{a}_{j} + \dots + 0 \cdot \mathbf{a}_{n} = \mathbf{a}_{j}.$$
(8)

Hence Fact 2 gives

$$\mathbf{AI} = \mathbf{A} \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \cdots & \mathbf{e}_n \end{bmatrix}$$
$$= \begin{bmatrix} \mathbf{A}\mathbf{e}_1 & \mathbf{A}\mathbf{e}_2 & \cdots & \mathbf{A}\mathbf{e}_n \end{bmatrix} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix};$$

that is, AI = A. The proof that IB = B is similar. (See Problems 41 and 42.)

The Inverse Matrix A⁻¹

If $a \neq 0$, then there is a number $b = a^{-1} = 1/a$ such that ab = ba = 1. Given a nonzero matrix **A**, we therefore wonder whether there is a matrix **B** such that AB = BA = I. The following two examples show that the answer to this question depends upon the particular matrix **A**.

Example 2

If

then

$$\mathbf{A} = \begin{bmatrix} 4 & 9 \\ 3 & 7 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} 7 & -9 \\ -3 & 4 \end{bmatrix},$$

 $\mathbf{AB} = \begin{bmatrix} 4 & 9 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} 7 & -9 \\ -3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \mathbf{I};$

BA = I by a similar computation.

Example 3 Let

$$\mathbf{A} = \begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

If the matrix **B** had the property that AB = BA = I, then

$$\mathbf{AB} = \begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$
$$= \begin{bmatrix} a - 3c & b - 3d \\ -2a + 6c & -2b + 6d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

But upon equating corresponding elements of **AB** and the 2×2 identity matrix in the last line, we find that

$$a - 3c = 1$$
 and $b - 3d = 0$
 $-2a + 6c = 0$ $-2b + 6d = 1.$

It is clear that these equations are inconsistent. Thus there can exist $no 2 \times 2$ matrix **B** such that AB = I.

DEFINITION Invertible Matrix

The square matrix **A** is called **invertible** if there exists a matrix **B** such that

$$AB = BA = I.$$

Thus the matrix **A** of Example 2 is invertible, whereas the matrix **A** of Example 3 is not invertible.

A matrix **B** such that AB = BA = I is called an **inverse matrix** of the matrix **A**. The following theorem says that no matrix can have two different inverse matrices.

THEOREM 1 Uniqueness of Inverse Matrices

If the matrix **A** is invertible, then there exists precisely one matrix **B** such that AB = BA = I.

Proof: If C is a (possibly different) matrix such that AC = CA = I as well, then the associative law of multiplication gives

$$\mathbf{C} = \mathbf{C}\mathbf{I} = \mathbf{C}(\mathbf{A}\mathbf{B}) = (\mathbf{C}\mathbf{A})\mathbf{B} = \mathbf{I}\mathbf{B} = \mathbf{B}.$$

Thus C is in fact the same matrix as **B**.

The unique inverse of an invertible matrix A is denoted by A^{-1} . Thus we say in Example 2 that

If
$$\mathbf{A} = \begin{bmatrix} 4 & 9 \\ 3 & 7 \end{bmatrix}$$
 then $\mathbf{A}^{-1} = \begin{bmatrix} 7 & -9 \\ -3 & 4 \end{bmatrix}$.

In the case of a 2×2 matrix **A**, it is easy to determine whether or not **A** is invertible and to find \mathbf{A}^{-1} if it exists. In Problems 36 and 37 we ask you to verify the following result.

THEOREM 2 Inverses of 2 x 2 Matrices

The 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

is invertible if and only if $ad - bc \neq 0$, in which case

$$\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$
 (9)

Equation (9) gives us the following prescription for writing the inverse of an invertible 2×2 matrix:

- First, interchange the two main diagonal entries.
- Then, change the signs of the two off-diagonal elements.
- Finally, divide each element of the resulting matrix by ad bc.

You might check that this is how $\mathbf{B} = \mathbf{A}^{-1}$ is obtained from A in Example 2 (in which ad - bc = 1).

Example 4

If

$$\mathbf{A} = \begin{bmatrix} 4 & 6\\ 5 & 9 \end{bmatrix},$$

then $ad - bc = 36 - 30 = 6 \neq 0$, so

 $\mathbf{A}^{-1} = \frac{1}{6} \begin{bmatrix} 9 & -6 \\ -5 & 4 \end{bmatrix} = \begin{bmatrix} \frac{3}{2} & -1 \\ -\frac{5}{6} & \frac{2}{3} \end{bmatrix}.$

Arbitrary integral powers of a square matrix A are defined as follows, though in the case of a negative exponent we must assume that A is also invertible. If n is a positive integer, we define

$$\mathbf{A}^{0} = \mathbf{I} \quad \text{and} \quad \mathbf{A}^{1} = \mathbf{A};$$
$$\mathbf{A}^{n+1} = \mathbf{A}^{n}\mathbf{A} \quad \text{for } n \ge 1;$$
$$\mathbf{A}^{-n} = (\mathbf{A}^{-1})^{n}.$$

In Problem 28 of Section 3.4, we asked you to verify the laws of exponents

$$\mathbf{A}^{r}\mathbf{A}^{s} = \mathbf{A}^{r+s}, \quad (\mathbf{A}^{r})^{s} = \mathbf{A}^{rs}$$
(10)

in the case of positive integral exponents, and Problem 31 of this section deals with the case of negative integral exponents. In Problem 29 we ask you to establish parts (a) and (b) of the following theorem.

THEOREM 3 Algebra of Inverse Matrices

If the matrices **A** and **B** of the same size are invertible, then

- (a) \mathbf{A}^{-1} is invertible and $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$;
- (b) If *n* is a nonnegative integer, then A^n is invertible and $(A^n)^{-1} = (A^{-1})^n$;
- (c) The product **AB** is invertible and

$$(\mathbf{A}\mathbf{B})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}.$$
 (11)

Proof of (c)

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I;$$

 $(B^{-1}A^{-1})(AB) = B^{-1}(A^{-1}A)B = B^{-1}IB = B^{-1}B = I.$

Thus we get **I** when we multiply **AB** on either side by $\mathbf{B}^{-1}\mathbf{A}^{-1}$. Because the inverse of the matrix **AB** is unique, this proves that **AB** is invertible and that its inverse matrix is $\mathbf{B}^{-1}\mathbf{A}^{-1}$.

In mathematics it is frequently important to note the surprises. The surprise in Eq. (11) is the reversal of the natural order of the factors in the right-hand side. You should now be able to show that

$$(ABC)^{-1} = C^{-1}B^{-1}A^{-1}.$$

In general, any product of invertible matrices of the same size is again invertible, and the inverse of a product of invertible matrices is the product *in reverse order* of their inverses.

THEOREM 4 Inverse Matrix Solution of Ax = b

If the $n \times n$ matrix **A** is invertible, then for any *n*-vector **b** the system

$$\mathbf{A}\mathbf{x} = \mathbf{b} \tag{12}$$

has the unique solution

$$\mathbf{x} = \mathbf{A}^{-1}\mathbf{b} \tag{13}$$

that is obtained by multiplying both sides in (12) on the left by the matrix A^{-1} .

Proof: We must show that $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$ is a solution and that it is the only solution of Eq. (12). First, the computation

$$\mathbf{A}(\mathbf{A}^{-1}\mathbf{b}) = (\mathbf{A}\mathbf{A}^{-1})\mathbf{b} = \mathbf{I}\mathbf{b} = \mathbf{b}$$

shows that $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$ is a solution. Second, if \mathbf{x}_1 is any (possibly different) solution, we observe that multiplication of each side of the equation $\mathbf{A}\mathbf{x}_1 = \mathbf{b}$ on the left by \mathbf{A}^{-1} yields $\mathbf{x}_1 = \mathbf{A}^{-1}\mathbf{b}$, and hence \mathbf{x}_1 is the same solution as \mathbf{x} after all.

Example 5 To solve the system

$$4x_1 + 6x_2 = 6$$

$$x_1 + 9x_2 = 18,$$

we use the inverse of the coefficient matrix

$$\mathbf{A} = \begin{bmatrix} 4 & 6\\ 5 & 9 \end{bmatrix}$$

that we found in Example 4. Then Eq. (13) yields

$$\mathbf{x} = \mathbf{A}^{-1}\mathbf{b} = \begin{bmatrix} 4 & 6\\ 5 & 9 \end{bmatrix}^{-1} \begin{bmatrix} 6\\ 18 \end{bmatrix}$$
$$= \begin{bmatrix} \frac{3}{2} & -1\\ -\frac{5}{6} & \frac{2}{3} \end{bmatrix} \begin{bmatrix} 6\\ 18 \end{bmatrix} = \begin{bmatrix} -9\\ 7 \end{bmatrix}.$$

Thus $x_1 = -9$, $x_2 = 7$ is the unique solution.

How to Find A⁻¹

Theorem 2 tells us only how to invert 2×2 matrices. The development of a method for inverting larger matrices involves a special class of matrices, which we define next.

DEFINITION Elementary Matrix

The $n \times n$ matrix **E** is called an **elementary matrix** if it can be obtained by performing a *single* elementary row operation on the $n \times n$ identity matrix **I**.

Example 6 We obtain some typical elementary matrices as follows.

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \xrightarrow{(3)R_1} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} = \mathbf{E}_1$$
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{(2)R_1 + R_3} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} = \mathbf{E}_2$$
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{\text{SWAP}(R_1, R_2)} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \mathbf{E}_3$$

The three elementary matrices E_1 , E_2 , and E_3 correspond to three typical elementary row operations.

Now, suppose that the $m \times m$ elementary matrix **E** corresponds to a certain elementary row operation. It turns out that if we perform this same operation on an arbitrary $m \times n$ matrix **A**, we get the product matrix **EA** that results upon multiplying **A** on the *left* by the matrix **E**. Thus we can carry out an elementary row operation by means of left multiplication by the corresponding elementary matrix. Problems 38–40 illustrate typical cases in the proof of the following theorem.

THEOREM 5 Elementary Matrices and Row Operations

If an elementary row operation is performed on the $m \times n$ matrix **A**, then the result is the product matrix **EA**, where **E** is the elementary matrix obtained by performing the same row operation on the $m \times m$ identity matrix.

Elementary Row Operation	Inverse Operation
$(c)R_i$	$\frac{1}{c}R_i$
$SWAP(R_i, R_j)$	$\text{SWAP}(R_i, R_j)$
$(c)R_i + R_j$	$(-c)R_i + R_j$

FIGURE 3.5.1. Inverse elementary row operations.

Elementary row operations are reversible. That is, to every elementary row operation there corresponds an *inverse* elementary row operation that cancels its effects (see Figure 3.5.1). It follows that *every elementary matrix is invertible*. To see why, let **E** be a given elementary matrix and let \mathbf{E}_1 be the elementary matrix corresponding to the inverse of the row operation that transforms **I** into **E**. Then the inverse operation transforms **E** to **I**, so Theorem 5 implies that $\mathbf{E}_1\mathbf{E} = \mathbf{I}$. We see similarly that $\mathbf{E}\mathbf{E}_1 = \mathbf{I}$. Hence, the elementary matrix **E** is invertible with $\mathbf{E}^{-1} = \mathbf{E}_1$.

Elementary matrices are not ordinarily used for computational purposes; it is simpler to carry out row operations directly than to multiply by elementary matrices. Instead, their principal role is in the proof of the following theorem, which leads in turn to a practical method for inverting matrices.

THEOREM 6 Invertible Matrices and Row Operations

The $n \times n$ matrix **A** is invertible if and only if it is row equivalent to the $n \times n$ identity matrix **I**.

Proof: Assume first that **A** is invertible. Then, by Theorem 4 (with $\mathbf{b} = \mathbf{0}$), it follows that $\mathbf{A}\mathbf{x} = \mathbf{0}$ has only the trivial solution $\mathbf{x} = \mathbf{0}$. But Theorem 4 in Section 3.3 implies that this is so (if and) only if **A** is row equivalent to **I**.

Now assume, conversely, that **A** is row equivalent to **I**. That is, there is a finite sequence of elementary row operations that transforms **A** into **I**. According to Theorem 5, each of these operations can be performed by multiplying on the left by the corresponding elementary matrix. If $\mathbf{E}_1, \mathbf{E}_2, \ldots, \mathbf{E}_k$ are the elementary matrices corresponding to these row operations, it follows that

$$\mathbf{E}_k \mathbf{E}_{k-1} \cdots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A} = \mathbf{I}. \tag{14}$$

If we now multiply each side in Eq. (14) by the inverse matrices $(\mathbf{E}_k)^{-1}$, $(\mathbf{E}_{k-1})^{-1}$, ..., $(\mathbf{E}_2)^{-1}$, $(\mathbf{E}_1)^{-1}$ in turn, we find that

$$\mathbf{A} = (\mathbf{E}_1)^{-1} (\mathbf{E}_2)^{-1} \cdots (\mathbf{E}_{k-1})^{-1} (\mathbf{E}_k)^{-1}.$$
 (15)

Thus \mathbf{A} is a product of invertible elementary matrices, and it follows from part (c) of Theorem 3 that \mathbf{A} is invertible.

The proof of Theorem 6 actually tells us *how to find* the inverse matrix of \mathbf{A} . If we invert each side in Eq. (15) (remembering to reverse the order on the right), we get

$$\mathbf{A}^{-1} = \mathbf{E}_k \mathbf{E}_{k-1} \cdots \mathbf{E}_2 \mathbf{E}_1 \mathbf{I}. \tag{16}$$

Because each left multiplication by an elementary matrix is equivalent to performing the corresponding row operation, we see by comparison of Eqs. (14) and (16) that the same sequence of elementary row operations that transforms **A** into **I** also transforms **I** into A^{-1} .

ALGORITHM Finding A⁻¹

To find the inverse A^{-1} of the invertible $n \times n$ matrix A, find a sequence of elementary row operations that reduces A to the $n \times n$ identity matrix I. Then apply the same sequence of operations in the same order to I to transform it into A^{-1} .

As a practical matter, it generally is more convenient to carry out the two reductions—from **A** to **I** and from **I** to \mathbf{A}^{-1} —in parallel, as illustrated in our next example.

Example 7 Find the inverse of the 3×3 matrix

$$\mathbf{A} = \begin{bmatrix} 4 & 3 & 2 \\ 5 & 6 & 3 \\ 3 & 5 & 2 \end{bmatrix}.$$

Solution We want to reduce **A** to the 3×3 identity matrix **I** while simultaneously performing the same sequence of row operations on **I** to obtain A^{-1} . In order to carry out this process efficiently, we adjoin **I** on the right of **A** to form the 3×6 matrix

Å

$$\begin{bmatrix} 4 & 3 & 2 & 1 & 0 & 0 \\ 5 & 6 & 3 & 0 & 1 & 0 \\ 3 & 5 & 2 & 0 & 0 & 1 \end{bmatrix}$$

We now apply the following sequence of elementary row operations to this 3×6 matrix (designed to transform its left half into the 3×3 identity matrix).

$\xrightarrow{(-1)R_3 + R_1}$	[1 5	-2_{6}	0 3	1 0	0 1	$\begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$
$\xrightarrow{(-1)R_3 + R_2}$	$\begin{bmatrix} 3 \\ 1 \\ 2 \\ 2 \end{bmatrix}$	-2 1	2 0 1	0 1 0	0 0 1	$\begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix}$
$\xrightarrow{(-2)R_1 + R_2}$	$\begin{bmatrix} 3 \\ 1 \\ 0 \\ 2 \end{bmatrix}$	5 -2 5	2 0 1	$1 \\ -2 \\ 0$	0 0 1	$\begin{bmatrix} 1 \\ -1 \\ 1 \\ 1 \end{bmatrix}$
$\xrightarrow{(-3)R_1 + R_3}$	$\begin{bmatrix} 3 \\ 1 \\ 0 \\ 0 \end{bmatrix}$	-2 5	2 0 1	$1 \\ -2 \\ 2$	0 1	$\begin{bmatrix} 1 \\ -1 \\ 1 \\ 4 \end{bmatrix}$
		11	2	-3	0	4
$\xrightarrow{(-2)R_2 + R_3}$	$\begin{bmatrix} 1\\ 0\\ 0 \end{bmatrix}$	-2 5	0 1	1 - 2	0 1	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$
$\xrightarrow{(-2)R_2 + R_3}$ $\xrightarrow{\text{SWAP}(R_2, R_3)}$	$\begin{bmatrix} 1\\0\\0\\\end{bmatrix}$	$ \begin{array}{r} -2 \\ 5 \\ 1 \\ -2 \\ 1 \end{array} $	0 1 0 0 0	$ \begin{array}{c} 1 \\ -2 \\ 1 \\ 1 \\ 1 \end{array} $	$ \begin{array}{c} 0 \\ 1 \\ -2 \\ 0 \\ -2 \end{array} $	$\begin{bmatrix} -1\\1\\2 \end{bmatrix}$ $\begin{bmatrix} -1\\2 \end{bmatrix}$
$\xrightarrow{(-2)R_2 + R_3}$ $\xrightarrow{\text{SWAP}(R_2, R_3)}$ $\xrightarrow{(2)R_2 + R_1}$	$\begin{bmatrix} 1\\0\\0\\\end{bmatrix}$	$ \begin{array}{c} -2 \\ 5 \\ 1 \\ -2 \\ 1 \\ 5 \\ 0 \\ 1 \end{array} $	0 1 0 0 0 1 0 0	$ \begin{array}{c} 1 \\ -2 \\ 1 \\ 1 \\ 1 \\ -2 \\ 3 \\ 1 \end{array} $	$\begin{array}{c} 0 \\ 1 \\ -2 \\ 0 \\ -2 \\ 1 \\ -4 \\ -2 \end{array}$	$ \begin{array}{c} -1\\1\\2\\-1\\2\\1\\3\\2\end{array} $
$(-2)R_2 + R_3 \rightarrow (-2)R_2 + R_3 \rightarrow (-5)R_2 + R_3 \rightarrow (-5)R_3 + (-5)R_$	$ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} $ $ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} $ $ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} $ $ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} $	$ \begin{array}{c} -2 \\ 5 \\ 1 \\ -2 \\ 1 \\ 5 \\ 0 \\ 1 \\ 5 \\ 0 \\ 1 \end{array} $	0 1 0 0 1 0 0 1 0 0	$ \begin{array}{c} 1 \\ -2 \\ 1 \\ 1 \\ -2 \\ 3 \\ 1 \\ -2 \\ 3 \\ 1 \end{array} $	$ \begin{array}{c} 0 \\ 1 \\ -2 \\ 0 \\ -2 \\ 1 \\ -4 \\ -2 \\ 1 \\ -4 \\ -2 \\ -4 \\ -2 \end{array} $	$ \begin{array}{c} -1 \\ 1 \\ 2 \\ -1 \\ 2 \\ 1 \\ 3 \\ 2 \\ 1 \\ 3 \\ 2 \\ 1 \end{array} $

Now that we have reduced the left half of the 3×6 matrix to **I**, we simply examine its right half to see that the inverse of **A** is

$$\mathbf{A}^{-1} = \begin{bmatrix} 3 & -4 & 3\\ 1 & -2 & 2\\ -7 & 11 & -9 \end{bmatrix}.$$

Remark Ordinarily, we do not know in advance whether a given square matrix is invertible or not. To find out, we attempt to carry out the reduction process illustrated in Example 7. If we succeed in reducing **A** to **I**, then **A** is invertible and thereby we find \mathbf{A}^{-1} . Otherwise—if, somewhere along the way, an all-zero row appears in the left half—we conclude that **A** is not row equivalent to **I**, and therefore (by Theorem 6) **A** is not invertible.

Matrix Equations

In certain applications, one needs to solve a system $A\mathbf{x} = \mathbf{b}$ of *n* equations in *n* unknowns several times in succession—with the same $n \times n$ coefficient matrix **A** each time, but with different constant vectors $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k$ on the right. Thus we want to find solution vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ such that

$$\mathbf{A}\mathbf{x}_1 = \mathbf{b}_1, \quad \mathbf{A}\mathbf{x}_2 = \mathbf{b}_2, \quad \dots, \quad \mathbf{A}\mathbf{x}_k = \mathbf{b}_k. \tag{17}$$

By Fact 2 at the beginning of this section,

$$\begin{bmatrix} \mathbf{A}\mathbf{x}_1 & \mathbf{A}\mathbf{x}_2 & \cdots & \mathbf{A}\mathbf{x}_k \end{bmatrix} = \mathbf{A}\begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix}.$$

So the k equations in (17) are equivalent to the single matrix equation

A

$$\mathbf{X} = \mathbf{B},\tag{18}$$

where

$$\mathbf{X} = \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_k \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \cdots & \mathbf{b}_k \end{bmatrix}.$$

If **A** is invertible and we know A^{-1} , we can find the $n \times k$ matrix of "unknowns" by multiplying each term in Equation (18) on the left by A^{-1} :

$$\mathbf{X} = \mathbf{A}^{-1}\mathbf{B}.\tag{19}$$

Note that this equation is a generalization of Eq. (13) in Theorem 4. If k = 1, it usually is simplest to solve the system by Gaussian elimination, but when several different solutions are sought, it may be simpler to find \mathbf{A}^{-1} first and then to apply (19).

Example 8	Find a 3×4 matrix X such that

 $\begin{bmatrix} 4 & 3 & 2 \\ 5 & 6 & 3 \\ 3 & 5 & 2 \end{bmatrix} \mathbf{X} = \begin{bmatrix} 3 & -1 & 2 & 6 \\ 7 & 4 & 1 & 5 \\ 5 & 2 & 4 & 1 \end{bmatrix}.$

Solution The coefficient matrix is the matrix **A** whose inverse we found in Example 7, so Eq. (19) yields

$$\mathbf{X} = \mathbf{A}^{-1}\mathbf{B} = \begin{bmatrix} 3 & -4 & 3\\ 1 & -2 & 2\\ -7 & 11 & -9 \end{bmatrix} \begin{bmatrix} 3 & -1 & 2 & 6\\ 7 & 4 & 1 & 5\\ 5 & 2 & 4 & 1 \end{bmatrix},$$

and hence

$$\mathbf{X} = \begin{bmatrix} -4 & -13 & 14 & 1\\ -1 & -5 & 8 & -2\\ 11 & 33 & -39 & 4 \end{bmatrix}.$$

By looking at the third columns of B and X, for instance, we see that the solution of

 $4x_1 + 3x_2 + 2x_3 = 2$ $5x_1 + 6x_2 + 3x_3 = 1$ $3x_1 + 5x_2 + 2x_3 = 4$

is $x_1 = 14$, $x_2 = 8$, $x_3 = -39$.

Nonsingular Matrices

Theorem 6 tells us that the square matrix **A** is invertible if and only if it is row equivalent to the identity matrix **I**, and Theorem 4 in Section 3.3 implies that the latter is true if and only if the system Ax = 0 has only the trivial solution x = 0. A square matrix having these equivalent properties is sometimes called a **nonsingular** matrix.

THEOREM 7 Properties of Nonsingular Matrices

The following properties of an $n \times n$ matrix **A** are equivalent.

- (a) A is invertible.
- (b) A is row equivalent to the $n \times n$ identity matrix I.
- (c) Ax = 0 has only the trivial solution.
- (d) For every *n*-vector **b**, the system Ax = b has a unique solution.
- (e) For every *n*-vector **b**, the system Ax = b is consistent.

Proof: By the remarks preceding the statement of Theorem 7, we already know that properties (a), (b), and (c) are equivalent—if **A** has any one of these properties, then it also has the other two. We can, therefore, complete the proof by establishing the chain of logical implications

$$(c) \Rightarrow (d) \Rightarrow (e) \Rightarrow (a).$$

That is, we need to show that if A has property (c), then it has property (d), and, similarly, that (d) implies (e) and that (e) implies (a).

(c) \Rightarrow (d): We already know that (c) implies (a), and Theorem 4 says that (a) implies (d). Therefore, (c) implies (d).

(d) \Rightarrow (e): This is obvious, because if the system Ax = b has a unique solution, then it certainly has a solution, and thus is consistent.

(e) \Rightarrow (a): Given the hypothesis that Ax = b is consistent for every **b**, we must prove that **A** is invertible. Let $\mathbf{b} = \mathbf{e}_j$, the *j* th column vector of the identity matrix **I**. Then the consistency of $Ax = \mathbf{e}_j$ yields an *n*-vector \mathbf{x}_j such that

$$\mathbf{A}\mathbf{x}_j = \mathbf{e}_j. \tag{20}$$

Let the vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ be obtained in this way for $j = 1, 2, \dots, n$, and let **B** be the $n \times n$ matrix with these vectors as its columns:

$$\mathbf{B} = \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \dots & \mathbf{x}_n \end{bmatrix}$$

Then

$$\mathbf{AB} = \mathbf{A} \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_n \end{bmatrix}$$
$$= \begin{bmatrix} \mathbf{A}\mathbf{x}_1 & \mathbf{A}\mathbf{x}_2 & \cdots & \mathbf{A}\mathbf{x}_n \end{bmatrix}$$
$$= \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \cdots & \mathbf{e}_n \end{bmatrix} \quad [by (20)].$$

Therefore, AB = I, and thus we have found a matrix **B** such that AB = I.

We next show that **B** is invertible by showing that $\mathbf{B}\mathbf{x} = \mathbf{0}$ has only the trivial solution [and by using the fact that property (c) implies property (a)]. But if $\mathbf{B}\mathbf{x} = \mathbf{0}$, then

$$\mathbf{A}(\mathbf{B}\mathbf{x}) = \mathbf{A}\mathbf{0} = \mathbf{0}$$

which implies that Ix = 0 and thus that x = 0. So **B** is indeed invertible. We can therefore multiply each term in the equation AB = I on the right by B^{-1} to get

$$\mathbf{A}\mathbf{B}\mathbf{B}^{-1} = \mathbf{I}\mathbf{B}^{-1}$$

so that $\mathbf{A} = \mathbf{B}^{-1}$. Thus \mathbf{A} is the inverse of an invertible matrix, and so is itself invertible. This establishes that property (e) implies property (a), and therefore we have completed the proof.

The proof of Theorem 7 is a bit long, but it summarizes most of the basic theory of Chapter 1 and is therefore well worth the effort. Indeed, this theorem is one of the central theorems of elementary linear algebra, and we will need to refer to it repeatedly in subsequent chapters.

3.5 **Problems**

In Problems 1–8, first apply the formulas in (9) to find A^{-1} . Then use A^{-1} (as in Example 5) to solve the system Ax = b.

- - 7

1.
$$\mathbf{A} = \begin{bmatrix} 3 & 2 \\ 4 & 3 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$$

2. $\mathbf{A} = \begin{bmatrix} 3 & 7 \\ 2 & 5 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$
3. $\mathbf{A} = \begin{bmatrix} 6 & 7 \\ 5 & 6 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$
4. $\mathbf{A} = \begin{bmatrix} 5 & 12 \\ 7 & 17 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 5 \\ 5 \end{bmatrix}$
5. $\mathbf{A} = \begin{bmatrix} 3 & 2 \\ 5 & 4 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 5 \\ 6 \end{bmatrix}$
6. $\mathbf{A} = \begin{bmatrix} 4 & 7 \\ 3 & 6 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 10 \\ 5 \end{bmatrix}$
7. $\mathbf{A} = \begin{bmatrix} 7 & 9 \\ 5 & 7 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$
8. $\mathbf{A} = \begin{bmatrix} 8 & 15 \\ 5 & 10 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 7 \\ 3 \end{bmatrix}$

In Problems 9–22, use the method of Example 7 to find the inverse \mathbf{A}^{-1} of each given matrix \mathbf{A} .

9.
$$\begin{bmatrix} 5 & 6 \\ 4 & 5 \end{bmatrix}$$
 10. $\begin{bmatrix} 5 & 7 \\ 4 & 6 \end{bmatrix}$

 11. $\begin{bmatrix} 1 & 5 & 1 \\ 2 & 5 & 0 \\ 2 & 7 & 1 \end{bmatrix}$
 12. $\begin{bmatrix} 1 & 3 & 2 \\ 2 & 8 & 3 \\ 3 & 10 & 6 \end{bmatrix}$

 13. $\begin{bmatrix} 2 & 7 & 3 \\ 1 & 3 & 2 \\ 3 & 7 & 9 \end{bmatrix}$
 14. $\begin{bmatrix} 3 & 5 & 6 \\ 2 & 4 & 3 \\ 2 & 3 & 5 \end{bmatrix}$

 15. $\begin{bmatrix} 1 & 1 & 5 \\ 1 & 4 & 13 \\ 3 & 2 & 12 \end{bmatrix}$
 16. $\begin{bmatrix} 1 & -3 & -3 \\ -1 & 1 & 2 \\ 2 & -3 & -3 \end{bmatrix}$

 17. $\begin{bmatrix} 1 & -3 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{bmatrix}$
 18. $\begin{bmatrix} 1 & -2 & 2 \\ 3 & 0 & 1 \\ 1 & -1 & 2 \end{bmatrix}$

 19. $\begin{bmatrix} 1 & 4 & 3 \\ 1 & 4 & 5 \\ 2 & 5 & 1 \end{bmatrix}$
 20. $\begin{bmatrix} 2 & 0 & -1 \\ 1 & 0 & 3 \\ 1 & 1 & 1 \end{bmatrix}$

 21. $\begin{bmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 3 & 0 & 0 & 1 \end{bmatrix}$
 22. $\begin{bmatrix} 4 & 0 & 1 & 1 \\ 3 & 1 & 3 & 1 \\ 0 & 1 & 2 & 0 \\ 3 & 2 & 4 & 1 \end{bmatrix}$

In Problems 23–28, use the method of Example 8 to find a matrix \mathbf{X} such that $\mathbf{A}\mathbf{X} = \mathbf{B}$.

23.
$$\mathbf{A} = \begin{bmatrix} 4 & 3 \\ 5 & 4 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 1 & 3 & -5 \\ -1 & -2 & 5 \end{bmatrix}$$

24. $\mathbf{A} = \begin{bmatrix} 7 & 6 \\ 8 & 7 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 2 & 0 & 4 \\ 0 & 5 & -3 \end{bmatrix}$
25. $\mathbf{A} = \begin{bmatrix} 1 & 4 & 1 \\ 2 & 8 & 3 \\ 2 & 7 & 4 \end{bmatrix}, \mathbf{B} = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 2 & 2 \\ -1 & 1 & 0 \end{bmatrix}$



Problems 30 through 37 explore the properties of matrix inverses.

- **30.** Suppose that **A**, **B**, and **C** are invertible matrices of the same size. Show that the product **ABC** is invertible and that $(\mathbf{ABC})^{-1} = \mathbf{C}^{-1}\mathbf{B}^{-1}\mathbf{A}^{-1}$.
- **31.** Suppose that **A** is an invertible matrix and that *r* and *s* are negative integers. Verify that $\mathbf{A}^r \mathbf{A}^s = \mathbf{A}^{r+s}$ and that $(\mathbf{A}^r)^s = \mathbf{A}^{rs}$.
- 32. Prove that if A is an invertible matrix and AB = AC, then B = C. Thus *invertible* matrices can be canceled.
- **33.** Let **A** be an $n \times n$ matrix such that $A\mathbf{x} = \mathbf{x}$ for every *n*-vector **x**. Show that $\mathbf{A} = \mathbf{I}$.
- **34.** Show that a diagonal matrix is invertible if and only if each diagonal element is nonzero. In this case, state concisely how the inverse matrix is obtained.
- **35.** Let **A** be an $n \times n$ matrix with either a row or a column consisting only of zeros. Show that **A** is not invertible.
- **36.** Show that $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is not invertible if ad bc = 0.
- **37.** Suppose that $ad bc \neq 0$ and \mathbf{A}^{-1} is defined as in Equation (9). Verify directly that $\mathbf{A}\mathbf{A}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$.

Problems 38 through 40 explore the effect of multiplying by an elementary matrix.

- 38. Let E be the elementary matrix E₁ of Example 6. If A is a 2 × 2 matrix, show that EA is the result of multiplying the first row of A by 3.
- **39.** Let **E** be the elementary matrix \mathbf{E}_2 of Example 6 and suppose that **A** is a 3×3 matrix. Show that **EA** is the result upon adding twice the first row of **A** to its third row.
- **40.** Let **E** be the elementary matrix E_3 of Example 6. Show that **EA** is the result of interchanging the first two rows of the matrix **A**.

Problems 41 and 42 complete the proof of Eq. (2).

- **41.** Show that the *i* th row of the product **AB** is **A**_{*i*}**B**, where **A**_{*i*} is the *i* th row of the matrix **A**.
- **42.** Apply the result of Problem 41 to show that if **B** is an $m \times n$ matrix and **I** is the $m \times m$ identity matrix, then **IB** = **B**.
- **43.** Suppose that the matrices **A** and **B** are row equivalent. Use Theorem 5 to prove that $\mathbf{B} = \mathbf{G}\mathbf{A}$, where **G** is a product of elementary matrices.

186 CHAPTER 3 LINEAR SYSTEMS AND MATRICES

- **44.** Show that every invertible matrix is a product of elementary matrices.
- **45.** Extract from the proof of Theorem 7 a self-contained proof of the following fact: If **A** and **B** are square matrices

3.5 Application Automated Solution of Linear Systems



Linear systems with more than two or three equations are most frequently solved with the aid of calculators or computers. If an $n \times n$ linear system is written in the matrix form $A\mathbf{x} = \mathbf{b}$, then we need to calculate first the inverse matrix \mathbf{A}^{-1} and then the matrix product $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$. Suppose the $n \times n$ matrix \mathbf{A} and the column vector \mathbf{b} have been entered (as illustrated in the 3.2 Application). If \mathbf{A} is invertible, then the inverse matrix \mathbf{A}^{-1} is calculated by the *Maple* command with(linalg): inverse(\mathbf{A}), the *Mathematica* command Inverse[\mathbf{A}], or the MATLAB command inv(\mathbf{A}). Consequently, the solution vector \mathbf{x} is calculated by the *Maple* command

such that AB = I, then A and B are invertible.

and **B** are themselves invertible.

46. Deduce from the result of Problem 45 that if A and B are

square matrices whose product AB is invertible, then A

with(linalg): x := multiply(inverse(A),b);

or the Mathematica command

or the MATLAB command

x = inv(A)*b

Figure 3.5.2 illustrates a similar calculator solution of the linear system



FIGURE 3.5.2. TI-89 solution of a linear system Ax = b.

 $3x_1 - 2x_2 + 7x_3 + 5x_4 = 505$ $2x_1 + 4x_2 - x_3 + 6x_4 = 435$ $5x_1 + x_2 + 7x_3 - 3x_4 = 286$ $4x_1 - 6x_2 - 8x_3 + 9x_4 = 445$

for the solution $x_1 = 59$, $x_2 = 13$, $x_3 = 17$, $x_4 = 47$. This solution is also given by the Wolfram Alpha query

Remark Whereas the preceding commands illustrate the handy use of conveniently available inverse matrices to solve linear systems, it might be mentioned that modern computer systems employ direct methods—involving Gaussian elimination and still more sophisticated techniques—that are more efficient and numerically reliable to solve a linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$ without first calculating the inverse matrix \mathbf{A}^{-1} .

Use an available calculator or computer system to solve the linear systems in Problems 1–6 of the 3.3 Application. The applied problems below are elementary in character—resembling the "word problems" of high school algebra—but might illustrate the practical advantages of automated solutions.

- 1. You are walking down the street minding your own business when you spot a small but heavy leather bag lying on the sidewalk. It turns out to contain U.S. Mint American Eagle gold coins of the following types:
 - One-half ounce gold coins that sell for \$285 each,
 - One-quarter ounce gold coins that sell for \$150 each, and

• One-tenth ounce gold coins that sell for \$70 each.

A bank receipt found in the bag certifies that it contains 258 such coins with a total weight of 67 ounces and a total value of exactly \$40,145. How many coins of each type are there?

- **2.** Now you really strike it rich! You find a bag containing one-ounce U.S. American Eagle gold coins valued at \$550 each, together with half-ounce and quarter-ounce coins valued as in the preceding problem. If this bag contains a total of 365 coins with a total weight of exactly 11 pounds and a total value of \$100,130, how many gold coins of each type are there?
- **3.** A commercial customer orders 81 gallons of paint that contains equal amounts of red paint, green paint, and blue paint—and, hence, could be prepared by mixing 27 gallons of each. However, the store wishes to prepare this order by mixing three types of paint that are already available in large quantity:
 - a *reddish* paint that is a mixture of 50% red, 25% green, and 25% blue paint;
 - a greenish paint that is 12.5% red, 75% green, and 12.5% blue paint; and
 - a *bluish* paint that is 20% red, 20% green, and 60% blue paint.

How many gallons of each are needed to prepare the customer's order?

- **4.** Now the paint store receives a really big order—for 244 gallons of paint that is 1/2 red paint, 1/4 green paint, and 1/4 blue paint. The store has three already-mixed types of paint available in large quantity—the greenish paint and the bluish paint of the preceding problem, plus a reddish paint that is 2/3 red paint, 1/6 green paint, and 1/6 blue paint. How many gallons of each must be mixed in order to fill this order?
- **5.** A tour busload of 45 people attended two Florida theme parks on successive days. On Day 1 the entrance fee was \$15 per adult, \$8 per child, \$12 per senior citizen and the total charge was \$558. On Day 2 the entrance fee was \$20 per adult, \$12 per child, \$17 per senior citizen and the total charge was \$771. How many adults, children, and senior citizens were on this tour bus?
- **6.** For some crazy reason, the lunches bought at the first theme park were totaled separately for the adults, children, and seniors. The adults ordered 34 hot dogs, 15 French fries, and 24 soft drinks for a total bill of \$70.85. The children ordered 20 hot dogs, 14 French fries, and 15 soft drinks for a total bill of \$46.65. The senior citizens ordered 11 hot dogs, 10 French fries, and 12 soft drinks for a total bill of \$30.05. What were the prices of a hot dog, an order of French fries, and a soft drink?
- 7. A fast-food restaurant sells four types of sandwiches—hamburgers, cheeseburgers, roast beef, and chicken—and has four cash registers. At the end of each day, each cash register tallies the number of each type of sandwich sold, and the total sandwich receipts for the day. The four cash register operators work at different speeds, and one day's totals were as follows:

	Hamburgers	Cheeseburgers	Roast Beef	Chicken	Receipts
Register 1	37	44	17	23	\$232.99
Register 2	28	35	13	17	\$178.97
Register 3	32	39	19	21	\$215.99
Register 4	47	51	25	29	\$294.38

What was the price of each of the four types of sandwiches?

	Hamburgers	Cheeseburgers	Roast Beef	Chicken	Ham	Receipts
Register 1	41	49	22	26	19	\$292.79
Register 2	34	39	18	20	16	\$236.73
Register 3	36	43	23	24	18	\$270.70
Register 4	49	52	26	31	24	\$340.19
Register 5	52	55	24	28	25	\$341.64

8. The fast-food restaurant of the preceding problem adds a ham sandwich to its menu and, because of increased business, it also adds a fifth cash register and reduces prices. After this expansion, one day's totals were as follows:

What were the new prices of the five types of sandwiches?

3.6 Determinants

In Theorem 2 of Section 3.5 we saw that the 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

is invertible if and only if $ad - bc \neq 0$. The number ad - bc is called the **determinant** of the 2 × 2 matrix **A**. There are several common notations for determinants:

$$\det \mathbf{A} = \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$
(1)

In particular, note the vertical bars that distinguish a determinant from a matrix. Sometimes we write det(A) to emphasize that (1) defines a function which associates a number |A| with each 2 × 2 matrix A.

Example 1

 $\begin{vmatrix} 3 & 7 \\ 4 & -6 \end{vmatrix} = (3)(-6) - (4)(7) = -46$

Determinants are often used in elementary algebra to solve a 2×2 system of the form

$$a_{11}x + a_{12}y = b_1$$

$$a_{21}x + a_{22}y = b_2.$$
(2)

It follows from Theorems 2 and 7 in Section 3.5 that this system has a unique solution if and only if its **coefficient determinant**—the determinant of its coefficient matrix **A**—is nonzero:

$$\det \mathbf{A} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \neq 0.$$
(3)

Then **Cramer's rule** says that this unique solution is given by

$$x = \frac{\begin{vmatrix} b_1 & a_{12} \\ b_2 & a_{22} \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}}, \qquad y = \frac{\begin{vmatrix} a_{11} & b_1 \\ a_{21} & b_2 \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}}.$$
 (4)

Thus Cramer's rule gives each of x and y as a quotient of two determinants, the denominator in each case being the determinant of the coefficient matrix. In the numerator for x the coefficients a_{11} and a_{21} of x are replaced with the right-side coefficients b_1 and b_2 , whereas in the numerator for y the coefficients of y are replaced with b_1 and b_2 .

The proof of Cramer's rule is a routine calculation—to verify that evaluation of the determinant in (4) gives the same expressions for x and y as are obtained by solution of (2) by the method of elimination.

Example 2 To apply Cramer's rule to solve the system

$$7x + 8y = 5$$
$$6x + 9y = 4$$

with coefficient determinant

$$\begin{vmatrix} 7 & 8 \\ 6 & 9 \end{vmatrix} = 15,$$

we simply substitute in the equations in (4) to get

$$x = \frac{\begin{vmatrix} 5 & 8 \\ 4 & 9 \end{vmatrix}}{15} = \frac{13}{15}, \qquad y = \frac{\begin{vmatrix} 7 & 5 \\ 6 & 4 \end{vmatrix}}{15} = -\frac{2}{15}.$$

Higher-Order Determinants

We define 3×3 determinants in terms of 2×2 determinants, 4×4 determinants in terms of 3×3 determinants, and so on. This type of definition—one dimension at a time, with the definition in each dimension depending on its meaning in lower dimensions—is called an *inductive definition*.

The determinant det $\mathbf{A} = |a_{ij}|$ of a 3 × 3 matrix $\mathbf{A} = [a_{ij}]$ is defined as follows:

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}.$$
 (5)

Note the single minus sign on the right-hand side. The three 2×2 determinants in (5) are multiplied by the elements a_{11} , a_{12} , and a_{13} along the *first row* of the matrix **A**. Each of these elements a_{1j} is multiplied by the determinant of the 2×2 submatrix of **A** that remains after the row and column containing a_{1j} are deleted.

Example 3

$$\begin{vmatrix} 5 & -2 & -3 \\ 4 & 0 & 1 \\ 3 & -1 & 2 \end{vmatrix} = (5) \begin{vmatrix} 0 & 1 \\ -1 & 2 \end{vmatrix} - (-2) \begin{vmatrix} 4 & 1 \\ 3 & 2 \end{vmatrix} + (-3) \begin{vmatrix} 4 & 0 \\ 3 & -1 \end{vmatrix}$$
$$= (5)(1) + (2)(5) - (3)(-4) = 27$$

The definition of higher-order determinants is simplified by the following notation and terminology.

DEFINITION Minors and Cofactors

Let $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ be an $n \times n$ matrix. The *ij*th **minor** of \mathbf{A} (also called the **minor** of a_{ij}) is the determinant M_{ij} of the $(n-1) \times (n-1)$ submatrix that remains after deleting the *i*th row and the *j*th column of \mathbf{A} . The *ij*th **cofactor** A_{ij} of \mathbf{A} (or the **cofactor** of a_{ij}) is defined to be

$$A_{ij} = (-1)^{i+j} M_{ij}.$$
 (6)

For example, the minor of a_{12} in a 3 × 3 matrix is

$$M_{12} = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}$$

The minor of a_{32} in a 4 × 4 matrix is

$$M_{32} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{vmatrix} = \begin{vmatrix} a_{11} & a_{13} & a_{14} \\ a_{21} & a_{23} & a_{24} \\ a_{41} & a_{43} & a_{44} \end{vmatrix}.$$

According to Eq. (6), the cofactor A_{ij} is obtained by attaching the sign $(-1)^{i+j}$ to the minor M_{ij} . This sign is most easily remembered as the one that appears in the *ij*th position in checkerboard arrays such as

$$\begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix} \text{ and } \begin{bmatrix} + & - & + & - \\ - & + & - & + \\ + & - & + & - \\ - & + & - & + \end{bmatrix}.$$

Note that a plus sign always appears in the upper left corner and that the signs alternate horizontally and vertically. In the 4×4 case, for instance,

$$A_{11} = +M_{11}, \quad A_{12} = -M_{12}, \quad A_{13} = +M_{13}, \quad A_{14} = -M_{14}, \\ A_{21} = -M_{21}, \quad A_{22} = +M_{22}, \quad A_{23} = -M_{23}, \quad A_{24} = +M_{24},$$

and so forth.

With this notation, the definition of 3×3 determinants in (5) can be rewritten as

$$\det \mathbf{A} = a_{11}M_{11} - a_{12}M_{12} + a_{13}M_{13}$$

= $a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13}.$ (7)

The last formula is the **cofactor expansion** of det A along the *first row* of A. Its natural generalization yields the definition of the determinant of an $n \times n$ matrix, under the inductive assumption that $(n - 1) \times (n - 1)$ determinants have already been defined.

DEFINITION n x n **Determinants**

The **determinant** det $\mathbf{A} = |a_{ij}|$ of an $n \times n$ matrix $\mathbf{A} = [a_{ij}]$ is defined as

$$\det \mathbf{A} = a_{11}A_{11} + a_{12}A_{12} + \dots + a_{1n}A_{1n}.$$
 (8)

Thus we multiply each element of the first row of \mathbf{A} by its cofactor and then add these *n* products to get det \mathbf{A} .

In numerical computations, it frequently is more convenient to work first with minors rather than with cofactors and then attach signs in accord with the checkerboard pattern illustrated previously. Note that determinants have been defined only for *square* matrices.

Example 4 To evaluate the determinant of

 $\mathbf{A} = \begin{bmatrix} 2 & 0 & 0 & -3 \\ 0 & -1 & 0 & 0 \\ 7 & 4 & 3 & 5 \\ -6 & 2 & 2 & 4 \end{bmatrix},$

we observe that there are only two nonzero terms in the cofactor expansion along the first row. We need not compute the cofactors of zeros, because they will be multiplied by zero in computing the determinant; hence

	-1	0	0		0	-1	0	
$\det \mathbf{A} = +(2)$	4	3	5	-(-3)	7	4	3	
	2	2	4		-6	2	2	

Each of the 3×3 determinants on the right-hand side has only a single nonzero term in its cofactor expansion along the first row, so

det
$$\mathbf{A} = (2)(-1) \begin{vmatrix} 3 & 5 \\ 2 & 4 \end{vmatrix} + (3)(+1) \begin{vmatrix} 7 & 3 \\ -6 & 2 \end{vmatrix}$$

= $(-2)(12 - 10) + (3)(14 + 18) = 92.$

Note that, if we could expand along the second row in Example 4, there would be only a single 3×3 determinant to evaluate. It is in fact true that a determinant can be evaluated by expansion along *any* row *or* column. The proof of the following theorem is included in Appendix B.

THEOREM 1 Cofactor Expansions of Determinants

The determinant of an $n \times n$ matrix $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ can be obtained by expansion along any row or column. The cofactor expansion along the *i*th row is

$$\det \mathbf{A} = a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}.$$
(9)

The cofactor expansion along the *j* th column is

$$\det \mathbf{A} = a_{1j}A_{1j} + a_{2j}A_{2j} + \dots + a_{nj}A_{nj}.$$
 (10)

The formulas in (9) and (10) provide 2n different cofactor expansions of an $n \times n$ determinant. For n = 3, for instance, we have

det
$$\mathbf{A} = a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13}$$

 $= a_{21}A_{21} + a_{22}A_{22} + a_{23}A_{23}$
 $= a_{31}A_{31} + a_{32}A_{32} + a_{33}A_{33}$ row expansions
 $= a_{11}A_{11} + a_{21}A_{21} + a_{31}A_{31}$
 $= a_{12}A_{12} + a_{22}A_{22} + a_{32}A_{32}$
 $= a_{13}A_{13} + a_{23}A_{23} + a_{33}A_{33}$.

In a specific example, we naturally attempt to choose the expansion that requires the least computational labor.

Example 5 To evaluate the determinant of

$$\mathbf{A} = \begin{bmatrix} 7 & 6 & 0 \\ 9 & -3 & 2 \\ 4 & 5 & 0 \end{bmatrix},$$

we expand along the third column, because it has only a single nonzero entry. Thus

det
$$\mathbf{A} = -(2) \begin{vmatrix} 7 & 6 \\ 4 & 5 \end{vmatrix} = (-2)(35 - 24) = -22.$$

In addition to providing ways of evaluating determinants, the theorem on cofactor expansions is a valuable tool for investigating the general properties of determinants. For instance, it follows immediately from the formulas in (9) and (10) that, *if the square matrix* **A** *has either an all-zero row or an all-zero column, then* det $\mathbf{A} = 0$. For example, by expanding along the second row we see immediately that

17	33	-24	
0	0	0	= 0.
80	-62	41	

Row and Column Properties

We now list seven properties of determinants that simplify their computation. Each of these properties is readily verified directly in the case of 2×2 determinants. Just as our definition of $n \times n$ determinants was inductive, the following discussion of Properties 1–7 of determinants is inductive. That is, we suppose that $n \ge 3$ and that these properties have already been verified for $(n - 1) \times (n - 1)$ determinants.

PROPERTY 1: If the $n \times n$ matrix **B** is obtained from **A** by multiplying a single row (or a column) of **A** by the constant *k*, then det **B** = *k* det **A**.

For instance, if the *i*th row of **A** is multiplied by *k*, then the elements *off* the *i*th row of **A** are unchanged. Hence for each j = 1, 2, ..., n, the *ij*th cofactors of **A** and **B** are equal: $A_{ij} = B_{ij}$. Therefore, expansion of **B** along the *i*th row gives

$$\det \mathbf{B} = (ka_{i1})B_{i1} + (ka_{i2})B_{i2} + \dots + (ka_{in})B_{in}$$

= $k(a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}),$

and thus det $\mathbf{B} = k \det \mathbf{A}$.

Property 1 implies simply that a constant can be factored out of a single row or column of a determinant. Thus we see that

$$\begin{vmatrix} 7 & 15 & -17 \\ -2 & 9 & 6 \\ 5 & -12 & 10 \end{vmatrix} = (3) \begin{vmatrix} 7 & 5 & -17 \\ -2 & 3 & 6 \\ 5 & -4 & 10 \end{vmatrix}$$

by factoring 3 out of the second column.

PROPERTY 2: If the $n \times n$ matrix **B** is obtained from **A** by interchanging two rows (or two columns), then det $\mathbf{B} = -\det \mathbf{A}$.

To see why this is so, suppose (for instance) that the first row is *not* one of the two that are interchanged (recall that $n \ge 3$). Then for each j = 1, 2, ..., n, the cofactor B_{1j} is obtained by interchanging two rows of the cofactor A_{1j} . Therefore, $B_{1j} = -A_{1j}$ by Property 2 for $(n - 1) \times (n - 1)$ determinants. Because $b_{1j} = a_{1j}$ for each j, it follows by expanding along the first row that

$$\det \mathbf{B} = b_{11}B_{11} + b_{12}B_{12} + \dots + b_{1n}B_{1n}$$

= $a_{11}(-A_{11}) + a_{12}(-A_{12}) + \dots + a_{1n}(-A_{1n})$
= $-(a_{11}A_{11} + a_{12}A_{12} + \dots + a_{1n}A_{1n}),$

and thus $\det \mathbf{B} = -\det \mathbf{A}$.

PROPERTY 3: If two rows (or two columns) of the $n \times n$ matrix **A** are identical, then det $\mathbf{A} = 0$.

To see why, let **B** denote the matrix obtained by interchanging the two identical rows of **A**. Then $\mathbf{B} = \mathbf{A}$, so det $\mathbf{B} = \det \mathbf{A}$. But Property 2 implies that det $\mathbf{B} = -\det \mathbf{A}$. Thus det $\mathbf{A} = -\det \mathbf{A}$, and it follows immediately that det $\mathbf{A} = 0$.

PROPERTY 4: Suppose that the $n \times n$ matrices A_1 , A_2 , and B are identical except for their *i* th rows—that is, the other n - 1 rows of the three matrices are identical—and that the *i* th row of **B** is the sum of the *i* th rows of A_1 and A_2 . Then

$$\det \mathbf{B} = \det \mathbf{A}_1 + \det \mathbf{A}_2.$$

This result also holds if columns are involved instead of rows.

Property 4 is readily established by expanding **B** along its *i* th row. In Problem 45 we ask you to supply the details for a typical case. The main importance (at this point) of Property 4 is that it implies the following property relating determinants and elementary row operations.

PROPERTY 5: If the $n \times n$ matrix **B** is obtained by adding a constant multiple of one row (or column) of **A** to another row (or column) of **A**, then det **B** = det **A**.

Thus the value of a determinant is *not* changed either by the type of elementary row operation described or by the corresponding type of elementary column operation. The following computation with 3×3 matrices illustrates the general proof of Property 5. Let

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} a_{11} & a_{12} & a_{13} + ka_{11} \\ a_{21} & a_{22} & a_{23} + ka_{21} \\ a_{31} & a_{32} & a_{33} + ka_{31} \end{bmatrix}$$

So **B** is the result of adding k times the first column of **A** to its third column. Then

$$\det \mathbf{B} = \begin{vmatrix} a_{11} & a_{12} & a_{13} + ka_{11} \\ a_{21} & a_{22} & a_{23} + ka_{21} \\ a_{31} & a_{32} & a_{33} + ka_{31} \end{vmatrix}$$
(11)
$$= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} + k \begin{vmatrix} a_{11} & a_{12} & a_{11} \\ a_{21} & a_{22} & a_{21} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} .$$

Here we first applied Property 4 and then factored k out of the second summand with the aid of Property 1. Now the first determinant on the right-hand side in (11) is simply det A, whereas the second determinant is zero (by Property 3—its first and third columns are identical). We therefore have shown that det $\mathbf{B} = \det \mathbf{A}$, as desired.

Although we use only elementary row operations in reducing a matrix to echelon form, Properties 1, 2, and 5 imply that we may use both elementary row operations and the analogous elementary column operations in simplifying the evaluation of determinants.

Example 6 The matrix

 $\mathbf{A} = \begin{bmatrix} 2 & -3 & -4 \\ -1 & 4 & 2 \\ 3 & 10 & 1 \end{bmatrix}$

has no zero elements to simplify the computation of its determinant *as it stands*, but we notice that we can "knock out" the first two elements of its third column by adding twice the first column to the third column. This gives

$$\det \mathbf{A} = \begin{vmatrix} 2 & -3 & -4 \\ -1 & 4 & 2 \\ 3 & 10 & 1 \end{vmatrix} = \begin{vmatrix} 2 & -3 & 0 \\ -1 & 4 & 0 \\ 3 & 10 & 7 \end{vmatrix}$$
$$= (+7) \begin{vmatrix} 2 & -3 \\ -1 & 4 \end{vmatrix} = 35.$$

The moral of the example is this: Evaluate determinants with your eyes open.

An **upper triangular matrix** is a square matrix having only zeros *below* its main diagonal. A **lower triangular matrix** is a square matrix having only zeros *above* its main diagonal. A **triangular matrix** is one that is either upper triangular or lower triangular, and thus looks like

[1]	6	10		1	0	0]
0	5	8	or	3	7	0	.
0	0	7		4	6	5_	

The next property tells us that determinants of triangular matrices are especially easy to evaluate.

PROPERTY 6: The determinant of a triangular matrix is equal to the product of its diagonal elements.

The reason is that the determinant of any triangular matrix can be evaluated as in the following computation:

$$\begin{vmatrix} 3 & 11 & 9 & 2 \\ 0 & -2 & 8 & -6 \\ 0 & 0 & 5 & 17 \\ 0 & 0 & 0 & -4 \end{vmatrix} = (3) \begin{vmatrix} -2 & 8 & -6 \\ 0 & 5 & 17 \\ 0 & 0 & -4 \end{vmatrix}$$
$$= (3)(-2) \begin{vmatrix} 5 & 17 \\ 0 & -4 \end{vmatrix} = (3)(-2)(5)(-4) = 120.$$

At each step we expand along the first column and pick up another diagonal element as a factor of the determinant.

Example 7 To evaluate

$$\begin{vmatrix} 2 & -1 & 3 \\ -2 & 1 & 5 \\ 4 & -2 & 10 \end{vmatrix},$$

we first add the first row to the second and then subtract twice the first row from the third. This yields

2	-1	3		2	-1	3	
-2	1	5	=	0	0	8	= 0,
4	-2	10		0	0	4	

because we produced a triangular matrix having a zero on its main diagonal. (Can you see an even quicker way to do it by keeping your eyes open?)

The Transpose of a Matrix

The transpose \mathbf{A}^T of a 2 × 2 matrix \mathbf{A} is obtained by interchanging its off-diagonal elements:

If
$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$
 then $\mathbf{A}^T = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$. (12)

More generally, the **transpose** of the $m \times n$ matrix $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ is the $n \times m$ matrix \mathbf{A}^T defined by

$$\mathbf{A}^T = \begin{bmatrix} a_{ji} \end{bmatrix}. \tag{13}$$

Note the reversal of subscripts; this means that the element of \mathbf{A}^T in its *j*th row and *i*th column is the element of \mathbf{A} in the *i*th row and the *j*th column of \mathbf{A} . Hence the rows of the transpose \mathbf{A}^T are (in order) the columns of \mathbf{A} , and the columns of \mathbf{A}^T are the rows of \mathbf{A} . Thus we obtain \mathbf{A}^T by changing the rows of \mathbf{A} into columns. For instance,

$$\begin{bmatrix} 2 & 0 & 5 \\ 4 & -1 & 7 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 \\ 0 & -1 \\ 5 & 7 \end{bmatrix}$$

and

$$\begin{bmatrix} 7 & -2 & 6\\ 1 & 2 & 3\\ 5 & 0 & 4 \end{bmatrix}^{T} = \begin{bmatrix} 7 & 1 & 5\\ -2 & 2 & 0\\ 6 & 3 & 4 \end{bmatrix}$$

If the matrix **A** is square, then we get \mathbf{A}^T by interchanging elements of **A** that are located symmetrically with respect to its main diagonal. Thus \mathbf{A}^T is the mirror reflection of **A** through its main diagonal.

In Problems 47 and 48 we ask you to verify the following properties of the transpose operation (under the assumption that **A** and **B** are matrices of appropriate sizes and c is a number):

(i) $(\mathbf{A}^T)^T = \mathbf{A};$ (ii) $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T;$ (iii) $(c\mathbf{A})^T = c\mathbf{A}^T;$ (iv) $(\mathbf{A}\mathbf{B})^T = \mathbf{B}^T\mathbf{A}^T.$

PROPERTY 7: If A is a square matrix, then $det(A^T) = det A$.

This property of determinants can be verified by writing the cofactor expansion of det A along its first *row* and the cofactor expansion of det(A^T) along its first *column*. When this is done, we see that the two expansions are identical, because the rows of A are the columns of A^T (as in Problem 49).

Determinants and Invertibility

We began this section with the remark that a 2×2 matrix **A** is invertible if and only if its determinant is nonzero: $|\mathbf{A}| \neq 0$. This result holds more generally for $n \times n$ matrices.

THEOREM 2 Determinants and Invertibility

The $n \times n$ matrix **A** is invertible if and only if det $\mathbf{A} \neq 0$.

This theorem (along with the others in this section) is proved in Appendix B. So now we can add the statement that det $\mathbf{A} \neq 0$ to the list of equivalent properties of nonsingular matrices stated in Theorem 7 of Section 3.5. Indeed, some texts *define* the square matrix \mathbf{A} to be nonsingular if and only if det $\mathbf{A} \neq 0$.

Example 8 According to Problem 61, the determinant of the matrix

$\mathbf{V} =$	1	a	a^2
	1	b	b^2
	1	С	c^2

is

$$|\mathbf{V}| = (b-a)(c-a)(c-b).$$

Note that $|\mathbf{V}| \neq 0$ if and only if the three numbers *a*, *b*, and *c* are *distinct*. Hence it follows from Theorem 2 that **V** is invertible if and only if *a*, *b*, and *c* are distinct. For instance, with a = -2, b = 3, and c = -4 we see that the matrix

[1	-2	4
1	3	9
1	-4	16

is invertible.

The connection between nonzero determinants and matrix invertibility is closely related to the fact that the determinant function "respects" matrix multiplication in the sense of the following theorem.

THEOREM 3 Determinants and Matrix Products

If **A** and **B** are $n \times n$ matrices, then

$$\mathbf{AB}| = |\mathbf{A}| \, |\mathbf{B}|. \tag{14}$$

The fact that $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$ seems so natural that we might fail to note that it is also quite remarkable. Contrast the simplicity of the equation $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$ with the complexity of the seemingly unrelated definitions of determinants and matrix products. For another contrast, we can mention that $|\mathbf{A} + \mathbf{B}|$ is generally *not* equal to $|\mathbf{A}| + |\mathbf{B}|$. (Pick a pair of 2 × 2 matrices and calculate their determinants and that of their sum.)

As a first application of Theorem 3, we can calculate the determinant of the *inverse* of an invertible matrix **A**:

$$\mathbf{A}\mathbf{A}^{-1}=\mathbf{I},$$

$$|\mathbf{A}| |\mathbf{A}^{-1}| = |\mathbf{A}\mathbf{A}^{-1}| = |\mathbf{I}| = 1,$$

and therefore

$$|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|}.\tag{15}$$

Now $|\mathbf{A}| \neq 0$ because **A** is invertible. Thus det(\mathbf{A}^{-1}) is the reciprocal of det **A**.

Example 9 The determinant of

$$\mathbf{A} = \begin{bmatrix} 2 & 2 & -2 & 0 \\ -4 & 2 & 3 & 1 \\ 0 & 4 & -1 & -4 \\ 3 & 1 & 3 & -1 \end{bmatrix}$$

happens to be $|\mathbf{A}| = 320 \neq 0$. Hence **A** is invertible, and *without* finding \mathbf{A}^{-1} we know from Eq. (15) that $|\mathbf{A}^{-1}| = 1/320$.

Cramer's Rule for $n \times n$ Systems

Suppose that we need to solve the $n \times n$ linear system

$$\mathbf{A}\mathbf{x} = \mathbf{b},\tag{16}$$

where

$$\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}.$$

We assume that the coefficient matrix **A** is invertible, so we know in advance that a unique solution **x** of (16) exists. The question is how to write **x** explicitly in terms of the coefficients a_{ij} and the constants b_i . In the following discussion, we think of **x** as a fixed (though as yet unknown) vector.

If we denote by $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ the column vectors of the $n \times n$ matrix **A**, then

$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}.$$

The desired formula for the *i*th unknown x_i involves the determinant of the matrix $\begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{b} & \cdots & \mathbf{a}_n \end{bmatrix}$ that is obtained upon replacing the *i*th column \mathbf{a}_i of \mathbf{A} with the constant vector \mathbf{b} .

THEOREM 4 Cramer's Rule

Consider the $n \times n$ linear system Ax = b, with

$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}.$$

If $|\mathbf{A}| \neq 0$, then the *i*th entry of the unique solution $\mathbf{x} = (x_1, x_2, \dots, x_n)$ is given by

$$x_{i} = \frac{|\mathbf{a}_{1} \cdots \mathbf{b} \cdots \mathbf{a}_{n}|}{|\mathbf{A}|}$$
$$= \frac{1}{|\mathbf{A}|} \begin{vmatrix} a_{11} \cdots b_{1} \cdots a_{1n} \\ a_{21} \cdots b_{2} \cdots a_{2n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} \cdots & b_{n} \cdots & a_{nn} \end{vmatrix},$$
(17)

where in the last expression the constant vector **b** replaces the *i*th column vector \mathbf{a}_i of **A**.

Example

Solution

For instance, the unique solution (x_1, x_2, x_3) of the 3 × 3 system

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3$$
(18)

with $|\mathbf{A}| = |a_{ij}| \neq 0$ is given by the formulas

$$x_{1} = \frac{1}{|\mathbf{A}|} \begin{vmatrix} b_{1} & a_{12} & a_{13} \\ b_{2} & a_{22} & a_{23} \\ b_{3} & a_{32} & a_{33} \end{vmatrix},$$

$$x_{2} = \frac{1}{|\mathbf{A}|} \begin{vmatrix} a_{11} & b_{1} & a_{13} \\ a_{21} & b_{2} & a_{23} \\ a_{31} & b_{3} & a_{33} \end{vmatrix},$$

$$x_{3} = \frac{1}{|\mathbf{A}|} \begin{vmatrix} a_{11} & a_{12} & b_{1} \\ a_{21} & a_{22} & b_{2} \\ a_{31} & a_{32} & b_{3} \end{vmatrix},$$
(19)

We find that

$$x_1 + 4x_2 + 5x_3 = 2$$

$$4x_1 + 2x_2 + 5x_3 = 3$$

$$-3x_1 + 3x_2 - x_3 = 1.$$

 $|\mathbf{A}| = \begin{vmatrix} 1 & 4 & 5 \\ 4 & 2 & 5 \\ -3 & 3 & -1 \end{vmatrix} = 29;$

then the formulas in (19) yield

$$x_{1} = \frac{1}{29} \begin{vmatrix} 2 & 4 & 5 \\ 3 & 2 & 5 \\ 1 & 3 & -1 \end{vmatrix} = \frac{33}{29},$$
$$x_{2} = \frac{1}{29} \begin{vmatrix} 1 & 2 & 5 \\ 4 & 3 & 5 \\ -3 & 1 & -1 \end{vmatrix} = \frac{35}{29},$$

and

$$x_3 = \frac{1}{29} \begin{vmatrix} 1 & 4 & 2 \\ 4 & 2 & 3 \\ -3 & 3 & 1 \end{vmatrix} = -\frac{23}{29}.$$

Inverses and the Adjoint Matrix

The inverse A^{-1} of the invertible $n \times n$ matrix A can be found by solving the matrix equation

$$\mathbf{A}\mathbf{X} = \mathbf{I}.\tag{20}$$

If we write the coefficient matrix $\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}$, the unknown matrix $\mathbf{X} = \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 & \cdots & \mathbf{x}_n \end{bmatrix}$, and the identity matrix $\mathbf{I} = \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \cdots & \mathbf{e}_n \end{bmatrix}$ in terms of their columns, then (20) says that

$$\mathbf{A}\mathbf{x}_j = \mathbf{e}_j \tag{21}$$

for j = 1, 2, ..., n. Each of these n equations can then be solved explicitly using Cramer's rule. In particular, Eq. (17) says that the *i*th element of the column matrix \mathbf{x}_i is given by

$$x_{ij} = \frac{\begin{vmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_{i-1} & \mathbf{e}_j & \mathbf{a}_{i+1} & \cdots & \mathbf{a}_n \end{vmatrix}}{|\mathbf{A}|}$$
(22)

for i, j = 1, 2, ..., n. When these results are assembled (Appendix B), we get an explicit formula for the inverse matrix.

THEOREM 5 The Inverse Matrix

The inverse of the invertible matrix A is given by the formula

$$\mathbf{A}^{-1} = \frac{\left[A_{ij}\right]^T}{|\mathbf{A}|},\tag{23}$$

where, as usual, A_{ij} denotes the *ij*th cofactor of **A**; that is, A_{ij} is the product of $(-1)^{i+j}$ and the *ij*th minor determinant of **A**.

We see in (23) the *transpose* of the **cofactor matrix** $\begin{bmatrix} A_{ij} \end{bmatrix}$ of the $n \times n$ matrix A. This transposed cofactor matrix is called the adjoint matrix of A and is denoted by

$$\operatorname{adj} \mathbf{A} = \begin{bmatrix} A_{ij} \end{bmatrix}^T = \begin{bmatrix} A_{ji} \end{bmatrix}.$$
(24)

Example 11 Apply the formula in (23) to find the inverse of the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 4 & 5 \\ 4 & 2 & 5 \\ -3 & 3 & -1 \end{bmatrix}$$

of Example 10, in which we saw that $|\mathbf{A}| = 29$.

Solution

First we calculate the cofactors of **A**, arranging our computations in a natural 3×3 array:

$$A_{11} = + \begin{vmatrix} 2 & 5 \\ 3 & -1 \end{vmatrix} = -17, \quad A_{12} = - \begin{vmatrix} 4 & 5 \\ -3 & -1 \end{vmatrix} = -11, \quad A_{13} = + \begin{vmatrix} 4 & 2 \\ -3 & 3 \end{vmatrix} = -18,$$

$$A_{21} = - \begin{vmatrix} 4 & 5 \\ 3 & -1 \end{vmatrix} = -19, \quad A_{22} = + \begin{vmatrix} 1 & 5 \\ -3 & -1 \end{vmatrix} = -14, \quad A_{23} = - \begin{vmatrix} 1 & 4 \\ -3 & 3 \end{vmatrix} = -15,$$

$$A_{31} = + \begin{vmatrix} 4 & 5 \\ 2 & 5 \end{vmatrix} = -10, \quad A_{32} = - \begin{vmatrix} 1 & 5 \\ 4 & 5 \end{vmatrix} = -15, \quad A_{33} = + \begin{vmatrix} 1 & 4 \\ 4 & 2 \end{vmatrix} = -14.$$

(Note the familiar checkerboard pattern of signs.) Thus the cofactor matrix of A is

$$\begin{bmatrix} A_{ij} \end{bmatrix} = \begin{bmatrix} -17 & -11 & 18 \\ 19 & 14 & -15 \\ 10 & 15 & -14 \end{bmatrix}.$$

Next, we interchange rows and columns to obtain the adjoint matrix

adj
$$\mathbf{A} = \begin{bmatrix} A_{ij} \end{bmatrix}^T = \begin{bmatrix} -17 & 19 & 10 \\ -11 & 14 & 15 \\ 18 & -15 & -14 \end{bmatrix}.$$

Finally, in accord with Eq. (23), we divide by $|\mathbf{A}| = 29$ to get the inverse matrix

$$\mathbf{A}^{-1} = \frac{1}{29} \begin{bmatrix} -17 & 19 & 10\\ -11 & 14 & 15\\ 18 & -15 & -14 \end{bmatrix}.$$

Computational Efficiency

The amount of labor required to complete a numerical calculation is measured by the number of arithmetical operations it involves. So let us count just the number of multiplications required to evaluate an $n \times n$ determinant using cofactor expansions. If n = 5, for instance, then the cofactor expansion along a row or column requires computations of *five* 4×4 determinants. A cofactor expansion of each of these five 4×4 determinants involves *four* 3×3 determinants. Each of these four 3×3 determinants leads to *three* 2×2 determinants, and finally each of these 2×2 determinants requires *two* multiplications for its evaluation. Hence the total number of multiplications required to evaluate the original 5×5 determinant is

$$5 \cdot 4 \cdot 3 \cdot 2 = 5! = 120$$

In general, the number of multiplications required to evaluate an $n \times n$ determinant completely by cofactor expansions is

$$n! = n \cdot (n-1) \cdots 3 \cdot 2 \cdot 1$$
 (*n* factorial).

Thus, if we ignore the smaller number of additions involved, then our *operations count* for an $n \times n$ determinant is n!. For instance, the evaluation of a 25 × 25 determinant by cofactor expansion would require about $25! \approx 1.55 \times 10^{25}$ operations. If we used a supercomputer capable of a billion operations per second this task would require about $(1.55 \times 10^{16})/(365.25 \times 24 \times 3600) \approx 492$ million years! This same supercomputer would require about 9.64×10^{47} years—incomparably longer than the estimated age of the universe (perhaps 20 billion years)—to evaluate a 50 × 50 determinant by cofactor expansion. Contemporary scientific and engineering applications routinely involve matrices of size 1000 × 1000 (or larger) whose cofactor expansions would require incomprehensibly long times with any conceivable computer.

However, a typical 2000-era desktop computer, using MATLAB and performing "only" about 100 million operations per second, calculates the determinant or inverse of a randomly generated 100×100 matrix almost instantaneously. How is this possible? Obviously not by using cofactor expansions!

The answer is that determinants are much more efficiently calculated by Gaussian elimination—that is, by reduction to triangular form, so that only the product of the remaining diagonal elements need be calculated. Similarly, the inverse of the invertible matrix **A** is much more efficiently calculated by reduction of the augmented matrix $\begin{bmatrix} A & I \end{bmatrix}$ to reduced echelon form (as in Section 3.5) than by use of Cramer's rule (as in Theorem 5 here). A careful count reveals that the number of arithmetic operations required to reduce an $n \times n$ matrix to echelon form is of the order of n^3 rather than n!. As indicated in the table of Fig. 3.6.1, n^3 is dramatically smaller than n! if n is fairly large. Consequently, cofactor expansions of determinants and Cramer's rule for the solution of systems are primarily of theoretical importance, and are seldom used in numerical problems where n is greater than 3 or 4.

n	5	10	25	50	100
$\frac{2}{3}n^{3}$	83	667	10,417	83,333	666,667
n!	120	3,628,800	1.55×10^{25}	3.04×10^{64}	9.33×10^{157}

FIGURE 3.6.1. Approximate operations counts for evaluating an $n \times n$ determinant by Gaussian elimination $(\frac{2}{3}n^3)$ and by cofactor expansion (*n*!).

3.6 **Problems**

Use cofactor expansions to evaluate the determinants in Problems 1–6. Expand along the row or column that minimizes the amount of computation that is required.

1.

$$\begin{pmatrix} 0 & 0 & 3 \\ 4 & 0 & 0 \\ 0 & 5 & 0 \end{pmatrix}$$
 2.
 $\begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$

 3.
 $\begin{pmatrix} 1 & 0 & 0 & 0 \\ 2 & 0 & 5 & 0 \\ 3 & 6 & 9 & 8 \\ 4 & 0 & 10 & 7 \end{pmatrix}$
 4.
 $\begin{pmatrix} 5 & 11 & 8 & 7 \\ 3 & -2 & 6 & 23 \\ 0 & 0 & 0 & -3 \\ 0 & 4 & 0 & 17 \end{pmatrix}$

 5.
 $\begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 4 \\ 0 & 5 & 0 & 0 & 0 \end{pmatrix}$
 6.

 6.
 $\begin{pmatrix} 3 & 0 & 11 & -5 & 0 \\ -2 & 4 & 13 & 6 & 5 \\ 0 & 0 & 5 & 0 & 0 \\ 7 & 6 & -9 & 17 & 7 \\ 0 & 0 & 8 & 2 & 0 \end{pmatrix}$

In Problems 7–12, evaluate each given determinant after first simplifying the computation (as in Example 6) by adding an appropriate multiple of some row or column to another.

7.	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	8.	$2 \\ -2 \\ 3$	3 - 3 2	4 1 7	
9.	$\begin{vmatrix} 3 & -2 & 5 \\ 0 & 5 & 17 \\ 6 & -4 & 12 \end{vmatrix}$	10.	-3 1 2	6 -2 -5	5 -4 12	
11.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	12.	$\begin{array}{c} 2\\ 0\\ 0\\ -4 \end{array}$	0 1 0 0	0 11 5 0	$\begin{array}{c c} -3 \\ 12 \\ 13 \\ 7 \end{array}$

Use the method of elimination to evaluate the determinants in Problems 13–20.

13.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$
15.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$16. \begin{vmatrix} 2 & 4 & -2 \\ -5 & -4 & -1 \\ -4 & 2 & 1 \end{vmatrix}$
17.	$\begin{vmatrix} 2 & 3 & 3 & 1 \\ 0 & 4 & 3 & -3 \\ 2 & -1 & -1 & -3 \\ 0 & -4 & -3 & 2 \end{vmatrix}$	
18.	$\begin{vmatrix} 1 & 4 & 4 & 1 \\ 0 & 1 & -2 & 2 \\ 3 & 3 & 1 & 4 \\ 0 & 1 & -3 & -2 \end{vmatrix}$	

	1	0	0	3
10	0	1	-2	0
19.	-2	3	-2	3
	0	-3	3	3
	1	2	1	-1
20	2	1	3	3
20.	0	1	-2	3
	-1	4	-2	4

Use Cramer's rule to solve the systems in Problems 21–32.

21.
$$3x + 4y = 2$$

 $5x + 7y = 1$ **22.** $5x + 8y = 3$
 $8x + 13y = 5$ **23.** $17x + 7y = 6$
 $12x + 5y = 4$ **24.** $11x + 15y = 10$
 $8x + 11y = 7$

- **25.** 5x + 6y = 123x + 4y = 6**26.** 6x + 7y = 38x + 9y = 4
- **27.** $5x_1 + 2x_2 2x_3 = 1$ $x_1 + 5x_2 - 3x_3 = -2$ $5x_1 - 3x_2 + 5x_3 = 2$
- **28.** $5x_1 + 4x_2 2x_3 = 4$ $2x_1 + 3x_3 = 2$ $2x_1 - x_2 + x_3 = 1$
- **29.** $3x_1 x_2 5x_3 = 3$ $4x_1 - 4x_2 - 3x_3 = -4$ $x_1 - 5x_3 = 2$
- **30.** $x_1 + 4x_2 + 2x_3 = 3$ $4x_1 + 2x_2 + x_3 = 1$ $2x_1 - 2x_2 - 5x_3 = -3$
- **31.** $2x_1 5x_3 = -3$ $4x_1 - 5x_2 + 3x_3 = -3$ $-2x_1 + x_2 + x_3 = -1$
- **32.** $3x_1 + 4x_2 3x_3 = 5$ $3x_1 - 2x_2 + 4x_3 = 7$ $3x_1 + 2x_2 - x_3 = 3$

Apply Theorem 5 to find the inverse A^{-1} of each matrix A given in Problems 33–40.

33.	$\begin{bmatrix} -5\\1\\5 \end{bmatrix}$	$-2 \\ 5 \\ -3$	$\begin{bmatrix} 2\\ -3\\ 1 \end{bmatrix}$	34.	$\begin{bmatrix} 2\\ -5\\ 2 \end{bmatrix}$	$0 \\ -4 \\ -1$	$\begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$
35.	$\begin{bmatrix} 3\\ -2\\ -5 \end{bmatrix}$	5 3 0	$\begin{bmatrix} 2\\ -4\\ -5 \end{bmatrix}$	36.	$\begin{bmatrix} -4\\ 3\\ 1 \end{bmatrix}$	$\begin{array}{c} 4\\ -1\\ 0 \end{array}$	$\begin{bmatrix} 3\\-5\\-5 \end{bmatrix}$
37.	$\begin{bmatrix} -4\\ -2\\ -3 \end{bmatrix}$	1 4 -3	$\begin{bmatrix} 5\\5\\-1 \end{bmatrix}$	38.	$\begin{bmatrix} 3\\ 3\\ -3 \end{bmatrix}$	4 2 2	$\begin{bmatrix} -3\\ -1\\ -4 \end{bmatrix}$
39.	$\begin{bmatrix} -3\\0\\2 \end{bmatrix}$	$-2 \\ 3 \\ 3$	$\begin{bmatrix} 3\\2\\-5 \end{bmatrix}$	40.	$\begin{bmatrix} 2\\ 2\\ -5 \end{bmatrix}$	$ \begin{array}{c} 4 \\ -3 \\ 0 \end{array} $	$\begin{bmatrix} -3\\ -1\\ -3 \end{bmatrix}$

41. Show that $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$ if **A** and **B** are arbitrary 2×2 matrices.
42. Consider the 2×2 matrices

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix}$$

where x and y denote the row vectors of **B**. Then the product **AB** can be written in the form

$$\mathbf{AB} = \begin{bmatrix} a\mathbf{x} + b\mathbf{y} \\ c\mathbf{x} + d\mathbf{y} \end{bmatrix}.$$

Use this expression and the properties of determinants to show that

det
$$\mathbf{AB} = (ad - bc) \begin{vmatrix} \mathbf{x} \\ \mathbf{y} \end{vmatrix} = (\det \mathbf{A})(\det \mathbf{B}).$$

Thus the determinant of a product of 2×2 matrices is equal to the product of their determinants.

Each of Problems 43-46 lists a special case of one of Property 1 through Property 5. Verify it by expanding the determinant on the left-hand side along an appropriate row or column.

.

$$43. \begin{vmatrix} ka_{11} & a_{12} & a_{13} \\ ka_{21} & a_{22} & a_{23} \\ ka_{31} & a_{32} & a_{33} \end{vmatrix} = k \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$44. \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{11} & a_{12} & a_{13} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = - \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$45. \begin{vmatrix} a_{1} & b_{1} & c_{1} + d_{1} \\ a_{2} & b_{2} & c_{2} + d_{2} \\ a_{3} & b_{3} & c_{3} + d_{3} \end{vmatrix} = \begin{vmatrix} a_{1} & b_{1} & c_{1} \\ a_{2} & b_{2} & c_{2} \\ a_{3} & b_{3} & c_{3} \end{vmatrix} + \begin{vmatrix} a_{1} & b_{1} & d_{1} \\ a_{2} & b_{2} & c_{2} \\ a_{3} & b_{3} & c_{3} \end{vmatrix}$$

$$46. \begin{vmatrix} a_{11} + ka_{12} & a_{12} & a_{13} \\ a_{21} + ka_{22} & a_{22} & a_{23} \\ a_{31} + ka_{32} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

Problems 47 through 49 develop properties of matrix transposes.

- 47. Suppose that A and B are matrices of the same size. Show that: (a) $(\mathbf{A}^T)^T = \mathbf{A}$; (b) $(c\mathbf{A})^T = c\mathbf{A}^T$; and (c) $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$.
- 48. Let A and B be matrices such that AB is defined. Show that $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$. Begin by recalling that the *ij*th element of **AB** is obtained by multiplying elements in the *i*th row of **A** with those in the *j*th column of **B**. What is the *ii*th element of $\mathbf{B}^T \mathbf{A}^T$?
- **49.** Let $\mathbf{A} = \begin{bmatrix} a_{ii} \end{bmatrix}$ be a 3 × 3 matrix. Show that det(\mathbf{A}^T) = det **A** by expanding det **A** along its first row and det(\mathbf{A}^T) along its first column.
- **50.** Suppose that $\mathbf{A}^2 = \mathbf{A}$. Prove that $|\mathbf{A}| = 0$ or $|\mathbf{A}| = 1$.
- 51. Suppose that $A^n = 0$ (the zero matrix) for some positive integer *n*. Prove that $|\mathbf{A}| = 0$.
- 52. The square matrix A is called orthogonal provided that $A^T = A^{-1}$. Show that the determinant of such a matrix must be either +1 or -1.
- 53. The matrices A and B are said to be similar provided that $\mathbf{A} = \mathbf{P}^{-1}\mathbf{B}\mathbf{P}$ for some invertible matrix **P**. Show that if **A** and **B** are similar, then $|\mathbf{A}| = |\mathbf{B}|$.

- **54.** Deduce from Theorems 2 and 3 that if **A** and **B** are $n \times n$ invertible matrices, then **AB** is invertible if and only if both A and B are invertible.
- 55. Let A and B be $n \times n$ matrices. Suppose it is known that either AB = I or BA = I. Use the result of Problem 54 to conclude that $\mathbf{B} = \mathbf{A}^{-1}$.
- **56.** Let **A** be an $n \times n$ matrix with det $\mathbf{A} = 1$ and with all elements of A integers.
 - (a) Show that A^{-1} has only integer entries.
 - (b) Suppose that **b** is an *n*-vector with only integer entries. Show that the solution vector \mathbf{x} of $\mathbf{A}\mathbf{x} = \mathbf{b}$ has only integer entries.
- **57.** Let A be a 3×3 upper triangular matrix with nonzero determinant. Show by explicit computation that A^{-1} is also upper triangular.
- 58. Figure 3.6.2 shows an acute triangle with angles A, B, and C and opposite sides a, b, and c. By dropping a perpendicular from each vertex to the opposite side, derive the equations

$$c \cos B + b \cos C = a$$

$$c \cos A + a \cos C = b$$

$$a \cos B + b \cos A = c.$$

Regarding these as linear equations in the unknowns $\cos A$, $\cos B$, and $\cos C$, use Cramer's rule to derive the law of cosines by solving for

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

Thus

$$a^2 = b^2 + c^2 - 2bc \cos A.$$

Note that the case $A = \pi/2$ (90°) reduces to the Pythagorean theorem.



FIGURE 3.6.2. The triangle of Problem 58.

59. Show that

$$\begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 3$$
 and $\begin{vmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{vmatrix} = 4.$

60. Consider the $n \times n$ determinant

$$B_n = \begin{vmatrix} 2 & 1 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 2 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 2 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 2 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 1 & 2 \end{vmatrix}$$

in which each entry on the main diagonal is a 2, each entry on the two adjacent diagonals is a 1, and every other entry is zero.

3.7 LINEAR EQUATIONS AND CURVE FITTING 203

(a) Expand along the first row to show that

$$B_n = 2B_{n-1} - B_{n-2}.$$

(b) Prove by induction on *n* that $B_n = n + 1$ for $n \ge 2$.

Problems 61-64 deal with the Vandermonde determinant

$$V(x_1, x_2, \dots, x_n) = \begin{vmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^{n-1} \end{vmatrix}$$

that will play an important role in Section 3.7.

61. Show by direct computation that V(a, b) = b - a and that

$$V(a,b,c) = \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} = (b-a)(c-a)(c-b).$$

62. The formulas in Problem 61 are the cases n = 2 and n = 3 of the general formula

$$V(x_1, x_2, \dots, x_n) = \prod_{\substack{i, j = 1 \\ i > j}}^n (x_i - x_j).$$
(25)

The case n = 4 is

$$V(x_1, x_2, x_3, x_4) = (x_2 - x_1)(x_3 - x_1)(x_3 - x_2)$$
$$\times (x_4 - x_1)(x_4 - x_2)(x_4 - x_3).$$

Prove this as follows. Given x_1 , x_2 , and x_3 , define the cubic polynomial P(y) to be

$$P(y) = \begin{vmatrix} 1 & x_1 & x_1^2 & x_1^3 \\ 1 & x_2 & x_2^2 & x_2^3 \\ 1 & x_3 & x_3^2 & x_3^3 \\ 1 & y & y^2 & y^3 \end{vmatrix}.$$
 (26)

Because $P(x_1) = P(x_2) = P(x_3) = 0$ (why?), the roots of P(y) are x_1, x_2 , and x_3 . It follows that

$$P(y) = k(y - x_1)(y - x_2)(y - x_3),$$

where k is the coefficient of y^3 in P(y). Finally, observe that expansion of the 4×4 determinant in (26) along its last row gives $k = V(x_1, x_2, x_3)$ and that $V(x_1, x_2, x_3, x_4) = P(x_4)$.

63. Generalize the argument in Problem 62 to prove the formula in (25) by induction on *n*. Begin with the (n - 1)st-degree polynomial

$$P(y) = \begin{vmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n-1} & x_{n-1}^2 & \cdots & x_{n-1}^{n-1} \\ 1 & y & y^2 & \cdots & y^{n-1} \end{vmatrix}$$

64. Use the formula in (25) to evaluate the two determinants given next.

(a)
$$\begin{vmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \end{vmatrix}$$

(b)
$$\begin{vmatrix} 1 & -1 & 1 & -1 \\ 1 & 2 & 4 & 8 \\ 1 & -2 & 4 & -8 \\ 1 & 3 & 9 & 27 \end{vmatrix}$$

3.7 Linear Equations and Curve Fitting



FIGURE 3.7.1. A curve y = f(x) interpolating (that is, passing through) given data points.

Linear algebra has important applications to the common scientific problem of representing empirical data by means of equations or functions of specified types. We give here only a brief introduction to this extensive subject.

Typically, we begin with a collection of given *data points* $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$ that are to be represented by a specific type of function y = f(x). For instance, y might be the volume of a sample of gas when its temperature is x. Thus the given data points are the results of experiment or measurement, and we want to determine the curve y = f(x) in the xy-plane so that it passes through each of these points; see Figure 3.7.1. Thus we speak of "fitting" the curve to the data points.

We will confine our attention largely to polynomial curves. A **polynomial of degree** *n* is a function of the form

$$f(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n,$$
(1)

where the coefficients $a_0, a_1, a_2, ..., a_n$ are constants. The data point (x_i, y_i) lies on the curve y = f(x) provided that $f(x_i) = y_i$. The condition that this be so for each

 $i = 0, 1, 2, \ldots, n$ yields the n + 1 equations

$$a_{0} + a_{1}x_{0} + a_{2}(x_{0})^{2} + \dots + a_{n}(x_{0})^{n} = y_{0}$$

$$a_{0} + a_{1}x_{1} + a_{2}(x_{1})^{2} + \dots + a_{n}(x_{1})^{n} = y_{1}$$

$$a_{0} + a_{1}x_{2} + a_{2}(x_{2})^{2} + \dots + a_{n}(x_{2})^{n} = y_{2}$$

$$\vdots$$

$$a_{0} + a_{1}x_{n} + a_{2}(x_{n})^{2} + \dots + a_{n}(x_{n})^{n} = y_{n}.$$
(2)

Because the numbers x_i and y_i are given, this is a system of n + 1 linear equations in the n + 1 unknowns $a_0, a_1, a_2, \ldots, a_n$ (the coefficients that determine the polynomial in (1)).

The $(n + 1) \times (n + 1)$ coefficient matrix of the system in (1) is the Vandermonde matrix

$$\mathbf{A} = \begin{bmatrix} 1 & x_0 & (x_0)^2 & \cdots & (x_0)^n \\ 1 & x_1 & (x_1)^2 & \cdots & (x_1)^n \\ 1 & x_2 & (x_2)^2 & \cdots & (x_2)^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & (x_n)^2 & \cdots & (x_n)^n \end{bmatrix},$$
(3)

whose determinant is discussed in Problems 61–63 of Section 3.6. It follows from Eq. (25) there that, if the *x*-coordinates $x_0, x_1, x_2, \ldots, x_n$ are distinct, then the matrix **A** is nonsingular. Hence Theorem 7 in Section 3.5 implies that the system in (2) has a unique solution for the coefficients $a_0, a_1, a_2, \ldots, a_n$ in (1). Thus there is a unique *n*th degree polynomial that fits the n + 1 given data points. We call it an **interpolating polynomial**, and say that it **interpolates** the given points.

Example 1 Find a cubic polynomial of the form

$$y = A + Bx + Cx^2 + Dx^3$$

that interpolates the data points (-1, 4), (1, 2), (2, 1), and (3, 16).

Solution

In a particular problem, it generally is simpler to use distinct capital letters rather than subscripted symbols to denote the coefficients. Here we want to find the values of *A*, *B*, *C*, and *D* so that y(-1) = 4, y(1) = 2, y(2) = 1, and y(3) = 16. These conditions yield the four linear equations

$$A - B + C - D = 4$$

$$A + B + C + D = 2$$

$$A + 2B + 4C + 8D = 1$$

$$A + 3B + 9C + 27D = 16.$$

We readily reduce this system to the echelon form

$$A - B + C - D = 4$$
$$B + D = -1$$
$$C + 2D = 0$$
$$D = 2,$$

and then back substitution yields A = 7, B = -3, C = -4, and D = 2. Thus the desired cubic polynomial is

$$y = 7 - 3x - 4x^2 + 2x^3$$

15

FIGURE 3.7.2. Cubic curve through the four data points of Example 1.

The graph of this cubic is shown in Fig. 3.7.2, along with the four original data points.

Modeling World Population Growth

As a concrete example of interpolation, we consider the growth of the world's human population. The table in Fig. 3.7.3 shows the total world population (in billions) at 5-year intervals. Actual populations are shown for the years 1975–2010. The figures listed for the years 2015–2040 are the world populations that were predicted by the United Nations on the basis of detailed demographic analysis of population trends during the 20th century on a country-by-country basis throughout the world. Each entry of the final column of this table gives the average annual percentage growth rate during the preceding 5-year period. For instance, $4.062(1.018)^5 \approx 4.44$ for the growth during the 5-year period 1975–1980, so the average annual growth during this period is about 1.8%.

	World Population	Percent
Year	(billions)	Growth
1975	4.062	
1980	4.440	1.80%
1985	4.853	1.79%
1990	5.310	1.82%
1995	5.735	1.55%
2000	6.127	1.33%
2005	6.520	1.25%
2010	6.930	1.23%
2015	7.349	1.18%
2020	7.758	1.09%
2025	8.142	0.97%
2030	8.501	0.87%
2035	8.839	0.78%
2040	9.157	0.71%

FIGURE 3.7.3. World population data.

We see that the world population grew at an annual rate of about 1.8% during the 1980s, but the rate of growth has slowed since then, and it is expected to slow even more during the coming decades of the 21st century. In particular, the growth of the world population at the present time in history is *not* natural or exponential in character—that characterization would imply a constant percentage rate of growth. We explore the possibility of interpolating world population data with polynomial models that might be usable to predict future populations. It seems natural to expect better results with higher-degree interpolating polynomials. Let's see whether this is so.

Example 2 First, we fit a linear polynomial $P_1(t) = a + bt$ (with t = 0 in 1900) to the 1995 and 2005 world population values. We need only solve the equations

$$a + 95b = 5.735$$

 $a + 105b = 6.520$

for a = -1.7225, b = 0.0785. Thus our linear interpolating polynomial is

$$P_1(t) = -1.7225 + 0.0785t.$$
⁽⁴⁾

206 CHAPTER 3 LINEAR SYSTEMS AND MATRICES

Example 3

1	95	95 ²	1	15.7	351	
1	100	100 ²	*	6.1	27	
1	105	105 ²		16.5	201	
					[-1.	523]
					0.0	745
					L.ZE	-5.1

FIGURE 3.7.4. TI-84 Plus CE calculator solution of the 3×3 system in Example 3.

Example 4

Now let's fit a quadratic polynomial $P_2(t) = a + bt + ct^2$ (with t = 0 in 1900) to the 1995, 2000, and 2005 world population values. With the three data points (95, 5.735), (100, 6.127), and (105, 6.520), the system in (2) yields the equations

$$a + 95b + 952c = 5.735$$

$$a + 100b + 1002c = 6.127$$

$$a + 105b + 1052c = 6.520$$

having the calculator solution a = -1.523, b = 0.0745, c = 0.00002 (Fig. 3.7.4). Thus our quadratic interpolating polynomial is

$$P_2(t) = -1.523 + 0.0745t + 0.00002t^2.$$
⁽⁵⁾

Next we fit a cubic polynomial $P_3(t) = a + bt + ct^2 + dt^3$ (with t = 0 in 1900) to the 1995, 2000, 2005, and 2010 world population values. With the four data points (95, 5.735), (100, 6.127), (105, 6.520), and (110, 6.930), the system in (2) yields the four equations

 $5.735 = a + 95b + 95^{2}c + 95^{3}d$ $6.127 = a + 100b + 100^{2}c + 100^{3}d$ $6.520 = a + 105b + 105^{2}c + 105^{3}d$ $6.930 = a + 110b + 110^{2}c + 110^{3}d.$

As in the 3.5 Application, a calculator or computer yields the solution

$\begin{bmatrix} a \\ b \end{bmatrix}$		1	95 100	95^2 100^2	$\begin{bmatrix} 95^3 \\ 100^3 \end{bmatrix}^{-1}$	-1	5.735 6.127		-22.803 0.713967	
$\begin{bmatrix} c \\ d \end{bmatrix}$	=	1 1	105 110	$105^2 \\ 110^2$	$105^3 \\ 110^3$		6.520 6.930	=	-0.00638 0.000021333	.

Thus our cubic interpolating polynomial is

$$P_3(t) = -22.803 + 0.713967t - 0.00638t^2 + 0.000021333t^3.$$
(6)

Example 5

In order to fit a fourth-degree population model of the form

$$P_4(t) = a + bt + ct^2 + dt^3 + et^4$$

to the 1990-1995-2000-2005-2010 world population data, we need to solve the linear system

$\begin{bmatrix} 1\\1\\1\\1\\1\\\end{bmatrix}$	90 95 100 105	90^{2} 95^{2} 100^{2} 105^{2}	90^{3} 95^{3} 100^{3} 105^{3}	90 ⁴ 95 ⁴ 100 ⁴ 105 ⁴	$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$	=	5.310 5.735 6.127 6.520	
	105 110	105^{2} 110^{2}	105^{3} 110^{3}	105^{+} 110^{4}	$\begin{bmatrix} a \\ e \end{bmatrix}$		6.520	

to find the values of the coefficients a, b, c, d, and e. The result is

$$P_4(t) = -154.473 + 5.867667t - 0.08195t^2 + 0.00051333t^3 - 0.0000012t^4.$$
⁽⁷⁾

The table in Fig. 3.7.5 compares our linear, quadratic, cubic, and quartic predictions with the "correct" United Nations prediction for the year 2030. Each "error" in the third column of this table is the amount by which the corresponding prediction undershoots (positive error) or overshoots (negative error) the U.N. prediction. We see that the quadratic prediction is better than the linear but also markedly

	Year 2030	
	Prediction	Error
Linear	8.482	+0.019
Quadratic	8.500	+0.001
Cubic	9.060	-0.559
Quartic	8.430	+0.071
United Nations	8.501	

FIGURE 3.7.5. Predictions of the 2030 world population.



FIGURE 3.7.6. Comparison of world population data and the interpolating polynomials $P_n(t)$ for n = 2, 3, 4.

better than the cubic prediction. The quartic prediction is an improvement over the cubic, yet still not as good as the quadratic. Thus there is at best an uncertain relationship between the degree of the polynomial model and the accuracy of its predictions.

Figure 3.7.6 shows the U.N. world population data points for the years 1975 through 2040, together with the plots of the quadratic, cubic, and quartic population functions of Examples 3, 4, and 5. (The plot of the linear population function of Example 2 is virtually indistinguishable from that of the quadratic function for the values of t shown in the figure.) It looks as though the more work we do to find a polynomial fitting selected data points, the less we get for our effort. It is certainly true in this figure that—outside the interval from 1990 to 2010—the higher the degree of the polynomial, the worse it appears to fit the given data points. The issue here is the difference between

- *interpolating* data points *within* the interval of given points being fitted, and
- *extrapolating* data points *outside* this interval.

All four of our polynomials appear to do a good job of interpolating but, somewhat paradoxically, the higher the degree, the worse the apparent accuracy of extrapolation. The highly questionable accuracy of data extrapolation outside the interval of interpolation has significant implications. For instance, consider a news report that when a certain alleged carcinogen was fed to mice in sufficient amounts to kill an elephant, the mice developed cancer. It is then argued that moderate amounts of this carcinogen may cause cancer in humans; or that if 1 part per billion of this carcinogen in the environment kills 1 person, then 1 part per million (a thousand times as much) will kill 1000 people. Such arguments are common, but they may well be cases of extrapolation beyond the range of accuracy. The bottom line is that interpolation is fairly safe—though hardly fail-safe—but extrapolation is risky.

Geometric Applications

In contrast with population prediction, there are interesting situations where curve fitting is exact. For instance, the fact that "two points determine a line" in the plane means that, when we fit the linear function y = a + bx to a given pair of points, we get precisely the one and only straight line in the plane that passes through these points. Similarly, "three points determine a circle," meaning that there is one and only one circle in the plane that passes through three given noncollinear points. In



FIGURE 3.7.7. The circle with center (h, k) and radius r.

Example 6

Solution

(F1700)	F2	*	Y F3+	Y FHT	Y F5	Y	F	67	7
-f-	Alge	bra	Calo	Other	PrgmI	0 0	lea	in l	JP
[-1	5	1	-26	ř.	[-1	Ĕ.	5	1	-26
• 5	-3	1	-34	÷a	5		-3	1	-34
6	4	1	-52	5.00	6		4	1	-52
					1	1	0	0	-4
= rre	f(a)					0	1	0	-2
2000205					1	0	0	1	-20
rref	(a)		020020	10 J	35655	0.0	500 C		
MAIN			RAD AUT	0	FUNC	22	30		

FIGURE 3.7.8. TI-89 calculator solution of the 3×3 system in Example 6.

order to find this particular circle, we recall that the equation of a circle with center (h, k) and radius r is

$$(x-h)^2 + (y-k)^2 = r^2$$
 (Fig. 3.7.7). (8)

Simplification gives

$$(x^{2} - 2hx + h^{2}) + (y^{2} - 2ky + k^{2}) = r^{2},$$

that is,

$$x^2 + y^2 + Ax + By + C = 0 (9)$$

(where A = -2h, B = -2k, and $C = h^2 + k^2 - r^2$) as the general equation of a circle in the plane.

Find the equation of the circle that is determined by the points P(-1, 5), Q(5, -3), and R(6, 4).

Substitution of the xy-coordinates of each of the three points P, Q, and R into (9) gives the three equations

$$-A + 5B + C = -26$$

 $5A - 3B + C = -34$
 $6A + 4B + C = -52.$

Reduction of the corresponding augmented coefficient matrix to reduced row-echelon form (Fig. 3.7.8) yields A = -4, B = -2, and C = -20. Thus the equation of the desired circle is

$$x^2 + y^2 - 4x - 2y - 20 = 0.$$

To find its center and radius, we complete the squares in x and y and get

$$(x-2)^2 + (y-1)^2 = 25$$

Thus the circle has center (2, 1) and radius 5 (Fig. 3.7.9).



FIGURE 3.7.9. The circle of Example 6.

FIGURE 3.7.10. A rotated central ellipse.

х

Three appropriate points in the plane also determine a **central conic** with equation of the form

$$Ax^2 + Bxy + Cy^2 = 1.$$
 (10)

This is a rotated conic section—an ellipse, parabola, or hyperbola—centered at the origin of the xy-coordinate system. Figure 3.7.10 shows a typical rotated ellipse in the plane.

Example 7

Solution

Find the equation of the central conic that passes through the same three points P(-1, 5), Q(5, -3), and R(6, 4) of Example 6.

Substitution of the *xy*-coordinates of each of the three points P, Q, and R into (10) gives the linear system of three equations

$$A - 5B + 25C = 1$$

$$25A - 15B + 9C = 1$$

$$36A + 24B + 16C = 1$$

(11)

in the three unknowns A, B, and C. Reduction of the corresponding augmented coefficient matrix to reduced row-echelon form (Fig. 3.7.11) yields the values

$$A = \frac{277}{14212}, \quad B = -\frac{172}{14212}, \quad \text{and} \quad C = \frac{523}{14212}$$

If we substitute these coefficient values in (10) and multiply the result by the common denominator 14212, we get the desired equation

$$277x^2 - 172xy + 523y^2 = 14212 \tag{12}$$

of our central conic. The computer plot in Fig. 3.7.12 verifies that this rotated ellipse does indeed pass through all three points P, Q, and R.



FIGURE 3.7.12. Central ellipse passing through the points *P*, *Q*, and *R* of Example 7.

FIGURE 3.7.11. TI-89 calculator solution of the reduced row-echelon form of the augmented coefficient matrix in (11).

3.7 **Problems**

In each of Problems 1–10, n + 1 data points are given. Find the nth degree polynomial y = f(x) that fits these points.

- **1.** (1, 1) and (3, 7)
- **2.** (-1, 11) and (2, -10)
- **3.** (0, 3), (1, 1), and (2, -5)
- **4.** (-1, 1), (1, 5), and (2, 16)
- **5.** (1, 3), (2, 3), and (3, 5)
- **6.** (-1, -1), (3, -13), and (5, 5)
- 7. (-1, 1), (0, 0), (1, 1), and (2, -4)
- **8.** (-1, 3), (0, 5), (1, 7), and (2, 3)
- **9.** (-2, -2), (-1, 2), (1, 10), and (2, 26)
- **10.** (-1, 27), (1, 13), (2, 3), and (3, -25)

Three points are given in each of Problems 11–14. Find the equation of the circle determined by these points, as well as its center and radius.

- **11.** (-1, -1), (6, 6), and (7, 5)
- **12.** (3, -4), (5, 10), and (-9, 12)
- **13.** (1,0), (0, -5), and (-5, -4)
- **14.** (0,0), (10,0), and (-7,7)

In Problems 15–18, find an equation of the central ellipse that passes through the three given points.

- **15.** (0, 5), (5, 0), and (5, 5)
- **16.** (0, 5), (5, 0), and (10, 10)
- **17.** (0, 1), (1, 0), and (10, 10)
- **18.** (0, 4), (3, 0), and (5, 5)
- **19.** Find a curve of the form y = A + (B/x) that passes through the points (1, 5) and (2, 4).
- **20.** Find a curve of the form $y = Ax + (B/x) + (C/x^2)$ that passes through the points (1, 2), (2, 20), and (4, 41).

A sphere in space with center (h, k, l) and radius r has equation

$$(x-h)^{2} + (y-k)^{2} + (z-l)^{2} = r^{2}.$$

Four given points in space suffice to determine the values of h, k, l, and r. In Problems 21 and 22, find the center and radius of the sphere that passes through the four given points P, Q, R, and S. Hint: Substitute each given triple of coordinates into the sphere equation above to obtain four equations that h, k, l, and r must satisfy. To solve these equations, first subtract the first one from each of the other three. How many unknowns are left in the three equations that result?

- **21.** *P*(4, 6, 15), *Q*(13, 5, 7), *R*(5, 14, 6), *S*(5, 5, -9)
- **22.** *P*(11, 17, 17), *Q*(29, 1, 15), *R*(13, -1, 33), *S*(-19, -13, 1)

Population Modeling

Problems 23–34 are intended as calculator or computer problems and are based on the U.S. census data in the table of Fig. 3.7.13, listed by national region in millions for the census years 1950–1990. See www.census.gov/population/ censusdata/table-16.pdf for further details.

In Problems 23–26, fit a quadratic function to the 1970, 1980, and 1990 population values for the indicated region.

- 23. The Northeast 24. The Midwest
- **25.** The South **26.** The West
- **27–30.** The same as Problems 23–26, except fit a cubic polynomial to the 1960, 1970, 1980, and 1990 population data for the indicated region.
- **31–34.** The same as Problems 23–26, except fit a quartic polynomial to the 1950, 1960, 1970, 1980, and 1990 population data for the indicated region.

Problems 35 through 40 illustrate the use of determinants in fitting polynomial curves to data points.

35. Explain why the determinant equation

$$\begin{vmatrix} y & x^2 & x & 1 \\ y_1 & x_1^2 & x_1 & 1 \\ y_2 & x_2^2 & x_2 & 1 \\ y_3 & x_3^2 & x_3 & 1 \end{vmatrix} = 0$$

fits a quadratic polynomial of the form $y = Ax^2 + Bx + C$ to the three given points (x_1, y_1) , (x_2, y_2) , and (x_3, y_3) .

- **36.** Expand the determinant in Problem 35 to find a parabola that interpolates the points (1, 3), (2, 3), and (3, 7).
- 37. Explain why the determinant equation

 $\begin{vmatrix} x^2 + y^2 & x & y & 1 \\ x_1^2 + y_1^2 & x_1 & y_1 & 1 \\ x_2^2 + y_2^2 & x_2 & y_2 & 1 \\ x_3^2 + y_3^2 & x_3 & y_3 & 1 \end{vmatrix} = 0$

fits a circle of the form $x^2 + y^2 + Ax + By + C = 0$ to the three given points $(x_1, y_1), (x_2, y_2)$, and (x_3, y_3) .

- **38.** Expand the determinant in Problem 37 to find the equation of a circle passing through the three points (3, -4), (5, 10), and (-9, 12). Then find its center and radius.
- 39. Explain why the determinant equation

 $\begin{vmatrix} x^2 & xy & y^2 & 1 \\ x_1^2 & x_1y_1 & y_1^2 & 1 \\ x_2^2 & x_2y_2 & y_2^2 & 1 \\ x_3^2 & x_3y_3 & y_3^2 & 1 \end{vmatrix} = 0$

fits a central conic equation of the form $Ax^2 + Bxy + Cy^2 = 1$ to the three given points (x_1, y_1) , (x_2, y_2) , and (x_3, y_3) .

40. Expand the determinant in Problem 39 to find the equation of the ellipse passing through the three points (0, 4), (3, 0), and (5, 5).

	1950	1960	1970	1980	1990
Northeast	30 / 78	14 678	/9.061	/0 137	50 809
Midwest	44.461	51.619	56.590	58.867	59.669
South	47.197	54.973	62.813	75.367	85.446
West	20.190	28.053	34.838	43.171	52.786
U.S.	151.326	179.323	203.302	226.542	248.710

FIGURE 3.7.13. Regional population data (in millions) for Problems 23–34.

4 Vector Spaces

4.1 The Vector Space R³



FIGURE 4.1.1. The arrow \overrightarrow{OP} representing the vector $\mathbf{v} = (a, b, c)$.

Here we take a fresh look—from the viewpoint of linear algebra—at the familiar 3-dimensional space of the physical world around us (and of multivariable calculus). This review, combining old and new ideas, will provide an introduction to basic concepts that are explored further in subsequent sections of the chapter.

We define **three-dimensional coordinate space** \mathbb{R}^3 to be the set of all ordered triples (a, b, c) of real numbers. In ordinary language, the elements of \mathbb{R}^3 are called **points**. The numbers *a*, *b*, and *c* are called the **coordinates** of the point P(a, b, c) and can be regarded as specifying the location of *P* in a fixed *xyz*-coordinate system.

The location of the point *P* can also be specified by means of the *arrow* or directed line segment (Fig. 4.1.1) that points from the origin (its *initial point*) to *P* (its *terminal point*). Arrows are often used in physics to represent *vector quantities*, such as force and velocity, that possess both magnitude and direction. For instance, the *velocity vector* \mathbf{v} of a point moving in space may be represented by an arrow that points in the direction of its motion, with the length of the arrow being equal to the speed of the moving point. If we locate this arrow with its initial point at the origin *O*, then its direction and length are determined by its terminal point (*a*, *b*, *c*). But the arrow is merely a pictorial object; the mathematical object associated with the vector \mathbf{v} is simply the point (*a*, *b*, *c*). For this reason, it is customary to use the words point and vector interchangeably for elements of 3-space \mathbf{R}^3 and thus to refer either to the *point* (*a*, *b*, *c*).

DEFINITION Vector

A vector v in 3-space \mathbb{R}^3 is simply an ordered triple (a, b, c) of real numbers. We write $\mathbf{v} = (a, b, c)$ and call the numbers a, b, and c the components of the vector v.

Thus "a vector is a point is a vector," but vector terminology often aids us in visualizing geometric relationships between different points. The point P(a, b, c)

determines the vector $\mathbf{v} = (a, b, c)$ in \mathbb{R}^3 , and \mathbf{v} is represented geometrically (as in Fig. 4.1.1) by the *position vector* \overrightarrow{OP} from the origin O(0, 0, 0) to *P*—or equally well by any parallel translate of this arrow. What is important about an arrow usually is not where it is, but how long it is and which way it points.

As in Section 3.4, we adopt the convention that the vector **v** with components v_1 , v_2 , and v_3 may be written interchangeably as either

$$\mathbf{v} = (v_1, v_2, v_3) \quad \text{or} \quad \mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}$$

with the column matrix regarded as just another symbol representing one and the same ordered triple of real numbers. Then the following definitions of addition of vectors and of multiplication of vectors by scalars are consistent with the matrix operations defined in Section 3.4.

DEFINITION Addition of Vectors

The sum $\mathbf{u} + \mathbf{v}$ of the two vectors $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{v} = (v_1, v_2, v_3)$ is the vector

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, u_3 + v_3) \tag{1}$$

that is obtained upon addition of respective components of \mathbf{u} and \mathbf{v} .

Thus we add vectors by adding corresponding components—that is, by *componentwise addition* (just as we add matrices). For instance, the sum of the vectors $\mathbf{u} = (4, 3, -5)$ and $\mathbf{v} = (-5, 2, 15)$ is the vector

$$\mathbf{u} + \mathbf{v} = (4, 3, -5) + (-5, 2, 15) = (4 - 5, 3 + 2, -5 + 15) = (-1, 5, 10).$$

The geometric representation of vectors as arrows often converts an algebraic relation into a picture that is readily understood and remembered. Addition of vectors is defined algebraically by Eq. (1). The geometric interpretation of vector addition is the *triangle law of addition* illustrated in Fig. 4.1.2 (for the case of 2-dimensional vectors in the plane), where the labeled lengths indicate why this interpretation is valid. An equivalent interpretation is the *parallelogram law of addition*, illustrated in Fig. 4.1.3.





FIGURE 4.1.2. The triangle law of vector addition.

FIGURE 4.1.3. The parallelogram law of vector addition.

Multiplication of a vector by a scalar (a real number) is also defined in a componentwise manner.

DEFINITION Multiplication of a Vector by a Scalar

If $\mathbf{v} = (v_1, v_2, v_3)$ is a vector and *c* is a real number, then the scalar multiple *c***v** is the vector

$$c\mathbf{v} = (cv_1, cv_2, cv_3) \tag{2}$$

that is obtained upon multiplying each component of \mathbf{v} by c.

The **length** $|\mathbf{v}|$ of the vector $\mathbf{v} = (a, b, c)$ is defined to be the distance of the point P(a, b, c) from the origin,

$$\mathbf{v}| = \sqrt{a^2 + b^2 + c^2}.$$
 (3)

The length of $c\mathbf{v}$ is |c| times the length of \mathbf{v} . For instance, if $\mathbf{v} = (4, 3, -12)$, then

$$(-7)\mathbf{v} = (-7 \cdot 4, -7 \cdot 3, -7 \cdot (-12)) = (-28, -21, 84),$$

 $|\mathbf{v}| = \sqrt{(4)^2 + (3)^2 + (-12)^2} = \sqrt{169} = 13,$ and
 $|-7\mathbf{v}| = |-7| \cdot |\mathbf{v}| = 7 \cdot 13 = 91.$

The geometric interpretation of scalar multiplication is that $c\mathbf{v}$ is a vector of length $|c| \cdot |\mathbf{v}|$, with the same direction as \mathbf{v} if c > 0 but the opposite direction if c < 0 (Fig. 4.1.4).

With vector addition and multiplication by scalars defined as in (1) and (2), \mathbf{R}^3 is a **vector space**. That is, these operations satisfy the conditions in (a)–(h) of the following theorem.

THEOREM 1 R³ as a Vector Space

If u ,	v , and w are vectors in \mathbf{R}^3 , and <i>r</i>	and <i>s</i> are real numbers, then
(a)	$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$	(commutativity)
(b)	$\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$	(associativity)
(c)	$\mathbf{u} + 0 = 0 + \mathbf{u} = \mathbf{u}$	(zero element)
(d)	u + (-u) = (-u) + u = 0	(additive inverse)
(e)	$r(\mathbf{u} + \mathbf{v}) = r\mathbf{u} + r\mathbf{v}$	(distributivity)
(f)	$(r+s)\mathbf{u} = r\mathbf{u} + s\mathbf{u}$	
(g)	$r(s\mathbf{u}) = (rs)\mathbf{u}$	
(h)	$1(\mathbf{u}) = \mathbf{u}$	(multiplicative identity).

Of course, $\mathbf{0} = (0, 0, 0)$ denotes the **zero vector** in (c) and (d), and $-\mathbf{u} = (-1)\mathbf{u}$ in (d). Each of properties (a)–(h) in Theorem 1 is readily verified in a componentwise manner. For instance, if $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{v} = (v_1, v_2, v_3)$, then use of the (ordinary) distributive law of real numbers gives

$$r(\mathbf{u} + \mathbf{v}) = r(u_1 + v_1, u_2 + v_2, u_3 + v_3)$$

= $(r(u_1 + v_1), r(u_2 + v_2), r(u_3 + v_3))$
= $(ru_1 + rv_1, ru_2 + rv_2, ru_3 + rv_3)$
= $(ru_1, ru_2, ru_3) + (rv_1, rv_2, rv_3)$
= $r(u_1, u_2, u_3) + r(v_1, v_2, v_3) = r\mathbf{u} + r\mathbf{v}$

so we have verified property (e).



FIGURE 4.1.4. The vector *c***u** may have the same direction as **u** or the opposite direction.

The Vector Space R²

The familiar coordinate plane \mathbf{R}^2 is the set of all ordered pairs (a, b) of real numbers. We may regard \mathbf{R}^2 as the *xy*-plane in \mathbf{R}^3 by identifying (a, b) with the vector (a, b, 0) in \mathbf{R}^3 . Then \mathbf{R}^2 is simply the set of all 3-dimensional vectors that have third component 0.

Clearly, the sum of any two vectors in \mathbf{R}^2 is again a vector in \mathbf{R}^2 , as is any scalar multiple of a vector in \mathbf{R}^2 . Indeed, vectors in \mathbf{R}^2 satisfy all the properties of a vector space enumerated in Theorem 1. Consequently, the plane \mathbf{R}^2 is a vector space in its own right.

The two vectors \mathbf{u} and \mathbf{v} are collinear—they lie on the same line through the origin and hence point either in the same direction (Fig. 4.1.5) or in opposite directions—if and only if one is a scalar multiple of the other; that is, either

$$\mathbf{u} = c\mathbf{v} \quad \text{or} \quad \mathbf{v} = c\mathbf{u},\tag{4}$$

for some scalar c. The scalar c merely adjusts the length and direction of one vector to fit the other. If **u** and **v** are nonzero vectors, then $c = \pm |\mathbf{u}|/|\mathbf{v}|$ in the first relation, with c being positive if the two vectors point in the same direction, negative otherwise.

If one of the relations in (4) holds for some scalar *c*, then we say that the two vectors are **linearly dependent**. Note that if $\mathbf{u} = \mathbf{0}$ while $\mathbf{v} \neq \mathbf{0}$, then $\mathbf{u} = 0\mathbf{v}$ but \mathbf{v} is not a scalar multiple of \mathbf{u} . (Why?) Thus, if precisely one of the two vectors \mathbf{u} and \mathbf{v} is the zero vector, then \mathbf{u} and \mathbf{v} are linearly dependent, but only one of the two relations in (4) holds.

If **u** and **v** are linearly dependent vectors with $\mathbf{u} = c\mathbf{v}$ (for instance), then $1 \cdot \mathbf{u} + (-c) \cdot \mathbf{v} = \mathbf{0}$. Thus there exist scalars *a* and *b* not both zero such that

$$a\mathbf{u} + b\mathbf{v} = \mathbf{0}.\tag{5}$$

Conversely, suppose that Eq. (5) holds with a and b not both zero. If $a \neq 0$ (for instance) then we can solve for

$$\mathbf{u} = -\frac{b}{a}\mathbf{v} = c\mathbf{v}$$

with c = -b/a, so it follows that **u** and **v** are linearly dependent. Therefore, we have proved the following theorem.

THEOREM 2 Two Linearly Dependent Vectors

The two vectors \mathbf{u} and \mathbf{v} are linearly dependent if and only if there exist scalars *a* and *b* not both zero such that

$$a\mathbf{u} + b\mathbf{v} = \mathbf{0}.\tag{5}$$

The most interesting pairs of vectors are those that are not linearly dependent. The two vectors \mathbf{u} and \mathbf{v} are said to be **linearly independent** provided that they are *not* linearly dependent. Thus \mathbf{u} and \mathbf{v} are linearly independent if and only if neither is a scalar multiple of the other. By Theorem 2 this is equivalent to the following



 (u_1, u_2)

 (v_1, v_2)

statement:

The two vectors \mathbf{u} and \mathbf{v} are linearly independent if and only if the relation

$$a\mathbf{u} + b\mathbf{v} = \mathbf{0} \tag{5}$$

implies that a = b = 0.

Thus the vectors \mathbf{u} and \mathbf{v} are linearly independent provided that no *nontrivial* linear combination of them is equal to the zero vector.

Example 1 If $\mathbf{u} = (3, -2)$, $\mathbf{v} = (-6, 4)$, and $\mathbf{w} = (5, -7)$, then \mathbf{u} and \mathbf{v} are linearly dependent, because $\mathbf{v} = -2\mathbf{u}$. On the other hand, \mathbf{u} and \mathbf{w} are linearly independent. Here is an argument to establish this fact: Suppose that there were scalars *a* and *b* such that

$$a\mathbf{u} + b\mathbf{w} = \mathbf{0}$$

Then

$$a(3,-2) + b(5,-7) = \mathbf{0},$$

and thus we get the simultaneous equations

$$3a + 5b = 0$$
$$-2a - 7b = 0$$

It is now easy to show that a = b = 0 is the (unique) solution of this system. This shows that whenever

$$a\mathbf{u} + b\mathbf{w} = \mathbf{0},$$

it follows that a = b = 0. Therefore, **u** and **w** are linearly independent.

Alternatively, we could prove that **u** and **w** are linearly independent by showing that neither is a scalar multiple of the other (because $\frac{5}{3} \neq \frac{7}{2}$).

The most important property of linearly independent pairs of plane vectors is this: If **u** and **v** are linearly independent vectors in the plane, then any third vector **w** in \mathbf{R}^2 can be expressed as a *linear combination* of **u** and **v**. That is, there exist scalars *a* and *b* such that $\mathbf{w} = a\mathbf{u} + b\mathbf{v}$ (Fig. 4.1.6). This is a statement that the two linearly independent vectors **u** and **v** suffice (in an obvious sense) to "generate" the whole plane \mathbf{R}^2 . This general fact—which Section 4.3 discusses in a broader context—is illustrated computationally by the following example.

Example 2 Express the vector $\mathbf{w} = (11, 4)$ as a linear combination of the vectors $\mathbf{u} = (3, -2)$ and $\mathbf{v} = (-2, 7)$.

Solution We want to find numbers a and b such that $a\mathbf{u} + b\mathbf{v} = \mathbf{w}$; that is,

$$a\begin{bmatrix}3\\-2\end{bmatrix}+b\begin{bmatrix}-2\\7\end{bmatrix}=\begin{bmatrix}11\\4\end{bmatrix}.$$

This vector equation is equivalent to the 2×2 linear system

$$\begin{bmatrix} 3 & -2 \\ -2 & 7 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 11 \\ 4 \end{bmatrix},$$

which (using Gaussian elimination or Cramer's rule) we readily solve for a = 5, b = 2. Thus $\mathbf{w} = 5\mathbf{u} + 2\mathbf{v}$.



FIGURE 4.1.6. The vector **w** as a linear combination of the two linearly independent vectors **u** and **v**.

Linear Independence in R³

We have said that the two vectors are linearly dependent provided that they lie on the same line through the origin. For *three* vectors $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$, and $\mathbf{w} = (w_1, w_2, w_3)$ in space, the analogous condition is that the three points (u_1, u_2, u_3) , (v_1, v_2, v_3) , and (w_1, w_2, w_3) lie in the same *plane* through the origin in \mathbf{R}^3 . Given \mathbf{u} , \mathbf{v} , and \mathbf{w} , how can we determine whether the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are *coplanar*? The key to the answer is the following observation: If r and s are scalars, then the parallelogram law of addition implies that the vectors \mathbf{u} , \mathbf{v} , and $r\mathbf{u} + s\mathbf{v}$ are coplanar; specifically, they lie in the plane through the origin that is determined by the parallelogram with vertices $\mathbf{0}$, $r\mathbf{u}$, $s\mathbf{v}$, and $r\mathbf{u} + s\mathbf{v}$. Thus any linear combination of \mathbf{u} and \mathbf{v} is coplanar with \mathbf{u} and \mathbf{v} . This is the motivation for our next definition.

DEFINITION Linearly Dependent Vectors in R³

The three vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in \mathbf{R}^3 are said to be **linearly dependent** provided that one of them is a linear combination of the other two—that is, either

 $\mathbf{w} = r\mathbf{u} + s\mathbf{v} \qquad \text{or} \\ \mathbf{u} = r\mathbf{v} + s\mathbf{w} \qquad \text{or} \qquad (6) \\ \mathbf{v} = r\mathbf{u} + s\mathbf{w}$

for appropriate scalars r and s.

Note that each of the three equations in (6) implies that there exist three scalars *a*, *b*, and *c* not all zero such that

$$a\mathbf{u} + b\mathbf{v} + c\mathbf{w} = \mathbf{0}.\tag{7}$$

For if $\mathbf{w} = r\mathbf{u} + s\mathbf{v}$ (for instance), then

$$r\mathbf{u} + s\mathbf{v} + (-1)\mathbf{w} = \mathbf{0}$$

so we can take a = r, b = s, and $c = -1 \neq 0$. Conversely, suppose that (7) holds with a, b, and c not all zero. If $c \neq 0$ (for instance), then we can solve for

$$\mathbf{w} = -\frac{a}{c}\mathbf{u} - \frac{b}{c}\mathbf{v} = r\mathbf{u} + s\mathbf{v}$$

with r = -a/c and s = -b/c, so it follows that the three vectors **u**, **v**, and **w** are linearly dependent. Therefore, we have proved the following theorem.

THEOREM 3 Three Linearly Dependent Vectors

The three vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in \mathbf{R}^3 are linearly dependent if and only if there exist scalars *a*, *b*, and *c* not all zero such that

$$a\mathbf{u} + b\mathbf{v} + c\mathbf{w} = \mathbf{0}.\tag{7}$$

The three vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are called **linearly independent** provided that they are *not* linearly dependent. Thus \mathbf{u} , \mathbf{v} , and \mathbf{w} are linearly independent if and

only if neither of them is a linear combination of the other two. As a consequence of Theorem 3, this is equivalent to the following statement:

The vectors **u**, **v**, and **w** are linearly independent if and only if the relation

$$a\mathbf{u} + b\mathbf{v} + c\mathbf{w} = \mathbf{0} \tag{7}$$

implies that a = b = c = 0.

Thus the three vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are linearly independent provided that no *non-trivial* linear combination of them is equal to the zero vector.

Given two vectors, we can see at a glance whether either is a scalar multiple of the other. By contrast, it is not evident at a glance whether or not three given vectors in \mathbf{R}^3 are linearly independent. The following theorem provides one way to resolve this question.

THEOREM 4 Three Linearly Independent Vectors

The vectors $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$, and $\mathbf{w} = (w_1, w_2, w_3)$ are linearly independent if and only if

$$\begin{vmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ u_3 & v_3 & w_3 \end{vmatrix} \neq 0.$$
(8)

Proof: We want to show that **u**, **v**, and **w** are linearly independent if and only if the matrix

$$\mathbf{A} = \begin{bmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ u_3 & v_3 & w_3 \end{bmatrix}$$

—with column vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} —has nonzero determinant. By Theorem 3, the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are linearly independent if and only if the equation

$$a\begin{bmatrix} u_1\\u_2\\u_3\end{bmatrix} + b\begin{bmatrix} v_1\\v_2\\v_3\end{bmatrix} + c\begin{bmatrix} w_1\\w_2\\w_3\end{bmatrix} = \mathbf{0}$$

implies that a = b = c = 0—that is, if and only if the system

$$\begin{bmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ u_3 & v_3 & w_3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$
(9)

with unknowns *a*, *b*, and *c* has only the trivial solution. But Theorem 7 in Section 3.5 implies that this is so if and only if the coefficient matrix **A** is invertible, and Theorem 2 in Section 3.6 implies that **A** is invertible if and only if $|\mathbf{A}| \neq 0$. Therefore, **u**, **v**, and **w** are linearly independent if and only if $|\mathbf{A}| \neq 0$.

Hence, in order to determine whether or not three given vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are linearly independent, we can calculate the determinant in (8). In practice, however, it is usually more efficient to set up and solve the linear system in (9). If we obtain only the trivial solution a = b = c = 0, then the three given vectors are linearly

independent. But if we find a nontrivial solution, we then can express one of the vectors as a linear combination of the other two and thus see *how* the three vectors are linearly dependent.

Example 3

To determine whether the three vectors $\mathbf{u} = (1, 2, -3)$, $\mathbf{v} = (3, 1, -2)$, and $\mathbf{w} = (5, -5, 6)$ are linearly independent or dependent, we need to solve the system

$$a\mathbf{u} + b\mathbf{v} + c\mathbf{w} = \begin{bmatrix} 1 & 3 & 5\\ 2 & 1 & -5\\ -3 & -2 & 6 \end{bmatrix} \begin{bmatrix} a\\ b\\ c \end{bmatrix} = \begin{bmatrix} 0\\ 0\\ 0 \end{bmatrix}.$$

By Gaussian elimination, we readily reduce this system to the echelon form

$$\begin{bmatrix} 1 & 3 & 5 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Therefore, we can choose c = 1, and it follows that b = -3c = -3 and a = -3b - 5c = 4. Therefore,

$$4\mathbf{u} + (-3)\mathbf{v} + \mathbf{w} = \mathbf{0}$$

and hence **u**, **v**, and **w** are linearly dependent, with $\mathbf{w} = -4\mathbf{u} + 3\mathbf{v}$.

Basis Vectors in R³

Perhaps the most familiar triple of linearly independent vectors in \mathbf{R}^3 consists of the **basic unit vectors**

$$\mathbf{i} = (1, 0, 0), \quad \mathbf{j} = (0, 1, 0), \text{ and } \mathbf{k} = (0, 0, 1).$$
 (10)

When represented by arrows with the initial points at the origin, these three vectors point in the positive directions along the three coordinate axes (Fig. 4.1.7). The expression

$$\mathbf{v} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k} = (a, b, c)$$

shows both that

- the three vectors **i**, **j**, and **k** are linearly independent (because **v** = **0** immediately implies *a* = *b* = *c* = 0), and that
- any vector in \mathbf{R}^3 can be expressed as a linear combination of \mathbf{i} , \mathbf{j} , and \mathbf{k} .

A **basis** for \mathbb{R}^3 is a triple **u**, **v**, **w** of vectors such that every vector **t** in \mathbb{R}^3 can be expressed as a linear combination

$$\mathbf{t} = a\mathbf{u} + b\mathbf{v} + c\mathbf{w} \tag{11}$$

of them. That is, given any vector \mathbf{t} in \mathbf{R}^3 , there exist scalars a, b, c such that Eq. (11) holds. Thus the unit vectors \mathbf{i} , \mathbf{j} , and \mathbf{k} constitute a basis for \mathbf{R}^3 . The following theorem says that any three *linearly independent* vectors constitute a basis for \mathbf{R}^3 .

THEOREM 5 Basis for R³

If the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in \mathbf{R}^3 are linearly independent, then they constitute a basis for \mathbf{R}^3 .



FIGURE 4.1.7. The basic unit vectors **i**, **j**, and **k**.

Proof: Let $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$, and $\mathbf{w} = (w_1, w_2, w_3)$. Given any fourth vector $\mathbf{t} = (t_1, t_2, t_3)$, we must show that there exist scalars a, b, and c such that

$$\mathbf{t} = a\mathbf{u} + b\mathbf{v} + c\mathbf{w} \tag{11}$$

—that is, such that the system

$$\begin{bmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ u_3 & v_3 & w_3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} t_1 \\ t_2 \\ t_3 \end{bmatrix}$$
(12)

has a solution (a, b, c). But by Theorem 4 the fact that **u**, **v**, and **w** are linearly independent implies that the coefficient matrix \mathbf{A} in (12) has nonzero determinant and is therefore invertible. Hence, there is a (unique) solution, which may be obtained by multiplying both sides in (12) by A^{-1} .

In order to express the vector $\mathbf{t} = (4, 20, 23)$ as a combination of the linearly independent vectors $\mathbf{u} = (1, 3, 2)$, $\mathbf{v} = (2, 8, 7)$, and $\mathbf{w} = (1, 7, 9)$, we need to solve the system

1	2	1	$\left[\begin{array}{c} a \end{array} \right]$		[4]
3	8	7	b	=	20
2	7	9	L c		_ 23 _

that we obtain by substitution in Eq. (12). The echelon form found by Gaussian elimination is

		t =	= 5 u -	- 2 v -	⊦3 w		
so $c = 3, b = 4 - 2c = -2$, and	a =	= 4 -	- 2b -	-c =	5. T	hus	
	0	0	1	_ c _		3	
	0	1	2	b	=	4	
	1	2				4	

Subspaces of R³

Up until this point, we have used the words *line* and *plane* only in an informal or intuitive way. It is now time for us to say precisely what is meant by a line or plane through the origin in \mathbb{R}^3 . Each is an example of a subspace of \mathbb{R}^3 .

The nonempty subset V of \mathbf{R}^3 is called a **subspace** of \mathbf{R}^3 provided that V itself is a vector space under the operations of vector addition and multiplication of vectors by scalars. Suppose that the nonempty subset V of \mathbb{R}^3 is closed under these operations—that is, that the sum of any two vectors in V is also in V and that every scalar multiple of a vector in V is also in V. Then the vectors in V automatically satisfy properties (a) through (h) of Theorem 1, because these properties are "inherited" from \mathbf{R}^3 ; they hold for all vectors in \mathbf{R}^3 , including those in V. Consequently, we see that a nonempty subset V of \mathbb{R}^3 is a subspace of \mathbb{R}^3 if and only if it satisfies the following two conditions:

- (i) If **u** and **v** are vectors in V, then $\mathbf{u} + \mathbf{v}$ is also in V (closure under addition).
- (ii) If **u** is a vector in V and c is a scalar, then c**u** is in V (closure under multiplication by scalars).

It is immediate that $V = \mathbf{R}^3$ is a subspace: \mathbf{R}^3 is a subspace of itself. At the opposite extreme, the subset $V = \{0\}$, containing only the zero vector, is also a subspace of \mathbb{R}^3 , because $\mathbf{0} + \mathbf{0} = \mathbf{0}$ and $c\mathbf{0} = \mathbf{0}$ for every scalar c. Thus $V = \{\mathbf{0}\}$ satisfies conditions (i) and (ii). The subspaces $\{0\}$ and \mathbb{R}^3 are sometimes called the trivial subspaces of \mathbf{R}^3 (because the verification that they *are* subspaces is quite trivial). All subspaces other than $\{0\}$ and \mathbb{R}^3 itself are called **proper subspaces** of **R**³.



Now we want to show that the proper subspaces of \mathbb{R}^3 are what we customarily call lines and planes through the origin. Let V be a subspace of \mathbb{R}^3 that is neither $\{0\}$ nor \mathbb{R}^3 itself. There are two cases to consider, depending on whether or not V contains two linearly independent vectors.

CASE 1: Suppose that V does *not* contain two linearly independent vectors. If **u** is a fixed nonzero vector in V, then, by condition (ii) above, every scalar multiple c**u** is also in V. Conversely, if **v** is any other vector in V, then **u** and **v** are linearly dependent, so it follows that $\mathbf{v} = c\mathbf{u}$ for some scalar c. Thus the subspace V is the set of all scalar multiples of the fixed nonzero vector **u** and is therefore what we call a **line** through the origin in **R**³. (See Fig. 4.1.8.)

CASE 2: Suppose that V contains two linearly independent vectors **u** and **v**. It then follows from conditions (i) and (ii) that V contains every linear combination $a\mathbf{u} + b\mathbf{v}$ of **u** and **v**. (See Problem 38.) Conversely, let **w** be any other vector in V. If **u**, **v**, **w** were linearly independent, then, by Theorem 4, V would be all of \mathbf{R}^3 . Therefore, **u**, **v**, **w** are linearly dependent, so it follows that there exist scalars a and b such that $\mathbf{w} = a\mathbf{u} + b\mathbf{v}$. (See Problem 40.) Thus the subspace V is the set of all linear combinations $a\mathbf{u} + b\mathbf{v}$ of the two linearly independent vectors **u** and **v** and is therefore what we call a **plane** through the origin in \mathbf{R}^3 . (See Fig. 4.1.9.)

Subspaces of the coordinate plane \mathbf{R}^2 are defined similarly—they are the nonempty subsets of \mathbf{R}^2 that are closed under addition and multiplication by scalars. In Problem 39 we ask you to show that every proper subspace of \mathbf{R}^2 is a line through the origin.

xample 5	Let V be the set of all vectors (x, y) in \mathbb{R}^2 such that $y = x$. Given u and v in V, we may write $\mathbf{u} = (u, u)$ and $\mathbf{v} = (v, v)$. Then $\mathbf{u} + \mathbf{v} = (u + v, u + v)$ and $c\mathbf{u} = (cu, cu)$ are in V. It
	follows that V is a subspace of \mathbb{R}^2 .
xample 6	Let V be the set of all vectors (x, y) in \mathbb{R}^2 such that $x + y = 1$. Thus V is the straight line
	that passes through the unit points on the x- and y-axes. Then $\mathbf{u} = (1,0)$ and $\mathbf{v} = (0,1)$ are
	in V, but the vector $\mathbf{u} + \mathbf{v} = (1, 1)$ is not. It follows that V is not a subspace of \mathbf{R}^2 .

Example 6 illustrates the fact that lines that do not pass through the origin are not subspaces of \mathbf{R}^2 . Because every subspace must contain the zero vector (per Problem 37), only lines and planes that pass through the origin are subspaces of \mathbf{R}^3 .

4.1 Problems

In Problems 1–4, find $|\mathbf{a} - \mathbf{b}|$, $2\mathbf{a} + \mathbf{b}$, and $3\mathbf{a} - 4\mathbf{b}$.

- **1.** $\mathbf{a} = (2, 5, -4), \mathbf{b} = (1, -2, -3)$
- **2.** $\mathbf{a} = (-1, 0, 2), \mathbf{b} = (3, 4, -5)$
- **3.** $\mathbf{a} = 2\mathbf{i} 3\mathbf{j} + 5\mathbf{k}, \mathbf{b} = 5\mathbf{i} + 3\mathbf{j} 7\mathbf{k}$
- 4. a = 2i j, b = j 3k

In Problems 5–8, determine whether the given vectors \mathbf{u} and \mathbf{v} are linearly dependent or linearly independent.

5. $\mathbf{u} = (0, 2), \mathbf{v} = (0, 3)$ 6. $\mathbf{u} = (0, 2), \mathbf{v} = (3, 0)$ 7. $\mathbf{u} = (2, 2), \mathbf{v} = (2, -2)$ 8. $\mathbf{u} = (2, -2), \mathbf{v} = (-2, 2)$

In Problems 9–14, express \mathbf{w} as a linear combination of \mathbf{u} and \mathbf{v} .

9. $\mathbf{u} = (1, -2), \mathbf{v} = (-1, 3), \mathbf{w} = (1, 0)$ 10. $\mathbf{u} = (3, 4), \mathbf{v} = (2, 3), \mathbf{w} = (0, -1)$ 11. $\mathbf{u} = (5, 7), \mathbf{v} = (2, 3), \mathbf{w} = (1, 1)$ 12. $\mathbf{u} = (4, 1), \mathbf{v} = (-2, -1), \mathbf{w} = (2, -2)$ 13. $\mathbf{u} = (7, 5), \mathbf{v} = (3, 4), \mathbf{w} = (5, -2)$ 14. $\mathbf{u} = (5, -2), \mathbf{v} = (-6, 4), \mathbf{w} = (5, 6)$

In Problems 15–18, apply Theorem 4 (that is, calculate a determinant) to determine whether the given vectors **u**, **v**, and **w** are linearly dependent or independent.

15. $\mathbf{u} = (3, -1, 2), \mathbf{v} = (5, 4, -6), \mathbf{w} = (8, 3, -4)$ **16.** $\mathbf{u} = (5, -2, 4), \mathbf{v} = (2, -3, 5), \mathbf{w} = (4, 5, -7)$ **17.** $\mathbf{u} = (1, -1, 2), \mathbf{v} = (3, 0, 1), \mathbf{w} = (1, -2, 2)$ **18.** $\mathbf{u} = (1, 1, 0), \mathbf{v} = (4, 3, 1), \mathbf{w} = (3, -2, -4)$

In Problems 19–24, use the method of Example 3 to determine whether the given vectors **u**, **v**, and **w** are linearly independent







or dependent. If they are linearly dependent, find scalars a, b, and c not all zero such that $a\mathbf{u} + b\mathbf{v} + c\mathbf{w} = \mathbf{0}$.

19. $\mathbf{u} = (2, 0, 1), \mathbf{v} = (-3, 1, -1), \mathbf{w} = (0, -2, -1)$ **20.** $\mathbf{u} = (5, 5, 4), \mathbf{v} = (2, 3, 1), \mathbf{w} = (4, 1, 5)$ **21.** $\mathbf{u} = (1, 1, -2), \mathbf{v} = (-2, -1, 6), \mathbf{w} = (3, 7, 2)$ **22.** $\mathbf{u} = (1, 1, 0), \mathbf{v} = (5, 1, 3), \mathbf{w} = (0, 1, 2)$ **23.** $\mathbf{u} = (2, 0, 3), \mathbf{v} = (5, 4, -2), \mathbf{w} = (2, -1, 1)$ **24.** $\mathbf{u} = (1, 4, 5), \mathbf{v} = (4, 2, 5), \mathbf{w} = (-3, 3, -1)$

In Problems 25–28, express the vector \mathbf{t} as a linear combination of the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} .

- **25.** $\mathbf{t} = (2, -7, 9), \mathbf{u} = (1, -2, 2), \mathbf{v} = (3, 0, 1), \mathbf{w} = (1, -1, 2)$ **26.** $\mathbf{t} = (5, 30, -21), \mathbf{u} = (5, 2, -2), \mathbf{v} = (1, 5, -3),$ $\mathbf{w} = (5, -3, 4)$
- **27.** $\mathbf{t} = (0, 0, 19), \mathbf{u} = (1, 4, 3), \mathbf{v} = (-1, -2, 2), \mathbf{w} = (4, 4, 1)$
- **28.** $\mathbf{t} = (7, 7, 7), \mathbf{u} = (2, 5, 3), \mathbf{v} = (4, 1, -1), \mathbf{w} = (1, 1, 5)$

In Problems 29–32, show that the given set V is closed under addition and under multiplication by scalars and is therefore a subspace of \mathbb{R}^3 .

- **29.** V is the set of all (x, y, z) such that x = 0.
- **30.** V is the set of all (x, y, z) such that x + y + z = 0.
- **31.** V is the set of all (x, y, z) such that 2x = 3y.
- **32.** V is the set of all (x, y, z) such that z = 2x + 3y.

In Problems 33–36, show that the given set V is not a subspace of \mathbb{R}^3 .

- **33.** V is the set of all (x, y, z) such that y = 1.
- **34.** V is the set of all (x, y, z) such that x + y + z = 3.
- **35.** V is the set of all (x, y, z) such that $z \ge 0$.
- **36.** V is the set of all (x, y, z) such that xyz = 1.
- 37. Show that every subspace V of R³ contains the zero vector0.
- 38. Suppose that V is a subspace of R³. Show that V is closed under the operation of taking linear combinations of pairs of vectors. That is, show that if u and v are in V and a and b are scalars, then au + bv is in V.
- **39.** Suppose that V is a proper subspace of **R**² and that **u** is a nonzero vector in V. Show that V is the set of all scalar multiples of **u** and therefore that V is a line through the origin.
- 40. Suppose that u, v, and w are vectors in R³ such that u and v are linearly independent but u, v, and w are linearly dependent. Show that there exist scalars a and b such that w = au + bv.
- **41.** Let V_1 and V_2 be subspaces of \mathbb{R}^3 . Their *intersection* $V = V_1 \cap V_2$ is the set of all vectors that lie both in V_1 and in V_2 . Show that V is a subspace of \mathbb{R}^3 .

4.2 The Vector Space R^{*n*} and Subspaces



FIGURE 4.2.1. Two points $P(x_1, x_2)$ and $Q(x_3, x_4)$ in **R**².

In Section 4.1 we defined 3-dimensional space \mathbf{R}^3 to be the set of all triples (x, y, z) of real numbers. This definition provides a *mathematical model* of the physical space in which we live, because geometric intuition and experience require that the location of every point be specified uniquely by *three* coordinates.

In science fiction, the fourth dimension often plays a rather exotic role. But there are common and ordinary situations where it is convenient to use four (or even more) coordinates rather than just two or three. For example, suppose we want to describe the motion of two points P and Q that are moving in the plane \mathbb{R}^2 under the action of some given physical law. (See Fig. 4.2.1.) In order to tell where P and Q are at a given instant, we need to give two coordinates for P and two coordinates for Q. So let us write $P(x_1, x_2)$ and $Q(x_3, x_4)$ to indicate these four coordinates. Then the two points P and Q determine a quadruple or 4-tuple (x_1, x_2, x_3, x_4) of real numbers, and any such 4-tuple determines a possible pair of locations of P and Q. In this way the set of all pairs of points P and Q in the plane corresponds to the set of all 4-tuples of real numbers. By analogy with our definition of \mathbb{R}^3 , we may define 4-dimensional space \mathbb{R}^4 to be the set of all such 4-tuples (x_1, x_2, x_3, x_4) . Then we can specify a pair of points P and Q in \mathbb{R}^2 by specifying a single point (x_1, x_2, x_3, x_4) in \mathbb{R}^4 , and this viewpoint may simplify our analysis of the motions of the original points P and Q. For instance, it may turn out that their coordinates satisfy some single equation such as

$$3x_1 - 4x_2 + 2x_3 - 5x_4 = 0$$

that is better understood in terms of a single point in \mathbb{R}^4 than in terms of the separate points *P* and *Q* in \mathbb{R}^2 . Finally, note that in this example the fourth dimension is quite tangible—it refers simply to the second coordinate x_4 of the point *Q*.

But the number 4 is no more special in this context than the numbers 2 and 3. To describe similarly the motion of three points in \mathbb{R}^3 , we would need 9 coordinates rather than 4. For instance, we might be studying the sun-earth-moon system in 3-dimensional space, with (x_1, x_2, x_3) denoting the coordinates of the sun, (x_4, x_5, x_6) the coordinates of the earth, and (x_7, x_8, x_9) the coordinates of the moon at a given instant. Then the single list $(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9)$ of 9 coordinates could be used to specify simultaneously the locations of the sun, earth, and moon. Application of Newton's second law of motion to each of these three bodies would then lead to a system of differential equations involving the dependent variables $x_i(t)$, $i = 1, 2, \ldots, 9$ as functions of time t. (We will discuss this very situation in Section 7.6.)

An efficient description of another situation might require fifty, a hundred, or even a thousand coordinates. For instance, an investor's stock portfolio might contain 50 different high-tech stocks. If x_i denotes the value of the *i*th stock on a particular day, then the state of the whole portfolio is described by the single list $(x_1, x_2, x_3, \ldots, x_{49}, x_{50})$ of 50 separate numbers. And the evolution of this portfolio with time could then be described by the motion of a single point in 50-dimensional space!

An *n*-tuple of real numbers is an (ordered) list $(x_1, x_2, x_3, ..., x_n)$ of *n* real numbers. Thus (1, 3, 4, 2) is a 4-tuple and (0, -3, 7, 5, 2, -1) is a 6-tuple. A 2-tuple is an ordered pair and a 3-tuple is an ordered triple of real numbers, so the following definition generalizes our earlier definitions of the plane \mathbb{R}^2 and 3-space \mathbb{R}^3 .

DEFINITION n-Space Rⁿ

The *n*-dimensional space \mathbb{R}^n is the set of all *n*-tuples $(x_1, x_2, x_3, ..., x_n)$ of real numbers.

The elements of *n*-space \mathbb{R}^n are called **points** or **vectors**, and we ordinarily use boldface letters to denote vectors. The *i* th entry of the vector $\mathbf{x} = (x_1, x_2, x_3, \dots, x_n)$ is called its *i* th **coordinate** or its *i* th **component**. For consistency with matrix operations, we agree that

$$\mathbf{x} = (x_1, x_2, x_3, \dots, x_n) = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix},$$

so that the $n \times 1$ matrix, or *column* vector, is simply another notation for the same ordered list of *n* real numbers.

If n > 3, we cannot visualize vectors in \mathbb{R}^n in the concrete way that we can "see" vectors in \mathbb{R}^2 and \mathbb{R}^3 . Nevertheless, we saw in Section 4.1 that the geometric properties of \mathbb{R}^2 and \mathbb{R}^3 stem ultimately from algebraic operations with vectors, and these algebraic operations can be defined for vectors in \mathbb{R}^n by analogy with their definitions in dimensions 2 and 3.

If $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ are vectors in \mathbf{R}^n , then their **sum** is the vector $\mathbf{u} + \mathbf{v}$ given by

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, \dots, u_n + v_n).$$
(1)

Thus addition of vectors in \mathbf{R}^n is defined in a componentwise manner, just as in \mathbf{R}^2 and \mathbf{R}^3 , and we can visualize $\mathbf{u} + \mathbf{v}$ as a diagonal vector in a parallelogram



FIGURE 4.2.2. The parallelogram law for addition of vectors, in \mathbf{R}^n just as in \mathbf{R}^2 or \mathbf{R}^3 .



FIGURE 4.2.3. Multiplication of the vector **u** by the scalar c, in \mathbf{R}^n just as in \mathbf{R}^2 or \mathbf{R}^3 .

determined by **u** and **v**, as in Fig. 4.2.2. If c is a scalar—a real number—then the scalar multiple c**u** is also defined in componentwise fashion:

$$c\mathbf{u} = (cu_1, cu_2, \dots, cu_n). \tag{2}$$

We can visualize $c\mathbf{u}$ as a vector collinear with \mathbf{u} , with only its length (and possibly its direction along the same line) altered by multiplication by the scalar c. See Fig. 4.2.3.

In Section 4.1 we saw that, for vectors in \mathbb{R}^2 and \mathbb{R}^3 , the operations in (1) and (2) satisfy a list of properties that constitute the definition of a *vector space*; the same is true for vectors in \mathbb{R}^n .

Definition of a Vector Space

Let V be a set of elements called *vectors*, in which the operations of addition of vectors and multiplication of vectors by scalars are defined. That is, given vectors **u** and **v** in V and a scalar c, the vectors $\mathbf{u} + \mathbf{v}$ and $c\mathbf{u}$ are also in V (so that V is *closed* under vector addition and multiplication by scalars). Then, with these operations, V is called a **vector space** provided that—given any vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and any scalars a and b—the following properties hold true:

(a) $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$		(commutativity)
(b) $u + (v + w) = (u + w)$	$(\mathbf{v} + \mathbf{v}) + \mathbf{w}$	(associativity)
(c) $u + 0 = 0 + u =$	u	(zero element)
(d) $u + (-u) = (-u)$	+ u = 0	(additive inverse)
(e) $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + \mathbf{v}$	av	(distributivity)
(f) $(a+b)\mathbf{u} = a\mathbf{u} + b$	bu	
(g) $a(b\mathbf{u}) = (ab)\mathbf{u}$		
(h) (1) u = u		

In property (c), it is meant that there exists a **zero vector 0** in V such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$. The zero vector in \mathbf{R}^n is

$$\mathbf{0} = (0, 0, \dots, 0).$$

Similarly, property (d) actually means that, given the vector **u** in V, there exists a vector $-\mathbf{u}$ in V such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$. In \mathbf{R}^n , we clearly have

$$-\mathbf{u} = (-u_1, -u_2, \ldots, -u_n).$$

The fact that $\mathbf{R} = \mathbf{R}^1$ satisfies properties (a)–(h), with the real numbers playing the dual roles of scalars *and* vectors, means that the real line may be regarded as a vector space. If n > 1, then each of properties (a)–(h) may be readily verified for \mathbf{R}^n by working with components and applying the corresponding properties of real numbers. For example, to verify the commutativity of vector addition in (a), we begin with Eq. (1) and write

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, \dots, u_n + v_n)$$

= $(v_1 + u_1, v_2 + u_2, \dots, v_n + u_n)$
= $\mathbf{v} + \mathbf{u}$

Thus *n*-space \mathbb{R}^n is a vector space with the operations defined in Eqs. (1) and (2). We will understand throughout that our scalars are the *real numbers*, though

in more advanced treatments of linear algebra other scalars (such as the complex numbers) are sometimes used.

Observe that, in the definition of a vector space, nothing is said about the elements of V (the "vectors") being *n*-tuples. They can be any objects for which addition and multiplication by scalars are defined and which satisfy properties (a)–(h). Although many of the vectors you will see in this text actually are *n*-tuples of real numbers, an example in which they are not may help to clarify the definition of a vector space.

Example 1 Let \mathcal{F} be the set of all real-valued functions defined on the real number line **R**. Then each vector in \mathcal{F} is a function **f** such that the real number $\mathbf{f}(x)$ is defined for all x in **R**. Given **f** and **g** in \mathcal{F} and a real number c, the functions $\mathbf{f} + \mathbf{g}$ and $c\mathbf{f}$ are defined in the natural way,

and

 $(\mathbf{f} + \mathbf{g})(x) = \mathbf{f}(x) + \mathbf{g}(x)$

 $(c\mathbf{f})(x) = c(\mathbf{f}(x)).$

Then each of properties (a)–(h) of a vector space follows readily from the corresponding property of the real numbers. For instance, if a is a scalar, then

 $[a(\mathbf{f} + \mathbf{g})](x) = a[(\mathbf{f} + \mathbf{g})(x)]$ = $a[\mathbf{f}(x) + \mathbf{g}(x)]$ = $a\mathbf{f}(x) + a\mathbf{g}(x)$ = $(a\mathbf{f} + a\mathbf{g})(x)$.

Thus $a(\mathbf{f} + \mathbf{g}) = a\mathbf{f} + a\mathbf{g}$, so \mathcal{F} enjoys property (e).

After verification of the other seven properties, we conclude that \mathcal{F} is, indeed, a vector space—one that differs in a fundamental way from each of the vector spaces \mathbf{R}^n . We naturally call \mathbf{R}^n an *n*-dimensional vector space, and in Section 4.4 we will define the word *dimension* in such a way that the dimension of \mathbf{R}^n actually *is n*. But the vector space \mathcal{F} of functions turns out not to have dimension *n* for any integer *n*; it is an example of an *infinite-dimensional* vector space. (See Section 4.4.)

Subspaces

Let $W = \{0\}$ be the subset of \mathbb{R}^n that contains only the zero vector $\mathbf{0}$. Then W satisfies properties (a)–(h) of a vector space, because each reduces trivially to the equation $\mathbf{0} = \mathbf{0}$. Thus W is a subset of the vector space $V = \mathbb{R}^n$ that is itself a vector space. According to the following definition, a subset of a vector space V that is itself a vector space is called a *subspace* of V.

DEFINITION Subspace

Let W be a nonempty subset of the vector space V. Then W is a **subspace** of V provided that W itself is a vector space with the operations of addition and multiplication by scalars as defined in V.

In order for the subset W to be a subspace of the vector space V, it first must be closed under the operations of vector addition and multiplication by scalars. Then it must satisfy properties (a)–(h) of a vector space. But W "inherits" all these properties from V, because the vectors in W all are vectors in V, and the vectors in V all satisfy properties (a)–(h). The following "subspace criterion" says that, in order to determine whether the subset W is a vector space, we need only check the two closure conditions.

THEOREM 1 Conditions for a Subspace

The nonempty subset W of the vector space V is a subspace of V if and only if it satisfies the following two conditions:

- (i) If **u** and **v** are vectors in W, then $\mathbf{u} + \mathbf{v}$ is also in W.
- (ii) If **u** is in W and c is a scalar, then the vector c**u** is also in W.

In Section 4.1 we saw that lines through the origin in \mathbb{R}^2 form subspaces of \mathbb{R}^2 and that lines and planes through the origin are subspaces of \mathbb{R}^3 . The subspace *W* of the following example may be regarded as a higher-dimensional "plane" (or "hyperplane") through the origin in \mathbb{R}^n .

Example 2 Let *W* be the subset of \mathbb{R}^n consisting of all those vectors $(x_1, x_2, ..., x_n)$ whose coordinates satisfy the single homogeneous linear equation

 $a_1x_1 + a_2x_2 + \dots + a_nx_n = 0,$

where the given coefficients $a_1, a_2, ..., a_n$ are not all zero. If $\mathbf{u} = (u_1, u_2, ..., u_n)$ and $\mathbf{v} = (v_1, v_2, ..., v_n)$ are vectors in W, then

$$a_1(u_1 + v_1) + a_2(u_2 + v_2) + \dots + a_n(u_n + v_n)$$

= $(a_1u_1 + a_2u_2 + \dots + a_nu_n) + (a_1v_1 + a_2v_2 + \dots + a_nv_n)$
= $0 + 0 = 0$,

so $\mathbf{u} + \mathbf{v} = (u_1 + v_1, \dots, u_n + v_n)$ is also in W. If c is a scalar, then

$$a_1(cu_1) + a_2(cu_2) + \dots + a_n(cu_n) = c(a_1u_1 + a_2u_2 + \dots + a_nu_n)$$

= (c)(0) = 0,

so $c\mathbf{u} = (cu_1, cu_2, \dots, cu_n)$ is in W. Thus we have shown that W satisfies conditions (i) and (ii) of Theorem 1 and is therefore a subspace of \mathbf{R}^n .

In order to apply Theorem 1 to show that W is a subspace of the vector space V, we must show that W satisfies *both* conditions in the theorem. But to apply Theorem 1 to show that W is *not* a subspace of V, we need only show *either* that condition (i) fails *or* that condition (ii) fails.

Example 3 Let *W* be the set of all those vectors (x_1, x_2, x_3, x_4) in \mathbb{R}^4 whose four coordinates are all nonnegative: $x_i \ge 0$ for i = 1, 2, 3, 4. Then it should be clear that the sum of two vectors in *W* is also a vector in *W*, because the sum of two nonnegative numbers is nonnegative. Thus *W* satisfies condition (i) of Theorem 1. But if we take $\mathbf{u} = (1, 1, 1, 1)$ in *W* and c = -1, then we find that the scalar multiple

$$c\mathbf{u} = (-1)(1, 1, 1, 1) = (-1, -1, -1, -1)$$

is not in W. Thus W fails to satisfy condition (ii) and therefore is not a subspace of \mathbb{R}^4 .

Example 4 Let *W* be the set of all those vectors (x_1, x_2, x_3, x_4) in \mathbb{R}^4 such that $x_1x_4 = 0$. Now *W* satisfies condition (ii) of Theorem 1, because $x_1x_4 = 0$ implies that $(cx_1)(cx_4) = 0$ for any scalar *c*. But if we take the vectors $\mathbf{u} = (1, 1, 0, 0)$ and $\mathbf{v} = (0, 0, 1, 1)$ in *W*, we see that their sum $\mathbf{u} + \mathbf{v} = (1, 1, 1, 1)$ is *not* in *W*. Thus *W* does not satisfy condition (i) and therefore is *not* a subspace of \mathbb{R}^4 .

Example 2 implies that the solution set of a homogeneous linear equation

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = 0$$

in *n* variables is always a *subspace* of \mathbb{R}^n . Theorem 2 further implies that the same is true of a *homogeneous* system of linear equations. Recall that any such system

of *m* equations in *n* unknowns can be written in the form $A\mathbf{x} = \mathbf{0}$, where **A** is the $m \times n$ coefficient matrix and $\mathbf{x} = (x_1, x_2, ..., x_n)$ is regarded as a column vector. The solution set of $A\mathbf{x} = \mathbf{0}$ is then the set of all vectors **x** in \mathbf{R}^n that satisfy this equation—that is, the set of all its solution vectors.

THEOREM 2 Solution Subspaces

If **A** is a (constant) $m \times n$ matrix, then the solution set of the homogeneous linear system

 $\mathbf{A}\mathbf{x} = \mathbf{0} \tag{3}$

is a subspace of \mathbf{R}^n .

Proof: Let W denote the solution set of Eq. (3). If **u** and **v** are vectors in W, then $A\mathbf{u} = A\mathbf{v} = \mathbf{0}$. Hence

$$A(u + v) = Au + Av = 0 + 0 = 0.$$

Thus the sum $\mathbf{u} + \mathbf{v}$ is also in *W*, and hence *W* satisfies condition (i) of Theorem 1. If *c* is a scalar, then

$$\mathbf{A}(c\mathbf{u}) = c(\mathbf{A}\mathbf{u}) = c\mathbf{0} = \mathbf{0},$$

so $c\mathbf{u}$ is in W if \mathbf{u} is in W. Thus W also satisfies condition (ii) of Theorem 1. It therefore follows that W is a subspace of \mathbf{R}^n .

Note that, in order to conclude by Theorem 2 that the solution set of a linear system is a subspace, it is necessary for the system to be *homogeneous*. Indeed, the solution set of a nonhomogeneous linear system

$$\mathbf{A}\mathbf{x} = \mathbf{b} \tag{4}$$

with $\mathbf{b} \neq \mathbf{0}$ is *never* a subspace. For if \mathbf{u} were a solution vector of the system in (4), then

$$\mathbf{A}(2\mathbf{u}) = 2(\mathbf{A}\mathbf{u}) = 2\mathbf{b} \neq \mathbf{b}$$

because $\mathbf{b} \neq \mathbf{0}$. Thus the scalar multiple 2**u** of the solution vector **u** is *not* a solution vector. Therefore, Theorem 1 implies that the solution set of (4) is *not* a subspace.

Because the solution set of a *homogeneous* linear system is a subspace, we often call it the **solution space** of the system. The subspaces of \mathbf{R}^n are the possible solution spaces of a homogeneous linear system with *n* unknowns, and this is one of the principal reasons for our interest in subspaces of \mathbf{R}^n .

At opposite extremes as subspaces of \mathbb{R}^n lie the **zero subspace** {0} and \mathbb{R}^n itself. Every other subspace of \mathbb{R}^n , each one that is neither {0} nor \mathbb{R}^n , is called a **proper subspace** of \mathbb{R}^n . The proper subspaces of \mathbb{R}^n play the same role in \mathbb{R}^n that lines and planes through the origin play in \mathbb{R}^3 . In the following two sections we develop the tools that are needed to analyze the structure of a given proper subspace of \mathbb{R}^n . In particular, given the homogeneous system $A\mathbf{x} = \mathbf{0}$, we ask how we can describe its solution space in a concise and illuminating way, beyond the mere statement that it is the set of all solution vectors of the system. Example 5 illustrates one possible way of doing this.

Example 5

In Example 4 of Section 3.4 we considered the homogeneous system

$$x_{1} + 3x_{2} - 15x_{3} + 7x_{4} = 0$$

$$x_{1} + 4x_{2} - 19x_{3} + 10x_{4} = 0$$

$$2x_{1} + 5x_{2} - 26x_{3} + 11x_{4} = 0.$$
(5)

4.2 THE VECTOR SPACEⁿ AND SUBSPACES 227

The reduced echelon form of the coefficient matrix of this system is

1	0	-3	-2	
0	1	-4	3	
0	0	0	0	

Hence x_1 and x_2 are the leading variables and x_3 and x_4 are free variables. Back substitution yields the general solution

$$x_3 = s$$
, $x_4 = t$, $x_2 = 4s - 3t$, $x_1 = 3s + 2t$

in terms of arbitrary parameters s and t. Thus a typical solution vector of the system in (5) has the form

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3s + 2t \\ 4s - 3t \\ s \\ t \end{bmatrix} = s \begin{bmatrix} 3 \\ 4 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} 2 \\ -3 \\ 0 \\ 1 \end{bmatrix}.$$

It follows that the solution space of the system in (5) can be described as the set of all linear combinations of the form

$$\mathbf{x} = s\mathbf{u} + t\mathbf{v},\tag{6}$$

where $\mathbf{u} = (3, 4, 1, 0)$ and $\mathbf{v} = (2, -3, 0, 1)$. Thus we have found two particular solution vectors \mathbf{u} and \mathbf{v} of our system that completely determine its solution space [by the formula in (6)].

4.2 **Problems**

In Problems 1–14, a subset W of some n-space \mathbb{R}^n is defined by means of a given condition imposed on the typical vector (x_1, x_2, \dots, x_n) . Apply Theorem 1 to determine whether or not W is a subspace of \mathbb{R}^n .

- **1.** W is the set of all vectors in \mathbf{R}^3 such that $x_3 = 0$.
- **2.** *W* is the set of all vectors in \mathbb{R}^3 such that $x_1 = 5x_2$.
- **3.** *W* is the set of all vectors in \mathbf{R}^3 such that $x_2 = 1$.
- 4. W is the set of all vectors in \mathbb{R}^3 such that $x_1 + x_2 + x_3 = 1$.
- 5. W is the set of all vectors in \mathbb{R}^4 such that $x_1 + 2x_2 + 3x_3 + 4x_4 = 0$.
- 6. W is the set of all vectors in \mathbb{R}^4 such that $x_1 = 3x_3$ and $x_2 = 4x_4$.
- 7. *W* is the set of all vectors in \mathbb{R}^2 such that $|x_1| = |x_2|$.
- 8. W is the set of all vectors in \mathbb{R}^2 such that $(x_1)^2 + (x_2)^2 = 0$.
- 9. W is the set of all vectors in \mathbb{R}^2 such that $(x_1)^2 + (x_2)^2 = 1$.
- **10.** W is the set of all vectors in \mathbf{R}^2 such that $|x_1| + |x_2| = 1$.
- **11.** W is the set of all vectors in \mathbf{R}^4 such that $x_1 + x_2 = x_3 + x_4$.
- **12.** *W* is the set of all vectors in \mathbf{R}^4 such that $x_1x_2 = x_3x_4$.
- **13.** W is the set of all vectors in \mathbf{R}^4 such that $x_1x_2x_3x_4 = 0$.
- 14. W is the set of all those vectors in \mathbf{R}^4 whose components are all nonzero.

In Problems 15–18, apply the method of Example 5 to find two solution vectors \mathbf{u} and \mathbf{v} such that the solution space is the set of all linear combinations of the form $s\mathbf{u} + t\mathbf{v}$.

15.
$$x_1 - 4x_2 + x_3 - 4x_4 = 0$$

 $x_1 + 2x_2 + x_3 + 8x_4 = 0$
 $x_1 + x_2 + x_3 + 6x_4 = 0$

- 16. $x_1 4x_2 3x_3 7x_4 = 0$ $2x_1 - x_2 + x_3 + 7x_4 = 0$ $x_1 + 2x_2 + 3x_3 + 11x_4 = 0$ 17. $x_1 + 3x_2 + 8x_3 - x_4 = 0$ $x_1 - 3x_2 - 10x_3 + 5x_5 = 0$ $x_1 + 4x_2 + 11x_3 - 2x_4 = 0$ 18. $x_1 + 3x_2 + 2x_3 + 5x_4 - x_5 = 0$
- $\begin{array}{rcrcrcrc} x_1 + 5x_2 + 2x_3 + 5x_4 x_5 = 0\\ 2x_1 + 7x_2 + 4x_3 + 11x_4 + 2x_5 = 0\\ 2x_1 + 6x_2 + 5x_3 + 12x_4 7x_5 = 0 \end{array}$

In Problems 19–22, reduce the given system to echelon form to find a single solution vector **u** such that the solution space is the set of all scalar multiples of **u**.

- **19.** $x_1 3x_2 5x_3 6x_4 = 0$ $2x_1 + x_2 + 4x_3 - 4x_4 = 0$ $x_1 + 3x_2 + 7x_3 + x_4 = 0$
- **20.** $x_1 + 5x_2 + x_3 8x_4 = 0$ $2x_1 + 5x_2 - 5x_4 = 0$
 - $2x_1 + 7x_2 + x_3 9x_4 = 0$
- **21.** $x_1 + 7x_2 + 2x_3 3x_4 = 0$ $2x_1 + 7x_2 + x_3 - 4x_4 = 0$ $3x_1 + 5x_2 - x_3 - 5x_4 = 0$
- **22.** $x_1 + 3x_2 + 3x_3 + 3x_4 = 0$ $2x_1 + 7x_2 + 5x_3 - x_4 = 0$ $2x_1 + 7x_2 + 4x_3 - 4x_4 = 0$
- **23.** Show that every subspace *W* of a vector space *V* contains the zero vector **0**.
- **24.** Apply the properties of a vector space *V* to show each of the following.
 - (a) $0\mathbf{u} = \mathbf{0}$ for every \mathbf{u} in V.
 - **(b)** $c\mathbf{0} = \mathbf{0}$ for every scalar *c*.

228 CHAPTER 4 VECTOR SPACES

(c) $(-1)\mathbf{u} = -\mathbf{u}$ for every \mathbf{u} in V.

Do *not* assume that the vectors in V are *n*-tuples of real numbers.

- **25.** Show that the nonempty subset W of a vector space V is a subspace of V if and only if for every pair of vectors \mathbf{u} and \mathbf{v} in W and every pair of scalars a and b, $a\mathbf{u} + b\mathbf{v}$ is also in W.
- 26. Prove: If u is a (fixed) vector in the vector space V, then the set W of all scalar multiples cu of u is a subspace of V.
- 27. Let u and v be (fixed) vectors in the vector space V. Show that the set W of all linear combinations au + bv of u and v is a subspace of V.
- **28.** Suppose that **A** is an $n \times n$ matrix and that k is a (constant) scalar. Show that the set of all vectors **x** such that $A\mathbf{x} = k\mathbf{x}$ is a subspace of \mathbf{R}^n .

- **29.** Let **A** be an $n \times n$ matrix, **b** be a nonzero vector, and \mathbf{x}_0 be a solution vector of the system $\mathbf{A}\mathbf{x} = \mathbf{b}$. Show that **x** is a solution of the nonhomogeneous system $\mathbf{A}\mathbf{x} = \mathbf{b}$ if and only if $\mathbf{y} = \mathbf{x} \mathbf{x}_0$ is a solution of the homogeneous system $\mathbf{A}\mathbf{y} = \mathbf{0}$.
- **30.** Let U and V be subspaces of the vector space W. Their intersection $U \cap V$ is the set of all vectors that are both in U and in V. Show that $U \cap V$ is a subspace of W. If U and V are two planes through the origin in \mathbb{R}^3 , what is $U \cap V$?
- **31.** Let *U* and *V* be subspaces of the vector space *W*. Their sum U + V is the set of all vectors **w** of the form

$$\mathbf{w} = \mathbf{u} + \mathbf{v},$$

where **u** is in U and **v** is in V. Show that U + V is a subspace of W. If U and V are lines through the origin in \mathbb{R}^3 , what is U + V?

4.3 Linear Combinations and Independence of Vectors

In Example 5 of Section 4.2 we solved the homogeneous linear system

$$x_1 + 3x_2 - 15x_3 + 7x_4 = 0$$

$$x_1 + 4x_2 - 19x_3 + 10x_4 = 0$$

$$2x_1 + 5x_2 - 26x_3 + 11x_4 = 0.$$
(1)

We found that its solution space W consists of all those vectors \mathbf{x} in \mathbf{R}^4 that have the form

$$\mathbf{x} = s(3, 4, 1, 0) + t(2, -3, 0, 1).$$
⁽²⁾

We therefore can visualize W as the plane in \mathbf{R}^4 determined by the vectors $\mathbf{v}_1 = (3, 4, 1, 0)$ and $\mathbf{v}_2 = (2, -3, 0, 1)$. The fact that every solution vector is a combination [as in (2)] of the particular solution vectors \mathbf{v}_1 and \mathbf{v}_2 gives us a tangible understanding of the solution space W of the system in (1).

More generally, we know from Theorem 2 in Section 4.2 that the solution set V of any $m \times n$ homogeneous linear system $\mathbf{Ax} = \mathbf{0}$ is a subspace of \mathbf{R}^n . In order to understand such a vector space V better, we would like to find a minimal set of vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ in V such that *every* vector in V is a sum of scalar multiples of these particular vectors.

The vector **w** is called a **linear combination** of the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ provided that there exist scalars c_1, c_2, \ldots, c_k such that

$$\mathbf{w} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_k \mathbf{v}_k. \tag{3}$$

Given a vector **w** in \mathbb{R}^n , the problem of determining whether or not **w** is a linear combination of the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ amounts to solving a linear system to see whether we can find scalars c_1, c_2, \ldots, c_k so that (3) holds.

Example 1

To determine whether the vector $\mathbf{w} = (2, -6, 3)$ in \mathbf{R}^3 is a linear combination of the vectors $\mathbf{v}_1 = (1, -2, -1)$ and $\mathbf{v}_2 = (3, -5, 4)$, we write the equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 = \mathbf{w}$ in matrix form:

$$c_1 \begin{bmatrix} 1\\-2\\-1 \end{bmatrix} + c_2 \begin{bmatrix} 3\\-5\\4 \end{bmatrix} = \begin{bmatrix} 2\\-6\\3 \end{bmatrix}$$

—that is,

$$c_1 + 3c_2 = 2$$

-2c_1 - 5c_2 = -6
-c_1 + 4c_2 = 3.

The augmented coefficient matrix

$$\begin{bmatrix} 1 & 3 & 2 \\ -2 & -5 & -6 \\ -1 & 4 & 3 \end{bmatrix}$$

can be reduced by elementary row operations to echelon form:

$\lceil 1 \rceil$	3	2]
0	1	-2
0	0	19

We see now, from the third row, that our system is inconsistent, so the desired scalars c_1 and c_2 do not exist. Thus w is *not* a linear combination of v_1 and v_2 .

Example 2

To express the vector $\mathbf{w} = (-7, 7, 11)$ as a linear combination of the vectors $\mathbf{v}_1 = (1, 2, 1)$, $\mathbf{v}_2 = (-4, -1, 2)$, and $\mathbf{v}_3 = (-3, 1, 3)$, we write the equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = \mathbf{w}$ in the form

$$c_1 \begin{bmatrix} 1\\2\\1 \end{bmatrix} + c_2 \begin{bmatrix} -4\\-1\\2 \end{bmatrix} + c_3 \begin{bmatrix} -3\\1\\3 \end{bmatrix} = \begin{bmatrix} -7\\7\\11 \end{bmatrix}$$

-that is,

$$c_1 - 4c_2 - 3c_3 = -7$$

$$2c_1 - c_2 + c_3 = 7$$

$$c_1 + 2c_2 + 3c_3 = 11.$$

The reduced echelon form of the augmented coefficient matrix of this system is

$$\begin{bmatrix} 1 & 0 & 1 & 5 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Thus c_3 is a free variable. With $c_3 = t$, back substitution yields $c_1 = 5 - t$ and $c_2 = 3 - t$. For instance, t = 1 gives $c_1 = 4$, $c_2 = 2$, and $c_3 = 1$, so

$$\mathbf{w} = 4\mathbf{v}_1 + 2\mathbf{v}_2 + \mathbf{v}_3.$$

But t = -2 yields $c_1 = 7$, $c_2 = 5$, and $c_3 = -2$, so w can also be expressed as

$$\mathbf{v} = 7\mathbf{v}_1 + 5\mathbf{v}_2 - 2\mathbf{v}_3.$$

We have found not only that w can be expressed as a linear combination of the vectors v_1 , v_2 , v_3 but also that this can be done in many different ways (one for each choice of the parameter t).

We began this section with the observation that every solution vector of the linear system in (1) is a linear combination of the vectors \mathbf{v}_1 and \mathbf{v}_2 that appear in the right-hand side of Eq. (2). A brief way of saying this is that the vectors \mathbf{v}_1 and \mathbf{v}_2 span the solution space. More generally, suppose that $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ are vectors in a vector space V. Then we say that the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ span the vector space V provided that every vector in V is a linear combination of these k vectors. We may also say that the set $S = {\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k}$ of vectors is a spanning set for V.

Example 3 The familiar unit vectors $\mathbf{i} = (1, 0, 0)$, $\mathbf{j} = (0, 1, 0)$, and $\mathbf{k} = (0, 0, 1)$ span \mathbf{R}^3 , because every vector $\mathbf{x} = (x_1, x_2, x_3)$ in \mathbf{R}^3 can be expressed as the linear combination

$$\mathbf{x} = x_1 \mathbf{i} + x_2 \mathbf{j} + x_3 \mathbf{k}$$

of these three vectors i, j, and k.

If the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ in the vector space V do not span V, we can ask about the subset of V consisting of all those vectors that are linear combinations of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. The following theorem implies that this subset is always a *subspace* of V.

THEOREM 1 The Span of a Set of Vectors

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ be vectors in the vector space V. Then the set W of all linear combinations of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ is a subspace of V.

Proof: We must show that W is closed under addition of vectors and multiplication by scalars. If

$$\mathbf{u} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_k \mathbf{v}_k$$

and

$$\mathbf{v} = b_1 \mathbf{v}_1 + b_2 \mathbf{v}_2 + \dots + b_k \mathbf{v}_k$$

are vectors in W, then

$$\mathbf{u} + \mathbf{v} = (a_1 + b_1)\mathbf{v}_1 + (a_2 + b_2)\mathbf{v}_2 + \dots + (a_k + b_k)\mathbf{v}_k$$

and

$$c\mathbf{u} = (ca_1)\mathbf{v}_1 + (ca_2)\mathbf{v}_2 + \dots + (ca_k)\mathbf{v}_k$$

for any scalar c. Thus $\mathbf{u} + \mathbf{v}$ and $c\mathbf{u}$ are linear combinations of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ and therefore are vectors in W. It is clear that W is nonempty, and hence W is a subspace of V.

We say that the subspace W of Theorem 1 is the space **spanned** by the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ (or is the **span** of the set $S = {\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k}$ of vectors). We sometimes write

$$W = \operatorname{span}(S) = \operatorname{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}.$$

Thus Example 3 implies that $\mathbf{R}^3 = \text{span}\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$. The question as to whether a given vector \mathbf{w} in \mathbf{R}^n lies in the subspace span $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ reduces to solving a linear system, as illustrated by Examples 1 and 2.

It is easy to verify that the space $W = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ of Theorem 1 is the smallest subspace of V that contains all the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ —meaning that every other subspace of V that contains these k vectors must also contain W (Problem 30).

Linear Independence

Henceforth, when we solve a homogeneous system of linear equations, we generally will seek a set $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ of solution vectors that span the solution space W of the system. Perhaps the most concrete way to describe a subspace W of a vector space V is to give explicitly a set $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ of vectors that span W. And this type of representation is most useful and desirable (as well as most aesthetically pleasing) when each vector \mathbf{w} in W is expressible in a *unique* way as a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$. (For instance, each vector in \mathbf{R}^3 is a unique linear combination of the vectors \mathbf{i}, \mathbf{j} , and \mathbf{k} of Example 3.) But Example 2 demonstrates that a vector \mathbf{w} may well be expressed in many different ways as a linear combination of given vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$.

Thus not all spanning sets enjoy the uniqueness property that we desire. Two questions arise:

- Given a subspace W of a vector space V, does there necessarily exist a spanning set with the uniqueness property that we desire?
- If so, how do we find such a spanning set for W?

The following definition provides the key to both answers.

DEFINITION Linear Independence

The vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ in a vector space V are said to be **linearly independent** provided that the equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0} \tag{4}$$

has only the trivial solution $c_1 = c_2 = \cdots = c_k = 0$. That is, the only linear combination of $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ that represents the zero vector **0** is the trivial combination $0\mathbf{v}_1 + 0\mathbf{v}_2 + \cdots + 0\mathbf{v}_k.$

Remark We can immediately verify that any vector w in the subspace W spanned by the linearly independent vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ is *uniquely* expressible as a linear combination of these vectors. For

$$\mathbf{w} = \sum_{i=1}^{k} a_i \mathbf{v}_i = \sum_{i=1}^{k} b_i \mathbf{v}_i \quad \text{implies that} \quad \sum_{i=1}^{k} (a_i - b_i) \mathbf{v}_i = \mathbf{0}.$$

Hence, with $c_i = a_i - b_i$ for each i = 1, ..., k, the linear independence of $\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_k$ implies that $c_1 = c_2 = \cdots = c_k = 0$. Thus

$$\sum_{i=1}^{k} a_i \mathbf{v}_i = \sum_{i=1}^{k} b_i \mathbf{v}_i \quad \text{implies that} \quad a_i = b_i \text{ for } i = 1, \dots, k,$$

so we see that w can be expressed in only *one* way as a combination of the linearly independent vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$.

Example 4	The standard unit vectors
	$\mathbf{e}_1 = (1, 0, 0, \dots, 0),$
	$\mathbf{e}_2 = (0, 1, 0, \dots, 0),$
	:
	$\mathbf{e}_n = (0, 0, 0, \dots, 1)$
	in \mathbf{R}^n are linearly independent. The reason is that the equation
	$c_1\mathbf{e}_1 + c_2\mathbf{e}_2 + \dots + c_n\mathbf{e}_n = 0$

evidently reduces to

 $(c_1, c_2, \ldots, c_n) = (0, 0, \ldots, 0)$ and thus has only the trivial solution $c_1 = c_2 = \cdots = c_n = 0$.

To determine whether the vectors $\mathbf{v}_1 = (1, 2, 2, 1)$, $\mathbf{v}_2 = (2, 3, 4, 1)$, and $\mathbf{v}_3 = (3, 8, 7, 5)$ in \mathbb{R}^4 Example 5 are linearly independent, we write the equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = \mathbf{0}$ as the linear system

> $c_1 + 2c_2 + 3c_3 = 0$ $2c_1 + 3c_2 + 8c_3 = 0$ $2c_1 + 4c_2 + 7c_3 = 0$ $c_1 + c_2 + 5c_3 = 0$

and then solve for c_1 , c_2 , and c_3 . The augmented coefficient matrix of this system reduces to the echelon form

2	3	0	
1	-2	0	
0	1	0	,
0	0	0	
	2 1 0 0	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

so we see that the only solution is $c_1 = c_2 = c_3 = 0$. Thus the vectors \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 are linearly independent.

Observe that linear independence of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ actually is a property of the set $S = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k}$ whose elements are these vectors. Occasionally the phraseology "the set $S = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k}$ is linearly independent" is more convenient. For instance, any subset of a linearly independent set $S = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k}$ is a linearly independent set of vectors (Problem 29).

Now we show that the coefficients in a linear combination of the linearly independent vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are unique. If both

$$\mathbf{w} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_k \mathbf{v}_k \tag{5}$$

and

$$\mathbf{w} = b_1 \mathbf{v}_1 + b_2 \mathbf{v}_2 + \dots + b_k \mathbf{v}_k,\tag{6}$$

then

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_k\mathbf{v}_k = b_1\mathbf{v}_1 + b_2\mathbf{v}_2 + \dots + b_k\mathbf{v}_k,$$

so it follows that

$$(a_1 - b_1)\mathbf{v}_1 + (a_2 - b_2)\mathbf{v}_2 + \dots + (a_k - b_k)\mathbf{v}_k = \mathbf{0}.$$
(7)

Because the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent, each of the coefficients in (7) must vanish. Therefore, $a_1 = b_1, a_2 = b_2, \dots, a_k = b_k$, so we have shown that the linear combinations in (5) and (6) actually are identical. Hence, if a vector \mathbf{w} is in the set span{ $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ }, then it can be expressed in only one way as a linear combination of these linearly independent vectors.

A set of vectors is called **linearly dependent** provided it is not linearly independent. Hence the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly dependent if and only if there exist scalars c_1, c_2, \dots, c_k not all zero such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0}.$$
 (8)

In short, a (finite) set of vectors is linearly dependent provided that some *nontrivial* linear combination of them equals the zero vector.

Example 6 Let $\mathbf{v}_1 = (2, 1, 3)$, $\mathbf{v}_2 = (5, -2, 4)$, $\mathbf{v}_3 = (3, 8, -6)$, and $\mathbf{v}_4 = (2, 7, -4)$. Then the equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + c_4\mathbf{v}_4 = \mathbf{0}$ is equivalent to the linear system

 $2c_1 + 5c_2 + 3c_3 + 2c_4 = 0$ $c_1 - 2c_2 + 8c_3 + 7c_4 = 0$ $3c_1 + 4c_2 - 6c_3 - 4c_4 = 0$

of three equations in four unknowns. Because this homogeneous system has more unknowns than equations, Theorem 3 in Section 3.3 implies that it has a nontrivial solution. Therefore we may conclude—without even solving explicitly for c_1 , c_2 , c_3 , and c_4 —that the vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 , and \mathbf{v}_4 are linearly dependent. (It happens that

$$2\mathbf{v}_1 - \mathbf{v}_2 + 3\mathbf{v}_3 - 4\mathbf{v}_4 = \mathbf{0},$$

4.3 LINEAR COMBINATIONS AND INDEPENDENCE OF VECTORS

The argument in Example 6 may be generalized in an obvious way to prove that *any set of more than n vectors in* \mathbb{R}^n *is linearly dependent*. For if k > n, then Eq. (8) is equivalent to a homogeneous linear system with more unknowns (k) than equations (n), so Theorem 3 in Section 3.3 yields a nontrivial solution.

We now look at the way in which the elements of a linearly dependent set of vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ "depend" on one another. We know that there exist scalars c_1, c_2, \ldots, c_k not all zero such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0}.$$
(9)

Suppose that the *p*th coefficient is nonzero: $c_p \neq 0$. Then we can solve Eq. (9) for $c_p \mathbf{v}_p$ and next divide by c_p to get

$$\mathbf{v}_{p} = a_{1}\mathbf{v}_{1} + \dots + a_{p-1}\mathbf{v}_{p-1} + a_{p+1}\mathbf{v}_{p+1} + \dots + a_{k}\mathbf{v}_{k}, \tag{10}$$

where $a_i = -c_i/c_p$ for $i \neq p$. Thus at least one of the linearly dependent vectors is a linear combination of the other k - 1. Conversely, suppose we are given a set of vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ with one of them dependent on the others as in Eq. (10). Then we can transpose all the terms to the left-hand side to get an equation of the form in (9) with $c_p = 1 \neq 0$. This shows that the vectors are linearly dependent. Therefore, we have proved that the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ are linearly dependent if and only if at least one of them is a linear combination of the others.

For instance (as we saw in Section 4.1), two vectors are linearly dependent if and only if one of them is a scalar multiple of the other, in which case the two vectors are collinear. Three vectors are linearly dependent if and only if one of them is a linear combination of the other two, in which case the three vectors are coplanar.

In Theorem 4 of Section 4.1 we saw that the *determinant* provides a criterion for deciding whether three vectors in \mathbf{R}^3 are linearly independent: The vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 in \mathbf{R}^3 are linearly independent if and only if the determinant of the 3 × 3 matrix

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 \end{bmatrix}$$

is nonzero. The proof given there in the three-dimensional case generalizes readily to the *n*-dimensional case. Given *n* vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ in \mathbf{R}^n , we consider the $n \times n$ matrix

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix}$$

having these vectors as its column vectors. Then, by Theorem 2 in Section 3.6, det $\mathbf{A} \neq 0$ if and only if \mathbf{A} is invertible, in which case the system $\mathbf{Ac} = \mathbf{0}$ has only the trivial solution $c_1 = c_2 = \cdots = c_n = 0$, so the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ must be linearly independent.

THEOREM 2 Independence of n Vectors in Rⁿ

The *n* vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ in \mathbf{R}^n are linearly independent if and only if the $n \times n$ matrix

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix}$$

having them as its column vectors has nonzero determinant.

We saw earlier that a set of *more* than *n* vectors in \mathbb{R}^n is always linearly dependent. The following theorem shows us how the determinant provides a criterion in the case of *fewer* than *n* vectors in \mathbb{R}^n .

THEOREM 3 Independence of Fewer Than n Vectors in Rⁿ

Consider k vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ in \mathbf{R}^n , with k < n. Let

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_k \end{bmatrix}$$

be the $n \times k$ matrix having them as its column vectors. Then the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ are linearly independent if and only if some $k \times k$ submatrix of **A** has nonzero determinant.

Rather than including a complete proof, we will simply illustrate the "if" part of Theorem 3 in the case n = 5, k = 3. Let $\mathbf{v}_1 = (a_1, a_2, a_3, a_4, a_5)$, $\mathbf{v}_2 = (b_1, b_2, b_3, b_4, b_5)$, and $\mathbf{v}_3 = (c_1, c_2, c_3, c_4, c_5)$ be three vectors in \mathbf{R}^5 such that the 5×3 matrix

 $\mathbf{A} = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \\ a_4 & b_4 & c_4 \\ a_5 & b_5 & c_5 \end{bmatrix}$

has a 3×3 submatrix with nonzero determinant. Suppose, for instance, that

a_1	b_1	c_1	
a_3	b_3	С3	$\neq 0$
a_5	b_5	c_5	

Then Theorem 2 implies that the three vectors $\mathbf{u}_1 = (a_1, a_3, a_5)$, $\mathbf{u}_2 = (b_1, b_3, b_5)$, and $\mathbf{u}_3 = (c_1, c_3, c_5)$ in \mathbb{R}^3 are linearly independent. Now suppose that $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = \mathbf{0}$. Then by deleting the second and fourth components of each vector in this equation, we find that $c_1\mathbf{u}_1 + c_2\mathbf{u}_2 + c_3\mathbf{u}_3 = \mathbf{0}$. But the fact that \mathbf{u}_1 , \mathbf{u}_2 , \mathbf{u}_3 are linearly independent implies that $c_1 = c_2 = c_3 = 0$, and it now follows that \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 are linearly independent.

4.3 **Problems**

In Problems 1–8, determine whether the given vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ are linearly independent or linearly dependent. Do this essentially by inspection—that is, without solving a linear system of equations.

1. $\mathbf{v}_1 = (4, -2, 6, -4), \mathbf{v}_2 = (6, -3, 9, -6)$

- **2.** $\mathbf{v}_1 = (3, 9, -3, 6), \mathbf{v}_2 = (2, 6, -2, 4)$
- **3.** $\mathbf{v}_1 = (3, 4), \mathbf{v}_2 = (6, -1), \mathbf{v}_3 = (7, 5)$
- **4.** $\mathbf{v}_1 = (4, -2, 2), \mathbf{v}_2 = (5, 4, -3), \mathbf{v}_3 = (4, 6, 5), \mathbf{v}_4 = (-7, 9, 3)$
- **5.** $\mathbf{v}_1 = (1, 0, 0), \mathbf{v}_2 = (0, -2, 0), \mathbf{v}_3 = (0, 0, 3)$
- **6.** $\mathbf{v}_1 = (1, 0, 0), \mathbf{v}_2 = (1, 1, 0), \mathbf{v}_3 = (1, 1, 1)$
- 7. $\mathbf{v}_1 = (2, 1, 0, 0), \mathbf{v}_2 = (3, 0, 1, 0), \mathbf{v}_3 = (4, 0, 0, 1)$
- 8. $\mathbf{v}_1 = (1, 0, 3, 0), \mathbf{v}_2 = (0, 2, 0, 4), \mathbf{v}_3 = (1, 2, 3, 4)$

In Problems 9–16, express the indicated vector \mathbf{w} as a linear combination of the given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ if this is possible. If not, show that it is impossible.

9. $\mathbf{w} = (1, 0, -7); \mathbf{v}_1 = (5, 3, 4), \mathbf{v}_2 = (3, 2, 5)$ **10.** $\mathbf{w} = (3, -1, -2); \mathbf{v}_1 = (-3, 1, -2), \mathbf{v}_2 = (6, -2, 3)$

- **11.** $\mathbf{w} = (1, 0, 0, -1); \mathbf{v}_1 = (7, -6, 4, 5), \mathbf{v}_2 = (3, -3, 2, 3)$
- **12.** $\mathbf{w} = (4, -4, 3, 3); \mathbf{v}_1 = (7, 3, -1, 9), \mathbf{v}_2 = (-2, -2, 1, -3)$
- **13.** $\mathbf{w} = (5, 2, -2); \mathbf{v}_1 = (1, 5, -3), \mathbf{v}_2 = (5, -3, 4)$
- **14.** $\mathbf{w} = (2, -3, 2, -3); \mathbf{v}_1 = (1, 0, 0, 3), \mathbf{v}_2 = (0, 1, -2, 0), \mathbf{v}_3 = (0, -1, 1, 1)$
- **15.** $\mathbf{w} = (4, 5, 6); \mathbf{v}_1 = (2, -1, 4), \mathbf{v}_2 = (3, 0, 1), \mathbf{v}_3 = (1, 2, -1)$
- **16.** $\mathbf{w} = (7, 7, 9, 11); \mathbf{v}_1 = (2, 0, 3, 1), \mathbf{v}_2 = (4, 1, 3, 2), \mathbf{v}_3 = (1, 3, -1, 3)$

In Problems 17–22, three vectors \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 are given. If they are linearly independent, show this; otherwise find a nontrivial linear combination of them that is equal to the zero vector.

17. $\mathbf{v}_1 = (1, 0, 1), \, \mathbf{v}_2 = (2, -3, 4), \, \mathbf{v}_3 = (3, 5, 2)$ **18.** $\mathbf{v}_1 = (2, 0, -3), \, \mathbf{v}_2 = (4, -5, -6), \, \mathbf{v}_3 = (-2, 1, 3)$ **19.** $\mathbf{v}_1 = (2, 0, 3, 0), \, \mathbf{v}_2 = (5, 4, -2, 1), \, \mathbf{v}_3 = (2, -1, 1, -1)$ **20.** $\mathbf{v}_1 = (1, 1, -1, 1), \, \mathbf{v}_2 = (2, 1, 1, 1), \, \mathbf{v}_3 = (3, 1, 4, 1)$

4.4 BASES AND DIMENSION FOR VECTOR SPACES 5

21. $\mathbf{v}_1 = (3, 0, 1, 2), \mathbf{v}_2 = (1, -1, 0, 1), \mathbf{v}_3 = (1, 2, 1, 0)$ **22.** $\mathbf{v}_1 = (3, 9, 0, 5), \mathbf{v}_2 = (3, 0, 9, -7), \mathbf{v}_3 = (4, 7, 5, 0)$

In Problems 23–26, the vectors $\{\mathbf{v}_i\}$ are known to be linearly independent. Apply the definition of linear independence to show that the vectors $\{\mathbf{u}_i\}$ are also linearly independent.

- **23.** $\mathbf{u}_1 = \mathbf{v}_1 + \mathbf{v}_2, \mathbf{u}_2 = \mathbf{v}_1 \mathbf{v}_2$
- **24.** $\mathbf{u}_1 = \mathbf{v}_1 + \mathbf{v}_2, \mathbf{u}_2 = 2\mathbf{v}_1 + 3\mathbf{v}_2$
- **25.** $\mathbf{u}_1 = \mathbf{v}_1, \mathbf{u}_2 = \mathbf{v}_1 + 2\mathbf{v}_2, \mathbf{u}_3 = \mathbf{v}_1 + 2\mathbf{v}_2 + 3\mathbf{v}_3$
- **26.** $\mathbf{u}_1 = \mathbf{v}_2 + \mathbf{v}_3$, $\mathbf{u}_2 = \mathbf{v}_1 + \mathbf{v}_3$, $\mathbf{u}_3 = \mathbf{v}_1 + \mathbf{v}_2$
- **27.** *Prove*: If the (finite) set *S* of vectors contains the zero vector, then *S* is linearly dependent.
- **28.** *Prove*: If the set *S* of vectors is linearly dependent and the (finite) set *T* contains *S*, then *T* is also linearly dependent. You may assume that $S = {\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_k}$ and that $T = {\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_m}$ with m > k.
- **29.** Show that if the (finite) set *S* of vectors is linearly independent, then any subset *T* of *S* is also linearly independent.
- **30.** Suppose that the subspace U of the vector space V contains the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. Show that U contains the subspace spanned by these vectors.

- **31.** Let *S* and *T* be sets of vectors in a vector space such that *S* is a subset of span(*T*). Show that span(*S*) is also a subset of span(*T*).
- **32.** Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ be linearly independent vectors in the set *S* of vectors. *Prove*: If no set of more than *k* vectors in *S* is linearly independent, then every vector in *S* is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$.

In Problems 33–35, let $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ be vectors in \mathbf{R}^n and let

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_k \end{bmatrix}$$

be the $n \times k$ matrix with these vectors as its column vectors.

- **33.** *Prove*: If some $k \times k$ submatrix of **A** is the $k \times k$ identity matrix, then $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$ are linearly independent.
- **34.** Suppose that k = n, that the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent, and that **B** is a nonsingular $n \times n$ matrix. Prove that the column vectors of the matrix **AB** are linearly independent.
- **35.** Suppose that k < n, that the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent, and that **B** is a nonsingular $k \times k$ matrix. Use Theorem 3 to show that the column vectors of **AB** are linearly independent.

4.4 Bases and Dimension for Vector Spaces

An especially useful way of describing the solution space of a homogeneous linear system is to list explicitly a set *S* of solution vectors such that *every* solution vector is a unique linear combination of these particular ones. The following definition specifies the properties of such a set *S* of "basic" solution vectors, and the concept is equally important for vector spaces other than solution spaces.

DEFINITION Basis

A finite set S of vectors in a vector space V is called a **basis** for V provided that (a) the vectors in S are linearly independent, and

(**b**) the vectors in *S* span *V*.

In short, a basis for the vector space V is a linearly independent spanning set of vectors in V. Thus, if $S = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n}$ is a basis for V, then any vector **w** in V can be expressed as a linear combination

$$\mathbf{w} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n \tag{1}$$

of the vectors in S, and we saw in Section 4.3 that the linear independence of S implies that the coefficients c_1, c_2, \ldots, c_n in (1) are *unique*. That is, w cannot be expressed differently as a linear combination of the basis vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$.

Example 1 The **standard basis** for \mathbf{R}^n consists of the unit vectors

$$\mathbf{e}_1 = (1, 0, 0, \dots, 0), \ \mathbf{e}_2 = (0, 1, 0, \dots, 0), \ \dots, \ \mathbf{e}_n = (0, 0, 0, \dots, 1)$$

If $\mathbf{x} = (x_1, x_2, \dots, x_n)$ is a vector in \mathbf{R}^n , then

$$\mathbf{x} = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n.$$

Thus the column vectors

$$\mathbf{e}_1 = \begin{bmatrix} 1\\0\\\vdots\\0 \end{bmatrix}, \quad \mathbf{e}_2 = \begin{bmatrix} 0\\1\\\vdots\\0 \end{bmatrix}, \quad \dots, \quad \mathbf{e}_n = \begin{bmatrix} 0\\0\\\vdots\\1 \end{bmatrix}$$

of the $n \times n$ identity matrix span \mathbb{R}^n , and we noted in Example 4 of Section 4.3 that these *standard unit vectors* are linearly independent.

Example 2 Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be *n* linearly independent vectors in \mathbf{R}^n . We saw in Section 4.3 that any set of more than *n* vectors in \mathbf{R}^n is linearly dependent. Hence, given a vector \mathbf{w} in \mathbf{R}^n , there exist scalars c, c_1, c_2, \dots, c_n not all zero such that

$$c\mathbf{w} + c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_n\mathbf{v}_n = \mathbf{0}.$$
 (2)

If *c* were zero, then (2) would imply that the vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ are linearly dependent. Hence $c \neq 0$, so Eq. (2) can be solved for **w** as a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$. Thus the linearly independent vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ also *span* \mathbf{R}^n and therefore constitute a basis for \mathbf{R}^n .

Example 2 shows that any set of n linearly independent vectors in \mathbb{R}^n is a basis for \mathbb{R}^n . By Theorem 2 in Section 4.3, we can therefore determine whether n given vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ form a basis for \mathbb{R}^n by calculating the determinant of the $n \times n$ matrix

$$\mathbf{A} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix}$$

with these vectors as its column vectors. They constitute a basis for \mathbf{R}^n if and only if det $\mathbf{A} \neq 0$.

Example 3 Let $\mathbf{v}_1 = (1, -1, -2, -3)$, $\mathbf{v}_2 = (1, -1, 2, 3)$, $\mathbf{v}_3 = (1, -1, -3, -2)$, and $\mathbf{v}_4 = (0, 3, -1, 2)$. Then we find that

$$\begin{vmatrix} 1 & 1 & 1 & 0 \\ -1 & -1 & -1 & 3 \\ -2 & 2 & -3 & -1 \\ -3 & 3 & -2 & 2 \end{vmatrix} = 30 \neq 0,$$

so it follows that $\{v_1, v_2, v_3, v_4\}$ is a basis for \mathbb{R}^4 .

Theorem 1 has the following import: Just as in \mathbb{R}^n , a basis for *any* vector space V contains the largest possible number of linearly independent vectors in V.

THEOREM 1 Bases as Maximal Linearly Independent Sets

Let $S = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n}$ be a basis for the vector space *V*. Then any set of more than *n* vectors in *V* is linearly dependent.

Proof: We need to show that if m > n, then any set $T = {\mathbf{w}_1, \mathbf{w}_2, ..., \mathbf{w}_m}$ of *m* vectors in *V* is linearly dependent. Because the basis *S* spans *V*, each vector \mathbf{w}_j in *T* can be expressed as a linear combination of $\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_n$:

$$\mathbf{w}_{1} = a_{11}\mathbf{v}_{1} + a_{21}\mathbf{v}_{2} + \dots + a_{n1}\mathbf{v}_{n}$$

$$\mathbf{w}_{2} = a_{12}\mathbf{v}_{1} + a_{22}\mathbf{v}_{2} + \dots + a_{n2}\mathbf{v}_{n}$$

$$\vdots$$

$$\mathbf{w}_{m} = a_{1m}\mathbf{v}_{1} + a_{2m}\mathbf{v}_{2} + \dots + a_{nm}\mathbf{v}_{n}.$$
(3)

Now we need to find scalars c_1, c_2, \ldots, c_m not all zero such that

$$c_1 \mathbf{w}_1 + c_2 \mathbf{w}_2 + \dots + c_m \mathbf{w}_m = \mathbf{0}. \tag{4}$$

4.4 BASES AND DIMENSION FOR VECTOR SPACES 7

Substituting in (4) the expressions in (3) for each \mathbf{w}_i , we get

$$c_{1}(a_{11}\mathbf{v}_{1} + a_{21}\mathbf{v}_{2} + \dots + a_{n1}\mathbf{v}_{n}) + c_{2}(a_{12}\mathbf{v}_{1} + a_{22}\mathbf{v}_{2} + \dots + a_{n2}\mathbf{v}_{n}) + \dots$$
(5)
+ $c_{m}(a_{1m}\mathbf{v}_{1} + a_{2m}\mathbf{v}_{2} + \dots + a_{nm}\mathbf{v}_{n}) = \mathbf{0}.$

Because the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent, the coefficient in (5) of each \mathbf{v}_i must vanish:

$$a_{11}c_{1} + a_{12}c_{2} + \dots + a_{1m}c_{m} = 0$$

$$a_{21}c_{1} + a_{22}c_{2} + \dots + a_{2m}c_{m} = 0$$

$$\vdots$$

$$a_{n1}c_{1} + a_{n2}c_{2} + \dots + a_{nm}c_{m} = 0.$$
(6)

But because m > n, this is a homogeneous linear system with more unknowns than equations, and therefore has a nontrivial solution (c_1, c_2, \ldots, c_m) . Then c_1, c_2, \ldots, c_m are scalars not all zero such that (4) holds, so we have shown that the set T of more than n vectors is linearly dependent.

Now let $S = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n}$ and $T = {\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_m}$ be two different bases for the same vector space V. Because S is a basis and T is linearly independent, Theorem 1 implies that $m \le n$. Next, reverse the roles of S and T; the fact that T is a basis and S is linearly independent implies similarly that $n \le m$. Hence m = n, so we have proved the following theorem for any vector space with a finite basis.

THEOREM 2 The Dimension of a Vector Space

Any two bases for a vector space consist of the same number of vectors.

A nonzero vector space V is called **finite dimensional** provided that there exists a basis for V consisting of a finite number of vectors from V. In this case the number n of vectors in each basis for V is called the **dimension** of V, denoted by $n = \dim V$. Then V is an *n*-dimensional vector space. The standard basis of Example 1 shows that \mathbf{R}^n is, indeed, an *n*-dimensional vector space.

Note that the zero vector space $\{0\}$ has no basis because it contains *no* linearly independent set of vectors. (Sometimes it is convenient to adopt the convention that the null set is a basis for $\{0\}$.) Here we define dim $\{0\}$ to be zero. A nonzero vector space that has no finite basis is called **infinite dimensional**. Infinite dimensional vector spaces are discussed in Section 4.5, but we include an illustrative example of one here.

Example 4 Let \mathcal{P} be the set of all polynomials of the form

$$p(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n,$$

where the largest exponent $n \ge 0$ that appears is the *degree* of the polynomial p(x), and the coefficients $a_0, a_1, a_2, \ldots, a_n$ are real numbers. We add polynomials in \mathcal{P} and multiply them by scalars in the usual way—that is, by collecting coefficients of like powers of x. For instance, if

$$p(x) = 3 + 2x + 5x^3$$
 and $q(x) = 7 + 4x + 3x^2 + 9x^4$

then

$$(p+q)(x) = (3+7) + (2+4)x + (0+3)x^2 + (5+0)x^3 + (0+9)x^4$$
$$= 10 + 6x + 3x^2 + 5x^3 + 9x^4$$
and

$$(7p)(x) = 7(3 + 2x + 5x^3) = 21 + 14x + 35x^3$$

It is readily verified that, with these operations, \mathcal{P} is a vector space. But \mathcal{P} has no finite basis. For if p_1, p_2, \ldots, p_n are elements of \mathcal{P} , then the degree of any linear combination of them is at most the maximum of *their* degrees. Hence no polynomial in \mathcal{P} of higher degree lies in span{ p_1, p_2, \ldots, p_n }. Thus no finite subset of \mathcal{P} spans \mathcal{P} , and therefore \mathcal{P} is an infinite dimensional vector space.

Here, our concern is with finite-dimensional vector spaces, and we note first that any proper subspace W of a finite-dimensional vector space V is itself finite dimensional, with dim $W < \dim V$. For if dim V = n, let $k \le n$ be the largest integer such that W contains k linearly independent vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k$. Then, by the same argument as in Example 2, we see that $\{\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k\}$ is a basis for W, so W is finite dimensional with dim W = k, and k < n because W is a proper subspace.

Moreover, an *n*-dimensional vector space V contains proper subspaces of each dimension k = 1, 2, ..., n - 1. For instance, if $\{\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_n\}$ is a basis for V and k < n, then $W = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_k\}$ is a subspace of dimension k. Thus \mathbf{R}^4 contains proper subspaces of dimensions 1, 2, and 3; \mathbf{R}^5 contains proper subspaces of dimensions 1, 2, 3, and 4; and so on. The proper subspaces of \mathbf{R}^n of dimensions 1, 2, ..., n - 1 are the higher-dimensional analogues of lines and planes through the origin in \mathbf{R}^3 .

Suppose that V is an n-dimensional vector space and that $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a set of n vectors in V. Then, in order to show that S is a basis for V, it is not necessary to prove *both* that S is linearly independent *and* that S spans V, because it turns out (for n vectors in an n-dimensional vector space) that either property of S implies the other. For instance, if $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a set of n linearly independent vectors in V, then Theorem 1 and the argument of Example 2 (once again) imply that S spans V and, hence, is a basis for V. This proves part (a) of Theorem 3. The remaining parts are left to the problems. (See Problems 27–32.)

THEOREM 3 Independent Sets, Spanning Sets, and Bases

- Let V be an n-dimensional vector space and let S be a subset of V. Then
- (a) If S is linearly independent and consists of n vectors, then S is a basis for V;
- (b) If S spans V and consists of n vectors, then S is a basis for V;
- (c) If S is linearly independent, then S is contained in a basis for V;
- (d) If S spans V, then S contains a basis for V.

Part (c) of Theorem 3 is often applied in the following form: If W is a kdimensional subspace of the *n*-dimensional vector space V, then any basis $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ for W can be "extended" to a basis $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ for V, consisting of the original basis vectors for W together with n - k additional vectors $\mathbf{v}_{k+1}, \mathbf{v}_{k+2}, \dots, \mathbf{v}_n$.

Bases for Solution Spaces

We consider now the homogeneous linear system

$$\mathbf{A}\mathbf{x} = \mathbf{0},\tag{7}$$

in which **A** is an $m \times n$ matrix, so the system consists of *m* equations in the *n* variables x_1, x_2, \ldots, x_n . Its solution space *W* is then a subspace of **R**^{*n*}. We want to determine the dimension of *W* and, moreover, to find an explicit basis for *W*. Thus we seek a maximal set of linearly independent solution vectors of (7).

4.4 BASES AND DIMENSION FOR VECTOR SPACES 9

Recall the Gaussian elimination method of Section 3.2. We use elementary row operations to reduce the coefficient matrix **A** to an echelon matrix **E** and note the (nonzero) *leading entries* in the rows of **E**. The *leading variables* are those that correspond to the columns of **E** containing the leading entries. The remaining variables (if any) are the *free variables*. If there are no free variables, then our system has only the trivial solution, so $W = \{0\}$. If there are free variables, we set each of them (separately) equal to a parameter and solve (by back substitution) for the leading variables as linear combinations of these parameters. The solution space W is then the set of all solution vectors obtained in this manner (for all possible values of the parameters).

To illustrate the general situation, let us suppose that the leading variables are the first *r* variables $x_1, x_2, ..., x_r$, so the k = n - r variables $x_{r+1}, x_{r+2}, ..., x_n$ are free variables. The reduced system **Ex** = **0** then takes the form

$$b_{11}x_{1} + b_{12}x_{2} + \dots + b_{1r}x_{r} + \dots + b_{1n}x_{n} = 0$$

$$b_{22}x_{2} + \dots + b_{2r}x_{r} + \dots + b_{2n}x_{n} = 0$$

$$\vdots$$

$$b_{rr}x_{r} + \dots + b_{rn}x_{n} = 0$$

$$0 = 0$$

$$\vdots$$

$$0 = 0$$

with the last m - r equations being "trivial." We set

 $x_{r+1} = t_1, \quad x_{r+2} = t_2, \dots, \quad x_n = t_k$ (9)

and then solve (by back substitution) the equations in (8) for the leading variables

$$x_{1} = c_{11}t_{1} + c_{12}t_{2} + \dots + c_{1k}t_{k}$$

$$x_{2} = c_{21}t_{1} + c_{22}t_{2} + \dots + c_{2k}t_{k}$$

$$\vdots$$

$$x_{r} = c_{r1}t_{1} + c_{r2}t_{2} + \dots + c_{rk}t_{k}.$$
(10)

The typical solution vector $(x_1, x_2, ..., x_n)$ is given in terms of the k parameters $t_1, t_2, ..., t_k$ by the equations in (9) and (10).

We now choose k particular solution vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ as follows: To get \mathbf{v}_j we set the *j* th parameter t_j equal to 1 and set all other parameters equal to zero. Then

$$\mathbf{w}_j = (c_{1j}, c_{2j}, \dots, c_{rj}, 0, \dots, 1, \dots, 0),$$
 (11)

with the 1 appearing as the (r + j)th entry. The vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are the column vectors of the $n \times k$ matrix

$$\begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1k} \\ c_{21} & c_{22} & \cdots & c_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ c_{r1} & c_{r2} & \cdots & c_{rk} \\ 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}.$$
(12)

Because of the presence of the lower $k \times k$ identity matrix, it is clear that the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent. (See Problem 36.) But Eqs. (9) and (10) show that the typical solution vector \mathbf{x} is a linear combination

$$\mathbf{x} = t_1 \mathbf{v}_1 + t_2 \mathbf{v}_2 + \dots + t_k \mathbf{v}_k. \tag{13}$$

Therefore, the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ defined in (11) form a basis for the solution space *W* of the original system in (7).

The following algorithm summarizes the steps in this procedure.

ALGORITHM A Basis for the Solution Space

To find a basis for the solution space W of the homogeneous linear system Ax = 0, carry out the following steps.

- 1. Reduce the coefficient matrix A to echelon form.
- 2. Identify the *r* leading variables and the k = n r free variables. If k = 0, then $W = \{0\}$.
- **3.** Set the free variables equal to parameters t_1, t_2, \ldots, t_k , and then solve by back substitution for the leading variables in terms of these parameters.
- **4.** Let \mathbf{v}_j be the solution vector obtained by setting t_j equal to 1 and the other parameters equal to zero. Then $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a basis for *W*.

Example 5 Find a basis for the solution space of the homogeneous linear system

$$3x_1 + 6x_2 - x_3 - 5x_4 + 5x_5 = 0$$

$$2x_1 + 4x_2 - x_3 - 3x_4 + 2x_5 = 0$$

$$3x_1 + 6x_2 - 2x_3 - 4x_4 + x_5 = 0.$$
(14)

Solution We readily reduce the coefficient matrix **A** to the echelon form

	1	2	0	-2	3]	
$\mathbf{E} =$	0	0	1	-1	4	
	0	0	0	0	0	

The leading entries are in the first and third columns, so the leading variables are x_1 and x_3 ; the free variables are x_2 , x_4 , and x_5 . To avoid subscripts, we use r, s, and t rather than t_1 , t_2 , and t_3 to denote the three parameters. Thus we set

$$x_2 = r, \quad x_4 = s, \quad \text{and} \quad x_5 = t.$$
 (15)

Then back substitution in the reduced system

$$x_1 + 2x_2 - 2x_4 + 3x_5 = 0$$

$$x_3 - x_4 + 4x_5 = 0$$

yields

$$x_1 = -2r + 2s - 3t$$
 and $x_3 = s - 4t$. (16)

The equations in (15) and (16) give the typical solution vector $(x_1, x_2, x_3, x_4, x_5)$ in terms of the parameters r, s, and t.

With r = 1 and s = t = 0,

we obtain
$$\mathbf{v}_1 = (-2, 1, 0, 0, 0).$$

With s = 1 and r = t = 0,

we obtain $\mathbf{v}_2 = (2, 0, 1, 1, 0).$

With t = 1 and r = s = 0,

we obtain $\mathbf{v}_3 = (-3, 0, -4, 0, 1)$.

Thus the solution space of the system in (14) is a 3-dimensional subspace of \mathbb{R}^5 with basis $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

4.4 **Problems**

In Problems 1–8, determine whether or not the given vectors in \mathbf{R}^n form a basis for \mathbf{R}^n .

- **1.** $\mathbf{v}_1 = (4, 7), \mathbf{v}_2 = (5, 6)$
- **2.** $\mathbf{v}_1 = (3, -1, 2), \mathbf{v}_2 = (6, -2, 4), \mathbf{v}_3 = (5, 3, -1)$
- **3.** $\mathbf{v}_1 = (1, 7, -3), \mathbf{v}_2 = (2, 1, 4), \mathbf{v}_3 = (6, 5, 1),$
- $\mathbf{v}_4 = (0, 7, 13)$
- **4.** $\mathbf{v}_1 = (3, -7, 5, 2), \mathbf{v}_2 = (1, -1, 3, 4), \mathbf{v}_3 = (7, 11, 3, 13)$
- **5.** $\mathbf{v}_1 = (0, 7, -3), \, \mathbf{v}_2 = (0, 5, 4), \, \mathbf{v}_3 = (0, 5, 10)$
- **6.** $\mathbf{v}_1 = (0, 0, 1), \mathbf{v}_2 = (0, 1, 2), \mathbf{v}_3 = (1, 2, 3)$
- 7. $\mathbf{v}_1 = (0, 0, 1), \mathbf{v}_2 = (7, 4, 11), \mathbf{v}_3 = (5, 3, 13)$
- 8. $\mathbf{v}_1 = (2, 0, 0, 0), \mathbf{v}_2 = (0, 3, 0, 0), \mathbf{v}_3 = (0, 0, 7, 6),$ $\mathbf{v}_4 = (0, 0, 4, 5)$

In Problems 9–11, find a basis for the indicated subspace of \mathbf{R}^3 .

- 9. The plane with equation x 2y + 5z = 0.
- **10.** The plane with equation y = z.
- **11.** The line of intersection of the planes described in Problems 9 and 10.

In Problems 12–14, find a basis for the indicated subspace of \mathbf{R}^4 .

- 12. The set of all vectors of the form (a, b, c, d) for which a = b + c + d.
- **13.** The set of all vectors of the form (a, b, c, d) such that a = 3c and b = 4d.
- 14. The set of all vectors of the form (a, b, c, d) for which a + 2b = c + 3d = 0.

In Problems 15–26, find a basis for the solution space of the given homogeneous linear system.

- **15.** $x_1 2x_2 + 3x_3 = 0$ $2x_1 - 3x_2 - x_3 = 0$ **16.** $x_1 + 3x_2 + 4x_3 = 0$
- $3x_1 + 8x_2 + 7x_3 = 0$
- **17.** $x_1 3x_2 + 2x_3 4x_4 = 0$ $2x_1 - 5x_2 + 7x_3 - 3x_4 = 0$
- **18.** $x_1 + 3x_2 + 4x_3 + 5x_4 = 0$ $2x_1 + 6x_2 + 9x_3 + 5x_4 = 0$
- **19.** $x_1 3x_2 9x_3 5x_4 = 0$ $2x_1 + x_2 - 4x_3 + 11x_4 = 0$ $x_1 + 3x_2 + 3x_3 + 13x_4 = 0$
- **20.** $x_1 3x_2 10x_3 + 5x_4 = 0$ $x_1 + 4x_2 + 11x_3 - 2x_4 = 0$ $x_1 + 3x_2 + 8x_3 - x_4 = 0$

21. $x_1 - 4x_2 - 3x_3 - 7x_4 = 0$ $2x_1 - x_2 + x_3 + 7x_4 = 0$ $x_1 + 2x_2 + 3x_3 + 11x_4 = 0$ **22.** $x_1 - 2x_2 - 3x_3 - 16x_4 = 0$ $2x_1 - 4x_2 + x_3 + 17x_4 = 0$ $x_1 - 2x_2 + 3x_3 + 26x_4 = 0$ **23.** $x_1 + 5x_2 + 13x_3 + 14x_4 = 0$ $2x_1 + 5x_2 + 11x_3 + 12x_4 = 0$ $2x_1 + 7x_2 + 17x_3 + 19x_4 = 0$ **24.** $x_1 + 3x_2 - 4x_3 - 8x_4 + 6x_5 = 0$ $x_1 + 2x_3 + x_4 + 3x_5 = 0$ $2x_1 + 7x_2 - 10x_3 - 19x_4 + 13x_5 = 0$ **25.** $x_1 + 2x_2 + 7x_3 - 9x_4 + 31x_5 = 0$ $2x_1 + 4x_2 + 7x_3 - 11x_4 + 34x_5 = 0$ $3x_1 + 6x_2 + 5x_3 - 11x_4 + 29x_5 = 0$

26. $3x_1 + x_2 - 3x_3 + 11x_4 + 10x_5 = 0$ $5x_1 + 8x_2 + 2x_3 - 2x_4 + 7x_5 = 0$ $2x_1 + 5x_2 - x_4 + 14x_5 = 0$

Problems 27 through 36 further explore independent sets, spanning sets, and bases.

- **27.** Suppose that *S* is a set of *n* linearly independent vectors in the *n*-dimensional vector space *V*. Prove that *S* is a basis for *V*.
- **28.** Suppose that S is a set of n vectors that span the n-dimensional vector space V. Prove that S is a basis for V.
- 29. Let {v₁, v₂,..., v_k} be a basis for the proper subspace W of the vector space V, and suppose that the vector v of V is not in W. Show that the vectors v₁, v₂,..., v_k, v are linearly independent.
- **30.** Use the result of Problem 29 to prove that every linearly independent set of vectors in a finite-dimensional vector space *V* is contained in a basis for *V*.
- **31.** Suppose that the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k, \mathbf{v}_{k+1}$ span the vector space *V* and that \mathbf{v}_{k+1} is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$. Show that the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ span *V*.
- **32.** Use the result of Problem 31 to prove that every spanning set for a finite-dimensional vector space *V* contains a basis for *V*.
- **33.** Let *S* be a linearly independent set of vectors in the finitedimensional vector space *V*. Then *S* is called a **maximal linearly independent set** provided that if any other vector is adjoined to *S*, then the resulting set is linearly dependent. Prove that every maximal linearly independent set in *V* is a basis for *V*.

- **34.** Let *S* be a finite set of vectors that span the vector space *V*. Then *S* is called a **minimal spanning set** provided that no proper subset of *S* spans *V*. Prove that every minimal spanning set in *V* is a basis for *V*.
- **35.** Let *S* be a finite set of vectors that span the vector space *V*. Then *S* is called a **uniquely spanning set** provided that

each vector in V can be expressed in one and only one way as a linear combination of the vectors in S. Prove that every uniquely spanning set in V is a basis for V.

36. Apply the definition of linear independence to show directly that the column vectors of the matrix in (12) are linearly independent.

4.5 Row and Column Spaces

In numerous examples we have observed the phenomenon of "disappearing equations" that sometimes occurs when we solve a linear system using the method of Gaussian elimination. The appearance in this process of a trivial equation 0 = 0means that one of the original equations was redundant. For instance, in the system

$$x - 2y + 2z = 0$$

$$x + 4y + 3z = 0$$

$$2x + 2y + 5z = 0$$

the third equation provides no additional information about a solution (x, y, z) because it is merely the sum of the first two equations.

Given a homogeneous linear system, it is natural to ask how many of the equations are "irredundant," and which ones they are. We will see that an answer to this question leads to a natural and simple relation between the number of irredundant equations, the number of unknowns, and the number of linearly independent solutions.

Row Space and Row Rank

The individual equations of the homogeneous linear system Ax = 0 correspond to the "row matrices"

 $\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \end{bmatrix} \\ \begin{bmatrix} a_{21} & a_{22} & \cdots & a_{2n} \end{bmatrix} \\ \vdots \\ \begin{bmatrix} a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$

of the $m \times n$ matrix $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$. The row vectors of \mathbf{A} are the *m* vectors

$$\mathbf{r}_{1} = (a_{11}, a_{12}, \dots, a_{1n})$$

$$\mathbf{r}_{2} = (a_{21}, a_{22}, \dots, a_{2n})$$

$$\vdots$$

$$\mathbf{r}_{m} = (a_{m1}, a_{m2}, \dots, a_{mn})$$
(1)

in \mathbb{R}^n . Recalling from Section 3.4 the convection that *n*-tuples denote column-vector elements of \mathbb{R}^n , we see that the row vectors of \mathbf{A} are the transposes of its row matrices; that is,

$$\mathbf{r}_{i} = \begin{bmatrix} a_{i1} \\ a_{i2} \\ \vdots \\ a_{in} \end{bmatrix} = \begin{bmatrix} a_{i1} & a_{i2} & \cdots & a_{in} \end{bmatrix}^{T}$$

for i = 1, 2, ..., m, and thus actually are column vectors (despite being called "row vectors"). The subspace of \mathbf{R}^n spanned by the *m* row vectors $\mathbf{r}_1, \mathbf{r}_2, ..., \mathbf{r}_m$ is called the **row space** Row(A) of the matrix A. The *dimension* of the row space Row(A) is called the **row rank** of the matrix A.

Example 1 Consider the 4×5 echelon matrix

 $\mathbf{A} = \begin{bmatrix} 1 & -3 & 2 & 5 & 3 \\ 0 & 0 & 1 & -4 & 2 \\ 0 & 0 & 0 & 1 & 7 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$

Its row vectors are $\mathbf{r}_1 = (1, -3, 2, 5, 3)$, $\mathbf{r}_2 = (0, 0, 1, -4, 2)$, $\mathbf{r}_3 = (0, 0, 0, 1, 7)$, and $\mathbf{r}_4 = (0, 0, 0, 0, 0, 0)$. We want to show that the row vectors that are not zero are linearly independent. To this end we calculate the linear combination

$$c_1\mathbf{r}_1 + c_2\mathbf{r}_2 + c_3\mathbf{r}_3 = (c_1, -3c_1, 2c_1 + c_2, 5c_1 - 4c_2 + c_3, 3c_1 + 2c_2 + 7c_3).$$

If $c_1\mathbf{r}_1 + c_2\mathbf{r}_2 + c_3\mathbf{r}_3 = \mathbf{0}$, then the first component gives $c_1 = 0$, next the third component $2c_1 + c_2 = 0$ yields $c_2 = 0$, and finally the fourth component $5c_1 - 4c_2 + c_3 = 0$ yields $c_3 = 0$. Thus the vectors \mathbf{r}_1 , \mathbf{r}_2 , and \mathbf{r}_3 are linearly independent and therefore form a basis for the row space Row(A). Hence Row(A) is a 3-dimensional subspace of \mathbf{R}^5 , and the row rank of A is 3.

It should be apparent that the method used in Example 1 applies to any *echelon* matrix $\mathbf{E} = [e_{ij}]$. If the first column of \mathbf{E} is not all zero, then its *nonzero* row vectors $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_k$ have the forms

$$\mathbf{r}_1 = (e_{11}, \dots, e_{1p}, \dots, e_{1q}, \dots),$$

$$\mathbf{r}_2 = (0, \dots, 0, e_{2p}, \dots, e_{2q}, \dots),$$

$$\mathbf{r}_3 = (0, \dots, 0, \dots, e_{3q}, \dots),$$

and so forth (where e_{11} , e_{2p} , and e_{3q} denote nonzero leading entries). Then the equation

$$c_1\mathbf{r}_1 + c_2\mathbf{r}_2 + \dots + c_k\mathbf{r}_k = \mathbf{0}$$

implies that

$$c_1e_{11} = 0$$
, $c_1e_{1p} + c_2e_{2p} = 0$, $c_1e_{1q} + c_2e_{2q} + c_3e_{3q} = 0$,

and so forth. We can therefore conclude in turn that $c_1 = 0, c_2 = 0, ..., c_k = 0$. Thus the row vectors $\mathbf{r}_1, \mathbf{r}_2, ..., \mathbf{r}_k$ are linearly independent, and hence we have the following result.

THEOREM 1 Row Space of an Echelon Matrix

The nonzero row vectors of an echelon matrix are linearly independent and therefore form a basis for its row space.

We investigate the row space of an arbitrary matrix **A** by reducing **A** to an echelon matrix **E**. The following theorem then guarantees that **A** and **E** have the same row space. Recall that two matrices are (row) *equivalent* provided that each can be transformed to the other by elementary row operations.

THEOREM 2 Row Spaces of Equivalent Matrices

If two matrices **A** and **B** are equivalent, then they have the same row space.

Proof: Because A can be transformed to B by elementary row operations, it follows that each row vector of B is a linear combination of the row vectors of A; the fact that this is obviously true if A is transformed into B by a single row operation implies its truth for any (finite) number of row operations. But then each vector in Row(B) is a linear combination of linear combinations of row vectors of A and hence is a linear combination of row vectors of A, which means that it is also a vector in Row(A). Therefore, the fact that A can be transformed to B implies that Row(A) contains Row(B).

But elementary row operations are "reversible," so **B** can also be transformed to **A** by row operations. Hence the same argument shows that Row(B) contains Row(A). Finally, the fact that each of the two row spaces Row(A) and Row(B) contains the other means that they are, indeed, identical.

Theorems 1 and 2 together provide a routine procedure for finding a basis for the row space of a given matrix, thereby determining its row rank.

ALGORITHM 1 A Basis for the Row Space

To find a basis for the row space of a matrix \mathbf{A} , use elementary row operations to reduce \mathbf{A} to an echelon matrix \mathbf{E} . Then the nonzero row vectors of \mathbf{E} form a basis for Row(\mathbf{A}).

Example 2	ple 2 To find a basis for the row space of the matrix						
	$\mathbf{A} =$	$\begin{bmatrix} 1\\ 3\\ 2\\ 2 \end{bmatrix}$	2 4 3 2	1 9 5 8	$3 \\ 0 \\ 1 \\ -3$	2 7 8 5	, (2)

we reduce it to its echelon form

$$\mathbf{E} = \begin{bmatrix} 1 & 2 & 1 & 3 & 2 \\ 0 & 1 & -3 & 5 & -4 \\ 0 & 0 & 0 & 1 & -7 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$
 (3)

Then the nonzero row vectors $\mathbf{v}_1 = (1, 2, 1, 3, 2)$, $\mathbf{v}_2 = (0, 1, -3, 5, -4)$, and $\mathbf{v}_3 = (0, 0, 0, 1, -7)$ form a basis for the row space of **A**. Thus Row(**A**) is a 3-dimensional subspace of \mathbf{R}^5 and the row rank of **A** is 3.

In Example 2, note that Algorithm 1 does *not* tell us that the first three row vectors of the matrix \mathbf{A} itself are linearly independent. We know that *some* three row vectors of \mathbf{A} span Row(\mathbf{A}), but without further investigation we do not know *which* three.

Column Space and Column Rank

Now we turn our attention from row vectors to column vectors. Given an $m \times n$ matrix $\mathbf{A} = [a_{ij}]$, the **column vectors** of \mathbf{A} are the *n* vectors

$$\mathbf{c}_{1} = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, \quad \mathbf{c}_{2} = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}, \quad \dots, \quad \mathbf{c}_{n} = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$
(4)

in \mathbb{R}^m . The subspace of \mathbb{R}^m spanned by the *n* column vectors $\mathbf{c}_1, \mathbf{c}_2, \ldots, \mathbf{c}_n$ is called the **column space** Col(A) of the matrix A. The *dimension* of the space Col(A) is called the **column rank** of the matrix A.

Example 3 Consider the 4×4

Consider the 4×5 echelon matrix

$$\mathbf{E} = \begin{bmatrix} 1 & 2 & 1 & 3 & 2 \\ 0 & 1 & -3 & 5 & -4 \\ 0 & 0 & 0 & 1 & -7 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$
 (5)

Its five column vectors \mathbf{c}_1 , \mathbf{c}_2 , \mathbf{c}_3 , \mathbf{c}_4 , \mathbf{c}_5 all lie in the subspace $\mathbf{R}^3 = \{(x_1, x_2, x_3, 0)\}$ of \mathbf{R}^4 . The column vectors that contain the leading entries in the nonzero rows of \mathbf{E} are

$$\mathbf{c}_1 = \begin{bmatrix} 1\\0\\0\\0 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} 2\\1\\0\\0 \end{bmatrix}, \quad \text{and} \quad \mathbf{c}_4 = \begin{bmatrix} 3\\5\\1\\0 \end{bmatrix}. \tag{6}$$

Note that

$$a_1\mathbf{c}_1 + a_2\mathbf{c}_2 + a_4\mathbf{c}_4 = (a_1 + 2a_2 + 3a_4, a_2 + 5a_4, a_4, 0)$$

Hence $a_1\mathbf{c}_1 + a_2\mathbf{c}_2 + a_4\mathbf{c}_4 = \mathbf{0}$ readily implies that $a_1 = a_2 = a_4 = 0$. Thus the vectors $\mathbf{c}_1, \mathbf{c}_2$, and \mathbf{c}_4 are linearly independent and therefore form a basis for $\mathbf{R}^3 = \{(x_1, x_2, x_3, 0)\}$. Hence Col(E) is a 3-dimensional subspace of \mathbf{R}^4 and the column rank of E is 3.

To find the column rank of an arbitrary $m \times n$ matrix **A**, we begin in the same way as when finding the row rank. First we use elementary row operations to reduce **A** to an echelon matrix **E**. But the relationship between the column spaces of **A** and **E** is far more subtle than between their row spaces; the reason is that elementary row operations generally do *not* preserve column spaces. (See Problem 36.)

To analyze this situation, let us denote by $\mathbf{c}_1, \mathbf{c}_2, \ldots, \mathbf{c}_n$ the column vectors of the original matrix **A**, and by $\mathbf{c}_1^*, \mathbf{c}_2^*, \ldots, \mathbf{c}_n^*$ the column vectors of the echelon matrix to which we have reduced **A**. Suppose that **E** has *k* nonzero rows. Then each column vector of **E** has the form $\mathbf{c}_j^* = (*, \ldots, *, 0, \ldots, 0)$ with m - k final zeros. Hence the column space Col(**E**) is contained in \mathbf{R}^k (considered as the subspace of \mathbf{R}^m for which $x_{k+1} = \cdots = x_m = 0$).

We are particularly interested in those columns of \mathbf{E} that contain the nonzero leading entries in the nonzero rows of \mathbf{E} . These *k* columns are called the **pivot** columns of \mathbf{E} . The echelon matrix \mathbf{E} has the general form

$$\mathbf{E} = \begin{bmatrix} d_1 & * & * & * & \cdots & * & * & * \\ 0 & d_2 & * & * & \cdots & * & * & * \\ 0 & 0 & 0 & d_3 & \cdots & * & * & * \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & d_k & * & * \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix},$$
(7)

where d_1, d_2, \ldots, d_k are the nonzero leading entries. Hence the k pivot column

vectors of **E** look like

Because of the upper triangular pattern visible here, it should be apparent that the argument in Example 3 can be used to show that these k pivot vectors form a basis for Col(**E**). Thus the column rank of the echelon matrix **E** is equal to the number k of its nonzero rows, and hence is equal to its row rank.

Now comes the subtle part. We want to show that the column rank of the original matrix A also is k. Note first that the homogeneous linear systems

$$\mathbf{A}\mathbf{x} = \mathbf{0} \quad \text{and} \quad \mathbf{E}\mathbf{x} = \mathbf{0} \tag{9}$$

have the same solution set because the matrices **A** and **E** are equivalent. If $\mathbf{x} = (x_1, x_2, ..., x_n)$, then the left-hand sides in (9) are linear combinations of column vectors:

$$\mathbf{A}\mathbf{x} = x_1\mathbf{c}_1 + x_2\mathbf{c}_2 + \dots + x_n\mathbf{c}_n,$$

$$\mathbf{E}\mathbf{x} = x_1\mathbf{c}_1^* + x_2\mathbf{c}_2^* + \dots + x_n\mathbf{c}_n^*.$$
(10)

Because \mathbf{x} satisfies *either* both *or* neither of the systems in (9), it follows that

$$x_1\mathbf{c}_1 + x_2\mathbf{c}_2 + \dots + x_n\mathbf{c}_n = \mathbf{0}$$

if and only if

$$x_1\mathbf{c}_1^* + x_2\mathbf{c}_2^* + \dots + x_n\mathbf{c}_n^* = \mathbf{0}.$$

Hence every linear dependence between column vectors of \mathbf{E} is mirrored in a linear dependence with the same coefficients between the corresponding column vectors of \mathbf{A} . In particular, because the k pivot column vectors of \mathbf{E} are linearly independent, it follows that the corresponding k column vectors of \mathbf{A} are linearly independent. And because the k pivot column vectors of \mathbf{E} span Col(\mathbf{E}), it follows that the corresponding k column vectors of \mathbf{E} span Col(\mathbf{E}), it follows that the corresponding k column vectors of \mathbf{A} span Col(\mathbf{A}). Thus the latter k vectors form a basis for the column space of \mathbf{A} , and the column rank of \mathbf{A} is k. Consequently we have established the following method for finding a basis for the column space of a given matrix.

ALGORITHM 2 A Basis for the Column Space

To find a basis for the column space of a matrix \mathbf{A} , use elementary row operations to reduce \mathbf{A} to an echelon matrix \mathbf{E} . Then the column vectors of \mathbf{A} that correspond to the pivot columns of \mathbf{E} form a basis for Col(\mathbf{A}).

Remark Since the pivot column vectors of the matrix **A**—that is, the column vectors corresponding to the pivot column vectors of the echelon matrix **E**—form a basis for the column space of **A**, it follows that every non-pivot column vector of **A** is a linear combination of its pivot column vectors.

Example 4 Consider again the 4×5 matrix

3 4 9 0 7	
$\mathbf{A} = \begin{bmatrix} 2 & 3 & 5 & 1 & 8 \end{bmatrix}$	
2 2 8 -3 5	

of Example 2, in which we reduced it to the echelon matrix

$$\mathbf{E} = \begin{bmatrix} 1 & 2 & 1 & 3 & 2 \\ 0 & 1 & -3 & 5 & -4 \\ 0 & 0 & 0 & 1 & -7 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The pivot columns of **E** are its first, second, and fourth columns, so the 3-dimensional column space of **A** has a basis consisting of its first, second, and fourth column vectors $\mathbf{c}_1 = (1, 3, 2, 2), \mathbf{c}_2 = (2, 4, 3, 2), \text{ and } \mathbf{c}_4 = (3, 0, 1, -3).$

The fact that Algorithm 2 provides a basis for Col(A) consisting of column vectors of A itself (rather than the echelon matrix E) enables us to apply it to the problem of extracting a maximal linearly independent subset from a given set of vectors in \mathbb{R}^n .

Example 5 Find a subset of the vectors $\mathbf{v}_1 = (1, -1, 2, 2)$, $\mathbf{v}_2 = (-3, 4, 1, -2)$, $\mathbf{v}_3 = (0, 1, 7, 4)$, and $\mathbf{v}_4 = (-5, 7, 4, -2)$ that forms a basis for the subspace *W* of \mathbf{R}^4 spanned by these four vectors. Solution It is not clear at the outset whether *W* is 2-dimensional, 3-dimensional, or 4-dimensional. To apply Algorithm 2, we arrange the given vectors as the *column* vectors of the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & -3 & 0 & -5 \\ -1 & 4 & 1 & 7 \\ 2 & 1 & 7 & 4 \\ 2 & -2 & 4 & -2 \end{bmatrix},$$

which reduces readily to the echelon matrix

	[1]	-3	0	-5	
E =	0	1	1	2	
	0	0	0	0	ŀ
	0	0	0	0	

The pivot columns of **E** are its first two columns, so the first two column vectors $\mathbf{v}_1 = (1, -1, 2, 2)$ and $\mathbf{v}_2 = (-3, 4, 1, -2)$ of **A** form a basis for the column space *W*. In particular, we see that *W* is 2-dimensional. (You can confirm that $\mathbf{v}_3 = 3\mathbf{v}_1 + \mathbf{v}_2$ and $\mathbf{v}_4 = \mathbf{v}_1 + 2\mathbf{v}_2$.)

Rank and Dimension

We have seen that the row rank *and* the column rank of an echelon matrix **E** *both* equal the number of rows of **E** that are not all zeros. But if **A** is any matrix that reduces by row operations to the echelon matrix **E**, then—according to Algorithm 1—the row rank of **A** is equal to the row rank of **E**, while Algorithm 2 implies that the column rank of **A** is equal to the column rank of **E**. We therefore have the following fundamental result.

THEOREM 3 Equality of Row Rank and Column Rank

The row rank and column rank of any matrix are equal.

For instance, if **A** is a 5×7 matrix and Row(**A**) is a 3-dimensional subspace of **R**⁷, then Theorem 3 tells us that Col(**A**) must be a 3-dimensional subspace of **R**⁵.

To see what an extraordinary theorem this is, think of a random 11×17 matrix **A**, possibly printed by a computer that uses a random number generator to produce independently the 187 entries in **A**. If it turns out that precisely 9 (no more) of the 11 row vectors of **A** are linearly independent, then Theorem 3 implies that precisely 9 (no more) of the 17 column vectors of **A** are linearly independent.

The common value of the row rank and the column rank of the matrix A is simply called the **rank** of A and is denoted by rank(A). The rank of A provides a meaning for the number of irredundant equations in the homogeneous linear system

$$\mathbf{A}\mathbf{x} = \mathbf{0}.\tag{11}$$

Indeed, the individual scalar equations in (11) correspond to the column vectors of the transpose matrix \mathbf{A}^T , whose rank *r* equals that of **A** (Problem 25). Then the *r* columns vectors of \mathbf{A}^T that form a basis for $\operatorname{Col}(\mathbf{A}^T)$ correspond to the *r* equations in (11) that are irredundant; the remaining equations are linear combinations of these irredundant ones and hence are redundant.

The solution space of the homogeneous system Ax = 0 is sometimes called the **null space** of A, denoted by Null(A). Note that if A is an $m \times n$ matrix, then Null(A) and Row(A) are subspaces of \mathbb{R}^n , whereas Col(A) is a subspace of \mathbb{R}^m . If $r = \operatorname{rank}(A)$, then the *r* column vectors of A that form a basis for Col(A) correspond to the *r* leading variables in a solution of Ax = 0 by Gaussian elimination. Moreover, we know from the algorithm in Section 4.4 that the dimension of the solution space Null(A) is equal to the number n - r of free variables. We therefore obtain the important identity

$$\operatorname{rank}(\mathbf{A}) + \operatorname{dim}\operatorname{Null}(\mathbf{A}) = n \tag{12}$$

for any $m \times n$ matrix **A**. For instance, if **A** is the matrix of Example 4 with rank 3 and n = 5, then the dimension of the null space of **A** is 5 - 3 = 2.

For another typical application of Equation (12), consider a homogeneous system of five linear equations in seven unknowns. If the 5×7 coefficient matrix of the system has rank 3, so only 3 of the 5 equations are irredundant, then (because n = 7) the system has 7 - 3 = 4 linearly independent solutions.

If **A** is an $m \times n$ matrix with rank m, then Equation (12) implies that the system $A\mathbf{x} = \mathbf{0}$ has n - m linearly independent solutions. Thus rank(**A**) = m is the condition under which the "conventional wisdom" about the relation between the numbers of equations, unknowns, and solutions is valid.

Nonhomogeneous Linear Systems

It follows immediately from Equation (10) and the definition of Col(A) that *the* nonhomogeneous linear system Ax = b is consistent if and only if the vector **b** is in the column space of **A**. In the problems below we outline some of the applications to nonhomogeneous systems of the results in this section.

If we can find a single particular solution \mathbf{x}_0 of the nonhomogeneous system

$$\mathbf{A}\mathbf{x} = \mathbf{b},\tag{13}$$

then the determination of its solution set reduces to solving the associated homogeneous system

$$\mathbf{A}\mathbf{x} = \mathbf{0}.\tag{14}$$

For then Problem 29 in Section 4.2 implies that the solution set of (13) is the set of all vectors **x** of the form

$$\mathbf{x} = \mathbf{x}_0 + \mathbf{x}_h$$

where \mathbf{x}_h is a solution of (14). Hence if $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_r$ is a basis for the solution space of (14), then the solution set of the nonhomogeneous system is the set of all vectors of the form

$$\mathbf{x} = \mathbf{x}_0 + c_1 \mathbf{x}_1 + c_2 \mathbf{x}_2 + \dots + c_r \mathbf{x}_r.$$

We may describe this solution set as the *translate* by the vector \mathbf{x}_0 of the *r*-dimensional solution space of the corresponding homogeneous system.



In Problems 13–16, a set S of vectors in \mathbb{R}^4 is given. Find a subset of S that forms a basis for the subspace of \mathbb{R}^4 spanned by S.

- **13.** $\mathbf{v}_1 = (1, 3, -2, 4), \mathbf{v}_2 = (2, -1, 3, 2), \mathbf{v}_3 = (5, 1, 4, 8)$
- **14.** $\mathbf{v}_1 = (1, -1, 2, 3), \mathbf{v}_2 = (2, 3, 4, 1), \mathbf{v}_3 = (1, 1, 2, 1), \mathbf{v}_4 = (4, 1, 8, 7)$
- **15.** $\mathbf{v}_1 = (3, 2, 2, 2), \, \mathbf{v}_2 = (2, 1, 2, 1), \, \mathbf{v}_3 = (4, 3, 2, 3), \, \mathbf{v}_4 = (1, 2, 3, 4)$
- **16.** $\mathbf{v}_1 = (5, 4, 2, 2), \, \mathbf{v}_2 = (3, 1, 2, 3), \, \mathbf{v}_3 = (7, 7, 2, 1), \, \mathbf{v}_4 = (1, -1, 2, 4), \, \mathbf{v}_5 = (5, 4, 6, 7)$

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ be a basis for the subspace W of \mathbb{R}^n . Then a basis T for \mathbb{R}^n that contains S can be found by applying the method of Example 5 to the vectors

$$\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_k, \mathbf{e}_1, \mathbf{e}_2, \ldots, \mathbf{e}_n$$

Do this in Problems 17–20.

- 17. Find a basis T for \mathbb{R}^3 that contains the vectors $\mathbf{v}_1 = (1, 2, 2)$ and $\mathbf{v}_2 = (2, 3, 3)$.
- **18.** Find a basis T for \mathbb{R}^3 that contains the vectors $\mathbf{v}_1 = (3, 2, -1)$ and $\mathbf{v}_2 = (2, -2, 1)$.
- **19.** Find a basis T for \mathbf{R}^4 that contains the vectors $\mathbf{v}_1 = (1, 1, 1, 1)$ and $\mathbf{v}_2 = (2, 3, 3, 3)$.
- **20.** Find a basis T for \mathbb{R}^4 that contains the vectors $\mathbf{v}_1 = (3, 2, 3, 3)$ and $\mathbf{v}_2 = (5, 4, 5, 5)$.

Given a homogeneous system Ax = 0 of (scalar) linear equations, we say that a subset of these equations is irredundant provided that the corresponding column vectors of the transpose A^T are linearly independent. In Problems 21–24, extract from each given system a maximal subset of irredundant equations.

- **21.** $x_1 3x_2 + 2x_3 = 0$ $2x_1 + 3x_2 + 2x_3 = 0$ $4x_1 - 3x_2 + 6x_3 = 0$
- **22.** $3x_1 + 2x_2 + 2x_3 + 2x_4 = 0$ $2x_1 + 3x_2 + 3x_3 + 3x_4 = 0$ $8x_1 + 7x_2 + 7x_3 + 7x_4 = 0$
- 23. $x_1 + 2x_2 x_3 + 2x_4 = 0$ $3x_1 - x_2 + 3x_3 + x_4 = 0$ $5x_1 + 3x_2 + x_3 + 5x_4 = 0$ $2x_1 + 5x_3 + 4x_4 = 0$
- **24.** $3x_1 + 2x_2 + 2x_3 = 0$ $2x_1 + 3x_2 + 3x_3 = 0$ $7x_1 + 8x_2 + 8x_3 = 0$ $8x_1 + 7x_2 + 7x_3 = 0$

$$5x_1 + 6x_2 + 5x_3 = 0$$

- **25.** Explain why the rank of a matrix \mathbf{A} is equal to the rank of its transpose \mathbf{A}^T .
- **26.** Explain why the $n \times n$ matrix **A** is invertible if and only if its rank is *n*.

Problems 27 through 32 explore the properties of nonhomogeneous systems.

- 27. Let **A** be a 3×5 matrix whose three row vectors are linearly independent. Prove that, for each **b** in **R**³, the non-homogeneous system **A** $\mathbf{x} = \mathbf{b}$ has a solution.
- **28.** Let **A** be a 5×3 matrix that has three linearly independent row vectors. Suppose that **b** is a vector in \mathbf{R}^5 such that the

nonhomogeneous system Ax = b has a solution. Prove that this solution is unique.

- **29.** Let **A** be an $m \times n$ matrix with m < n. Given **b** in \mathbb{R}^m , note that any solution of $A\mathbf{x} = \mathbf{b}$ expresses **b** as a linear combination of the column vectors of **A**. Then prove that no such solution is unique.
- **30.** Given the $m \times n$ matrix **A** with m > n, show that there exists a vector **b** in \mathbf{R}^m such that the system $\mathbf{A}\mathbf{x} = \mathbf{b}$ has no solution.
- **31.** Existence of Solutions Let A be an $m \times n$ matrix. Prove that the system Ax = b is consistent for every b in \mathbb{R}^m if and only if the rank of A is equal to m.
- **32.** Uniqueness of Solutions Let A be an $m \times n$ matrix and suppose that the system Ax = b is consistent. Prove that

4.6 Orthogonal Vectors in \mathbb{R}^n

its solution is unique if and only if the rank of \mathbf{A} is equal to n.

- **33.** Prove that the pivot column vectors in (8) are linearly independent.
- **34.** Let **A** be a matrix with rank *r*, and suppose that **A** can be reduced to echelon form *without row interchanges*. Show that the first *r* row vectors of **A** are linearly independent.
- **35.** Deduce from Theorem 3 in Section 4.3 that the rank of the matrix **A** is the largest integer r such that **A** has a nonsingular $r \times r$ submatrix.
- **36.** Let **A** be the 3×2 matrix whose columns are the unit basis vectors $\mathbf{i} = (1, 0, 0)$ and $\mathbf{j} = (0, 1, 0)$. If **B** is the row-equivalent matrix obtained by adding the first and second rows of **A** to its third row, show that **A** and **B** do *not* have the same column space.

In this section we show that the geometrical concepts of *distance* and *angle* in *n*-dimensional space can be based on the definition of the *dot product* of two vectors in \mathbf{R}^n . Recall from elementary calculus that the dot product $\mathbf{u} \cdot \mathbf{v}$ of two 3-dimensional vectors $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{v} = (v_1, v_2, v_3)$ is (by definition) the sum

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + u_3 v_3$$

of the products of corresponding scalar components of the two vectors.

Similarly, the **dot product** $\mathbf{u} \cdot \mathbf{v}$ of two *n*-dimensional vectors $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ is defined by

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n \tag{1}$$

(with one additional scalar product term for each additional dimension). And just as in \mathbb{R}^3 , it follows readily from the formula in (1) that if \mathbf{u} , \mathbf{v} , and \mathbf{w} are vectors in \mathbb{R}^n and *c* is a scalar, then

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u} \qquad (symmetry) \qquad (2)$$

$$\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w} \qquad (\text{distributivity}) \qquad (3)$$

$$(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v}) \qquad (\text{homogeneity}) \qquad (4)$$

Therefore, the dot product in
$$\mathbf{R}^n$$
 is an example of an *inner product*.

 $\mathbf{u} = \mathbf{0}$.

DEFINITION Inner Product

An **inner product** on a vector space V is a function that associates with each pair of vectors **u** and **v** in V a scalar $\langle \mathbf{u}, \mathbf{v} \rangle$ such that, if **u**, **v**, and **w** are vectors and c is a scalar, then

- (i) $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle;$
- (ii) $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle;$
- (iii) $\langle c\mathbf{u}, \mathbf{v} \rangle = c \langle \mathbf{u}, \mathbf{v} \rangle;$
- (iv) $\langle \mathbf{u}, \mathbf{u} \rangle \ge 0$; $\langle \mathbf{u}, \mathbf{u} \rangle = 0$ if and only if $\mathbf{u} = \mathbf{0}$.

The dot product on \mathbb{R}^n is sometimes called the Euclidean inner product, and with this inner product \mathbb{R}^n is sometimes called Euclidean *n*-dimensional space. We can use any of the notations in

$$\mathbf{u} \cdot \mathbf{v} = \langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u}^T \mathbf{v}$$

for the dot product of the two $n \times 1$ column vectors **u** and **v**. (Note in the last expression that $\mathbf{u}^T = (u_1, u_2, \dots, u_n)^T$ is the $1 \times n$ row vector with the indicated entries, so the 1×1 matrix product $\mathbf{u}^T \mathbf{v}$ is simply a scalar.) Here we will ordinarily use the notation $\mathbf{u} \cdot \mathbf{v}$.

The **length** $|\mathbf{u}|$ of the vector $\mathbf{u} = (u_1, u_2, \dots, u_n)$ is defined as follows:

$$|\mathbf{u}| = \sqrt{(\mathbf{u} \cdot \mathbf{u})} = \left(u_1^2 + u_2^2 + \dots + u_n^2\right)^{1/2}.$$
 (6)

Note that the case n = 2 is a consequence of the familiar Pythagorean formula in the plane.

Theorem 1 gives one of the most important inequalities in mathematics. Many proofs are known, but none of them seems direct and well motivated.

THEOREM 1 The Cauchy-Schwarz Inequality

If **u** and **v** are vectors in \mathbf{R}^n , then

$$|\mathbf{u} \cdot \mathbf{v}| \le |\mathbf{u}| |\mathbf{v}|. \tag{7}$$

Proof: If $\mathbf{u} = \mathbf{0}$, then $|\mathbf{u} \cdot \mathbf{v}| = |\mathbf{u}| = 0$, so the inequality is satisfied trivially. If $\mathbf{u} \neq \mathbf{0}$, then we let $a = \mathbf{u} \cdot \mathbf{u}$, $b = 2\mathbf{u} \cdot \mathbf{v}$, and $c = \mathbf{v} \cdot \mathbf{v}$. For any real number x, the distributivity and positivity properties of the dot product then yield

$$0 \leq (x\mathbf{u} + \mathbf{v}) \cdot (x\mathbf{u} + \mathbf{v})$$

= $(\mathbf{u} \cdot \mathbf{u})x^2 + 2(\mathbf{u} \cdot \mathbf{v})x + (\mathbf{v} \cdot \mathbf{v}),$

so that

 \succ

$$0 \leq ax^2 + bx + c.$$

Thus the quadratic equation $ax^2 + bx + c = 0$ either has no real roots or has a repeated real root. Hence the quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

implies that the discriminant $b^2 - 4ac$ cannot be positive; that is, $b^2 \leq 4ac$, so

$$4(\mathbf{u} \cdot \mathbf{v})^2 \leq 4(\mathbf{u} \cdot \mathbf{u})(\mathbf{v} \cdot \mathbf{v}).$$

We get the Cauchy-Schwarz inequality in (7) when we take square roots, remembering that the numbers $|\mathbf{u}| = (\mathbf{u} \cdot \mathbf{u})^{1/2}$ and $|\mathbf{v}| = (\mathbf{v} \cdot \mathbf{v})^{1/2}$ are nonnegative.



FIGURE 4.6.1. The angle θ between the vectors **u** and **v**.

The Cauchy-Schwarz inequality enables us to *define* the angle θ between the nonzero vectors **u** and **v**. (See Figure 4.6.1.) Division by the positive number $|\mathbf{u}||\mathbf{v}|$ in (7) yields $|\mathbf{u} \cdot \mathbf{v}|/(|\mathbf{u}||\mathbf{v}|) \leq 1$, so

$$-1 \leq \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}||\mathbf{v}|} \leq +1.$$
(8)

Hence there is a unique angle θ between 0 and π radians, inclusive (that is, between 0° and 180°), such that

$$\cos\theta = \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}||\mathbf{v}|}.\tag{9}$$

Thus we obtain the same geometric interpretation

$$\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}| |\mathbf{v}| \cos \theta \tag{10}$$

of the dot product in \mathbb{R}^n as one sees (for 3-dimensional vectors) in elementary calculus textbooks—for instance, see Section 11.2 of Edwards and Penney, *Calculus: Early Transcendentals*, 7th edition, Hoboken, NJ: Pearson, 2008.

On the basis of (10) we call the vectors **u** and **v orthogonal** provided that

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{0}. \tag{11}$$

If **u** and **v** are nonzero vectors this means that $\cos \theta = 0$, so $\theta = \pi/2$ (90°). Note that **u** = **0** satisfies (11) for all **v**, so the zero vector is orthogonal to *every* vector.

Example 1 Find the angle θ_n in \mathbb{R}^n between the x_1 -axis and the line through the origin and the point (1, 1, ..., 1).

Solution We take $\mathbf{u} = (1, 0, 0, ..., 0)$ on the x_1 -axis and $\mathbf{v} = (1, 1, ..., 1)$. Then $|\mathbf{u}| = 1$, $|\mathbf{v}| = \sqrt{n}$, and $\mathbf{u} \cdot \mathbf{v} = 1$, so the formula in (9) gives

$$\cos \theta_n = \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}||\mathbf{v}|} = \frac{1}{\sqrt{n}}$$

For instance, if

>

>

$$n = 3, \quad \text{then} \quad \theta_3 = \cos^{-1} \left(\frac{1}{\sqrt{3}}\right) \approx 0.9553 \quad (55^\circ);$$

$$n = 4, \quad \text{then} \quad \theta_4 = \cos^{-1} \left(\frac{1}{\sqrt{4}}\right) \approx 1.0472 \quad (60^\circ);$$

$$n = 5, \quad \text{then} \quad \theta_5 = \cos^{-1} \left(\frac{1}{\sqrt{5}}\right) \approx 1.1071 \quad (63^\circ);$$

$$n = 100, \quad \text{then} \quad \theta_{100} = \cos^{-1} \left(\frac{1}{10}\right) \approx 1.4706 \quad (84^\circ).$$

It is interesting to note that θ_n increases as *n* increases. Indeed, θ_n approaches $\cos^{-1}(0) = \pi/2$ (90°) as *n* increases without bound (so that $1/\sqrt{n}$ approaches zero).

In addition to angles, the dot product provides a definition of distance in \mathbb{R}^n . The **distance** $d(\mathbf{u}, \mathbf{v})$ between the points (vectors) $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ is defined to be

$$d(\mathbf{u}, \mathbf{v}) = |\mathbf{u} - \mathbf{v}|$$

= $[(u_1 - v_1)^2 + (u_2 - v_2)^2 + \dots + (u_n - v_n)^2]^{1/2}$. (12)

Example 2 The distance between the points $\mathbf{u} = (1, -1, -2, 3, 5)$ and $\mathbf{v} = (4, 3, 4, 5, 9)$ in \mathbb{R}^5 is

$$|\mathbf{u} - \mathbf{v}| = \sqrt{3^2 + 4^2 + 6^2 + 2^2 + 4^2} = \sqrt{81} = 9.$$

u + v o u

FIGURE 4.6.2. The "triangle" of the triangle inequality.

The *triangle inequality* of Theorem 2 relates the three sides of the triangle shown in Figure 4.6.2.

If **u** and **v** are vectors in \mathbf{R}^n , then

$$|\mathbf{u} + \mathbf{v}| \le |\mathbf{u}| + |\mathbf{v}|. \tag{13}$$

Proof: We apply the Cauchy-Schwarz inequality to find that

 $|\mathbf{u} + \mathbf{v}|^2 = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$ = $\mathbf{u} \cdot \mathbf{u} + 2\mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{v}$ $\leq \mathbf{u} \cdot \mathbf{u} + 2|\mathbf{u}||\mathbf{v}| + \mathbf{v} \cdot \mathbf{v}$ = $|\mathbf{u}|^2 + 2|\mathbf{u}||\mathbf{v}| + |\mathbf{v}|^2$,

and therefore

$$|\mathbf{u} + \mathbf{v}|^2 \leq (|\mathbf{u}| + |\mathbf{v}|)^2.$$

We now get (13) when we take square roots.

The vectors **u** and **v** are orthogonal if and only if $\mathbf{u} \cdot \mathbf{v} = 0$, so line (14) in the proof of the triangle inequality yields the fact that the **Pythagorean formula**

$$|\mathbf{u} + \mathbf{v}|^2 = |\mathbf{u}|^2 + |\mathbf{v}|^2 \tag{14}$$

holds if and only if the triangle with "adjacent side vectors" \mathbf{u} and \mathbf{v} is a right triangle with hypotenuse vector $\mathbf{u} + \mathbf{v}$ (see Figure 4.6.3).

The following theorem states a simple relationship between orthogonality and linear independence.

THEOREM 3 Orthogonality and Linear Independence

If the nonzero vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are mutually orthogonal—that is, each two of them are orthogonal—then they are linearly independent.

Proof: Suppose that

 $c_1\mathbf{v}_1+c_2\mathbf{v}_2+\cdots+c_k\mathbf{v}_k=\mathbf{0},$

where, as usual, c_1, c_2, \ldots, c_k are scalars. When we take the dot product of each side of this equation with \mathbf{v}_i , we find that

$$c_i \mathbf{v}_i \cdot \mathbf{v}_i = c_i |\mathbf{v}_i|^2 = 0.$$

Now $|\mathbf{v}_i| \neq 0$ because \mathbf{v}_i is a nonzero vector. It follows that $c_i = 0$. Thus $c_1 = c_2 = \cdots = c_k = 0$, and therefore the mutually orthogonal nonzero vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent.

In particular, any set of *n* mutually orthogonal nonzero vectors in \mathbb{R}^n constitutes a basis for \mathbb{R}^n . Such a basis is called an **orthogonal basis**. For instance, the standard unit vectors $\mathbf{e}_1, \mathbf{e}_2, \ldots, \mathbf{e}_n$ form an orthogonal basis for \mathbb{R}^n .



FIGURE 4.6.3. A right triangle in \mathbf{R}^n .

Orthogonal Complements

Now we want to relate orthogonality to the solution of systems of linear equations. Consider the homogeneous linear system

$$\mathbf{x} = \mathbf{0} \tag{15}$$

of *m* equations in *n* unknowns. If $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are the row vectors of the $m \times n$ coefficient matrix **A**, then the system looks like



Consequently, it is clear that x is a solution vector of Ax = 0 if and only if x is orthogonal to each row vector of A. But in the latter event x is orthogonal to every linear combination of row vectors of A because

$$\mathbf{x} \cdot (c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_m \mathbf{v}_m)$$

= $c_1 \mathbf{x} \cdot \mathbf{v}_1 + c_2 \mathbf{x} \cdot \mathbf{v}_2 + \dots + c_m \mathbf{x} \cdot \mathbf{v}_m$
= $(c_1)(0) + (c_2)(0) + \dots + (c_m)(0) = 0$

Thus we have shown that the vector \mathbf{x} in \mathbf{R}^n is a solution vector of $\mathbf{A}\mathbf{x} = \mathbf{0}$ if and only if \mathbf{x} is orthogonal to each vector in the row space $\text{Row}(\mathbf{A})$ of the matrix \mathbf{A} . This situation motivates the following definition.

DEFINITION The Orthogonal Complement of a Subspace

The vector **u** is **orthogonal** to the subspace V of \mathbb{R}^n provided that **u** is orthogonal to every vector in V. The **orthogonal complement** V^{\perp} (read "V perp") of V is the set of all those vectors in \mathbb{R}^n that are orthogonal to the subspace V.

If \mathbf{u}_1 and \mathbf{u}_2 are vectors in V^{\perp} , **v** is in V, and c_1 and c_2 are scalars, then

$$(c_1\mathbf{u}_1 + c_2\mathbf{u}_2) \cdot \mathbf{v} = c_1\mathbf{u}_1 \cdot \mathbf{v} + c_2\mathbf{u}_2 \cdot \mathbf{v}$$
$$= (c_1)(0) + (c_2)(0) = 0.$$

Thus any linear combination of vectors in V^{\perp} is orthogonal to every vector in V and hence is a vector in V^{\perp} . Therefore *the orthogonal complement* V^{\perp} *of a subspace* V *is itself a subspace of* \mathbf{R}^n . The standard picture of two complementary subspaces V and V^{\perp} consists of an orthogonal line and plane through the origin in \mathbf{R}^3 (see Fig. 4.6.4). The proofs of the remaining parts of Theorem 4 are left to the problems.

THEOREM 4 Properties of Orthogonal Complements

Let *V* be a subspace of \mathbf{R}^n . Then

- **1.** Its orthogonal complement V^{\perp} is also a subspace of \mathbf{R}^{n} ;
- 2. The only vector that lies in both V and V^{\perp} is the zero vector;
- 3. The orthogonal complement of V^{\perp} is V—that is, $(V^{\perp})^{\perp} = V$;
- **4.** If S is a spanning set for V, then the vector **u** is in V^{\perp} if and only if **u** is orthogonal to every vector in S.



FIGURE 4.6.4. Orthogonal complements.

In our discussion of the homogeneous linear system Ax = 0 in (16), we showed that a vector space lies in the **null space** Null(A) of A—that is, in the solution space of Ax = 0—if and only if it is orthogonal to each vector in the row space of A. In the language of orthogonal complements, this proves Theorem 5.

THEOREM 5 The Row Space and the Null Space

Let **A** be an $m \times n$ matrix. Then the row space Row(**A**) and the null space Null(**A**) are orthogonal complements in **R**^{*n*}. That is,

If $V = \operatorname{Row}(\mathbf{A})$, then $V^{\perp} = \operatorname{Null}(\mathbf{A})$. (16)

Now suppose that a subspace V of \mathbb{R}^n is given, with $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_m$ a set of vectors that span V. For instance, these vectors may form a given basis for V. Then the implication in (17) provides the following algorithm for finding a basis for the orthogonal complement V^{\perp} of V.

- **1.** Let **A** be the $m \times n$ matrix with row vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_m$.
- 2. Reduce A to echelon form and use the algorithm of Section 4.4 to find a basis $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ for the solution space Null(A) of $\mathbf{A}\mathbf{x} = \mathbf{0}$. Because $V^{\perp} =$ Null(A), this will be a basis for the orthogonal complement of V.

Example 3 Let V be the 1-dimensional subspace of \mathbf{R}^3 spanned by the vector $\mathbf{v}_1 = (1, -3, 5)$. Then $\mathbf{A} = \begin{bmatrix} 1 & -3 & 5 \end{bmatrix}$

and our linear system Ax = 0 consists of the single equation

$$x_1 - 3x_2 + 5x_3 = 0.$$

If $x_2 = s$ and $x_3 = t$, then $x_1 = 3s - 5t$. With s = 1 and t = 0 we get the solution vector $\mathbf{u}_1 = (3, 1, 0)$, whereas with s = 0 and t = 1 we get the solution vector $\mathbf{u}_2 = (-5, 0, 1)$. Thus the orthogonal complement V^{\perp} is the 2-dimensional subspace of \mathbf{R}^3 having $\mathbf{u}_1 = (3, 1, 0)$ and $\mathbf{u}_2 = (-5, 0, 1)$ as basis vectors.

Example 4 Let V be the 2-dimensional subspace of \mathbb{R}^5 that has $\mathbf{v}_1 = (1, 2, 1, -3, -3)$ and $\mathbf{v}_2 = (2, 5, 6, -10, -12)$ as basis vectors. The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 1 & -3 & -3 \\ 2 & 5 & 6 & -10 & -12 \end{bmatrix}$$

with row vectors v_1 and v_2 has reduced echelon form

$$\mathbf{E} = \begin{bmatrix} 1 & 0 & -7 & 5 & 9 \\ 0 & 1 & 4 & -4 & -6 \end{bmatrix}.$$

Hence the solution space of Ax = 0 is described parametrically by

$$x_3 = r, \quad x_4 = s, \quad x_5 = t,$$

 $x_2 = -4r + 4s + 6t$
 $x_1 = 7r - 5s - 9t.$

Then the choice

r = 1, s = 0, t = 0 yields $\mathbf{u}_1 = (7, -4, 1, 0, 0)$; r = 0, s = 1, t = 0 yields $\mathbf{u}_2 = (-5, 4, 0, 1, 0)$; r = 0, s = 0, t = 1 yields $\mathbf{u}_3 = (-9, 6, 0, 0, 1)$.

Thus the orthogonal complement V^{\perp} is the 3-dimensional subspace of \mathbb{R}^5 with basis $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$.

Observe that dim $V + \dim V^{\perp} = 3$ in Example 3, but dim $V + \dim V^{\perp} = 5$ in Example 4. It is no coincidence that in each case the dimensions of V and V^{\perp} add up to the dimension n of the Euclidean space containing them. To see why, suppose that V is a subspace of \mathbb{R}^n and let \mathbf{A} be an $m \times n$ matrix whose row vectors span V. Then Equation (12) in Section 4.5 implies that

$$rank(\mathbf{A}) + \dim Null(\mathbf{A}) = n$$

But

 $\dim V = \dim \operatorname{Row}(\mathbf{A}) = \operatorname{rank}(\mathbf{A})$

and

>

$$\dim V^{\perp} = \dim \operatorname{Null}(\mathbf{A})$$

by Theorem 5, so it follows that

$$\dim V + \dim V^{\perp} = n. \tag{17}$$

Moreover, it should be apparent intuitively that if

 $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$ is a basis for *V*

and

 $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ is a basis for V^{\perp} ,

then

 $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m, \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ is a basis for \mathbf{R}^n .

That is, *the union of a basis for V and a basis for V^{\perp} is a basis for R^{<i>n*}. In Problem 34 of this section we ask you to prove that this is so.

4.6 **Problems**

In Problems 1–4, determine whether the given vectors are mutually orthogonal.

- **1.** $\mathbf{v}_1 = (2, 1, 2, 1), \mathbf{v}_2 = (3, -6, 1, -2), \mathbf{v}_3 = (3, -1, -5, 5)$
- **2.** $\mathbf{v}_1 = (3, -2, 3, -4), \mathbf{v}_2 = (6, 3, 4, 6), \mathbf{v}_3 = (17, -12, -21, 3)$
- **3.** $\mathbf{v}_1 = (5, 2, -4, -1), \mathbf{v}_2 = (3, -5, 1, 1),$ $\mathbf{v}_3 = (3, 0, 8, -17)$
- **4.** $\mathbf{v}_1 = (1, 2, 3, -2, 1), \mathbf{v}_2 = (3, 2, 3, 6, -4), \mathbf{v}_3 = (6, 2, -4, 1, 4)$

In Problems 5–8, the three vertices A, B, and C of a triangle are given. Prove that each triangle is a right triangle by showing that its sides a, b, and c satisfy the Pythagorean relation $a^2 + b^2 = c^2$.

- **5.** A(6, 6, 5, 8), B(6, 8, 6, 5), C(5, 7, 4, 6)
- **6.** A(3, 5, 1, 3), B(4, 2, 6, 4), C(1, 3, 4, 2)
- 7. A(4, 5, 3, 5, -1), B(3, 4, -1, 4, 4), C(1, 3, 1, 3, 1)
- **8.** A(2, 8, -3, -1, 2), B(-2, 5, 6, 2, 12), C(-5, 3, 2, -3, 5)

9–12. Find the acute angles (in degrees) of each of the right triangles of Problems 5–8, respectively.

In Problems 13–22, the given vectors span a subspace V of the indicated Euclidean space. Find a basis for the orthogonal complement V^{\perp} of V.

- **13.** $\mathbf{v}_1 = (1, -2, 3)$ **14.** $\mathbf{v}_1 = (1, 5, -3)$ **15.** $\mathbf{v}_1 = (1, -2, -3, 5)$ **16.** $\mathbf{v}_1 = (1, 7, -6, -9)$ **17.** $\mathbf{v}_1 = (1, 3, 2, 4), \mathbf{v}_2 = (2, 7, 7, 3)$ **18.** $\mathbf{v}_1 = (1, -3, 3, 5), \mathbf{v}_2 = (2, -5, 9, 3)$ **19.** $\mathbf{v}_1 = (1, 2, 5, 2, 3), \mathbf{v}_2 = (3, 7, 11, 9, 5)$ **20.** $\mathbf{v}_1 = (2, 5, 5, 4, 3), \mathbf{v}_2 = (3, 7, 8, 8, 8)$ **21.** $\mathbf{v}_1 = (1, 2, 3, 1, 3), \mathbf{v}_2 = (1, 3, 4, 3, 6), \mathbf{v}_3 = (2, 2, 4, 3, 5)$ **22.** $\mathbf{v}_1 = (1, 1, 1, 1, 3), \mathbf{v}_2 = (2, 3, 1, 4, 7), \mathbf{v}_3 = (5, 3, 7, 1, 5)$ **23.** Prove: For arbitrary vectors **u** and **v**,
 - (a) $|\mathbf{u} + \mathbf{v}|^2 + |\mathbf{u} \mathbf{v}|^2 = 2|\mathbf{u}|^2 + 2|\mathbf{v}|^2;$
 - **(b)** $|\mathbf{u} + \mathbf{v}|^2 |\mathbf{u} \mathbf{v}|^2 = 4\mathbf{u} \cdot \mathbf{v}.$

24. Given mutually orthogonal vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$, show that

$$|\mathbf{v}_1 + \mathbf{v}_2 + \dots + \mathbf{v}_k|^2 = |\mathbf{v}_1|^2 + |\mathbf{v}_2|^2 + \dots + |\mathbf{v}_k|^2.$$

- **25.** Suppose that *A*, *B*, and *C* are the unit points on distinct coordinate axes in \mathbb{R}^n (for example, perhaps $A = \mathbf{e}_1$, $B = \mathbf{e}_2$, and $C = \mathbf{e}_3$). Show that *ABC* is an equilateral triangle.
- **26.** Prove that $|\mathbf{u} \cdot \mathbf{v}| = |\mathbf{u}||\mathbf{v}|$ if and only if the vectors \mathbf{u} and \mathbf{v} are linearly dependent.

In Problems 27–29, V denotes a subspace of \mathbb{R}^n .

- **27.** Show that the only vector that lies in both V and V^{\perp} is the zero vector.
- **28.** Prove that if $W = V^{\perp}$ then $W^{\perp} = V$.
- **29.** Let *S* be a spanning set for *V*. Show that the vector **u** is in V^{\perp} if and only if **u** is orthogonal to every vector in *S*.
- **30.** Suppose that the vectors **u** and **v** are orthogonal and that $\mathbf{u} + \mathbf{v} = \mathbf{0}$. Show that $\mathbf{u} = \mathbf{v} = \mathbf{0}$.
- **31.** Let *S* and *T* be two sets of vectors in \mathbb{R}^n such that each vector in *S* is orthogonal to every vector in *T*. Prove that each vector in span(*S*) is orthogonal to every vector in span(*T*).
- **32.** Let $S = {\mathbf{u}_1, \mathbf{u}_2}$ and $T = {\mathbf{v}_1, \mathbf{v}_2}$ be linearly independent sets of vectors such that each \mathbf{u}_i in S is orthogonal

4.7 General Vector Spaces

to every vector \mathbf{v}_j in *T*. Then use the results of Problems 30 and 31 to show that the four vectors \mathbf{u}_1 , \mathbf{u}_2 , \mathbf{v}_1 , \mathbf{v}_2 are linearly independent.

- **33.** Let $S = {\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k}$ and $T = {\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m}$ be linearly independent sets of vectors such that each \mathbf{u}_i in *S* is orthogonal to every vector \mathbf{v}_j in *T*. Then generalize Problem 32 to show that the k + m vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are linearly independent.
- 34. Deduce from the result of Problem 33 that if V is a subspace of Rⁿ, then the union of a basis for V and a basis for V[⊥] is a basis for Rⁿ.
- **35.** Let **A** be an $m \times n$ matrix and **b** a vector in **R**^{*m*}. Combine the observations listed below to prove that *the nonhomogeneous linear system* $\mathbf{A}\mathbf{x} = \mathbf{b}$ *is consistent if and only if* **b** *is orthogonal to the solution space of the homogeneous system* $\mathbf{A}^T \mathbf{y} = \mathbf{0}$.
 - (a) Ax = b is consistent if and only if b is in Col(A).
 - (b) **b** is in Col(A) if and only if **b** is orthogonal to ${Col(A)}^{\perp} = {Row(A^T)}^{\perp}$.
 - (c) $\{\operatorname{Row}(\mathbf{A}^T)\}^{\perp} = \operatorname{Null}(\mathbf{A}^T).$

In the previous six sections of this chapter, almost all the specific vector spaces appearing in our examples and problems have been vector spaces of n-tuples of real numbers. We have thus confined our attention largely to Euclidean spaces and their subspaces. In this section we discuss examples of some other types of vector spaces that play important roles in various branches of mathematics and their applications.

Example 1 Given fixed positive integers m and n, let M_{mn} denote the set of all $m \times n$ matrices with real number entries. Then M_{mn} is a vector space with the usual operations of addition of matrices and multiplication of matrices by scalars (real numbers). That is, these operations satisfy properties (a)–(h) in the definition of a vector space (Section 4.2). In particular, the zero element in M_{mn} is the $m \times n$ matrix **0** matrix whose elements are all zeros, and the negative $-\mathbf{A}$ of the matrix \mathbf{A} is (as usual) the matrix whose elements are the negatives of the corresponding entries of \mathbf{A} .

Given positive integers *i* and *j* with $1 \le i \le m$ and $1 \le j \le n$, let \mathbf{E}_{ij} denote the $m \times n$ matrix whose only nonzero entry is the number 1 in the *i*th row and the *j*th column. Then it should be clear that the mn elements { \mathbf{E}_{ij} } of M_{mn} form a basis for M_{mn} , and so M_{mn} is a finite-dimensional vector space of dimension mn.

For instance, in the case of the vector space M_{22} of all 2×2 matrices, these alleged basis elements are the four matrices

$$\mathbf{E}_{11} = \begin{bmatrix} 1 & 0\\ 0 & 0 \end{bmatrix}, \quad \mathbf{E}_{12} = \begin{bmatrix} 0 & 1\\ 0 & 0 \end{bmatrix},$$
$$\mathbf{E}_{21} = \begin{bmatrix} 0 & 0\\ 1 & 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{E}_{22} = \begin{bmatrix} 0 & 0\\ 0 & 1 \end{bmatrix}.$$

Then any matrix **A** in M_{22} can be expressed as

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a\mathbf{E}_{11} + b\mathbf{E}_{12} + c\mathbf{E}_{21} + d\mathbf{E}_{22},$$

so the set $\{E_{11}, E_{12}, E_{21}, E_{22}\}$ spans M_{22} . Moreover,

$$a\mathbf{E}_{11} + b\mathbf{E}_{12} + c\mathbf{E}_{21} + d\mathbf{E}_{22} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

implies immediately that a = b = c = d = 0, so the set { $\mathbf{E}_{11}, \mathbf{E}_{12}, \mathbf{E}_{21}, \mathbf{E}_{22}$ } is also linearly independent and therefore forms a basis for M_{22} . Thus, we have shown that M_{22} is a 4-dimensional vector space.

Example 2 Let C denote the subset of M_{22} consisting of all 2×2 matrices of the form

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}.$$
 (1)

Obviously the sum of any two such matrices is such a matrix, as is any scalar multiple of such a matrix. Thus *C* is closed under the operations of addition of matrices and multiplication by scalars and is therefore a *subspace* of M_{22} .

To determine the *dimension* of the subspace C, we consider the two special matrices

$$\mathbf{Re} = \begin{bmatrix} 1 & 0\\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \mathbf{Im} = \begin{bmatrix} 0 & -1\\ 1 & 0 \end{bmatrix}$$
(2)

in *C*. Here **Re** is another notation for the 2×2 identity matrix **I**, whereas the matrix **Im** is not so familiar. The matrices **Re** and **Im** are linearly independent—obviously neither is a scalar multiple of the other—and

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} = a\mathbf{Re} + b\mathbf{Im},\tag{3}$$

so these two matrices span *C*. Thus {**Re**, **Im**} is a basis for *C*, and so *C* is a 2-dimensional subspace of the 4-dimensional vector space M_{22} of 2×2 matrices.

Now observe that

$$(\mathbf{Im})^2 = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = -\mathbf{Re}.$$
 (4)

It follows that the matrix product of any two elements $a\mathbf{Re} + b\mathbf{Im}$ and $c\mathbf{Re} + d\mathbf{Im}$ of C is given by

$$(a\mathbf{Re} + b\mathbf{Im})(c\mathbf{Re} + d\mathbf{Im}) = ac(\mathbf{Re})^2 + bc\mathbf{Im}\,\mathbf{Re} + ad\,\mathbf{Re}\,\mathbf{Im} + bd(\mathbf{Im})^2$$
$$= (ac - bd)\mathbf{Re} + (ad + bc)\mathbf{Im}.$$
(5)

This explicit formula for the product of two matrices in *C* shows that, in addition to being closed under matrix addition and multiplication by scalars, the subspace *C* of M_{22} also is *closed under matrix multiplication*.

If you feel that the idea of a space of matrices being closed under matrix multiplication is a rather complex one, you are correct! Indeed, the vector space *C* of Example 2 can be taken as a "model" for the set of all *complex numbers* of the form a + bi, where *a* and *b* are real numbers and $i = \sqrt{-1}$ denotes the "imaginary" square root of -1 (so that $i^2 = -1$). Because $(\mathbf{Im})^2 = -\mathbf{Re}$, the matrix \mathbf{Im} plays the role of *i* and the matrix

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} = a\mathbf{R}\mathbf{e} + b\mathbf{I}\mathbf{m}$$

corresponds to the complex number a + bi. And the matrix multiplication formula in (5) is then analogous to the formula

$$(a+bi)(c+di) = (ac-bd) + (ad+bc)i$$

for the multiplication of complex numbers. In addition to providing a concrete interpretation of complex numbers in terms of real numbers, the matrix model *C* provides a convenient way to manipulate complex numbers in computer programming languages in which matrix operations are "built in" but complex operations are not.

Function Spaces

In Example 1 of Section 4.2 we introduced the vector space \mathcal{F} of all real-valued functions defined on the real line **R**. If **f** and **g** are elements of \mathcal{F} and *c* is a scalar, then

$$(\mathbf{f} + \mathbf{g})(x) = \mathbf{f}(x) + \mathbf{g}(x)$$

and

$$(c\mathbf{f})(x) = c(\mathbf{f}(x))$$

for all x in **R**. The zero element in \mathcal{F} is the function **0** such that $\mathbf{0}(x) = 0$ for all x. We frequently refer to a function in \mathcal{F} by simply specifying its formula. For instance, by "the function x^2 " we mean that function whose value at each x in **R** is x^2 .

The functions $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_k$ in \mathcal{F} are linearly dependent provided that there exist scalars c_1, c_2, \dots, c_k , not all zero, such that

$$c_1\mathbf{f}_1(x) + c_2\mathbf{f}_2(x) + \dots + c_k\mathbf{f}_k(x) = 0$$

for all x in **R**. We often can determine whether two given functions are linearly dependent simply by observing whether one is a scalar multiple of the other.

Example 3 The functions e^x and e^{-2x} are linearly independent because either of the equations $e^x = ae^{-2x}$ or $e^{-2x} = be^x$ would imply that e^{3x} is a constant, which obviously is not so. By contrast, the functions $\sin 2x$ and $\sin x \cos x$ are linearly dependent, because of the trigonometric identity $\sin 2x = 2 \sin x \cos x$. The three functions $1, \cos^2 x$, and $\sin^2 x$ are linearly dependent, because the fundamental identity $\cos^2 x + \sin^2 x = 1$ can be written in the form

$$(-1)(1) + (1)(\cos^2 x) + (1)(\sin^2 x) = 0.$$

A subspace of \mathcal{F} is called a **function space**. An example of a function space is the vector space \mathcal{P} of all polynomials in \mathcal{F} (Example 4 in Section 4.4). Recall that a function $\mathbf{p}(x)$ in \mathcal{F} is called a **polynomial of degree** $n \ge 0$ if it can be expressed in the form

$$\mathbf{p}(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n \tag{7}$$

with $a_n \neq 0$. Polynomials are added and multiplied by scalars in the usual manner by collecting coefficients of like powers of x. Clearly any linear combination of polynomials is a polynomial, so \mathcal{P} is, indeed, a subspace of \mathcal{F} . (Recall the subspace criterion of Theorem 1 in Section 4.2.)

Example 4 Given $n \ge 0$, denote by \mathcal{P}_n the set of all polynomials of degree at most n. That is, \mathcal{P}_n consists of all polynomials of degrees 0, 1, 2, ..., n. Any linear combination of two polynomials of degree at most n is again a polynomial of degree at most n, so \mathcal{P}_n is a subspace of \mathcal{P} (and of \mathcal{F}). The formula in (7) shows that the n + 1 polynomials

$$1, x, x^2, x^3, \dots, x^n$$
 (8)

span \mathcal{P}_n . To show that these "monomials" are linearly independent, suppose that

$$c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n = 0 \tag{9}$$

for all x in **R**. If a_0, a_1, \ldots, a_n are fixed *distinct* real numbers, then each a_i satisfies Eq. (9).

(6)

In matrix notation this means that

$$\begin{bmatrix} 1 & a_0 & a_0^2 & \cdots & a_0^n \\ 1 & a_1 & a_1^2 & \cdots & a_1^n \\ 1 & a_2 & a_2^2 & \cdots & a_2^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & a_n & a_n^2 & \cdots & a_n^n \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$
(10)

The coefficient matrix in (10) is a Vandermonde matrix **V**, and Problems 61–63 in Section 3.6 imply that any Vandermonde matrix is nonsingular. Therefore the system $\mathbf{Vc} = \mathbf{0}$ has only the trivial solution $\mathbf{c} = \mathbf{0}$, so it follows from (10) that $c_0 = c_1 = c_2 = \cdots = c_n = 0$. Thus we have proved that the n + 1 monomials $1, x, x^2, \dots, x^n$ are linearly independent and hence constitute a basis for \mathcal{P}_n . Consequently \mathcal{P}_n is an (n + 1)-dimensional vector space.

The fact that the monomials in (8) are linearly independent implies that the coefficients in a polynomial are unique. That is, if

$$a_0 + a_1 x + \dots + a_n x^n = b_0 + b_1 x + \dots + b_n x^n$$

for all x, then $a_0 = b_0$, $a_1 = b_1$, ..., and $a_n = b_n$. (See the remark that follows the definition of linear independence in Section 4.3.) This *identity principle* for polynomials is often used without proof in elementary algebra. A typical application is the method of *partial-fraction decomposition* illustrated in Example 5.

Example 5 Find constants *A*, *B*, and *C* such that

$$\frac{6x}{(x-1)(x+1)(x+2)} = \frac{A}{x-1} + \frac{B}{x+1} + \frac{C}{x+2}$$
(11)

Solution

for all x (other than x = 1, -1, or -2).

Multiplication of each side of the equation in (11) by the denominator on the left-hand side yields

$$6x = A(x + 1)(x + 2) + B(x - 1)(x + 2) + C(x - 1)(x + 1);$$

$$6x = (2A - 2B - C) + (3A + B)x + (A + B + C)x^{2}.$$

Then the identity principle for polynomials yields the linear equations

$$2A - 2B - C = 0$$

$$3A + B = 6$$

$$A + B + C = 0$$

which we readily solve for A = 1, B = 3, and C = -4. Therefore

$$\frac{6x}{(x-1)(x+1)(x+2)} = \frac{1}{x-1} + \frac{3}{x+1} - \frac{4}{x+2}$$

if $x \neq 1, -1, -2$.

Example 6

Show that the four polynomials

1,
$$x$$
, $3x^2 - 1$, and $5x^3 - 3x$ (12)

form a basis for \mathcal{P}_3 .

Solution

In order to show simultaneously that these four polynomials span \mathcal{P}_3 and are linearly independent, it suffices to see that every polynomial

$$p(x) = b_0 + b_1 x + b_2 x^2 + b_3 x^3$$
(13)

in \mathcal{P}_3 can be expressed uniquely as a linear combination

$$c_0(1) + c_1(x) + c_2(3x^2 - 1) + c_3(5x^3 - 3x)$$

= $(c_0 - c_2) + (c_1 - 3c_3)x + (3c_2)x^2 + (5c_3)x^3$ (14)

of the polynomials in (12). But upon comparing the coefficients in (13) and (14), we see that we need only observe that the linear system

[1]	0	-1	0]	$\begin{bmatrix} c_0 \end{bmatrix}$		$\begin{bmatrix} b_0 \end{bmatrix}$
0	1	0	-3	<i>c</i> ₁		b_1
0	0	3	0	<i>c</i> ₂	=	b_2
0	0	0	5	<i>c</i> ₃		b_3

obviously has a unique solution for c_0, c_1, c_2, c_3 in terms of b_0, b_1, b_2, b_3 .

Because the vector space \mathcal{P} of all polynomials contains n + 1 linearly independent functions for every integer $n \ge 0$, it follows that \mathcal{P} is an infinite-dimensional vector space. So too is any function space, such as \mathcal{F} itself, that contains \mathcal{P} . Another important function space is the set $\mathbb{C}^{(0)}$ of all continuous functions on \mathbf{R} ; $\mathbb{C}^{(0)}$ is a subspace of \mathcal{F} because every linear combination of continuous functions is again a continuous function. Every polynomial is a continuous function, so $\mathbb{C}^{(0)}$ contains \mathcal{P} and is therefore an infinite-dimensional vector space. Similarly, the set $\mathbb{C}^{(k)}$ of all functions in \mathcal{F} that have continuous *k*th-order derivatives is an infinite-dimensional function space.

Solution Spaces of Differential Equations

In Section 1.5 we saw that the (now familiar) first-order differential equation

$$y' = ky$$
 (k constant) (15)

has the general solution $y(x) = Ce^{kx}$. Thus the "solution space" of (15) is the set of all constant multiples of the single function e^{kx} , and it is therefore the 1-dimensional function space with basis $\{e^{kx}\}$.

In Section 5.1 we will see that the set of all solutions y(x) of a linear secondorder differential equation of the form

$$ay'' + by' + cy = 0 (16)$$

(with constant coefficients a, b, and c) is a 2-dimensional function space ϑ , called the **solution space** of the differential equation. Here we will illustrate this general fact with three simple examples of second-order equations that can be solved in an elementary manner.

Example 7 With a = 1 and b = c = 0 in (16), we get the differential equation

$$y'' = 0.$$
 (17)

If y(x) is any solution of this equation, so y''(x) = 0, then two integrations give first

$$y'(x) = \int y''(x) \, dx = \int (0) \, dx = A$$

and then

$$y(x) = \int y'(x) \, dx = \int A \, dx = Ax + B,$$

where *A* and *B* are arbitrary constants of integration. Thus every solution of (17) is of the form y(x) = Ax + B, and clearly every such function is a solution of the differential equation. Thus the solution space of (17) is simply the 2-dimensional space \mathcal{P}_1 of linear polynomials generated by the basis $\{1, x\}$.

Example 8

With a = 1, b = -2, and c = 0 in (16), we get the differential equation

$$y'' - 2y' = 0. (18)$$

If y(x) is any solution of this equation and if v(x) = y'(x), then (18) says that

$$v'(x) = 2v(x).$$

But this is a familiar first-order differential equation with general solution $v(x) = Ce^{2x}$. Then integration gives

$$y(x) = \int y'(x) \, dx = \int v(x) \, dx = \int C e^{2x} \, dx = \frac{1}{2} C e^{2x} + A.$$

Thus every solution of (18) is of the form $y(x) = A + Be^{2x}$ (where $B = \frac{1}{2}C$), and you should show by substitution in (18) that every function of this form is a solution of the differential equation. Therefore, the solution space of (18) is the 2-dimensional function space that is generated by the basis $\{1, e^{2x}\}$.

Example 9 With a = 1, b = 0, and c = -1 in (16), we get the differential equation

$$y'' - y = 0. (19)$$

If y(x) is any solution of this equation and if v(x) = y'(x), then (19) says that

$$y'' = \frac{dv}{dx} = \frac{dv}{dy}\frac{dy}{dx} = v\frac{dv}{dy} = y;$$

so v dv = y dy. Integration therefore gives

$$\frac{1}{2}v^2 = \frac{1}{2}y^2 + C$$
, so $v^2 = y^2 \pm a^2$, (20)

where the final sign depends on the sign of C. The final result is independent of this choice (Problem 27); we'll take the negative sign for the sake of illustration. Since v = dy/dx, it then follows from (20) that

$$x = \pm \int \frac{dy}{\sqrt{y^2 - a^2}}.$$
(21)

Here again the final result is independent of the choice of sign; this time we'll take the positive sign. Then the substitution y = au and the standard integral

$$\int \frac{du}{\sqrt{u^2 - 1}} = \cosh^{-1} u + C$$

(see Section 7.6 of Edwards and Penney, *Calculus: Early Transcendentals*, 7th edition, Hoboken, NJ: Pearson, 2008) yield

$$x = \cosh^{-1}\frac{y}{a} + b$$

so finally,

$$y = a\cosh(x - b) = a(\cosh x \cosh b - \sinh x \sinh b),$$

which can be written as

$$y(x) = A\cosh x + B\sinh x,$$
(22)

with $A = a \cosh b$ and $B = -a \sinh b$. Thus every solution of the differential equation y'' - y = 0 in (19) is of the form $y(x) = A \cosh x + B \sinh x$, and you should show by substitution in (19) that every function of this form is a solution of the differential equation. Therefore, the solution space & of (19) is the 2-dimensional function space that is generated by the basis { $\cosh x$, $\sinh x$ }. If we substitute into (22) the relations

$$\cosh x = \frac{e^x + e^{-x}}{2}, \quad \sinh x = \frac{e^x - e^{-x}}{2}$$

that define the hyperbolic cosine and sine functions in terms of exponentials, we see that every solution of y'' - y = 0 can also be written in the form of the linear combination

$$y(x) = Ce^x + De^{-x}$$

Consequently, $\{e^x, e^{-x}\}$ is a second basis for the 2-dimensional solution space \mathscr{S} .

In Section 5.3 we will learn a much quicker way to solve linear second-order differential equations like the equation y'' - y = 0 of Example 9 and the equation y'' + y = 0 of Problem 28.

4.7 **Problems**

In Problems 1–4, determine whether or not the indicated set of 3×3 matrices is a subspace of M_{33} .

- **1.** The set of all diagonal 3×3 matrices.
- **2.** The set of all symmetric 3×3 matrices (that is, matrices $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ such that $a_{ij} = a_{ji}$ for $1 \le i \le 3, 1 \le j \le 3$).
- **3.** The set of all nonsingular 3×3 matrices.
- **4.** The set of all singular 3×3 matrices.

In Problems 5–8, determine whether or not each indicated set of functions is a subspace of the space \mathcal{F} of all real-valued functions on **R**.

- 5. The set of all f such that f(0) = 0.
- **6.** The set of all f such that $f(x) \neq 0$ for all x.
- 7. The set of all f such that f(0) = 0 and f(1) = 1.
- 8. The set of all f such that f(-x) = -f(x) for all x.

In Problems 9–12, a condition on the coefficients of a polynomial $a_0 + a_1x + a_2x^2 + a_3x^3$ is given. Determine whether or not the set of all such polynomials satisfying this condition is a subspace of the space \mathcal{P} of all polynomials.

9. $a_3 \neq 0$

- **10.** $a_0 = a_1 = 0$
- **11.** $a_0 + a_1 + a_2 + a_3 = 0$
- 12. a_0, a_1, a_2 , and a_3 are all integers

In Problems 13–18, determine whether the given functions are linearly independent.

13. $\sin x$ and $\cos x$ **14.** e^x and xe^x **15.** 1 + x, 1 - x, and $1 - x^2$ **16.** 1 + x, $x + x^2$, and $1 - x^2$ **17.** $\cos 2x$, $\sin^2 x$, and $\cos^2 x$ **18.** $2\cos x + 3\sin x$ and $4\cos x + 5\sin x$

In Problems 19–22, use the method of Example 5 to find the constants A, B, and C in the indicated partial-fraction decompositions.

19.
$$\frac{x-5}{(x-2)(x-3)} = \frac{A}{x-2} + \frac{B}{x-3}$$

20.
$$\frac{2}{x(x^2 - 1)} = \frac{A}{x} + \frac{B}{x - 1} + \frac{C}{x + 1}$$

21.
$$\frac{8}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 4}$$

22.
$$\frac{2x}{(x + 1)(x + 2)(x + 3)} = \frac{A}{x + 1} + \frac{B}{x + 2} + \frac{C}{x + 3}$$

In Problems 23 and 24, use the method of Example 7 to find a basis for the solution space of the given differential equation. (It's 3-dimensional in Problem 23 and 4-dimensional in Problem 24.)

23.
$$y''' = 0$$
 24. $y^{(4)} = 0$

In Problems 25 and 26, use the method of Example 8 to find a basis for the 2-dimensional solution space of the given differential equation.

- **25.** y'' 5y' = 0 **26.** y'' + 10y' = 0
- **27.** Take the positive sign in Eq. (20), and then use the standard integral

$$\int \frac{du}{\sqrt{u^2 + 1}} = \sinh^{-1} u + C$$

to derive the same general solution $y(x) = A \cosh x + B \sinh x$ given in Eq. (22).

28. Use the method of Example 9 and the standard integral

$$\int \frac{du}{\sqrt{1-u^2}} = \sin^{-1}u + C$$

to derive the general solution $y(x) = A \cos x + B \sin x$ of the second-order differential equation y'' + y = 0. Thus its solution space has basis { $\cos x$, $\sin x$ }.

29. Let *V* be the set of all infinite sequences $\{x_n\} = \{x_1, x_2, x_3, ...\}$ of real numbers. Let addition of elements of *V* and multiplication by scalars be defined as follows:

$$\{x_n\} + \{y_n\} = \{x_n + y_n\}$$

and

- $c\{x_n\} = \{cx_n\}.$
- (a) Show that V is a vector space with these operations.
- (b) Prove that V is infinite dimensional.

264 CHAPTER 4 VECTOR SPACES

30. Let *V* be the vector space of Problem 29 and let the subset *W* consist of those elements $\{x_n\}$ of *V* such that $x_n = x_{n-1} + x_{n-2}$ for $n \ge 2$. Thus a typical element of *W* is the *Fibonacci sequence*

$$\{1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \ldots\}.$$

- (a) Show that W is a subspace of V.
- (b) Prove that W is 2-dimensional.

Problems 31 through 33 develop a method of multiplying matrices whose entries are complex numbers.

31. Motivated by Example 2, let us define a function or transformation *T* from the set of all complex numbers to the space of all 2×2 matrices of the form in (1) as follows. Given a complex number z = a + bi, let

$$T(z) = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}.$$

(a) Suppose that z_1 and z_2 are complex numbers and c_1 and c_2 are real numbers. Show that

$$T(c_1z_1 + c_2z_2) = c_1T(z_1) + c_2T(z_2).$$

(b) Show that

$$T(z_1 z_2) = T(z_1)T(z_2)$$

for all complex numbers z_1 and z_2 . Note the complex multiplication on the left in contrast with the matrix multiplication on the right.

(c) Prove that if z is an arbitrary nonzero complex number, then

$$T(z^{-1}) = \{T(z)\}^{-1}$$

where $z^{-1} = 1/z$ and $\{T(z)\}^{-1}$ is the inverse of the matrix T(z).

32. Let **A** and **B** be 4×4 (real) matrices partitioned into 2×2 submatrices or "blocks":

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{bmatrix}.$$

Then verify that **AB** can be calculated in "blockwise" fashion:

$$\mathbf{AB} = \begin{bmatrix} \mathbf{A}_{11}\mathbf{B}_{11} + \mathbf{A}_{12}\mathbf{B}_{21} & \mathbf{A}_{11}\mathbf{B}_{12} + \mathbf{A}_{12}\mathbf{B}_{22} \\ \mathbf{A}_{21}\mathbf{B}_{11} + \mathbf{A}_{22}\mathbf{B}_{21} & \mathbf{A}_{21}\mathbf{B}_{12} + \mathbf{A}_{22}\mathbf{B}_{22} \end{bmatrix}.$$

33. Given a 2×2 matrix $\mathbf{M} = \begin{bmatrix} z_{ij} \end{bmatrix}$ whose entries are complex numbers, let $T(\mathbf{M})$ denote the 4×4 matrix of real numbers given in block form by

$$T(\mathbf{M}) = \begin{bmatrix} T(z_{11}) & T(z_{12}) \\ T(z_{21}) & T(z_{22}) \end{bmatrix}$$

The transformation T on the right-hand side is the one we defined in Problem 31. Suppose that **M** and **N** are 2×2 complex matrices. Use Problems 31(b) and 32 to show that

$$T(\mathbf{MN}) = T(\mathbf{M})T(\mathbf{N}).$$

Hence one can find the product of the 2×2 complex matrices **M** and **N** by calculating the product of the 4×4 real matrices $T(\mathbf{M})$ and $T(\mathbf{N})$. For instance, if

$$\mathbf{M} = \begin{bmatrix} 1+i & 2+i \\ 1+2i & 1+3i \end{bmatrix}$$

and

$$\mathbf{N} = \begin{bmatrix} 3-i & 1-2i \\ 2-i & 3-2i \end{bmatrix},$$

then

$$T(\mathbf{M})T(\mathbf{N}) = \begin{bmatrix} 1 & -1 & 2 & -1 \\ 1 & 1 & 1 & 2 \\ \hline 1 & -2 & 1 & -3 \\ 2 & 1 & 3 & 1 \end{bmatrix} \times \begin{bmatrix} 3 & 1 & 1 & 2 \\ -1 & 3 & -2 & 1 \\ \hline 2 & 1 & 3 & 2 \\ -1 & 2 & -2 & 3 \end{bmatrix}$$
$$= \begin{bmatrix} 9 & -2 & 11 & 2 \\ \hline 2 & 9 & -2 & 11 \\ \hline 10 & -10 & 14 & -7 \\ 10 & 10 & 7 & 14 \end{bmatrix}.$$

Therefore,

$$\mathbf{MN} = \begin{bmatrix} 9+2i & 11-2i \\ 10+10i & 14+7i \end{bmatrix}.$$

If **M** is a nonsingular 2×2 complex matrix, it can be shown that $T(\mathbf{M}^{-1}) = \{T(\mathbf{M})\}^{-1}$. The $n \times n$ versions of these results are sometimes used to carry out complex matrix operations in computer languages that support only real arithmetic.

5 Higher-Order Linear Differential Equations

5.1 Introduction: Second-Order Linear Equations

In Chapters 1 and 2 we investigated first-order differential equations. We now turn to equations of higher order $n \ge 2$, beginning in this chapter with equations that are linear. The general theory of linear differential equations parallels the second-order case (n = 2), which we outline in this initial section.

Recall that a second-order differential equation in the (unknown) function y(x) is one of the form

$$G(x, y, y', y'') = 0.$$
 (1)

This differential equation is said to be **linear** provided that G is linear in the dependent variable y and its derivatives y' and y''. Thus a linear second-order equation takes (or can be written in) the form

$$A(x)y'' + B(x)y' + C(x)y = F(x).$$
 (2)

Unless otherwise noted, we will always assume that the (known) coefficient functions A(x), B(x), C(x), and F(x) are continuous on some open interval I (perhaps unbounded) on which we wish to solve this differential equation, but we do *not* require that they be linear functions of x. Thus the differential equation

$$e^{x}y'' + (\cos x)y' + (1 + \sqrt{x})y = \tan^{-1}x$$

is linear because the dependent variable y and its derivatives y' and y'' appear linearly. By contrast, the equations

$$y'' = yy'$$
 and $y'' + 3(y')^2 + 4y^3 = 0$

are not linear because products and powers of y or its derivatives appear.

If the function F(x) on the right-hand side of Eq. (2) vanishes identically on I, then we call Eq. (2) a **homogeneous** linear equation; otherwise, it is **nonhomo-geneous**. For example, the second-order equation

$$x^2y'' + 2xy' + 3y = \cos x$$

is nonhomogeneous; its associated homogeneous equation is

$$x^2y'' + 2xy' + 3y = 0.$$

In general, the homogeneous linear equation associated with Eq. (2) is

$$A(x)y'' + B(x)y' + C(x)y = 0.$$
 (3)

In case the differential equation in (2) models a physical system, the nonhomogeneous term F(x) frequently corresponds to some *external* influence on the system.

Remark Note that the meaning of the term "homogeneous" for a second-order linear differential equation is quite different from its meaning for a first-order differential equation (as in Section 1.6). Of course, it is not unusual—either in mathematics or in the English language more generally—for the same word to have different meanings in different contexts.

A Typical Application

Linear differential equations frequently appear as mathematical models of mechanical systems and electrical circuits. For example, suppose that a mass m is attached both to a spring that exerts on it a force F_S and to a dashpot (shock absorber) that exerts a force F_R on the mass (Fig. 5.1.1). Assume that the restoring force F_S of the spring is proportional to the *displacement* x of the mass from its equilibrium position and acts opposite to the direction of displacement. Then

$$F_S = -kx$$
 (with $k > 0$)

so $F_S < 0$ if x > 0 (spring stretched) while $F_S > 0$ if x < 0 (spring compressed). We assume that the dashpot force F_R is proportional to the *velocity* v = dx/dt of the mass and acts opposite to the direction of motion. Then

$$F_R = -cv = -c\frac{dx}{dt} \quad \text{(with } c > 0\text{)},$$

so $F_R < 0$ if v > 0 (motion to the right) while $F_R > 0$ if v < 0 (motion to the left).

If F_R and F_S are the only forces acting on the mass *m* (Fig. 5.1.2) and its resulting acceleration is a = dv/dt, then Newton's law F = ma gives

$$mx'' = F_S + F_R; (4)$$

that is,

$$m\frac{d^2x}{dt^2} + c\frac{dx}{dt} + kx = 0.$$
(5)

Thus we have a differential equation satisfied by the position function x(t) of the mass *m*. This homogeneous second-order linear equation governs the *free vibrations* of the mass; we will return to this problem in detail in Section 5.4.

If, in addition to F_S and F_R , the mass *m* is acted on by an external force F(t)—which must then be added to the right-hand side in Eq. (4)—the resulting equation is

$$m\frac{d^2x}{dt^2} + c\frac{dx}{dt} + kx = F(t).$$
(6)

This nonhomogeneous linear differential equation governs the *forced vibrations* of the mass under the influence of the external force F(t).



FIGURE 5.1.1. A mass–spring– dashpot system.



FIGURE 5.1.2. Directions of the forces acting on *m*.

Homogeneous Second-Order Linear Equations

Consider the general second-order linear equation

$$A(x)y'' + B(x)y' + C(x)y = F(x),$$
(7)

where the coefficient functions A, B, C, and F are continuous on the open interval I. Here we assume in addition that $A(x) \neq 0$ at each point of I, so we can divide each term in Eq. (7) by A(x) and write it in the form

$$y'' + p(x)y' + q(x)y = f(x).$$
(8)

We will discuss first the associated homogeneous equation

$$y'' + p(x)y' + q(x)y = 0.$$
 (9)

A particularly useful property of this *homogeneous* linear equation is the fact that the sum of any two solutions of Eq. (9) is again a solution, as is any constant multiple of a solution. This is the central idea of the following theorem.

THEOREM 1 Principle of Superposition for Homogeneous Equations

Let y_1 and y_2 be two solutions of the homogeneous linear equation in (9) on the interval *I*. If c_1 and c_2 are constants, then the linear combination

$$v = c_1 y_1 + c_2 y_2 \tag{10}$$

is also a solution of Eq. (9) on *I*.

]

Proof: The conclusion follows almost immediately from the linearity of the operation of differentiation, which gives

$$y' = c_1 y'_1 + c_2 y'_2$$
 and $y'' = c_1 y''_1 + c_2 y''_2$.

Then

$$y'' + py' + qy = (c_1y_1 + c_2y_2)'' + p(c_1y_1 + c_2y_2)' + q(c_1y_1 + c_2y_2)$$

= $(c_1y_1'' + c_2y_2'') + p(c_1y_1' + c_2y_2') + q(c_1y_1 + c_2y_2)$
= $c_1(y_1'' + py_1' + qy_1) + c_2(y_2'' + py_2' + qy_2)$
= $c_1 \cdot 0 + c_2 \cdot 0 = 0$

because y_1 and y_2 are solutions. Thus $y = c_1y_1 + c_2y_2$ is also a solution.

Remark According to Theorem 1 in Section 4.2, Theorem 1 here implies that the set \mathscr{S} of all solutions of the homogeneous linear second-order equation (9) is a subspace of the vector space of all functions on the real line **R**—and hence is itself a vector space. We therefore call *S* the **solution space** of the differential equation.

Example 1 We can see by inspection that

$$y_1(x) = \cos x$$
 and $y_2(x) = \sin x$

are two solutions of the equation

$$y'' + y = 0.$$

 $y(x) = c_1 \cos x$ of y'' + y = 0.

Theorem 1 tells us that any linear combination of these solutions, such as

$$y(x) = 3y_1(x) - 2y_2(x) = 3\cos x - 2\sin x,$$

is also a solution. We will see later that, conversely, *every* solution of y'' + y = 0 is a linear combination of these two particular solutions y_1 and y_2 . Thus a general solution of y'' + y = 0 is given by

$$y(x) = c_1 \cos x + c_2 \sin x$$

It is important to understand that this single formula for the general solution encompasses a "twofold infinity" of particular solutions, because the two coefficients c_1 and c_2 can be selected independently. Figures 5.1.3 through 5.1.5 illustrate some of the possibilities, with either c_1 or c_2 set equal to zero, or with both nonzero.



FIGURE 5.1.4. Solutions $y(x) = c_2 \sin x$ of y'' + y = 0.



Earlier in this section we gave the linear equation mx'' + cx' + kx = F(t) as a mathematical model of the motion of the mass shown in Fig. 5.1.1. Physical considerations suggest that the motion of the mass should be determined by its initial position and initial velocity. Hence, given any preassigned values of x(0) and x'(0), Eq. (6) ought to have a *unique* solution satisfying these initial conditions. More generally, in order to be a "good" mathematical model of a deterministic physical situation, a differential equation must have unique solutions satisfying any appropriate initial conditions. The following existence and uniqueness theorem (proved in the Appendix) gives us this assurance for the general second-order equation.

THEOREM 2 Existence and Uniqueness for Linear Equations

Suppose that the functions p, q, and f are continuous on the open interval I containing the point a. Then, given any two numbers b_0 and b_1 , the equation

$$y'' + p(x)y' + q(x)y = f(x)$$
(8)

has a unique (that is, one and only one) solution on the entire interval I that satisfies the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$
 (11)

Remark 1 Equation (8) and the conditions in (11) constitute a second-order linear **initial** value problem. Theorem 2 tells us that any such initial value problem has a unique solution on the *whole* interval I where the coefficient functions in (8) are continuous. Recall from

Section 1.3 that a *nonlinear* differential equation generally has a unique solution on only a smaller interval.

Remark 2 Whereas a *first-order* differential equation dy/dx = F(x, y) generally admits only a single solution curve y = y(x) passing through a given initial point (a, b), Theorem 2 implies that the *second-order* equation in (8) has infinitely many solution curves passing through the point (a, b_0) —namely, one for each (real number) value of the initial slope $y'(a) = b_1$. That is, instead of there being only one line through (a, b_0) tangent to a solution curve, *every* nonvertical straight line through (a, b_0) is tangent to some solution curve of Eq. (8). Figure 5.1.6 shows a number of solution curves of the equation y'' + 3y' + 2y = 0 all having the same initial value y(0) = 1, while Fig. 5.1.7 shows a number of solution curves all having the same initial slope y'(0) = 1. The application at the end of this section suggests how to construct such families of solution curves for a given homogeneous second-order linear differential equation.



y'' + 3y' + 2y = 0 with the same initial value y(0) = 1 but different initial slopes.



Example 1 Continued We saw in the first part of Example 1 that $y(x) = 3 \cos x - 2 \sin x$ is a solution (on the entire real line) of y'' + y = 0. It has the initial values y(0) = 3, y'(0) = -2. Theorem 2 tells us that this is the *only* solution with these initial values. More generally, the solution

$$y(x) = b_0 \cos x + b_1 \sin x$$

satisfies the *arbitrary* initial conditions $y(0) = b_0$, $y'(0) = b_1$; this illustrates the *existence* of such a solution, also as guaranteed by Theorem 2.

Example 1 suggests how, given a *homogeneous* second-order linear equation, we might actually find the solution y(x) whose existence is assured by Theorem 2. First, we find two "essentially different" solutions y_1 and y_2 ; second, we attempt to impose on the general solution

$$y = c_1 y_1 + c_2 y_2 \tag{12}$$

the initial conditions $y(a) = b_0$, $y'(a) = b_1$. That is, we attempt to solve the simultaneous equations

$$c_1 y_1(a) + c_2 y_2(a) = b_0,$$

$$c_1 y_1'(a) + c_2 y_2'(a) = b_1$$
(13)

for the coefficients c_1 and c_2 .

Example 2

 $y_1(x) = e^x$ and $y_2(x) = xe^x$

are solutions of the differential equation

$$y'' - 2y' + y = 0,$$

and then find a solution satisfying the initial conditions y(0) = 3, y'(0) = 1.

Solution The verification is routine; we omit it. We impose the given initial conditions on the general solution

$$y(x) = c_1 e^x + c_2 x e^x,$$

for which



FIGURE 5.1.8. Different solutions $y(x) = 3e^x + c_2xe^x$ of y'' - 2y' + y = 0 with the same initial value y(0) = 3.

$$y'(x) = (c_1 + c_2)e^x + c_2xe^x$$

to obtain the simultaneous equations

$$y(0) = c_1 = 3$$

 $y'(0) = c_1 + c_2 = 1$

The resulting solution is $c_1 = 3$, $c_2 = -2$. Hence the solution of the original initial value problem is

$$y(x) = 3e^x - 2xe^x$$

Figure 5.1.8 shows several additional solutions of y'' - 2y' + y = 0, all having the same initial value y(0) = 3.

Linearly Independent Solutions

In order for the procedure of Example 2 to succeed, the two solutions y_1 and y_2 must have the property that the equations in (13) can always be solved for c_1 and c_2 , no matter what the initial values $y(a) = b_0$ and $y'(a) = b_1$ might be. This is so, provided that the two functions y_1 and y_2 are linearly independent. According to the following definition, linear independence of functions defined on an interval I is analogous to linear independence of ordinary vectors.

DEFINITION Linear Independence of Two Functions

Two functions defined on an open interval *I* are said to be **linearly independent** on *I* provided that neither is a constant multiple of the other.

Two functions are said to be **linearly dependent** on an open interval provided that they are not linearly independent there; that is, one of them is a constant multiple of the other. We can always determine whether two given functions f and g are linearly dependent on an interval I by noting at a glance whether either of the two quotients f/g or g/f is a constant-valued function on I.

Example 3 Thus it is clear that the following pairs of functions are linearly independent on the entire real line:

 $\begin{array}{rcl}
\sin x & \text{and} & \cos x;\\
e^x & \text{and} & e^{-2x};\\
e^x & \text{and} & xe^x;\\
x+1 & \text{and} & x^2;\\
x & \text{and} & |x|.\\
\end{array}$

That is, neither $\sin x/\cos x = \tan x$ nor $\cos x/\sin x = \cot x$ is a constant-valued function; neither $e^x/e^{-2x} = e^{3x}$ nor e^{-2x}/e^x is a constant-valued function; and so forth. But the

identically zero function $f(x) \equiv 0$ and any other function g are linearly dependent on every interval, because $0 \cdot g(x) = 0 = f(x)$. Also, the functions

$$f(x) = \sin 2x$$
 and $g(x) = \sin x \cos x$

are linearly dependent on any interval because f(x) = 2g(x) is the familiar trigonometric identity $\sin 2x = 2 \sin x \cos x$.

General Solutions

But does the homogeneous equation y'' + py' + qy = 0 always have two linearly independent solutions? Theorem 2 says yes! We need only choose y_1 and y_2 so that

$$y_1(a) = 1$$
, $y'_1(a) = 0$ and $y_2(a) = 0$, $y'_2(a) = 1$.

It is then impossible that either $y_1 = ky_2$ or $y_2 = ky_1$ because $k \cdot 0 \neq 1$ for any constant k. Theorem 2 tells us that two such linearly independent solutions *exist*; actually finding them is a crucial matter that we will discuss briefly at the end of this section, and in greater detail beginning in Section 5.3.

We want to show, finally, that given *any* two linearly independent solutions y_1 and y_2 of the homogeneous equation

$$y''(x) + p(x)y'(x) + q(x)y(x) = 0,$$
(9)

every solution y of Eq. (9) can be expressed as a linear combination

$$y = c_1 y_1 + c_2 y_2 \tag{12}$$

of y_1 and y_2 . This means that the function in (12) is a *general solution* of Eq. (9)—it provides all possible solutions of the differential equation.

As suggested by the equations in (13), the determination of the constants c_1 and c_2 in (12) depends on a certain 2×2 determinant of values of y_1 , y_2 , and their derivatives. Given two functions f and g, the **Wronskian** of f and g is the determinant

$$\blacktriangleright \qquad \qquad W = \begin{vmatrix} f & g \\ f' & g' \end{vmatrix} = fg' - f'g.$$

We write either W(f, g) or W(x), depending on whether we wish to emphasize the two functions or the point x at which the Wronskian is to be evaluated. For example,

$$W(\cos x, \sin x) = \begin{vmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{vmatrix} = \cos^2 x + \sin^2 x = 1$$

and

$$W(e^x, xe^x) = \begin{vmatrix} e^x & xe^x \\ e^x & e^x + xe^x \end{vmatrix} = e^{2x}.$$

These are examples of linearly *independent* pairs of solutions of differential equations (see Examples 1 and 2). Note that in both cases the Wronskian is everywhere *nonzero*.

On the other hand, if the functions f and g are linearly dependent, with f = kg (for example), then

$$W(f,g) = \begin{vmatrix} kg & g \\ kg' & g' \end{vmatrix} = kgg' - kg'g \equiv 0.$$

Thus the Wronskian of two linearly *dependent* functions is identically zero. In Section 5.2 we will prove that, if the two functions y_1 and y_2 are solutions of a homogeneous second-order linear equation, then the strong converse stated in part (b) of Theorem 3 holds.

THEOREM 3 Wronskians of Solutions

Suppose that y_1 and y_2 are two solutions of the homogeneous second-order linear equation (Eq. (9))

$$y'' + p(x)y' + q(x)y = 0$$

on an open interval I on which p and q are continuous.

(a) If y_1 and y_2 are linearly dependent, then $W(y_1, y_2) \equiv 0$ on *I*.

(b) If y_1 and y_2 are linearly independent, then $W(y_1, y_2) \neq 0$ at each point of *I*.

Thus, given two solutions of Eq. (9), there are just two possibilities: The Wronskian W is identically zero if the solutions are linearly dependent; the Wronskian is never zero if the solutions are linearly independent. The latter fact is what we need to show that $y = c_1y_1 + c_2y_2$ is the general solution of Eq. (9) if y_1 and y_2 are linearly independent solutions.

THEOREM 4 General Solutions of Homogeneous Equations

Let y_1 and y_2 be two linearly independent solutions of the homogeneous equation (Eq. (9))

$$y'' + p(x)y' + q(x)y = 0$$

with p and q continuous on the open interval I. If Y is any solution whatsoever of Eq. (9) on I, then there exist numbers c_1 and c_2 such that

$$Y(x) = c_1 y_1(x) + c_2 y_2(x)$$

for all x in I.

Remark In essence, Theorem 4 tells us that, when we have found *two* linearly independent solutions y_1 and y_2 of the homogeneous equation in (9), we have then found *all* of its solutions. Specifically, $\{y_1, y_2\}$ is then a basis for the solution space of the differential equation. We therefore call the linear combination $Y = c_1y_1 + c_2y_2$ a *general solution* of the differential equation.

Proof of Theorem 4: Choose a point a of I, and consider the simultaneous equations

$$c_1 y_1(a) + c_2 y_2(a) = Y(a),$$

$$c_1 y_1'(a) + c_2 y_2'(a) = Y'(a).$$
(14)

The determinant of the coefficients in this system of linear equations in the unknowns c_1 and c_2 is simply the Wronskian $W(y_1, y_2)$ evaluated at x = a. By Theorem 3, this determinant is nonzero, so by elementary algebra it follows that the equations in (14) can be solved for c_1 and c_2 . With these values of c_1 and c_2 , we define the solution

$$G(x) = c_1 y_1(x) + c_2 y_2(x)$$

of Eq. (9); then

$$G(a) = c_1 y_1(a) + c_2 y_2(a) = Y(a)$$

and

$$G'(a) = c_1 y'_1(a) + c_2 y'_2(a) = Y'(a).$$

Thus the two solutions Y and G have the same initial values at a; likewise, so do Y' and G'. By the uniqueness of a solution determined by such initial values (Theorem 2), it follows that Y and G agree on I. Thus we see that

$$Y(x) \equiv G(x) = c_1 y_1(x) + c_2 y_2(x),$$

as desired. If $y_1(x) = e^{2x}$ and $y_2(x) = e^{-2x}$, then

Example 4

$$y_1'' = (2)(2)e^{2x} = 4e^{2x} = 4y_1$$
 and $y_2'' = (-2)(-2)e^{-2x} = 4e^{-2x} = 4y_2$.

Therefore, y_1 and y_2 are linearly independent solutions of

$$y'' - 4y = 0. (15)$$

But $y_3(x) = \cosh 2x$ and $y_4(x) = \sinh 2x$ are also solutions of Eq. (15), because

$$\frac{d^2}{dx^2}(\cosh 2x) = \frac{d}{dx}(2\sinh 2x) = 4\cosh 2x$$

and, similarly, $(\sinh 2x)'' = 4 \sinh 2x$. It therefore follows from Theorem 4 that the functions $\cosh 2x$ and $\sinh 2x$ can be expressed as linear combinations of $y_1(x) = e^{2x}$ and $y_2(x) = e^{-2x}$. Of course, this is no surprise, because

$$\cosh 2x = \frac{1}{2}e^{2x} + \frac{1}{2}e^{-2x}$$
 and $\sinh 2x = \frac{1}{2}e^{2x} - \frac{1}{2}e^{-2x}$

by the definitions of the hyperbolic cosine and hyperbolic sine. Thus the solution space of the differential equation y'' - 4y = 0 has the two different bases $\{e^{2x}, e^{-2x}\}$ and $\{\cosh 2x, \sinh 2x\}$.

Remark Because e^{2x} , e^{-2x} and $\cosh x$, $\sinh x$ are two different pairs of linearly independent solutions of the equation y'' - 4y = 0 in (15), Theorem 4 implies that every particular solution Y(x) of this equation can be written both in the form

$$Y(x) = c_1 e^{2x} + c_2 e^{-2x}$$

and in the form

 \succ

$$Y(x) = a\cosh 2x + b\sinh 2x.$$

Thus these two different linear combinations (with arbitrary constant coefficients) provide two different descriptions of the set of all solutions of the same differential equation y'' - 4y = 0. Hence each of these two linear combinations is a general solution of the equation. Indeed, this is why it is accurate to refer to a specific such linear combination as "*a* general solution" rather than as "*the* general solution."

Linear Second-Order Equations with Constant Coefficients

As an illustration of the general theory introduced in this section, we discuss the homogeneous second-order linear differential equation

$$ay'' + by' + cy = 0 (16)$$

with *constant* coefficients a, b, and c. We first look for a *single* solution of Eq. (16) and begin with the observation that

$$(e^{rx})' = re^{rx}$$
 and $(e^{rx})'' = r^2 e^{rx}$, (17)

so any derivative of e^{rx} is a constant multiple of e^{rx} . Hence, if we substituted $y = e^{rx}$ in Eq. (16), then each term would be a constant multiple of e^{rx} , with the constant coefficients dependent on r and the coefficients a, b, and c. This suggests that we try to find a value of r so that these multiples of e^{rx} will have sum zero. If we succeed, then $y = e^{rx}$ will be a solution of Eq. (16).
For example, if we substitute $y = e^{rx}$ in the equation

$$y'' - 5y' + 6y = 0,$$

we obtain

$$r^2 e^{rx} - 5r e^{rx} + 6e^{rx} = 0.$$

Thus

 \succ

$$(r^2 - 5r + 6)e^{rx} = 0;$$
 $(r - 2)(r - 3)e^{rx} = 0.$

Hence $y = e^{rx}$ will be a solution if either r = 2 or r = 3. So, in searching for a single solution, we actually have found two solutions: $y_1(x) = e^{2x}$ and $y_2(x) = e^{3x}$.

To carry out this procedure in the general case, we substitute $y = e^{rx}$ in Eq. (16). With the aid of the equations in (17), we find the result to be

$$ar^2e^{rx} + bre^{rx} + ce^{rx} = 0.$$

Because e^{rx} is never zero, we conclude that $y(x) = e^{rx}$ will satisfy the differential equation in (16) precisely when *r* is a root of the *algebraic* equation

$$ar^2 + br + c = 0. (18)$$

This quadratic equation is called the **characteristic equation** of the homogeneous linear differential equation

$$ay'' + by' + cy = 0.$$
 (16)

If Eq. (18) has two *distinct* (unequal) roots r_1 and r_2 , then the corresponding solutions $y_1(x) = e^{r_1 x}$ and $y_2(x) = e^{r_2 x}$ of (16) are linearly independent. (Why?) This gives the following result.

THEOREM 5 Distinct Real Roots

If the roots r_1 and r_2 of the characteristic equation in (18) are real and distinct, then

$$y(x) = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$
(19)

is a general solution of Eq. (16). Thus the solution space of the equation ay'' + by' + cy = 0 has basis $\{e^{r_1x}, e^{r_2x}\}$.

Example 5 Find the general solution of

>

$$2y'' - 7y' + 3y = 0.$$

Solution

$$2r^2 - 7r + 3 = 0$$

by factoring:

$$(2r-1)(r-3) = 0$$

The roots $r_1 = \frac{1}{2}$ and $r_2 = 3$ are real and distinct, so Theorem 5 yields the general solution

$$y(x) = c_1 e^{x/2} + c_2 e^{3x}.$$

Example 6

The differential equation y'' + 2y' = 0 has characteristic equation

$$r^2 + 2r = r(r+2) = 0$$

with distinct real roots $r_1 = 0$ and $r_2 = -2$. Because $e^{0 \cdot x} \equiv 1$, we get the general solution

$$y(x) = c_1 + c_2 e^{-2x}$$

Figure 5.1.9 shows several different solution curves with $c_1 = 1$, all appearing to approach the solution curve $y(x) \equiv 1$ (with $c_2 = 0$) as $x \to +\infty$.

Remark Note that Theorem 5 changes a problem involving a differential equation into one involving only the solution of an *algebraic* equation.

If the characteristic equation in (18) has equal roots $r_1 = r_2$, we get (at first) only the single solution $y_1(x) = e^{r_1 x}$ of Eq. (16). The problem in this case is to produce the "missing" second solution of the differential equation.

A double root $r = r_1$ will occur precisely when the characteristic equation is a constant multiple of the equation

$$(r - r_1)^2 = r^2 - 2r_1r + r_1^2 = 0.$$

Any differential equation with this characteristic equation is equivalent to

$$y'' - 2r_1y' + r_1^2y = 0. (20)$$

But it is easy to verify by direct substitution that $y = xe^{r_1x}$ is a second solution of Eq. (20). It is clear (but you should verify) that

$$y_1(x) = e^{r_1 x}$$
 and $y_2(x) = x e^{r_1 x}$

are linearly independent functions, so the general solution of the differential equation in (20) is

$$y(x) = c_1 e^{r_1 x} + c_2 x e^{r_1 x}.$$

THEOREM 6 Repeated Roots

If the characteristic equation in (18) has equal (necessarily real) roots $r_1 = r_2$, then

$$y(x) = (c_1 + c_2 x)e^{r_1 x}$$
(21)

is a general solution of Eq. (16). In this case the solution space of the equation ay'' + by' + cy = 0 has basis $\{e^{r_1x}, xe^{r_1x}\}$.

Example 7 To solve the initial value problem

$$y'' + 2y' + y = 0;$$

 $y(0) = 5, y'(0) = -3,$

we note first that the characteristic equation

$$r^2 + 2r + 1 = (r+1)^2 = 0$$

has equal roots $r_1 = r_2 = -1$. Hence the general solution provided by Theorem 6 is

$$y(x) = c_1 e^{-x} + c_2 x e^{-x}.$$



FIGURE 5.1.9. Solutions $y(x) = 1 + c_2e^{-2x}$ of y'' + 2y' = 0 with different values of c_2 .



FIGURE 5.1.10. Solutions $y(x) = c_1 e^{-x} + 2x e^{-x}$ of y'' + 2y' + y = 0 with different values of c_1 .

Problems 5.1

In Problems 1 through 16, a homogeneous second-order linear differential equation, two functions y_1 and y_2 , and a pair of initial conditions are given. First verify that y_1 and y_2 are solutions of the differential equation. Then find a particular solution of the form $y = c_1y_1 + c_2y_2$ that satisfies the given initial conditions. Primes denote derivatives with respect to x.

- **1.** y'' y = 0; $y_1 = e^x$, $y_2 = e^{-x}$; y(0) = 0, y'(0) = 5
- **2.** y'' 9y = 0; $y_1 = e^{3x}$, $y_2 = e^{-3x}$; y(0) = -1, y'(0) = 15
- **3.** y'' + 4y = 0; $y_1 = \cos 2x$, $y_2 = \sin 2x$; y(0) = 3, y'(0) = 8
- **4.** y'' + 25y = 0; $y_1 = \cos 5x$, $y_2 = \sin 5x$; y(0) = 10, v'(0) = -10
- 5. v'' 3v' + 2v = 0; $v_1 = e^x$, $v_2 = e^{2x}$; v(0) = 1, v'(0) = 0
- **6.** $y'' + y' 6y = 0; y_1 = e^{2x}, y_2 = e^{-3x}; y(0) = 7,$ v'(0) = -1
- 7. y'' + y' = 0; $y_1 = 1$, $y_2 = e^{-x}$; y(0) = -2, y'(0) = 8
- 8. y'' 3y' = 0; $y_1 = 1$, $y_2 = e^{3x}$; y(0) = 4, y'(0) = -2
- 9. $y'' + 2y' + y = 0; y_1 = e^{-x}, y_2 = xe^{-x}; y(0) = 2,$ v'(0) = -1
- **10.** y'' 10y' + 25y = 0; $y_1 = e^{5x}$, $y_2 = xe^{5x}$; y(0) = 3, v'(0) = 13
- **11.** y'' 2y' + 2y = 0; $y_1 = e^x \cos x$, $y_2 = e^x \sin x$; y(0) = 0, y'(0) = 5
- **12.** y'' + 6y' + 13y = 0; $y_1 = e^{-3x} \cos 2x$, $y_2 = e^{-3x} \sin 2x$; y(0) = 2, y'(0) = 0
- **13.** $x^2y'' 2xy' + 2y = 0$; $y_1 = x$, $y_2 = x^2$; y(1) = 3, y'(1) = 1
- **14.** $x^2y'' + 2xy' 6y = 0$; $y_1 = x^2$, $y_2 = x^{-3}$; y(2) = 10, v'(2) = 15
- **15.** $x^2y'' xy' + y = 0; y_1 = x, y_2 = x \ln x; y(1) = 7,$ v'(1) = 2
- **16.** $x^2y'' + xy' + y = 0$; $y_1 = \cos(\ln x)$, $y_2 = \sin(\ln x)$; v(1) = 2, v'(1) = 3

The following three problems illustrate the fact that the superposition principle does not generally hold for nonlinear equations.

Differentiation yields

$$y'(x) = -c_1 e^{-x} + c_2 e^{-x} - c_2 x e^{-x},$$

so the initial conditions yield the equations

$$y(0) = c_1 = 5,$$

 $y'(0) = -c_1 + c_2 = -3,$

which imply that $c_1 = 5$ and $c_2 = 2$. Thus the desired particular solution of the initial value problem is

$$y(x) = 5e^{-x} + 2xe^{-x}.$$

This particular solution, together with several others of the form $y(x) = c_1 e^{-x} + 2x e^{-x}$, is illustrated in Fig. 5.1.10.

The characteristic equation in (18) may have either real or complex roots. The case of complex roots will be discussed in Section 5.3.

- 17. Show that y = 1/x is a solution of $y' + y^2 = 0$, but that if $c \neq 0$ and $c \neq 1$, then y = c/x is not a solution.
- **18.** Show that $y = x^3$ is a solution of $yy'' = 6x^4$, but that if $c^2 \neq 1$, then $y = cx^3$ is not a solution.
- **19.** Show that $y_1 \equiv 1$ and $y_2 = \sqrt{x}$ are solutions of yy'' + y'' = 1 $(y')^2 = 0$, but that their sum $y = y_1 + y_2$ is not a solution.

Determine whether the pairs of functions in Problems 20 through 26 are linearly independent or linearly dependent on the real line.

- **20.** $f(x) = \pi$, $g(x) = \cos^2 x + \sin^2 x$
- **21.** $f(x) = x^3$, $g(x) = x^2|x|$
- **22.** f(x) = 1 + x, g(x) = 1 + |x|
- **23.** $f(x) = xe^x$, $g(x) = |x|e^x$ **24.** $f(x) = \sin^2 x$, $g(x) = 1 \cos 2x$
- **25.** $f(x) = e^x \sin x, g(x) = e^x \cos x$
- **26.** $f(x) = 2\cos x + 3\sin x, g(x) = 3\cos x 2\sin x$
- **27.** Let y_p be a particular solution of the nonhomogeneous equation y'' + py' + qy = f(x) and let y_c be a solution of its associated homogeneous equation. Show that $y = y_c + y_p$ is a solution of the given nonhomogeneous equation.
- **28.** With $y_p = 1$ and $y_c = c_1 \cos x + c_2 \sin x$ in the notation of Problem 27, find a solution of y'' + y = 1 satisfying the initial conditions y(0) = -1 = y'(0).

Problems 29 through 32 explore the propeties of the Wronskian.

- **29.** Show that $y_1 = x^2$ and $y_2 = x^3$ are two different solutions of $x^2y'' - 4xy' + 6y = 0$, both satisfying the initial conditions y(0) = 0 = y'(0). Explain why these facts do not contradict Theorem 2 (with respect to the guaranteed uniqueness).
- **30.** (a) Show that $y_1 = x^3$ and $y_2 = |x^3|$ are linearly independent solutions on the real line of the equation $x^2 y''$ – 3xy' + 3y = 0. (b) Verify that $W(y_1, y_2)$ is identically zero. Why do these facts not contradict Theorem 3?

- **31.** Show that $y_1 = \sin x^2$ and $y_2 = \cos x^2$ are linearly independent functions, but that their Wronskian vanishes at x = 0. Why does this imply that there is *no* differential equation of the form y'' + p(x)y' + q(x)y = 0, with both *p* and *q* continuous everywhere, having both y_1 and y_2 as solutions?
- **32.** Let y_1 and y_2 be two solutions of A(x)y'' + B(x)y' + C(x)y = 0 on an open interval *I* where *A*, *B*, and *C* are continuous and A(x) is never zero. (a) Let $W = W(y_1, y_2)$. Show that

$$A(x)\frac{dW}{dx} = (y_1)(Ay_2'') - (y_2)(Ay_1'').$$

Then substitute for Ay_2'' and Ay_1'' from the original differential equation to show that

$$A(x)\frac{dW}{dx} = -B(x)W(x).$$

(b) Solve this first-order equation to deduce Abel's formula

$$W(x) = K \exp\left(-\int \frac{B(x)}{A(x)} dx\right)$$

where *K* is a constant. (c) Why does Abel's formula imply that the Wronskian $W(y_1, y_2)$ is either zero everywhere or nonzero everywhere (as stated in Theorem 3)?

Apply Theorems 5 and 6 to find general solutions of the differential equations given in Problems 33 through 42. Primes denote derivatives with respect to x.

33. $y'' - 3y' + 2y = 0$	34. $y'' + 2y' - 15y = 0$
35. $y'' + 5y' = 0$	36. $2y'' + 3y' = 0$
37. $2y'' - y' - y = 0$	38. $4y'' + 8y' + 3y = 0$
39. $4y'' + 4y' + y = 0$	40. $9y'' - 12y' + 4y = 0$
41. $6y'' - 7y' - 20y = 0$	42. $35y'' - y' - 12y = 0$

Each of Problems 43 through 48 gives a general solution y(x) of a homogeneous second-order differential equation ay'' + by' + cy = 0 with constant coefficients. Find such an equation.

43. $y(x) = c_1 + c_2 e^{-10x}$ 44. $y(x) = c_1 e^{10x} + c_2 e^{-10x}$ 45. $y(x) = c_1 e^{-10x} + c_2 x e^{-10x}$ 46. $y(x) = c_1 e^{10x} + c_2 e^{100x}$ 47. $y(x) = c_1 + c_2 x$ 48. $y(x) = e^x \left(c_1 e^{x\sqrt{2}} + c_2 e^{-x\sqrt{2}} \right)$

Problems 49 and 50 deal with the solution curves of y'' + 3y' + 2y = 0 shown in Figs. 5.1.6 and 5.1.7.

- **49.** Find the highest point on the solution curve with y(0) = 1 and y'(0) = 6 in Fig. 5.1.6.
- **50.** Figure 5.1.7 suggests that the solution curves shown all meet at a common point in the third quadrant. Assuming that this is indeed the case, find the coordinates of that point.
- 51. A second-order Euler equation is one of the form

$$ax^{2}y'' + bxy' + cy = 0$$
(22)

where *a*, *b*, *c* are constants. (a) Show that if x > 0, then the substitution $v = \ln x$ transforms Eq. (22) into the constant-coefficient linear equation

$$a\frac{d^{2}y}{dv^{2}} + (b-a)\frac{dy}{dv} + cy = 0$$
 (23)

with independent variable v. (b) If the roots r_1 and r_2 of the characteristic equation of Eq. (23) are real and distinct, conclude that a general solution of the Euler equation in (22) is $y(x) = c_1 x^{r_1} + c_2 x^{r_2}$.

Make the substitution $v = \ln x$ of Problem 51 to find general solutions (for x > 0) of the Euler equations in Problems 52–56.

52. $x^2y'' + xy' - y = 0$ **53.** $x^2y'' + 2xy' - 12y = 0$ **54.** $4x^2y'' + 8xy' - 3y = 0$ **55.** $x^2y'' + xy' = 0$ **56.** $x^2y'' - 3xy' + 4y = 0$

5.1 Application Plotting Second-Order Solution Families

Go to goo.gl/jRVU01 to download this application's computing resources including *Maple/Mathematica*/MATLAB. This application deals with the plotting by computer of families of solutions such as those illustrated in Figs. 5.1.6 and 5.1.7. Show first that the general solution of the differential equation

$$y'' + 3y' + 2y = 0 \tag{1}$$

is

$$y(x) = c_1 e^{-x} + c_2 e^{-2x}.$$
 (2)

This is equivalent to the graphing calculator result shown in Figure 5.1.11, and to the Wolfram Alpha output generated by the simple query

$$y'' + 3y' + 2y = 0$$



FIGURE 5.1.11. TI-Nspire CX CAS screen showing the general solution of y'' + 3y' + 2y = 0.



FIGURE 5.1.12. MATLAB graph of Eq. (3) with a = 2 and b = 1. Using the sliders, a and b can be changed interactively.



FIGURE 5.1.13. TI-Nspire CX CAS graph of Eq. (3) with a = 2 and b = 1. The arrows allow a and b to be changed interactively.

Next show that the particular solution of Eq. (1) satisfying y(0) = a, y'(0) = b corresponds to $c_1 = 2a + b$ and $c_2 = -a - b$, that is

$$y(x) = (2a+b)e^{-x} - (a+b)e^{-2x}.$$
(3)

For Fig. 5.1.6, we fix a = 1, leading to the particular solution

$$y(x) = (b+2)e^{-x} - (b+1)e^{-2x}.$$
(4)

The MATLAB loop

x = -1 : 0.02 : 5 % x-vector from x = -1 to x = 5
for b = -6 : 2 : 6 % for b = -6 to 6 with db = 2 do
 y = (b + 2)*exp(-x) - (b + 1)*exp(-2*x);
 plot(x,y)
end

was used to generate Fig. 5.1.6.

For Fig. 5.1.7, we instead fix b = 1, leading to the particular solution

$$y(x) = (2a+1)e^{-x} - (a+1)e^{-2x}.$$
(5)

The MATLAB loop

```
x = -2 : 0.02 : 4 % x-vector from x = -2 to x = 4
for a = -3 : 1 : 3 % for a = -3 to 3 with da = 1 do
    y = (2*a + 1)*exp(-x) - (a + 1)*exp(-2*x);
    plot(x,y)
end
```

was used to generate Fig. 5.1.7.

Computer systems, such as *Maple* and *Mathematica*, as well as graphing calculators, have commands to carry out for-loops such as the two shown here. Moreover, such systems often allow for *interactive* investigation, in which the solution curve is immediately redrawn in response to on-screen input. For example, Fig. 5.1.12 was generated using MATLAB's uicontrol command; moving the sliders allows the user to experiment with various combinations of the initial conditions y(0) = a and y'(0) = b.

The *Mathematica* command

```
Manipulate[
    Plot[(2*a+b)*Exp[-x]+(-b-a)*Exp[-2*x],
    {x,-1,5}, PlotRange -> {-5,5}],
    {a,-3,3}, {b,-6,6}]
```

produces a similar display, as does the Maple command

(after first bringing up a dialog box in which the ranges of values for a and b are specified). Likewise some graphing calculators feature a touchpad or other screen navigation method allowing the user to vary specified parameters in real time (see Fig. 5.1.13).

Begin by either reproducing Figs. 5.1.6 and 5.1.7 or by creating an interactive display that shows the graph of (3) for any desired combination of *a* and *b*. Then, for each of the following differential equations, modify your commands to examine the family of solution curves satisfying y(0) = 1, as well as the family of solution curves satisfying the initial condition y'(0) = 1.

1.
$$y'' - y = 0$$

2. $y'' - 3y' + 2y = 0$
3. $2y'' + 3y' + y = 0$
4. $y'' + y = 0$ (see Example 1)
5. $y'' + 2y' + 2y = 0$, which has general solution $y(x) = e^{-x}(c_1 \cos x + c_2 \sin x)$

5.2 General Solutions of Linear Equations

>

We now show that our discussion in Section 5.1 of second-order linear equations generalizes in a very natural way to the general *n*th-order linear differential equation of the form

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \dots + P_{n-1}(x)y' + P_n(x)y = F(x).$$
(1)

Unless otherwise noted, we will always assume that the coefficient functions $P_i(x)$ and F(x) are continuous on some open interval I (perhaps unbounded) where we wish to solve the equation. Under the additional assumption that $P_0(x) \neq 0$ at each point of I, we can divide each term in Eq. (1) by $P_0(x)$ to obtain an equation with leading coefficient 1, of the form

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = f(x).$$
(2)

The homogeneous linear equation associated with Eq. (2) is

>
$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0.$$
 (3)

Just as in the second-order case, a *homogeneous n*th-order linear differential equation has the valuable property that any superposition, or *linear combination*, of solutions of the equation is again a solution. The proof of the following theorem is essentially the same—a routine verification—as that of Theorem 1 of Section 5.1.

THEOREM 1 Principle of Superposition for Homogeneous Equations

Let y_1, y_2, \ldots, y_n be *n* solutions of the homogeneous linear equation in (3) on the interval *I*. If c_1, c_2, \ldots, c_n are constants, then the linear combination

$$y = c_1 y_1 + c_2 y_2 + \dots + c_n y_n \tag{4}$$

is also a solution of Eq. (3) on *I*.

Example 1 It is easy to verify that the three functions

 $y_1(x) = e^{-3x}$, $y_2(x) = \cos 2x$, and $y_3(x) = \sin 2x$

are all solutions of the homogeneous third-order equation

 $y^{(3)} + 3y'' + 4y' + 12y = 0$

on the entire real line. Theorem 1 tells us that any linear combination of these solutions, such as

$$y(x) = -3y_1(x) + 3y_2(x) - 2y_3(x) = -3e^{-3x} + 3\cos 2x - 2\sin 2x,$$

is also a solution on the entire real line. We will see that, conversely, every solution of the differential equation of this example is a linear combination of the three particular solutions y_1 , y_2 , and y_3 . Thus a general solution is given by

$$y(x) = c_1 e^{-3x} + c_2 \cos 2x + c_3 \sin 2x.$$

Existence and Uniqueness of Solutions

We saw in Section 5.1 that a particular solution of a *second-order* linear differential equation is determined by *two* initial conditions. Similarly, a particular solution of an *n*th-order linear differential equation is determined by n initial conditions. The following theorem, proved in the Appendix, is the natural generalization of Theorem 2 of Section 5.1.

THEOREM 2 Existence and Uniqueness for Linear Equations

Suppose that the functions $p_1, p_2, ..., p_n$, and f are continuous on the open interval I containing the point a. Then, given n numbers $b_0, b_1, ..., b_{n-1}$, the *n*th-order linear equation (Eq. (2))

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = f(x)$$

has a unique (that is, one and only one) solution on the entire interval I that satisfies the n initial conditions

$$y(a) = b_0, \quad y'(a) = b_1, \quad \dots, \quad y^{(n-1)}(a) = b_{n-1}.$$
 (5)

Equation (2) and the conditions in (5) constitute an *n*th-order **initial value problem**. Theorem 2 tells us that any such initial value problem has a unique solution on the *whole* interval I where the coefficient functions in (2) are continuous. It tells us nothing, however, about how to find this solution. In Section 5.3 we will see how to construct explicit solutions of initial value problems in the *constant*-coefficient case that occurs often in applications.

Example 1

Continued We saw earlier that



FIGURE 5.2.1. The particular solution y(x) = $-3e^{-3x} + 3\cos 2x - 2\sin 2x$.

$$y(x) = -3e^{-3x} + 3\cos 2x - 2\sin 2x$$

is a solution of

$$y^{(3)} + 3y'' + 4y' + 12y = 0$$

on the entire real line. This particular solution has initial values y(0) = 0, y'(0) = 5, and y''(0) = -39, and Theorem 2 implies that there is no other solution with these same initial values. Note that its graph (in Fig. 5.2.1) looks periodic on the right. Indeed, because of the negative exponent, we see that $y(x) \approx 3 \cos 2x - 2 \sin 2x$ for large positive x.

Remark Because its general solution involves the three arbitrary constants c_1 , c_2 , and c_3 , the third-order equation in Example 1 has a "threefold infinity" of solutions, including three families of especially simple solutions:

- $y(x) = c_1 e^{-3x}$ (obtained from the general solution with $c_2 = c_3 = 0$),
- $y(x) = c_2 \cos 2x$ (with $c_1 = c_3 = 0$), and
- $y(x) = c_3 \sin 2x$ (with $c_1 = c_2 = 0$).

Alternatively, Theorem 2 suggests a threefold infinity of particular solutions corresponding to independent choices of the three initial values $y(0) = b_0$, $y'(0) = b_1$, and $y''(0) = b_2$. Figures 5.2.2 through 5.2.4 illustrate three corresponding families of solutions—for each of which, two of these three initial values are zero.



FIGURE 5.2.2. Solutions of $y^{(3)} + 3y'' + 4y' + 12y = 0$ with y'(0) = y''(0) = 0 but with different values for y(0).



FIGURE 5.2.3. Solutions of $y^{(3)} + 3y'' + 4y' + 12y = 0$ with y(0) = y''(0) = 0 but with different values for y'(0).



FIGURE 5.2.4. Solutions of $y^{(3)} + 3y'' + 4y' + 12y = 0$ with y(0) = y'(0) = 0 but with different values for y''(0).

Note that Theorem 2 implies that the *trivial* solution $y(x) \equiv 0$ is the only solution of the *homogeneous* equation

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$
(3)

that satisfies the trivial initial conditions

$$y(a) = y'(a) = \dots = y^{(n-1)}(a) = 0.$$

Example 2 It is easy to verify that

$$y_1(x) = x^2$$
 and $y_2(x) = x^3$

are two different solutions of

$$x^2y'' - 4xy' + 6y = 0,$$

and that both satisfy the initial conditions y(0) = y'(0) = 0. Why does this not contradict the uniqueness part of Theorem 2? It is because the leading coefficient in this differential equation vanishes at x = 0, so this equation cannot be written in the form of Eq. (3) with coefficient functions *continuous* on an open interval containing the point x = 0.

Linearly Independent Solutions

On the basis of our knowledge of general solutions of second-order linear equations, we anticipate that a general solution of the *homogeneous n*th-order linear equation

>
$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$
 (3)

will be a linear combination

$$y = c_1 y_1 + c_2 y_2 + \dots + c_n y_n, \tag{4}$$

where y_1, y_2, \ldots, y_n are particular solutions of Eq. (3). But these *n* particular solutions must be "sufficiently independent" that we can always choose the coefficients c_1, c_2, \ldots, c_n in (4) to satisfy arbitrary initial conditions of the form in (5). The question is this: What should be meant by *independence* of three or more functions?

Recall that *two* functions f_1 and f_2 are linearly *dependent* if one is a constant multiple of the other; that is, if either $f_1 = kf_2$ or $f_2 = kf_1$ for some constant k. If we write these equations as

$$(1)f_1 + (-k)f_2 = 0$$
 or $(k)f_1 + (-1)f_2 = 0$,

we see that the linear dependence of f_1 and f_2 implies that there exist two constants c_1 and c_2 not both zero such that

$$c_1 f_1 + c_2 f_2 = 0. (6)$$

Conversely, if c_1 and c_2 are not both zero, then Eq. (6) certainly implies that f_1 and f_2 are linearly dependent.

In analogy with Eq. (6), we say that *n* functions f_1, f_2, \ldots, f_n are *linearly dependent* provided that some *nontrivial* linear combination

$$c_1f_1 + c_2f_2 + \dots + c_nf_n$$

of them vanishes identically; *nontrivial* means that *not all* of the coefficients c_1, c_2, \ldots, c_n are zero (although some of them may be zero).

DEFINITION Linear Dependence of Functions

The *n* functions f_1, f_2, \ldots, f_n are said to be **linearly dependent** on the interval *I* provided that there exist constants c_1, c_2, \ldots, c_n not all zero such that

$$c_1 f_1 + c_2 f_2 + \dots + c_n f_n = 0 \tag{7}$$

on *I*; that is,

$$c_1 f_1(x) + c_2 f_2(x) + \dots + c_n f_n(x) = 0$$

for all x in I.

Thus linear dependence of functions on an interval I is precisely analogous to linear dependence of ordinary vectors (Section 4.3).

If not all the coefficients in Eq. (7) are zero, then clearly we can solve for at least one of the functions as a linear combination of the others, and conversely. Thus the functions f_1, f_2, \ldots, f_n are linearly dependent if and only if at least one of them is a linear combination of the others.

Example 3 The functions

$$f_1(x) = \sin 2x$$
, $f_2(x) = \sin x \cos x$, and $f_3(x) = e^x$

are linearly dependent on the real line because

$$(1) f_1 + (-2) f_2 + (0) f_3 = 0$$

(by the familiar trigonometric identity $\sin 2x = 2 \sin x \cos x$).

The *n* functions $f_1, f_2, ..., f_n$ are called **linearly independent** on the interval *I* provided that they are not linearly dependent there. Equivalently, they are linearly independent on *I* provided that the identity

$$c_1 f_1 + c_2 f_2 + \dots + c_n f_n = 0 \tag{7}$$

holds on *I* only in the trivial case

$$c_1 = c_2 = \cdots = c_n = 0;$$

that is, *no* nontrivial linear combination of these functions vanishes on *I*. Put yet another way, the functions f_1, f_2, \ldots, f_n are linearly independent if no one of them is a linear combination of the others. (Why?)

Sometimes one can show that n given functions are linearly dependent by finding, as in Example 3, nontrivial values of the coefficients so that Eq. (7) holds. But in order to show that n given functions are linearly independent, we must prove that nontrivial values of the coefficients *cannot* be found, and this is seldom easy to do in any direct or obvious manner.

Fortunately, in the case of *n* solutions of a homogeneous *n*th-order linear equation, there is a tool that makes the determination of their linear dependence or independence a routine matter in many examples. This tool is the Wronskian determinant, which we introduced (for the case n = 2) in Section 5.1. Suppose that the *n* functions f_1, f_2, \ldots, f_n are each n - 1 times differentiable. Then their **Wronskian** is the $n \times n$ determinant

$$W = \begin{vmatrix} f_1 & f_2 & \cdots & f_n \\ f'_1 & f'_2 & \cdots & f'_n \\ \vdots & \vdots & & \vdots \\ f_1^{(n-1)} & f_2^{(n-1)} & \cdots & f_n^{(n-1)} \end{vmatrix}.$$
(8)

We write $W(f_1, f_2, ..., f_n)$ or W(x), depending on whether we wish to emphasize the functions or the point x at which their Wronskian is to be evaluated. The Wronskian is named after the Polish mathematician J. M. H. Wronski (1778–1853).

We saw in Section 5.1 that the Wronskian of two linearly dependent functions vanishes identically. More generally, the Wronskian of n linearly dependent functions f_1, f_2, \ldots, f_n is identically zero. To prove this, assume that Eq. (7) holds on the interval I for some choice of the constants c_1, c_2, \ldots, c_n not all zero. We then differentiate this equation n - 1 times in succession, obtaining the n equations

$$c_{1}f_{1}(x) + c_{2}f_{2}(x) + \dots + c_{n}f_{n}(x) = 0,$$

$$c_{1}f_{1}'(x) + c_{2}f_{2}'(x) + \dots + c_{n}f_{n}'(x) = 0,$$

$$\vdots$$

$$c_{1}f_{1}^{(n-1)}(x) + c_{2}f_{2}^{(n-1)}(x) + \dots + c_{n}f_{n}^{(n-1)}(x) = 0,$$

(9)

which hold for all x in I. We recall from Theorem 7 in Section 3.5 that a *homo*geneous $n \times n$ linear system of equations has a nontrivial solution if and only if its coefficient matrix is *not* invertible, which by Theorem 2 in Section 3.6 is so if and only if the coefficient determinant vanishes. In Eq. (9) the unknowns are the constants c_1, c_2, \ldots, c_n and the determinant of coefficients is simply the Wronskian $W(f_1, f_2, \ldots, f_n)$ evaluated at the typical point x of I. Because we know that the c_i are not all zero, it follows that $W(x) \equiv 0$, as we wanted to prove.

Therefore, to show that the functions f_1, f_2, \ldots, f_n are *linearly independent* on the interval *I*, it suffices to show that their Wronskian is nonzero at just one point of *I*.

Example 4

Show that the functions $y_1(x) = e^{-3x}$, $y_2(x) = \cos 2x$, and $y_3(x) = \sin 2x$ (of Example 1) are linearly independent.

Solution Their Wronskian is

$$W = \begin{vmatrix} e^{-3x} & \cos 2x & \sin 2x \\ -3e^{-3x} & -2\sin 2x & 2\cos 2x \\ 9e^{-3x} & -4\cos 2x & -4\sin 2x \end{vmatrix}$$
$$= e^{-3x} \begin{vmatrix} -2\sin 2x & 2\cos 2x \\ -4\cos 2x & -4\sin 2x \end{vmatrix} + 3e^{-3x} \begin{vmatrix} \cos 2x & \sin 2x \\ -4\cos 2x & -4\sin 2x \end{vmatrix}$$
$$+ 9e^{-3x} \begin{vmatrix} \cos 2x & \sin 2x \\ -2\sin 2x & 2\cos 2x \end{vmatrix} = 26e^{-3x} \neq 0.$$

Because $W \neq 0$ everywhere, it follows that y_1 , y_2 , and y_3 are linearly independent on any open interval (including the entire real line).

Example 5

Show first that the three solutions

$$y_1(x) = x$$
, $y_2(x) = x \ln x$, and $y_3(x) = x^2$

of the third-order equation

$$x^{3}y^{(3)} - x^{2}y'' + 2xy' - 2y = 0$$
⁽¹⁰⁾

are linearly independent on the open interval x > 0. Then find a particular solution of Eq. (10) that satisfies the initial conditions

$$y(1) = 3, y'(1) = 2, y''(1) = 1.$$
 (11)

Solution Note that for x > 0, we could divide each term in (10) by x^3 to obtain a homogeneous linear equation of the standard form in (3). When we compute the Wronskian of the three given solutions, we find that

$$W = \begin{vmatrix} x & x \ln x & x^2 \\ 1 & 1 + \ln x & 2x \\ 0 & \frac{1}{x} & 2 \end{vmatrix} = x$$

Thus $W(x) \neq 0$ for x > 0, so y_1 , y_2 , and y_3 are linearly independent on the interval x > 0. To find the desired particular solution, we impose the initial conditions in (11) on

$$y(x) = c_1 x + c_2 x \ln x + c_3 x^2,$$

$$y'(x) = c_1 + c_2 (1 + \ln x) + 2c_3 x,$$

$$y''(x) = 0 + \frac{c_2}{x} + 2c_3.$$

This yields the simultaneous equations

$$y(1) = c_1 + c_3 = 3,$$

$$y'(1) = c_1 + c_2 + 2c_3 = 2,$$

$$y''(1) = c_2 + 2c_3 = 1;$$

we solve to find $c_1 = 1$, $c_2 = -3$, and $c_3 = 2$. Thus the particular solution in question is

$$y(x) = x - 3x \ln x + 2x^2.$$

Provided that $W(y_1, y_2, ..., y_n) \neq 0$, it turns out (Theorem 4) that we can always find values of the coefficients in the linear combination

$$y = c_1 y_1 + c_2 y_2 + \dots + c_n y_n$$

that satisfy any given initial conditions of the form in (5). Theorem 3 provides the necessary nonvanishing of W in the case of linearly independent solutions.

THEOREM 3 Wronskians of Solutions

Suppose that $y_1, y_2, ..., y_n$ are *n* solutions of the homogeneous *n*th-order linear equation

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$
(3)

on an open interval I, where each p_i is continuous. Let

$$W = W(y_1, y_2, \ldots, y_n).$$

(a) If y_1, y_2, \ldots, y_n are linearly dependent, then $W \equiv 0$ on *I*.

(b) If y_1, y_2, \ldots, y_n are linearly independent, then $W \neq 0$ at each point of *I*.

Thus there are just two possibilities: Either W = 0 everywhere on I, or $W \neq 0$ everywhere on I.

Proof: We have already proven part (a). To prove part (b), it is sufficient to assume that W(a) = 0 at some point of *I*, and show this implies that the solutions y_1, y_2, \ldots, y_n are linearly dependent. But W(a) is simply the determinant of coefficients of the system of *n* homogeneous linear equations

$$c_{1}y_{1}(a) + c_{2}y_{2}(a) + \dots + c_{n}y_{n}(a) = 0,$$

$$c_{1}y'_{1}(a) + c_{2}y'_{2}(a) + \dots + c_{n}y'_{n}(a) = 0,$$

$$\vdots$$
(12)

$$c_1 y_1^{(n-1)}(a) + c_2 y_2^{(n-1)}(a) + \dots + c_n y_n^{(n-1)}(a) = 0$$

in the *n* unknowns c_1, c_2, \ldots, c_n . Because W(a) = 0, the basic fact from linear algebra quoted just after (9) implies that the equations in (12) have a nontrivial solution. That is, the numbers c_1, c_2, \ldots, c_n are not all zero.

We now use these values to define the particular solution

$$Y(x) = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x)$$
(13)

of Eq. (3). The equations in (12) then imply that Y satisfies the trivial initial conditions

$$Y(a) = Y'(a) = \dots = Y^{(n-1)}(a) = 0.$$

Theorem 2 (uniqueness) therefore implies that $Y(x) \equiv 0$ on *I*. In view of (13) and the fact that c_1, c_2, \ldots, c_n are not all zero, this is the desired conclusion that the solutions y_1, y_2, \ldots, y_n are linearly dependent. This completes the proof of Theorem 3.

Remark According to Problem 35, the Wronskian of Theorem 3 satisfies the first-order equation $W' = -p_1(x)W$. Solution of this equation yields *Abel's formula*

$$W(x) = K \exp\left(-\int p_1(x) \, dx\right)$$

for the homogeneous linear equation in (3). According as the constant K = 0 or $K \neq 0$, it follows immediately from Abel's formula that either W vanishes everywhere or W is nonzero everywhere.

General Solutions

We can now show that, given any fixed set of *n* linearly independent solutions of a *homogeneous n*th-order equation, *every* (other) solution of the equation can be expressed as a linear combination of those *n* particular solutions. Using the fact from Theorem 3 that the Wronskian of *n* linearly independent solutions is nonzero, the proof of the following theorem is essentially the same as the proof of Theorem 4 of Section 5.1 (the case n = 2).

THEOREM 4 General Solutions of Homogeneous Equations

Let y_1, y_2, \ldots, y_n be *n* linearly independent solutions of the homogeneous equation

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$
(3)

on an open interval I where the p_i are continuous. If Y is any solution whatsoever of Eq. (3), then there exist numbers c_1, c_2, \ldots, c_n such that

$$Y(x) = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x)$$

for all x in I.

Remark Theorem 4 tells us that, once we have found *n* linearly independent solutions y_1, y_2, \ldots, y_n of the *n*th-order homogeneous linear equation in (3), we really have found *all* of its solutions. For then the solution space \mathscr{S} of the equation is an *n*-dimensional vector space with basis $\{y_1, y_2, \ldots, y_n\}$. Because *every* solution of (3) can be expressed as a linear combination of the form

$$y = c_1 y_1 + c_2 y_2 + \dots + c_n y_n, \tag{14}$$

we call such a linear combination of *n* linearly independent particular solutions a **general** solution of the homogeneous linear differential equation.

Example 6 According to Example 4, the particular solutions $y_1(x) = e^{-3x}$, $y_2(x) = \cos 2x$, and $y_3(x) = \sin 2x$ of the linear differential equation $y^{(3)} + 3y'' + 4y' + 12y = 0$ are linearly independent. Now Theorem 2 says that—given b_0 , b_1 , and b_2 —there exists a particular solution y(x) satisfying the initial conditions $y(0) = b_0$, $y'(0) = b_1$, and $y''(0) = b_2$. Hence Theorem 4 implies that *this* particular solution is a linear combination of y_1 , y_2 , and y_3 . That is, there exist coefficients c_1 , c_2 , and c_3 such that

$$y(x) = c_1 e^{-3x} + c_2 \cos 2x + c_3 \sin 2x.$$

Upon successive differentiation and substitution of x = 0, we discover that to find these coefficients, we need only solve the three linear equations

$$c_1 + c_2 = b_0,$$

 $-3c_1 + 2c_3 = b_1,$
 $9c_1 - 4c_2 = b_2.$

(See the application for this section.)

Nonhomogeneous Equations

We now consider the nonhomogeneous nth-order linear differential equation

>
$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = f(x)$$
 (2)

with associated homogeneous equation

>
$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0.$$
 (3)

Suppose that a single fixed particular solution y_p of the nonhomogeneous equation in (2) is known, and that Y is any other solution of Eq. (2). If $y_c = Y - y_p$, then subsitution of y_c in the differential equation gives (using the linearity of differentiation)

$$y_{c}^{(n)} + p_{1}y_{c}^{(n-1)} + \dots + p_{n-1}y_{c}' + p_{n}y_{c}$$

$$= \left[(Y^{(n)} + p_{1}Y^{(n-1)} + \dots + p_{n-1}Y' + p_{n}Y \right]$$

$$- \left[(y_{p}^{(n)} + p_{1}y_{p}^{(n-1)} + \dots + p_{n-1}y_{p}' + p_{n}y_{p} \right]$$

$$= f(x) - f(x) = 0.$$

Thus $y_c = Y - y_p$ is a solution of the associated homogeneous equation in (3). Then

$$Y = y_c + y_p, \tag{14}$$

and it follows from Theorem 4 that

$$y_c = c_1 y_1 + c_2 y_2 + \dots + c_n y_n, \tag{15}$$

where y_1, y_2, \ldots, y_n are linearly independent solutions of the associated *homogeneous* equation. We call y_c a **complementary function** of the nonhomogeneous equation and have thus proved that a *general solution* of the nonhomogeneous equation in (2) is the sum of its complementary function y_c and a single particular solution y_p of Eq. (2).

THEOREM 5 Solutions of Nonhomogeneous Equations

Let y_p be a particular solution of the nonhomogeneous equation in (2) on an open interval *I* where the functions p_i and *f* are continuous. Let y_1, y_2, \ldots, y_n be linearly independent solutions of the associated homogeneous equation in (3). If *Y* is any solution whatsoever of Eq. (2) on *I*, then there exist numbers c_1, c_2, \ldots, c_n such that

$$Y(x) = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x) + y_p(x)$$
(16)

for all x in I.

Example 7 It is evident that $y_p = 3x$ is a particular solution of the equation

$$y'' + 4y = 12x,$$
 (17)

and that $y_c(x) = c_1 \cos 2x + c_2 \sin 2x$ is its complementary solution. Find a solution of Eq. (17) that satisfies the initial conditions y(0) = 5, y'(0) = 7.

Solution The general solution of Eq. (17) is

$$y(x) = c_1 \cos 2x + c_2 \sin 2x + 3x.$$

Now

$$y'(x) = -2c_1 \sin 2x + 2c_2 \cos 2x + 3.$$

Hence the initial conditions give

$$y(0) = c_1 = 5,$$

 $y'(0) = 2c_2 + 3 = 7.$

We find that $c_1 = 5$ and $c_2 = 2$. Thus the desired solution is

 $y(x) = 5\cos 2x + 2\sin 2x + 3x.$

5.2 Problems

In Problems 1 through 6, show directly that the given functions are linearly dependent on the real line. That is, find a nontrivial linear combination of the given functions that vanishes identically.

- 1. f(x) = 2x, $g(x) = 3x^2$, $h(x) = 5x 8x^2$
- **2.** f(x) = 5, $g(x) = 2 3x^2$, $h(x) = 10 + 15x^2$
- 3. $f(x) = 0, g(x) = \sin x, h(x) = e^x$
- 4. $f(x) = 17, g(x) = 2\sin^2 x, h(x) = 3\cos^2 x$
- 5. $f(x) = 17, g(x) = \cos^2 x, h(x) = \cos 2x$
- 6. $f(x) = e^x$, $g(x) = \cosh x$, $h(x) = \sinh x$

In Problems 7 through 12, use the Wronskian to prove that the given functions are linearly independent on the indicated interval.

- 7. f(x) = 1, g(x) = x, $h(x) = x^2$; the real line
- 8. $f(x) = e^x$, $g(x) = e^{2x}$, $h(x) = e^{3x}$; the real line
- **9.** $f(x) = e^x$, $g(x) = \cos x$, $h(x) = \sin x$; the real line
- **10.** $f(x) = e^x$, $g(x) = x^{-2}$, $h(x) = x^{-2} \ln x$; x > 0
- **11.** $f(x) = x, g(x) = xe^x, h(x) = x^2e^x$; the real line
- 12. $f(x) = x, g(x) = \cos(\ln x), h(x) = \sin(\ln x); x > 0$

In Problems 13 through 20, a third-order homogeneous linear equation and three linearly independent solutions are given. Find a particular solution satisfying the given initial conditions.

- **13.** $y^{(3)} + 2y'' y' 2y = 0$; y(0) = 1, y'(0) = 2, y''(0) = 0; $y_1 = e^x$, $y_2 = e^{-x}$, $y_3 = e^{-2x}$
- **14.** $y^{(3)} 6y'' + 11y' 6y = 0$; y(0) = 0, y'(0) = 0, y''(0) = 3; $y_1 = e^x$, $y_2 = e^{2x}$, $y_3 = e^{3x}$
- **15.** $y^{(3)} 3y'' + 3y' y = 0; y(0) = 2, y'(0) = 0, y''(0) = 0;$ $y_1 = e^x, y_2 = xe^x, y_3 = x^2e^x$
- **16.** $y^{(3)} 5y'' + 8y' 4y = 0; y(0) = 1, y'(0) = 4, y''(0) = 0;$ $y_1 = e^x, y_2 = e^{2x}, y_3 = xe^{2x}$
- **17.** $y^{(3)} + 9y' = 0$; y(0) = 3, y'(0) = -1, y''(0) = 2; $y_1 = 1$, $y_2 = \cos 3x$, $y_3 = \sin 3x$
- **18.** $y^{(3)} 3y'' + 4y' 2y = 0; y(0) = 1, y'(0) = 0, y''(0) = 0;$ $y_1 = e^x, y_2 = e^x \cos x, y_3 = e^x \sin x.$

19. $x^{3}y^{(3)} - 3x^{2}y'' + 6xy' - 6y = 0; y(1) = 6, y'(1) = 14,$ $y''(1) = 22; y_{1} = x, y_{2} = x^{2}, y_{3} = x^{3}$ **20.** $x^{3}y^{(3)} + 6x^{2}y'' + 4xy' - 4y = 0; y(1) = 1, y'(1) = 5,$ $y''(1) = -11; y_{1} = x, y_{2} = x^{-2}, y_{3} = x^{-2} \ln x$

In Problems 21 through 24, a nonhomogeneous differential equation, a complementary solution y_c , and a particular solution y_p are given. Find a solution satisfying the given initial conditions.

- **21.** y'' + y = 3x; y(0) = 2, y'(0) = -2; $y_c = c_1 \cos x + c_2 \sin x$; $y_p = 3x$
- **22.** y'' 4y = 12; y(0) = 0, y'(0) = 10; $y_c = c_1 e^{2x} + c_2 e^{-2x}; y_p = -3$
- **23.** y'' 2y' 3y = 6; y(0) = 3, y'(0) = 11; $y_c = c_1 e^{-x} + c_2 e^{3x}$; $y_p = -2$
- **24.** y'' 2y' + 2y = 2x; y(0) = 4, y'(0) = 8; $y_c = c_1 e^x \cos x + c_2 e^x \sin x$; $y_p = x + 1$
- **25.** Let Ly = y'' + py' + qy. Suppose that y_1 and y_2 are two functions such that

$$Ly_1 = f(x)$$
 and $Ly_2 = g(x)$.

Show that their sum $y = y_1 + y_2$ satisfies the nonhomogeneous equation Ly = f(x) + g(x).

26. (a) Find by inspection particular solutions of the two nonhomogeneous equations

$$y'' + 2y = 4$$
 and $y'' + 2y = 6x$.

(b) Use the method of Problem 25 to find a particular solution of the differential equation y'' + 2y = 6x + 4.

27. Prove directly that the functions

 $f_1(x) \equiv 1$, $f_2(x) = x$, and $f_3(x) = x^2$

are linearly independent on the whole real line. (Suggestion: Assume that $c_1 + c_2x + c_3x^2 = 0$. Differentiate this equation twice, and conclude from the equations you get that $c_1 = c_2 = c_3 = 0$.)

28. Generalize the method of Problem 27 to prove directly that the functions

$$f_0(x) \equiv 1, \ f_1(x) = x, \ f_2(x) = x^2, \dots, \ f_n(x) = x^n$$

are linearly independent on the real line.

29. Use the result of Problem 28 and the definition of linear independence to prove directly that, for any constant r, the functions

$$f_0(x) = e^{rx}, \quad f_1(x) = xe^{rx}, \quad \dots, \quad f_n(x) = x^n e^{rx}$$

are linearly independent on the whole real line.

30. Verify that $y_1 = x$ and $y_2 = x^2$ are linearly independent solutions on the entire real line of the equation

$$x^2y'' - 2xy' + 2y = 0,$$

but that $W(x, x^2)$ vanishes at x = 0. Why do these observations not contradict part (b) of Theorem 3?

31. This problem indicates why we can impose *only n* initial conditions on a solution of an *n*th-order linear differential equation. (a) Given the equation

$$y'' + py' + qy = 0,$$

explain why the value of y''(a) is determined by the values of y(a) and y'(a). (b) Prove that the equation

$$y'' - 2y' - 5y = 0$$

has a solution satisfying the conditions

$$y(0) = 1$$
, $y'(0) = 0$, and $y''(0) = C$

if and only if C = 5.

32. Prove that an *n*th-order homogeneous linear differential equation satisfying the hypotheses of Theorem 2 has *n* linearly independent solutions y_1, y_2, \ldots, y_n . (*Suggestion:* Let y_i be the unique solution such that

$$y_i^{(i-1)}(a) = 1$$
 and $y_i^{(k)}(a) = 0$ if $k \neq i - 1$.)

33. Suppose that the three numbers r_1 , r_2 , and r_3 are distinct. Show that the three functions $\exp(r_1x)$, $\exp(r_2x)$, and $\exp(r_3x)$ are linearly independent by showing that their Wronskian

$$W = \exp[(r_1 + r_2 + r_3)x] \cdot \begin{vmatrix} 1 & 1 & 1 \\ r_1 & r_2 & r_3 \\ r_1^2 & r_2^2 & r_3^2 \end{vmatrix}$$

is nonzero for all x.

34. Assume as known that the Vandermonde determinant

$$V = \begin{vmatrix} 1 & 1 & \cdots & 1 \\ r_1 & r_2 & \cdots & r_n \\ r_1^2 & r_2^2 & \cdots & r_n^2 \\ \vdots & \vdots & & \vdots \\ r_1^{n-1} & r_2^{n-1} & \cdots & r_n^{n-1} \end{vmatrix}$$

is nonzero if the numbers r_1, r_2, \ldots, r_n are distinct. Prove by the method of Problem 33 that the functions

$$f_i(x) = \exp(r_i x), \quad 1 \le i \le n$$

are linearly independent.

35. According to Problem 32 of Section 5.1, the Wronskian $W(y_1, y_2)$ of two solutions of the second-order equation

$$y'' + p_1(x)y' + p_2(x)y = 0$$

is given by Abel's formula

$$W(x) = K \exp\left(-\int p_1(x) \, dx\right)$$

for some constant *K*. It can be shown that the Wronskian of *n* solutions y_1, y_2, \ldots, y_n of the *n*th-order equation

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$

satisfies the same identity. Prove this for the case n = 3 as follows: (a) The derivative of a determinant of functions is the sum of the determinants obtained by separately differentiating the rows of the original determinant. Conclude that

$$W' = \begin{vmatrix} y_1 & y_2 & y_3 \\ y'_1 & y'_2 & y'_3 \\ y_1^{(3)} & y_2^{(3)} & y_3^{(3)} \end{vmatrix}$$

(**b**) Substitute for $y_1^{(3)}$, $y_2^{(3)}$, and $y_3^{(3)}$ from the equation

$$y^{(3)} + p_1 y'' + p_2 y' + p_3 y = 0,$$

and then show that $W' = -p_1 W$. Integration now gives Abel's formula.

36. Suppose that one solution $y_1(x)$ of the homogeneous second-order linear differential equation

$$y'' + p(x)y' + q(x)y = 0$$
 (18)

is known (on an interval *I* where *p* and *q* are continuous functions). The method of **reduction of order** consists of substituting $y_2(x) = v(x)y_1(x)$ in (18) and attempting to determine the function v(x) so that $y_2(x)$ is a second linearly independent solution of (18). After substituting $y = v(x)y_1(x)$ in Eq. (18), use the fact that $y_1(x)$ is a solution to deduce that

$$y_1v'' + (2y_1' + py_1)v' = 0.$$
(19)

If $y_1(x)$ is known, then (19) is a separable equation that is readily solved for the derivative v'(x) of v(x). Integration of v'(x) then gives the desired (nonconstant) function v(x). **37.** Before applying Eq. (19) with a given homogeneous second-order linear differential equation and a known solution $y_1(x)$, the equation must first be written in the form of (18) with leading coefficient 1 in order to correctly determine the coefficient function p(x). Frequently it is more convenient to simply substitute $y = v(x)y_1(x)$ in the given differential equation and then proceed directly to find v(x). Thus, starting with the readily verified solution $y_1(x) = x^3$ of the equation

$$x^2y'' - 5xy' + 9y = 0 \quad (x > 0),$$

substitute $y = vx^3$ and deduce that xv'' + v' = 0. Thence solve for $v(x) = C \ln x$, and thereby obtain (with C = 1) the second solution $y_2(x) = x^3 \ln x$.

In each of Problems 38 through 42, a differential equation and one solution y_1 are given. Use the method of reduction of order as in Problem 37 to find a second linearly independent solution y_2 .

38.
$$x^2y'' + xy' - 9y = 0$$
 (x > 0); $y_1(x) = x^3$

- **39.** $4y'' 4y' + y = 0; y_1(x) = e^{x/2}$
- **40.** $x^2y'' x(x+2)y' + (x+2)y = 0$ (x > 0); $y_1(x) = x$
- **41.** (x + 1)y'' (x + 2)y' + y = 0 $(x > -1); y_1(x) = e^x$
- **42.** $(1-x^2)y'' + 2xy' 2y = 0$ (-1 < x < 1); $y_1(x) = x$
- **43.** First note that $y_1(x) = x$ is one solution of Legendre's equation of order 1,

$$(1 - x^2)y'' - 2xy' + 2y = 0.$$

Then use the method of reduction of order to derive the second solution

$$y_2(x) = 1 - \frac{x}{2} \ln \frac{1+x}{1-x}$$
 (for $-1 < x < 1$).

44. First verify by substitution that $y_1(x) = x^{-1/2} \cos x$ is one solution (for x > 0) of Bessel's equation of order $\frac{1}{2}$,

$$x^{2}y'' + xy' + (x^{2} - \frac{1}{4})y = 0.$$

Then derive by reduction of order the second solution $y_2(x) = x^{-1/2} \sin x$.

5.2 Application Plotting Third-Order Solution Families

Go to goo.gl/rj9zB2 to download this application's computing resources including *Maple/Mathematica*/MATLAB. This application deals with the plotting by computer of families of solutions such as those illustrated in Figs. 5.2.2 through 5.2.4. We know from Example 6 that the general solution of

$$y^{(3)} + 3y'' + 4y' + 12y = 0$$
(1)

is

$$y(x) = c_1 e^{-3x} + c_2 \cos 2x + c_3 \sin 2x.$$
 (2)

For Fig. 5.2.2, use the method of Example 6 to show that the particular solution of Eq. (1) satisfying the initial conditions y(0) = a, y'(0) = 0, and y''(0) = 0 is given by

$$y(x) = \frac{a}{13} \left(4e^{-3x} + 9\cos 2x + 6\sin 2x \right).$$
(3)

The MATLAB loop

x = -1.5 : 0.02 : 5 % x-vector from x = -1.5 to x = 5
for a = -3 : 1 : 3 % for a = -3 to 3 with da = 1 do
 c1 = 4*a/13;
 c2 = 9*a/13;
 c3 = 6*a/13;
 y = c1*exp(-3*x) + c2*cos(2*x) + c3*sin(2*x);
 plot(x,y)
end

was used to generate Fig. 5.2.2.

For Fig. 5.2.3, show that the particular solution of Eq. (1) satisfying the initial conditions y(0) = 0, y'(0) = b, and y''(0) = 0 is given by

$$y(x) = \frac{b}{2}\sin 2x,\tag{4}$$

and alter the preceding for-loop accordingly.

For Fig. 5.2.4, show that the particular solution of Eq. (1) satisfying the initial conditions y(0) = 0, y'(0) = 0, and y''(0) = c is given by

$$y(x) = \frac{c}{26} \left(2e^{-3x} - 2\cos 2x + 3\sin 2x \right).$$
 (5)

Computer algebra systems such as *Maple* and *Mathematica*, as well as graphing calculators, have commands to carry out for-loops such as the one shown here. Begin by reproducing Figs. 5.2.2 through 5.2.4. Then plot similar families of solution curves for the differential equations in Problems 13 through 20.

5.3 Homogeneous Equations with Constant Coefficients

In Section 5.2 we saw that a general solution of an *n*th-order homogeneous linear equation is a linear combination of *n* linearly independent particular solutions, but we said little about how actually to find even a single solution. The solution of a linear differential equation with *variable* coefficients ordinarily requires numerical methods (Chapter 2) or infinite series methods (Chapter 8). But we can now show how to find, explicitly and in a rather straightforward way, *n* linearly independent solutions of a given *n*th-order linear equation if it has *constant* coefficients. The general such equation may be written in the form

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_2 y'' + a_1 y' + a_0 y = 0,$$
 (1)

where the coefficients $a_0, a_1, a_2, \ldots, a_n$ are real constants with $a_n \neq 0$.

The Characteristic Equation

We first look for a single solution of Eq. (1), and begin with the observation that

$$\frac{d^k}{dx^k}(e^{rx}) = r^k e^{rx},\tag{2}$$

so any derivative of e^{rx} is a constant multiple of e^{rx} . Hence, if we substituted $y = e^{rx}$ in Eq. (1), each term would be a constant multiple of e^{rx} , with the constant coefficients depending on r and the coefficients a_k . This suggests that we try to find r so that all these multiples of e^{rx} will have sum zero, in which case $y = e^{rx}$ will be a solution of Eq. (1).

For example, in Section 5.1 we substituted $y = e^{rx}$ in the second-order equation

$$ay'' + by' + cy = 0$$

to derive the characteristic equation

$$ar^2 + br + c = 0$$

that r must satisfy.

To carry out this technique in the general case, we substitute $y = e^{rx}$ in Eq. (1), and with the aid of Eq. (2) we find the result to be

$$a_n r^n e^{rx} + a_{n-1} r^{n-1} e^{rx} + \dots + a_2 r^2 e^{rx} + a_1 r e^{rx} + a_0 e^{rx} = 0;$$

that is,

 $\mathbf{>}$

$$e^{rx} \left(a_n r^n + a_{n-1} r^{n-1} + \dots + a_2 r^2 + a_1 r + a_0 \right) = 0$$

Because e^{rx} is never zero, we see that $y = e^{rx}$ will be a solution of Eq. (1) precisely when *r* is a root of the equation

$$a_n r^n + a_{n-1} r^{n-1} + \dots + a_2 r^2 + a_1 r + a_0 = 0.$$
(3)

This equation is called the **characteristic equation** or **auxiliary equation** of the differential equation in (1). Our problem, then, is reduced to the solution of this purely algebraic equation.

According to the fundamental theorem of algebra, every *n*th-degree polynomial—such as the one in Eq. (3)—has *n* zeros, though not necessarily distinct and not necessarily real. Finding the exact values of these zeros may be difficult or even impossible; the quadratic formula is sufficient for second-degree equations, but for equations of higher degree we may need either to spot a fortuitous factorization or to apply a numerical technique such as Newton's method (or use a calculator/computer solve command).

Distinct Real Roots

Whatever the method we use, let us suppose that we have solved the characteristic equation. Then we can always write a general solution of the differential equation. The situation is slightly more complicated in the case of repeated roots or complex roots of Eq. (3), so let us first examine the simplest case—in which the characteristic equation has *n* distinct (no two equal) *real* roots r_1, r_2, \ldots, r_n . Then the functions

$$e^{r_1x}, e^{r_2x}, \ldots, e^{r_nx}$$

are all solutions of Eq. (1), and (by Problem 34 of Section 5.2) these *n* solutions are linearly independent on the entire real line. In summary, we have proved Theorem 1.

THEOREM 1 Distinct Real Roots

If the roots r_1, r_2, \ldots, r_n of the characteristic equation in (3) are real and distinct, then

is a general solution of Eq. (1). Thus the *n* linearly independent functions $\{e^{r_1x}, e^{r_2x}, \ldots, e^{r_nx}\}$ constitute a basis for the *n*-dimensional solution space of Eq. (1).

$$y^{(3)} + 3y'' - 10y' = 0;$$

 $y(0) = 7, y'(0) = 0, y''(0) = 70$

Solution The characteristic equation of the given differential equation is

$$r^3 + 3r^2 - 10r = 0.$$

We solve by factoring:

$$r(r^{2} + 3r - 10) = r(r + 5)(r - 2) = 0,$$

and so the characteristic equation has the three distinct real roots r = 0, r = -5, and r = 2. Because $e^0 = 1$, Theorem 1 gives the general solution

$$y(x) = c_1 + c_2 e^{-5x} + c_3 e^{2x}.$$

Then the given initial conditions yield the linear equations

$$y(0) = c_1 + c_2 + c_3 = 7,$$

$$y'(0) = -5c_2 + 2c_3 = 0,$$

$$y''(0) = 25c_2 + 4c_3 = 70$$

in the coefficients c_1 , c_2 , and c_3 . The last two equations give $y''(0) - 2y'(0) = 35c_2 = 70$, so $c_2 = 2$. Then the second equation gives $c_3 = 5$, and finally the first equation gives $c_1 = 0$. Thus the desired particular solution is

$$y(x) = 2e^{-5x} + 5e^{2x}.$$

Polynomial Differential Operators

If the roots of the characteristic equation in (3) are *not* distinct—there are repeated roots—then we cannot produce *n* linearly independent solutions of Eq. (1) by the method of Theorem 1. For example, if the roots are 1, 2, 2, and 2, we obtain only the *two* functions e^x and e^{2x} . The problem, then, is to produce the missing linearly independent solutions. For this purpose, it is convenient to adopt "operator notation" and write Eq. (1) in the form Ly = 0, where the **operator**

$$L = a_n \frac{d^n}{dx^n} + a_{n-1} \frac{d^{n-1}}{dx^{n-1}} + \dots + a_2 \frac{d^2}{dx^2} + a_1 \frac{d}{dx} + a_0$$
(5)

operates on the *n*-times differentiable function y(x) to produce the linear combination

$$Ly = a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_2 y^{(2)} + a_1 y' + a_0 y$$

of y and its first n derivatives. We also denote by D = d/dx the operation of differentiation with respect to x, so that

$$Dy = y', \quad D^2y = y'', \quad D^3y = y^{(3)},$$

and so on. In terms of D, the operator L in (5) may be written

$$L = a_n D^n + a_{n-1} D^{n-1} + \dots + a_2 D^2 + a_1 D + a_0,$$
(6)

and we will find it useful to think of the right-hand side in Eq. (6) as a (formal) *n*th-degree polynomial in the "variable" *D*; it is a **polynomial differential operator**.

A first-degree polynomial operator with leading coefficient 1 has the form D - a, where a is a real number. It operates on a function y = y(x) to produce

$$(D-a)y = Dy - ay = y' - ay.$$

The important fact about such operators is that any two of them *commute*:

$$(D-a)(D-b)y = (D-b)(D-a)y$$
(7)

for any twice differentiable function y = y(x). The proof of the formula in (7) is the following computation:

$$(D-a)(D-b)y = (D-a)(y'-by) = D(y'-by) - a(y'-by) = y'' - (b+a)y' + aby = y'' - (a+b)y' + bay = D(y'-ay) - b(y'-ay) = (D-b)(y'-ay) = (D-b)(D-a)y.$$

We see here also that $(D - a)(D - b) = D^2 - (a + b)D + ab$. Similarly, it can be shown by induction on the number of factors that an operator product of the form $(D - a_1)(D - a_2)\cdots(D - a_n)$ expands—by multiplying out and collecting coefficients—in the same way as does an ordinary product $(x - a_1)(x - a_2)\cdots(x - a_n)$ of linear factors, with x denoting a real variable. Consequently, the algebra of polynomial differential operators closely resembles the algebra of ordinary real polynomials.

Repeated Real Roots

Let us now consider the possibility that the characteristic equation

$$a_n r^n + a_{n-1} r^{n-1} + \dots + a_2 r^2 + a_1 r + a_0 = 0$$
(3)

has *repeated* roots. For example, suppose that Eq. (3) has only two distinct roots, r_0 of multiplicity 1 and r_1 of multiplicity k = n - 1 > 1. Then (after dividing by a_n) Eq. (3) can be rewritten in the form

$$(r - r_1)^k (r - r_0) = (r - r_0)(r - r_1)^k = 0.$$
(8)

Similarly, the corresponding operator L in (6) can be written as

$$L = (D - r_1)^k (D - r_0) = (D - r_0)(D - r_1)^k,$$
(9)

the order of the factors making no difference because of the formula in (7).

Two solutions of the differential equation Ly = 0 are certainly $y_0 = e^{r_0 x}$ and $y_1 = e^{r_1 x}$. This is, however, not sufficient; we need k + 1 linearly independent solutions in order to construct a general solution, because the equation is of order k + 1. To find the missing k - 1 solutions, we note that

$$Ly = (D - r_0)[(D - r_1)^k y] = 0.$$

Consequently, every solution of the kth-order equation

$$(D - r_1)^k y = 0 (10)$$

will also be a solution of the original equation Ly = 0. Hence our problem is reduced to that of finding a general solution of the differential equation in (10).

The fact that $y_1 = e^{r_1 x}$ is one solution of Eq. (10) suggests that we try the substitution

$$y(x) = u(x)y_1(x) = u(x)e^{r_1x},$$
(11)

where u(x) is a function yet to be determined. Observe that

$$(D - r_1) [ue^{r_1 x}] = (Du)e^{r_1 x} + u(r_1 e^{r_1 x}) - r_1(ue^{r_1 x}) = (Du)e^{r_1 x}.$$
 (12)

Upon k applications of this fact, it follows that

$$(D - r_1)^k \left[u e^{r_1 x} \right] = (D^k u) e^{r_1 x}$$
(13)

for any sufficiently differentiable function u(x). Hence $y = ue^{r_1 x}$ will be a solution of Eq. (10) if and only if $D^k u = u^{(k)} = 0$. But this is so if and only if

$$u(x) = c_1 + c_2 x + c_3 x^2 + \dots + c_k x^{k-1},$$

a polynomial of degree at most k - 1. Hence our desired solution of Eq. (10) is

$$y(x) = ue^{r_1x} = (c_1 + c_2x + c_3x^2 + \dots + c_kx^{k-1})e^{r_1x}.$$

In particular, we see here the additional solutions xe^{r_1x} , $x^2e^{r_1x}$, ..., $x^{k-1}e^{r_1x}$ of the original differential equation Ly = 0.

The preceding analysis can be carried out with the operator $D - r_1$ replaced with an arbitrary polynomial operator. When this is done, the result is a proof of the following theorem.

THEOREM 2 Repeated Roots

If the characteristic equation in (3) has a repeated root r of multiplicity k, then the part of a general solution of the differential equation in (1) corresponding to r is of the form

$$(c_1 + c_2 x + c_3 x^2 + \dots + c_k x^{k-1}) e^{rx}.$$
 (14)

Remark According to Problem 29 of Section 5.2, the k functions e^{rx} , xe^{rx} , x^2e^{rx} , ..., $x^{k-1}e^{rx}$ involved in (14) are linearly independent. Thus a root of multiplicity $k \ge 1$ corresponds to k linearly independent "basis solutions" that generate a k-dimensional subspace of the *n*-dimensional solution space \mathscr{S} of the *n*th-order homogeneous linear differential equation. Because the sum of the multiplicities of the separate roots of the *n*th-degree characteristic equation is n, we get a basis $\{y_1, y_2, \ldots, y_n\}$ for \mathscr{S} when we assemble all the basis solutions corresponding to these separate roots. We can then write a general solution $y = c_1y_1 + c_2y_2 + \cdots + c_ny_n$ of the differential equation. In effect, we thereby construct a general solution by combining different "parts" corresponding to the separate roots of the separate roots of the characteristic equation.

Example 2 Find a general solution of the fifth-order differential equation

$$9y^{(5)} - 6y^{(4)} + y^{(3)} = 0.$$

Solution The characteristic equation is

$$9r^{5} - 6r^{4} + r^{3} = r^{3}(9r^{2} - 6r + 1) = r^{3}(3r - 1)^{2} = 0$$

It has the triple root r = 0 and the double root $r = \frac{1}{3}$. The triple root r = 0 contributes

$$c_1 e^{0 \cdot x} + c_2 x e^{0 \cdot x} + c_3 x^2 e^{0 \cdot x} = c_1 + c_2 x + c_3 x^2$$

to the solution, while the double root $r = \frac{1}{3}$ contributes $c_4 e^{x/3} + c_5 x e^{x/3}$. Hence a general solution of the given differential equation is

$$y(x) = c_1 + c_2 x + c_3 x^2 + c_4 e^{x/3} + c_5 x e^{x/3}.$$

Complex-Valued Functions and Euler's Formula

Because we have assumed that the coefficients of the differential equation and its characteristic equation are real, any complex (nonreal) roots will occur in complex conjugate pairs $a \pm bi$ where a and b are real and $i = \sqrt{-1}$. This raises the question of what might be meant by an exponential such as $e^{(a+bi)x}$.

To answer this question, we recall from elementary calculus the Taylor (or MacLaurin) series for the exponential function

$$e^{t} = \sum_{n=0}^{\infty} \frac{t^{n}}{n!} = 1 + t + \frac{t^{2}}{2!} + \frac{t^{3}}{3!} + \frac{t^{4}}{4!} + \cdots$$

If we substitute $t = i\theta$ in this series and recall that $i^2 = -1$, $i^3 = -i$, $i^4 = 1$, and so on, we get

$$e^{i\theta} = \sum_{n=0}^{\infty} \frac{(i\theta)^n}{n!}$$

= $1 + i\theta - \frac{\theta^2}{2!} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} - \cdots$
= $\left(1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \cdots\right) + i\left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \cdots\right).$

Because the two real series in the last line are the Taylor series for $\cos \theta$ and $\sin \theta$, respectively, this implies that

$$e^{i\theta} = \cos\theta + i\sin\theta. \tag{15}$$

This result is known as **Euler's formula**. Because of it, we *define* the exponential function e^z , for z = x + iy an arbitrary complex number, to be

$$e^{z} = e^{x+iy} = e^{x}e^{iy} = e^{x}(\cos y + i\sin y).$$
 (16)

Thus it appears that complex roots of the characteristic equation will lead to complex-valued solutions of the differential equation. A **complex-valued function** F of the real variable x associates with each real number x (in its domain of definition) the complex number

$$F(x) = f(x) + ig(x).$$
 (17)

The real-valued functions f and g are called the **real** and **imaginary** parts, respectively, of F. If they are differentiable, we define the **derivative** F' of F by

$$F'(x) = f'(x) + ig'(x).$$
(18)

Thus we simply differentiate the real and imaginary parts of *F* separately.

We say that the complex-valued function F(x) satisfies the homogeneous linear differential equation L[F(x)] = 0 provided that its real and imaginary parts in (17) separately satisfy this equation—so L[F(x)] = L[f(x)] + iL[g(x)] = 0.

The particular complex-valued functions of interest here are of the form $F(x) = e^{rx}$, where $r = a \pm bi$. We note from Euler's formula that

$$e^{(a+bi)x} = e^{ax}e^{ibx} = e^{ax}(\cos bx + i\sin bx)$$
 (19a)

and

$$e^{(a-bi)x} = e^{ax}e^{-ibx} = e^{ax}(\cos bx - i\sin bx).$$
 (19b)

The most important property of e^{rx} is that

$$D_x(e^{rx}) = re^{rx}, (20)$$

if *r* is a complex number. The proof of this assertion is a straightforward computation based on the definitions and formulas given earlier:

$$D_x(e^{rx}) = D_x(e^{ax}\cos bx) + iD_x(e^{ax}\sin bx)$$

= $(ae^{ax}\cos bx - be^{ax}\sin bx) + i(ae^{ax}\sin bx + be^{ax}\cos bx)$
= $(a + bi)(e^{ax}\cos bx + ie^{ax}\sin bx) = re^{rx}.$

Complex Roots

It follows from Eq. (20) that when r is complex (just as when r is real), e^{rx} will be a solution of the differential equation in (1) if and only if r is a root of its characteristic equation. If the complex conjugate pair of roots $r_1 = a + bi$ and $r_2 = a - bi$ are simple (nonrepeated), then the corresponding part of a general solution of Eq. (1) is

$$y(x) = C_1 e^{r_1 x} + C_2 e^{r_2 x} = C_1 e^{(a+bi)x} + C_2 e^{(a-bi)x}$$

= $C_1 e^{ax} (\cos bx + i \sin bx) + C_2 e^{ax} (\cos bx - i \sin bx)$
 $y(x) = (C_1 + C_2) e^{ax} \cos bx + i (C_1 - C_2) e^{ax} \sin bx,$

where the arbitrary constants C_1 and C_2 can be complex. For instance, the choice $C_1 = C_2 = \frac{1}{2}$ gives the real-valued solution $y_1(x) = e^{ax} \cos bx$, while the choice $C_1 = -\frac{1}{2}i$, $C_2 = \frac{1}{2}i$ gives the independent real-valued solution $y_2(x) = e^{ax} \sin bx$. This yields the following result.

THEOREM 3 Complex Roots

If the characteristic equation in (3) has an unrepeated pair of complex conjugate roots $a \pm bi$ (with $b \neq 0$), then the corresponding part of a general solution of Eq. (1) has the form

$$e^{ax}(c_1\cos bx + c_2\sin bx). \tag{21}$$

Thus the linearly independent solutions $e^{ax} \cos bx$ and $e^{ax} \sin bx$ (corresponding to the complex conjugate characteristic roots $a \pm bi$) generate a 2-dimensional subspace of the solution space of the differential equation.

Example 3 The characteristic equation of $y'' + b^2 y = 0$ (b > 0) is $r^2 + b^2 = 0$, with roots $r = \pm bi$. So Theorem 3 (with a = 0) gives the general solution

$$y(x) = c_1 \cos bx + c_2 \sin bx.$$

Completion

Example 4

Find the particular solution of

Solution

for which
$$y(0) = 1$$
 and $y'(0) = 5$.
Completion of the square in the characteristic equation yields

$$r^{2} - 4r + 5 = (r - 2)^{2} + 1 = 0,$$

y'' - 4y' + 5y = 0

so $r-2 = \pm \sqrt{-1} = \pm i$. Thus we obtain the complex conjugate roots $2 \pm i$ (which could also be found directly using the quadratic formula). Hence Theorem 3 with a = 2 and b = 1gives the general solution

$$y(x) = e^{2x}(c_1 \cos x + c_2 \sin x).$$

Then

$$y'(x) = 2e^{2x}(c_1\cos x + c_2\sin x) + e^{2x}(-c_1\sin x + c_2\cos x)$$

so the initial conditions give

$$y(0) = c_1 = 1$$
 and $y'(0) = 2c_1 + c_2 = 5$.

It follows that $c_2 = 3$, and so the desired particular solution is

$$y(x) = e^{2x}(\cos x + 3\sin x).$$

In Example 5 below we employ the **polar form**

$$z = x + iy = re^{i\theta} \tag{22}$$



FIGURE 5.3.1. Modulus and argument of the complex number x + iy.

of the complex number z. This form follows from Euler's formula upon writing

$$z = r\left(\frac{x}{r} + i\frac{y}{r}\right) = r(\cos\theta + i\sin\theta) = re^{i\theta}$$

in terms of the modulus $r = \sqrt{x^2 + y^2} > 0$ of the number z and its argument θ indicated in Fig. 5.3.1. For instance, the imaginary number *i* has modulus 1 and argument $\pi/2$, so $i = e^{i\pi/2}$. Similarly, $-i = e^{3\pi/2}$. Another consequence is the fact that the nonzero complex number $z = re^{i\theta}$ has the two square roots

$$\sqrt{z} = \pm (re^{i\theta})^{1/2} = \pm \sqrt{r}e^{i\theta/2},\tag{23}$$

where \sqrt{r} denotes (as usual for a positive real number) the positive square root of the modulus of z.

Example 5 Solution Find a general solution of $y^{(4)} + 4y = 0$. The characteristic equation is

$$r^{4} + 4 = (r^{2})^{2} - (2i)^{2} = (r^{2} + 2i)(r^{2} - 2i) = 0$$

and its four roots are $\pm \sqrt{\pm 2i}$. Since $i = e^{i\pi/2}$ and $-i = e^{i3\pi/2}$, we find that

$$\sqrt{2i} = (2e^{i\pi/2})^{1/2} = \sqrt{2}e^{i\pi/4} = \sqrt{2}\left(\cos\frac{\pi}{4} + i\sin\frac{\pi}{4}\right) = 1 + i$$

and

$$\sqrt{-2i} = (2e^{i3\pi/2})^{1/2} = \sqrt{2}e^{i3\pi/4} = \sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right) = -1 + i$$

Thus the four (distinct) roots of the characteristic equation are $r = \pm (1 + i)$ and r = $\pm(-1+i)$. Grouping these four into the complex conjugate pairs $1 \pm i$ and $-1 \pm i$ yields the general solution

$$y(x) = e^{x}(c_1 \cos x + c_2 \sin x) + e^{-x}(c_3 \cos x + c_4 \sin x)$$

of the differential equation $y^{(4)} + 4y = 0$.

Repeated Complex Roots

Theorem 2 holds for repeated complex roots. If the conjugate pair $a \pm bi$ has multiplicity k, then the corresponding part of the general solution has the form

$$(A_1 + A_2 x + \dots + A_k x^{k-1})e^{(a+bi)x} + (B_1 + B_2 x + \dots + B_k x^{k-1})e^{(a-bi)x}$$
$$= \sum_{p=0}^{k-1} x^p e^{ax} (c_p \cos bx + d_p \sin bx).$$
(24)

It can be shown that the 2k functions

 $x^{p}e^{ax}\cos bx, \quad x^{p}e^{ax}\sin bx, \quad 0 \le p \le k-1$

that appear in Eq. (24) are linearly independent.

Example 6

Find a general solution of $(D^2 + 6D + 13)^2 v = 0$. Solution By completing the square, we see that the characteristic equation

 $(r^{2} + 6r + 13)^{2} = [(r + 3)^{2} + 4]^{2} = 0$

has as its roots the conjugate pair $-3 \pm 2i$ of multiplicity k = 2. Hence Eq. (24) gives the general solution

$$y(x) = e^{-3x}(c_1\cos 2x + d_1\sin 2x) + xe^{-3x}(c_2\cos 2x + d_2\sin 2x).$$

In applications we are seldom presented in advance with a factorization as convenient as the one in Example 6. Often the most difficult part of solving a homogeneous linear equation is finding the roots of its characteristic equation. Example 7 illustrates an approach that may succeed when a root of the characteristic equation can be found by inspection. The application material for this section illustrates other possibilities.

Example 7 The characteristic equation of the differential equation

$$y^{(3)} + y' - 10y = 0$$

is the cubic equation

$$r^3 + r - 10 = 0$$

By a standard theorem of elementary algebra, the only possible rational roots are the factors ± 1 , ± 2 , ± 5 , and ± 10 of the constant term 10. By trial and error (if not by inspection) we discover the root 2. The factor theorem of elementary algebra implies that r-2 is a factor of $r^3 + r - 10$, and division of the former into the latter produces as quotient the quadratic polynomial

$$r^2 + 2r + 5 = (r+1)^2 + 4.$$

The roots of this quotient are the complex conjugates $-1 \pm 2i$. The three roots we have found now yield the general solution

$$y(x) = c_1 e^{2x} + e^{-x} (c_2 \cos 2x + c_3 \sin 2x).$$

Example 8

-5, $2 \pm 3i$, and $2 \pm 3i$. Write a general solution of this homogeneous differential equation. The solution can be read directly from the list of roots. It is Solution

$$y(x) = c_1 + c_2 x + c_3 x^2 + c_4 x^3 + c_5 e^{3x} + c_6 e^{-5x} + c_7 x e^{-5x} + e^{2x} (c_8 \cos 3x + c_9 \sin 3x) + x e^{2x} (c_{10} \cos 3x + c_{11} \sin 3x)$$

5.3 **Problems**

Find the general solutions of the differential equations in Problems 1 through 20.

- **2.** 2v'' 3v' = 01. v'' - 4v = 04. 2v'' - 7v' + 3v = 03. v'' + 3v' - 10v = 05. y'' + 6y' + 9y = 06. y'' + 5y' + 5y = 07. 4y'' - 12y' + 9y = 08. y'' - 6y' + 13y = 010. $5y^{(4)} + 3y^{(3)} = 0$ 9. y'' + 8y' + 25y = 0**11.** $v^{(4)} - 8v^{(3)} + 16v'' = 0$ 12. $v^{(4)} - 3v^{(3)} + 3v'' - v' = 0$ **13.** $9y^{(3)} + 12y'' + 4y' = 0$ 14. $y^{(4)} + 3y'' - 4y = 0$ 15. $v^{(4)} - 8v'' + 16v = 0$ **16.** $v^{(4)} + 18v'' + 81v = 0$ 17. $6y^{(4)} + 11y'' + 4y = 0$ 18. $v^{(4)} = 16v$ **19.** $v^{(3)} + v'' - v' - v = 0$
- **20.** $y^{(4)} + 2y^{(3)} + 3y'' + 2y' + y = 0$ (Suggestion: Expand $(r^2 + r + 1)^2$.)

Solve the initial value problems given in Problems 21 through 26.

- **21.** y'' 4y' + 3y = 0; y(0) = 7, y'(0) = 11
- **22.** 9y'' + 6y' + 4y = 0; y(0) = 3, y'(0) = 4
- **23.** y'' 6y' + 25y = 0; y(0) = 3, y'(0) = 1
- **24.** $2y^{(3)} 3y'' 2y' = 0$; y(0) = 1, y'(0) = -1, y''(0) = 3
- **25.** $3y^{(3)} + 2y'' = 0$; y(0) = -1, y'(0) = 0, y''(0) = 1
- **26.** $y^{(3)} + 10y'' + 25y' = 0; y(0) = 3, y'(0) = 4, y''(0) = 5$

Find general solutions of the equations in Problems 27 through 32. First find a small integral root of the characteristic equation by inspection; then factor by division.

27.
$$y^{(3)} + 3y'' - 4y = 0$$

28. $2y^{(3)} - y'' - 5y' - 2y = 0$
29. $y^{(3)} + 27y = 0$
30. $y^{(4)} - y^{(3)} + y'' - 3y' - 6y = 0$
31. $y^{(3)} + 3y'' + 4y' - 8y = 0$
32. $y^{(4)} + y^{(3)} - 3y'' - 5y' - 2y = 0$

In Problems 33 through 36, one solution of the differential equation is given. Find the general solution.

- **33.** $y^{(3)} + 3y'' 54y = 0; y = e^{3x}$
- **34.** $3y^{(3)} 2y'' + 12y' 8y = 0; y = e^{2x/3}$
- **35.** $6y^{(4)} + 5y^{(3)} + 25y'' + 20y' + 4y = 0; y = \cos 2x$
- **36.** $9y^{(3)} + 11y'' + 4y' 14y = 0; y = e^{-x} \sin x$
- **37.** Find a function y(x) such that $y^{(4)}(x) = y^{(3)}(x)$ for all x and y(0) = 18, y'(0) = 12, y''(0) = 13, and $y^{(3)}(0) = 7$.
- 38. Solve the initial value problem

$$y^{(3)} - 5y'' + 100y' - 500y = 0;$$

 $y(0) = 0, \quad y'(0) = 10, \quad y''(0) = 250$

given that $y_1(x) = e^{5x}$ is one particular solution of the differential equation.

In Problems 39 through 42, find a linear homogeneous constant-coefficient equation with the given general solution.

- **39.** $y(x) = (A + Bx + Cx^2)e^{2x}$
- **40.** $y(x) = Ae^{2x} + B\cos 2x + C\sin 2x$
- **41.** $y(x) = A \cos 2x + B \sin 2x + C \cosh 2x + D \sinh 2x$
- 42. $y(x) = (A + Bx + Cx^2)\cos 2x + (D + Ex + Fx^2)\sin 2x$

Problems 43 through 47 pertain to the solution of differential equations with complex coefficients.

- **43.** (a) Use Euler's formula to show that every complex number can be written in the form $re^{i\theta}$, where $r \ge 0$ and $-\pi < \theta \le \pi$. (b) Express the numbers 4, -2, 3*i*, 1 + *i*, and $-1 + i\sqrt{3}$ in the form $re^{i\theta}$. (c) The two square roots of $re^{i\theta}$ are $\pm \sqrt{r}e^{i\theta/2}$. Find the square roots of the numbers $2 2i\sqrt{3}$ and $-2 + 2i\sqrt{3}$.
- **44.** Use the quadratic formula to solve the following equations. Note in each case that the roots are not complex conjugates.
 - (a) $x^2 + ix + 2 = 0$ (b) $x^2 2ix + 3 = 0$
- **45.** Find a general solution of y'' 2iy' + 3y = 0.
- **46.** Find a general solution of y'' iy' + 6y = 0.
- **47.** Find a general solution of $y'' = (-2 + 2i\sqrt{3})y$.
- **48.** Solve the initial value problem

$$y^{(3)} = y; \quad y(0) = 1, \quad y'(0) = y''(0) = 0.$$

(Suggestion: Impose the given initial conditions on the general solution

$$y(x) = Ae^x + Be^{\alpha x} + Ce^{\beta x},$$

where α and β are the complex conjugate roots of $r^3 - 1 = 0$, to discover that

$$y(x) = \frac{1}{3} \left(e^x + 2e^{-x/2} \cos \frac{x\sqrt{3}}{2} \right)$$

is a solution.)

49. Solve the initial value problem

$$y^{(4)} = y^{(3)} + y'' + y' + 2y;$$

$$y(0) = y'(0) = y''(0) = 0, 2y^{(3)}(0) = 30$$

50. The differential equation

$$y'' + (\operatorname{sgn} x)y = 0 \tag{25}$$

has the discontinuous coefficient function

$$\operatorname{sgn} x = \begin{cases} +1 & \text{if } x > 0, \\ -1 & \text{if } x < 0. \end{cases}$$

Show that Eq. (25) nevertheless has two linearly independent solutions $y_1(x)$ and $y_2(x)$ defined for all x such that

- Each satisfies Eq. (25) at each point $x \neq 0$;
- Each has a continuous derivative at x = 0;
- $y_1(0) = y'_2(0) = 1$ and $y_2(0) = y'_1(0) = 0$.

(*Suggestion:* Each $y_i(x)$ will be defined by one formula for x < 0 and by another for $x \ge 0$.) The graphs of these two solutions are shown in Fig. 5.3.2.



FIGURE 5.3.2. Graphs of $y_1(x)$ and $y_2(x)$ in Problem 50.

51. According to Problem 51 in Section 5.1, the substitution $v = \ln x (x > 0)$ transforms the second-order Euler equation $ax^2y'' + bxy' + cy = 0$ to a constant-coefficient homogeneous linear equation. Show similarly that this same

substitution transforms the third-order Euler equation

$$ax^{3}y''' + bx^{2}y'' + cxy' + dy = 0$$

(where a, b, c, d are constants) into the constantcoefficient equation

$$a\frac{d^3y}{dv^3} + (b-3a)\frac{d^2y}{dv^2} + (c-b+2a)\frac{dy}{dv} + dy = 0.$$

Make the substitution $v = \ln x$ of Problem 51 to find general solutions (for x > 0) of the Euler equations in Problems 52 through 58.

52. $x^{2}y'' + xy' + 9y = 0$ 53. $x^{2}y'' + 7xy' + 25y = 0$ 54. $x^{3}y''' + 6x^{2}y'' + 4xy' = 0$ 55. $x^{3}y''' - x^{2}y'' + xy' = 0$ 56. $x^{3}y''' + 3x^{2}y'' + xy' = 0$ 57. $x^{3}y''' - 3x^{2}y'' + xy' = 0$ 58. $x^{3}y''' + 6x^{2}y'' + 7xy' + y = 0$

5.3 Application Approximate Solutions of Linear Equations



To meet the needs of applications such as those of this section, polynomial-solving utilities are now a common feature of calculator and computer systems and can be used to solve a characteristic equation numerically even when no simple factorization is evident or even possible. For instance, suppose that we want to solve the homogeneous linear differential equation

$$y^{(3)} - 3y'' + y = 0 \tag{1}$$

with characteristic equation

$$r^3 - 3r^2 + 1 = 0. (2)$$

A typical graphing calculator has a **solve** command that can be used to find the approximate roots of a polynomial equation. As indicated in Figs. 5.3.3 and 5.3.4, we find that the roots of Eq. (2) are given by $r \approx -0.5321, 0.6527, \text{ and } 2.8794$. Some analogous computer algebra system commands are

solv	e(R ³ -	3R ² +1	.R	-1)	
			-0.5	3208	88862
solv	e(R ³ -	3R ² +1	R.1	.)	
			0.6	5270	36447
solv	e(R ³ -	3R ² +1	.,R,3	3)	
			. 2.	8793	85242

FIGURE 5.3.3. Solving the equation $r^3 - 3r^2 + 1 = 0$ with a TI-84 Plus CE calculator that requires an estimate of each root.



FIGURE 5.3.4. Solving the characteristic equation with a TI-89 calculator having a more sophisticated **solve** facility.

$fsolve(r^3 - 3*r^2 + 1 = 0, r);$	(Maple)
$NSolve[r^3 - 3*r^2 + 1 == 0, r]$	(Mathematica)
$r^3 - 3r^2 + 1 = 0$	(Wolfram Alpha)
roots([1 -3 0 1])	(MATLAB)

(In the MATLAB command, one enters the polynomial's vector [1 -3 0 1] of coefficients, listed in descending order.) However we find these approximate roots, it follows that a general solution of the differential equation in (1) is given (approximately) by

$$y(x) = c_1 e^{-(0.5321)x} + c_2 e^{(0.6527)x} + c_3 e^{(2.8794)x}.$$
(3)

Use calculator or computer methods like those indicated here to find general solutions (in approximate numerical form) of the following differential equations.

1. $y^{(3)} - 3y' + y = 0$ 2. $y^{(3)} + 3y'' - 3y = 0$ 3. $y^{(3)} + y' + y = 0$ 4. $y^{(3)} + 3y' + 5y = 0$ 5. $y^{(4)} + 2y^{(3)} - 3y = 0$ 6. $y^{(4)} + 3y' - 4y = 0$

5.4 Mechanical Vibrations



```
FIGURE 5.4.1. A mass–spring–dashpot system.
```

The motion of a mass attached to a spring serves as a relatively simple example of the vibrations that occur in more complex mechanical systems. For many such systems, the analysis of these vibrations is a problem in the solution of linear differential equations with constant coefficients.

We consider a body of mass *m* attached to one end of an ordinary spring that resists compression as well as stretching; the other end of the spring is attached to a fixed wall, as shown in Fig. 5.4.1. Assume that the body rests on a frictionless horizontal plane, so that it can move only back and forth as the spring compresses and stretches. Denote by *x* the distance of the body from its **equilibrium position**— its position when the spring is unstretched. We take x > 0 when the spring is stretched, and thus x < 0 when it is compressed.

According to Hooke's law, the restorative force F_S that the spring exerts on the mass is proportional to the distance x that the spring has been stretched or compressed. Because this is the same as the displacement x of the mass m from its equilibrium position, it follows that

$$F_S = -kx. \tag{1}$$

The positive constant of proportionality k is called the **spring constant**. Note that F_S and x have opposite signs: $F_S < 0$ when x > 0, $F_S > 0$ when x < 0.

Figure 5.4.1 shows the mass attached to a dashpot—a device, like a shock absorber, that provides a force directed opposite to the instantaneous direction of motion of the mass m. We assume the dashpot is so designed that this force F_R is proportional to the velocity v = dx/dt of the mass; that is,

$$F_R = -cv = -c\frac{dx}{dt}.$$
(2)

The positive constant c is the **damping constant** of the dashpot. More generally, we may regard Eq. (2) as specifying frictional forces in our system (including air resistance to the motion of m).

If, in addition to the forces F_S and F_R , the mass is subjected to a given **exter**nal force $F_E = F(t)$, then the total force acting on the mass is $F = F_S + F_R + F_E$. Using Newton's law

$$F = ma = m\frac{d^2x}{dt^2} = mx'',$$

we obtain the second-order linear differential equation

$$mx'' + cx' + kx = F(t) \tag{3}$$

that governs the motion of the mass.

If there is no dashpot (and we ignore all frictional forces), then we set c = 0 in Eq. (3) and call the motion **undamped**; it is **damped** motion if c > 0. If there is no external force, we replace F(t) with 0 in Eq. (3). We refer to the motion as **free** in this case and **forced** in the case $F(t) \neq 0$. Thus the homogeneous equation

$$mx'' + cx' + kx = 0 (4)$$

describes free motion of a mass on a spring with dashpot but with no external forces applied. We will defer discussion of forced motion until Section 5.6.

For an alternative example, we might attach the mass to the lower end of a spring that is suspended vertically from a fixed support, as in Fig. 5.4.2. In this case the weight W = mg of the mass would stretch the spring a distance s_0 determined by Eq. (1) with $F_S = -W$ and $x = s_0$. That is, $mg = ks_0$, so that $s_0 = mg/k$. This gives the **static** equilibrium position of the mass. If y denotes the displacement of the mass in motion, measured downward from its static equilibrium position, then we ask you to show in Problem 9 that y satisfies Eq. (3); specifically, that

$$my'' + cy' + ky = F(t)$$
⁽⁵⁾

if we include damping and external forces (meaning those other than gravity).

The Simple Pendulum

The importance of the differential equation that appears in Eqs. (3) and (5) stems from the fact that it describes the motion of many other simple mechanical systems. For example, a **simple pendulum** consists of a mass *m* swinging back and forth on the end of a string (or better, a *massless rod*) of length *L*, as shown in Fig. 5.4.3. We may specify the position of the mass at time *t* by giving the counterclockwise angle $\theta = \theta(t)$ that the string or rod makes with the vertical at time *t*. To analyze the motion of the mass *m*, we will apply the law of the *conservation of mechanical energy*, according to which the sum of the kinetic energy and the potential energy of *m* remains constant.

The distance along the circular arc from 0 to m is $s = L\theta$, so the velocity of the mass is $v = ds/dt = L(d\theta/dt)$, and therefore its kinetic energy is

$$T = \frac{1}{2}mv^2 = \frac{1}{2}m\left(\frac{ds}{dt}\right)^2 = \frac{1}{2}mL^2\left(\frac{d\theta}{dt}\right)^2.$$

We next choose as reference point the lowest point *O* reached by the mass (see Fig. 5.4.3). Then its potential energy *V* is the product of its weight mg and its vertical height $h = L(1 - \cos \theta)$ above *O*, so

$$V = mgL(1 - \cos\theta).$$



FIGURE 5.4.2. A mass suspended vertically from a spring.



FIGURE 5.4.3. The simple pendulum.

The fact that the sum of T and V is a constant C therefore gives

$$\frac{1}{2}mL^2\left(\frac{d\theta}{dt}\right)^2 + mgL(1-\cos\theta) = C.$$

We differentiate both sides of this identity with respect to t to obtain

$$mL^{2}\left(\frac{d\theta}{dt}\right)\left(\frac{d^{2}\theta}{dt^{2}}\right) + mgL(\sin\theta)\frac{d\theta}{dt} = 0,$$

so

$$\frac{d^2\theta}{dt^2} + \frac{g}{L}\sin\theta = 0 \tag{6}$$

after removal of the common factor $mL^2(d\theta/dt)$. This differential equation can be derived in a seemingly more elementary manner using the familiar second law F = ma of Newton (applied to tangential components of the acceleration of the mass and the force acting on it). However, derivations of differential equations based on conservation of energy are often seen in more complex situations where Newton's law is not so directly applicable, and it may be instructive to see the energy method in a simpler application like the pendulum.

Now recall that if θ is small, then $\sin \theta \approx \theta$ (this approximation obtained by retaining just the first term in the Taylor series for $\sin \theta$). In fact, $\sin \theta$ and θ agree to two decimal places when $|\theta|$ is at most $\pi/12$ (that is, 15°). In a typical pendulum clock, for example, θ would never exceed 15°. It therefore seems reasonable to simplify our mathematical model of the simple pendulum by replacing $\sin \theta$ with θ in Eq. (6). If we also insert a term $c\theta'$ to account for the frictional resistance of the surrounding medium, the result is an equation in the form of Eq. (4):

$$\theta'' + c\theta' + k\theta = 0, \tag{7}$$

where k = g/L. Note that this equation is independent of the mass *m* on the end of the rod. We might, however, expect the effects of the discrepancy between θ and $\sin \theta$ to accumulate over a period of time, so that Eq. (7) will probably not describe accurately the actual motion of the pendulum over a long period of time.

In the remainder of this section, we first analyze free undamped motion and then free damped motion.

Free Undamped Motion

If we have only a mass on a spring, with neither damping nor external force, then Eq. (3) takes the simpler form

$$mx'' + kx = 0. ag{8}$$

It is convenient to define

$$\omega_0 = \sqrt{\frac{k}{m}} \tag{9}$$

and rewrite Eq. (8) as

$$x'' + \omega_0^2 x = 0. (8')$$

The general solution of Eq. (8') is

$$x(t) = A\cos\omega_0 t + B\sin\omega_0 t.$$
(10)



FIGURE 5.4.4. The angle α .

To analyze the motion described by this solution, we choose constants C and α so that

$$C = \sqrt{A^2 + B^2}, \quad \cos \alpha = \frac{A}{C}, \quad \text{and} \quad \sin \alpha = \frac{B}{C},$$
 (11)

as indicated in Fig. 5.4.4. Note that, although $\tan \alpha = B/A$, the angle α is *not* given by the principal branch of the inverse tangent function (which gives values only in the interval $-\pi/2 < x < \pi/2$). Instead, α is the angle between 0 and 2π whose cosine and sine have the signs given in (11), where either A or B or both may be negative. Thus

$$\alpha = \begin{cases} \tan^{-1}(B/A) & \text{if } A > 0, B > 0 \text{ (first quadrant),} \\ \pi + \tan^{-1}(B/A) & \text{if } A < 0 \text{ (second or third quadrant),} \\ 2\pi + \tan^{-1}(B/A) & \text{if } A > 0, B < 0 \text{ (fourth quadrant),} \end{cases}$$

where $\tan^{-1}(B/A)$ is the angle in $(-\pi/2, \pi/2)$ given by a calculator or computer. In any event, from (10) and (11) we get

$$x(t) = C\left(\frac{A}{C}\cos\omega_0 t + \frac{B}{C}\sin\omega_0 t\right) = C(\cos\alpha\cos\omega_0 t + \sin\alpha\sin\omega_0 t).$$

With the aid of the cosine addition formula, we find that

$$x(t) = C\cos(\omega_0 t - \alpha). \tag{12}$$

Thus the mass oscillates to and fro about its equilibrium position with

1. AmplitudeC,2. Circular frequency ω_0 , and3. Phase angle α .

Such motion is called **simple harmonic motion**.

If time *t* is measured in seconds, the circular frequency ω_0 has dimensions of radians per second (rad/s). The **period** of the motion is the time required for the system to complete one full oscillation, so is given by

$$T = \frac{2\pi}{\omega_0} \tag{13}$$

 $C \xrightarrow{x} x(t) = C \cos(\omega_0 t - \alpha)$

FIGURE 5.4.5. Simple harmonic motion.

seconds; its **frequency** is

$$\nu = \frac{1}{T} = \frac{\omega_0}{2\pi} \tag{14}$$

in hertz (Hz), which measures the number of complete cycles per second. Note that frequency is measured in cycles per second, whereas circular frequency has the dimensions of radians per second.

A typical graph of a simple harmonic position function

$$x(t) = C\cos(\omega_0 t - \alpha) = C\cos\left(\omega_0\left(t - \frac{\alpha}{\omega_0}\right)\right) = C\cos(\omega_0(t - \delta))$$

is shown in Fig. 5.4.5, where the geometric significance of the amplitude C, the period T, and the **time lag**

$$\delta = \frac{\alpha}{\omega_0}$$

are indicated.

If the initial position $x(0) = x_0$ and initial velocity $x'(0) = v_0$ of the mass are given, we first determine the values of the coefficients *A* and *B* in Eq. (10), then find the amplitude *C* and phase angle α by carrying out the transformation of x(t) to the form in Eq. (12), as indicated previously.

Example 1 Undamped mass-spring system A body with mass $m = \frac{1}{2}$ kilogram (kg) is attached to the end of a spring that is stretched 2 meters (m) by a force of 100 newtons (N). It is set in motion with initial position $x_0 = 1$ (m) and initial velocity $v_0 = -5$ (m/s). (Note that these initial conditions indicate that the body is displaced to the right and is moving to the left at time t = 0.) Find the position function of the body as well as the amplitude, frequency, period of oscillation, and time lag of its motion.

Solution The spring constant is k = (100 N)/(2 m) = 50 (N/m), so Eq. (8) yields $\frac{1}{2}x'' + 50x = 0$; that is,

$$x'' + 100x = 0$$

Consequently, the circular frequency of the resulting simple harmonic motion of the body will be $\omega_0 = \sqrt{100} = 10$ (rad/s). Hence it will oscillate with period

$$T = \frac{2\pi}{\omega_0} = \frac{2\pi}{10} \approx 0.6283 \text{ s}$$

and with frequency

$$v = \frac{1}{T} = \frac{\omega_0}{2\pi} = \frac{10}{2\pi} \approx 1.5915 \text{ Hz}.$$

We now impose the initial conditions x(0) = 1 and x'(0) = -5 on the position function

 $x(t) = A\cos 10t + B\sin 10t$ with $x'(t) = -10A\sin 10t + 10B\cos 10t$.

It follows readily that A = 1 and $B = -\frac{1}{2}$, so the position function of the body is

$$x(t) = \cos 10t - \frac{1}{2}\sin 10t.$$

Hence its amplitude of motion is

$$C = \sqrt{(1)^2 + (-\frac{1}{2})^2} = \frac{1}{2}\sqrt{5} \text{ m}.$$

To find the time lag, we write

$$x(t) = \frac{\sqrt{5}}{2} \left(\frac{2}{\sqrt{5}} \cos 10t - \frac{1}{\sqrt{5}} \sin 10t \right) = \frac{\sqrt{5}}{2} \cos(10t - \alpha),$$

where the phase angle α satisfies

$$\cos \alpha = \frac{2}{\sqrt{5}} > 0$$
 and $\sin \alpha = -\frac{1}{\sqrt{5}} < 0.$

Hence α is the fourth-quadrant angle

$$\alpha = 2\pi + \tan^{-1}\left(\frac{-1/\sqrt{5}}{2/\sqrt{5}}\right) = 2\pi - \tan^{-1}(\frac{1}{2}) \approx 5.8195,$$

and the time lag of the motion is

$$\delta = \frac{\alpha}{\omega_0} \approx 0.5820 \text{ s}$$

With the amplitude and approximate phase angle shown explicitly, the position function of the body takes the form

$$x(t) \approx \frac{1}{2}\sqrt{5}\cos(10t - 5.8195),$$

and its graph is shown in Fig. 5.4.6.



FIGURE 5.4.6. Graph of the position function $x(t) = C \cos(\omega_0 t - \alpha)$ in Example 1, with amplitude $C \approx 1.118$, period $T \approx 0.628$, and time lag $\delta \approx 0.582$.

Free Damped Motion

With damping but no external force, the differential equation we have been studying takes the form mx'' + cx' + kx = 0; alternatively,

$$x'' + 2px' + \omega_0^2 x = 0, (15)$$

where $\omega_0 = \sqrt{k/m}$ is the corresponding *undamped* circular frequency and

$$p = \frac{c}{2m} > 0. \tag{16}$$

The characteristic equation $r^2 + 2pr + \omega_0^2 = 0$ of Eq. (15) has roots

$$r_1, r_2 = -p \pm (p^2 - \omega_0^2)^{1/2}$$
(17)

that depend on the sign of

$$p^{2} - \omega_{0}^{2} = \frac{c^{2}}{4m^{2}} - \frac{k}{m} = \frac{c^{2} - 4km}{4m^{2}}.$$
 (18)

The **critical damping** c_{cr} is given by $c_{cr} = \sqrt{4km}$, and we distinguish three cases, according as $c > c_{cr}$, $c = c_{cr}$, or $c < c_{cr}$.

OVERDAMPED CASE: $c > c_{cr}$ ($c^2 > 4km$). Because *c* is relatively large in this case, we are dealing with a strong resistance in comparison with a relatively weak spring or a small mass. Then (17) gives distinct real roots r_1 and r_2 , both of which are negative. The position function has the form

$$x(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t}.$$
(19)

It is easy to see that $x(t) \to 0$ as $t \to +\infty$ and that the body settles to its equilibrium position without any oscillations (Problem 29). Figure 5.4.7 shows some typical graphs of the position function for the overdamped case; we chose x_0 a fixed positive number and illustrated the effects of changing the initial velocity v_0 . In every case the would-be oscillations are damped out.



FIGURE 5.4.7. Overdamped motion: $x(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t}$ with $r_1 < 0$ and $r_2 < 0$. Solution curves are graphed with the same initial position x_0 and different initial velocities.

CRITICALLY DAMPED CASE: $c = c_{cr} (c^2 = 4km)$. In this case, (17) gives equal roots $r_1 = r_2 = -p$ of the characteristic equation, so the general solution is

$$x(t) = e^{-pt}(c_1 + c_2 t).$$
(20)

Because $e^{-pt} > 0$ and $c_1 + c_2t$ has at most one positive zero, the body passes through its equilibrium position at most once, and it is clear that $x(t) \rightarrow 0$ as $t \rightarrow +\infty$. Some graphs of the motion in the critically damped case appear in Fig. 5.4.8, and they resemble those of the overdamped case (Fig. 5.4.7). In the critically damped case, the resistance of the dashpot is just large enough to damp out any oscillations, but even a slight reduction in resistance will bring us to the remaining case, the one that shows the most dramatic behavior.

UNDERDAMPED CASE: $c < c_{\rm cr}$ ($c^2 < 4km$). The characteristic equation now has two complex conjugate roots $-p \pm i \sqrt{\omega_0^2 - p^2}$, and the general solution is

$$x(t) = e^{-pt} (A \cos \omega_1 t + B \sin \omega_1 t), \qquad (21)$$

where

 \succ

$$\omega_1 = \sqrt{\omega_0^2 - p^2} = \frac{\sqrt{4km - c^2}}{2m}.$$
(22)

(23)

Using the cosine addition formula as in the derivation of Eq. (12), we may rewrite Eq. (20) as

 $x(t) = Ce^{-pt}\cos(\omega_1 t - \alpha)$

 $x(t) = Ce^{-pt} \left(\frac{A}{C}\cos\omega_1 t + \frac{B}{C}\sin\omega_1 t\right),$

so

where

$$C = \sqrt{A^2 + B^2}, \quad \cos \alpha = \frac{A}{C}, \quad \text{and} \quad \sin \alpha = \frac{B}{C}.$$

The solution in (21) represents exponentially damped oscillations of the body around its equilibrium position. The graph of x(t) lies between the "amplitude envelope" curves $x = -Ce^{-pt}$ and $x = Ce^{-pt}$ and touches them when $\omega_1 t - \alpha$ is an integral multiple of π . The motion is not actually periodic, but it is nevertheless useful to call ω_1 its **circular frequency** (more properly, its **pseudofrequency**), $T_1 = 2\pi/\omega_1$ its **pseudoperiod** of oscillation, and Ce^{-pt} its **time-varying amplitude**. Most of these quantities are shown in the typical graph of underdamped motion in Fig. 5.4.9. Note from Eq. (22) that in this case ω_1 is less than the undamped circular frequency ω_0 , so T_1 is larger than the period T of oscillation of the same mass without damping on the same spring. Thus the action of the dashpot has at least two effects:

- **1.** It exponentially damps the oscillations, in accord with the time-varying amplitude.
- 2. It slows the motion; that is, the dashpot decreases the frequency of the motion.

As the following example illustrates, damping typically also delays the motion further—that is, increases the time lag—as compared with undamped motion with the same initial conditions.



FIGURE 5.4.9. Underdamped oscillations: $x(t) = Ce^{-pt} \cos(\omega_1 t - \alpha).$



FIGURE 5.4.8. Critically damped motion: $x(t) = (c_1 + c_2t)e^{-pt}$ with p > 0. Solution curves are graphed

with the same initial position x_0 and

different initial velocities.

Example 2 Damped system The mass and spring of Example 1 are now attached also to a dashpot that provides 1 N of resistance for each meter per second of velocity. The mass is set in motion with the same initial position x(0) = 1 and initial velocity x'(0) = -5 as in Example 1. Now find the position function of the mass, its new frequency and pseudoperiod of motion, its new time lag, and the times of its first four passages through the initial position x = 0.

Solution Rather than memorizing the various formulas given in the preceding discussion, it is better practice in a particular case to set up the differential equation and then solve it directly. Recall that $m = \frac{1}{2}$ and k = 50; we are now given c = 1 in mks units. Hence Eq. (4) is $\frac{1}{2}x'' + x' + 50x = 0$; that is,

$$x'' + 2x' + 100x = 0.$$

The characteristic equation $r^2 + 2r + 100 = (r+1)^2 + 99 = 0$ has roots $r_1, r_2 = -1 \pm \sqrt{99}i$, so the general solution is

$$x(t) = e^{-t} (A \cos \sqrt{99}t + B \sin \sqrt{99}t).$$
(24)

Consequently, the new circular (pseudo)frequency is $\omega_1 = \sqrt{99} \approx 9.9499$ (as compared with $\omega_0 = 10$ in Example 1). The new (pseudo)period and frequency are

$$T_1 = \frac{2\pi}{\omega_1} = \frac{2\pi}{\sqrt{99}} \approx 0.6315 \text{ s}$$

and

$$v_1 = \frac{1}{T_1} = \frac{\omega_1}{2\pi} = \frac{\sqrt{99}}{2\pi} \approx 1.5836 \,\mathrm{Hz}$$

(as compared with $T \approx 0.6283 < T_1$ and $\nu \approx 1.5915 > \nu_1$ in Example 1).

We now impose the initial conditions x(0) = 1 and x'(0) = -5 on the position function in (23) and the resulting velocity function

$$x'(t) = -e^{-t}(A\cos\sqrt{99}t + B\sin\sqrt{99}t) + \sqrt{99}e^{-t}(-A\sin\sqrt{99}t + B\cos\sqrt{99}t).$$

It follows that

$$x(0) = A = 1$$
 and $x'(0) = -A + B\sqrt{99} = -5$,

whence we find that A = 1 and $B = -4/\sqrt{99}$. Thus the new position function of the body is

$$x(t) = e^{-t} \left(\cos \sqrt{99} t - \frac{4}{\sqrt{99}} \sin \sqrt{99} t \right).$$

Hence its time-varying amplitude of motion is

$$C_1 e^{-t} = \sqrt{(1)^2 + \left(-\frac{4}{\sqrt{99}}\right)^2} e^{-t} = \sqrt{\frac{115}{99}} e^{-t}.$$

We therefore write

$$\begin{aligned} x(t) &= \frac{\sqrt{115}}{\sqrt{99}} e^{-t} \left(\frac{\sqrt{99}}{\sqrt{115}} \cos \sqrt{99} t - \frac{4}{\sqrt{115}} \sin \sqrt{99} t \right) \\ &= \sqrt{\frac{115}{99}} e^{-t} \cos(\sqrt{99} t - \alpha_1), \end{aligned}$$

where the phase angle α_1 satisfies

$$\cos \alpha_1 = \frac{\sqrt{99}}{\sqrt{115}} > 0$$
 and $\sin \alpha_1 = -\frac{4}{\sqrt{115}} < 0.$

Hence α_1 is the fourth-quadrant angle

$$\alpha_1 = 2\pi + \tan^{-1}\left(\frac{-4/\sqrt{115}}{\sqrt{99}/\sqrt{115}}\right) = 2\pi - \tan^{-1}\left(\frac{4}{\sqrt{99}}\right) \approx 5.9009,$$
and the time lag of the motion is

$$\delta_1 = \frac{\alpha_1}{\omega_1} \approx 0.5931 \text{ s}$$

(as compared with $\delta \approx 0.5820 < \delta_1$ in Example 1). With the time-varying amplitude and approximate phase angle shown explicitly, the position function of the mass takes the form

$$x(t) \approx \sqrt{\frac{115}{99}} e^{-t} \cos(\sqrt{99}t - 5.9009),$$
 (25)

and its graph is the damped exponential that is shown in Fig. 5.4.10 (in comparison with the undamped oscillations of Example 1).

From (24) we see that the mass passes through its equilibrium position x = 0 when $\cos(\omega_1 t - \alpha_1) = 0$, and thus when

$$\omega_1 t - \alpha_1 = -\frac{3\pi}{2}, \quad -\frac{\pi}{2}, \quad \frac{\pi}{2}, \quad \frac{3\pi}{2}, \quad \dots$$

х

that is, when

$$t = \delta_1 - \frac{3\pi}{2\omega_1}, \quad \delta_1 - \frac{\pi}{2\omega_1}, \quad \delta_1 + \frac{\pi}{2\omega_1}, \quad \delta_1 + \frac{3\pi}{2\omega_1}, \quad \dots$$

We see similarly that the undamped mass of Example 1 passes through equilibrium when

$$t = \delta_0 - \frac{3\pi}{2\omega_0}, \quad \delta_0 - \frac{\pi}{2\omega_0}, \quad \delta_0 + \frac{\pi}{2\omega_0}, \quad \delta_0 + \frac{3\pi}{2\omega_0}, \quad \dots$$

The following table compares the first four values t_1, t_2, t_3, t_4 we calculate for the undamped and damped cases, respectively.

n	1	2	3	4	
t_n (undamped)	0.1107	0.4249	0.7390	1.0532	
t_n (damped)	0.1195	0.4352	0.7509	1.0667	

Accordingly, in Fig. 5.4.11 (where only the first three equilibrium passages are shown) we see the damped oscillations lagging slightly behind the undamped ones.



FIGURE 5.4.10. Graphs of the position function $x(t) = C_1 e^{-t} \cos(\omega_1 t - \alpha_1)$ of Example 2 (damped oscillations), the position function $x(t) = C \cos(\omega_0 t - \alpha)$ of Example 1 (undamped oscillations), and the envelope curves $x(t) = \pm C_1 e^{-t}$.



FIGURE 5.4.11. Graphs on the interval $0 \le t \le 0.8$ illustrating the additional delay associated with damping.

5.4 Problems

- **1.** Determine the period and frequency of the simple harmonic motion of a 4-kg mass on the end of a spring with spring constant 16 N/m.
- **2.** Determine the period and frequency of the simple harmonic motion of a body of mass 0.75 kg on the end of a spring with spring constant 48 N/m.
- **3.** A mass of 3 kg is attached to the end of a spring that is stretched 20 cm by a force of 15 N. It is set in motion with initial position $x_0 = 0$ and initial velocity $v_0 = -10$ m/s. Find the amplitude, period, and frequency of the resulting motion.
- 4. A body with mass 250 g is attached to the end of a spring that is stretched 25 cm by a force of 9 N. At time t = 0 the body is pulled 1 m to the right, stretching the spring, and set in motion with an initial velocity of 5 m/s to the left.
 (a) Find x(t) in the form C cos(ω₀t α). (b) Find the amplitude and period of motion of the body.

Simple Pendulum

In Problems 5 through 8, assume that the differential equation of a simple pendulum of length L is $L\theta'' + g\theta = 0$, where $g = GM/R^2$ is the gravitational acceleration at the location of the pendulum (at distance R from the center of the earth; M denotes the mass of the earth).

5. Two pendulums are of lengths L_1 and L_2 and—when located at the respective distances R_1 and R_2 from the center of the earth—have periods p_1 and p_2 . Show that

$$\frac{p_1}{p_2} = \frac{R_1\sqrt{L_1}}{R_2\sqrt{L_2}}$$

- 6. A certain pendulum keeps perfect time in Paris, where the radius of the earth is R = 3956 (mi). But this clock loses 2 min 40 s per day at a location on the equator. Use the result of Problem 5 to find the amount of the equatorial bulge of the earth.
- 7. A pendulum of length 100.10 in., located at a point at sea level where the radius of the earth is R = 3960 (mi), has the same period as does a pendulum of length 100.00



FIGURE 5.4.12. The buoy of Problem 10.

in. atop a nearby mountain. Use the result of Problem 5 to find the height of the mountain.

- **8.** Most grandfather clocks have pendulums with adjustable lengths. One such clock loses 10 min per day when the length of its pendulum is 30 in. With what length pendulum will this clock keep perfect time?
- **9.** Derive Eq. (5) describing the motion of a mass attached to the bottom of a vertically suspended spring. (*Suggestion:* First denote by x(t) the displacement of the mass below the unstretched position of the spring; set up the differential equation for x. Then substitute $y = x s_0$ in this differential equation.)
- **10.** Floating buoy Consider a floating cylindrical buoy with radius *r*, height *h*, and uniform density $\rho \leq 0.5$ (recall that the density of water is 1 g/cm³). The buoy is initially suspended at rest with its bottom at the top surface of the water and is released at time t = 0. Thereafter it is acted on by two forces: a downward gravitational force equal to its weight $mg = \rho \pi r^2 hg$ and (by Archimedes' principle of buoyancy) an upward force equal to the weight $\pi r^2 xg$ of water displaced, where x = x(t) is the depth of the bottom of the buoy beneath the surface at time t (Fig. 5.4.12). Assume that friction is negligible. Conclude that the buoy undergoes simple harmonic motion around its equilibrium position $x_e = \rho h$ with period $p = 2\pi \sqrt{\rho h/g}$. Compute p and the amplitude of the motion if $\rho = 0.5$ g/cm³, h = 200 cm, and g = 980 cm/s².
- 11. Floating buoy A cylindrical buoy weighing 100 lb (thus of mass m = 3.125 slugs in ft-lb-s (fps) units) floats in water with its axis vertical (as in Problem 10). When depressed slightly and released, it oscillates up and down four times every 10 s. Find the radius of the buoy.
- 12. Hole through the earth Assume that the earth is a solid sphere of uniform density, with mass M and radius R = 3960 (mi). For a particle of mass m within the earth at distance r from the center of the earth, the gravitational force attracting m toward the center is $F_r = -GM_rm/r^2$, where M_r is the mass of the part of the earth within a sphere of radius r (Fig. 5.4.13). (a) Show that $F_r = -GMmr/R^3$.



FIGURE 5.4.13. A mass *m* falling down a hole through the center of the earth (Problem 12).

(b) Now suppose that a small hole is drilled straight through the center of the earth, thus connecting two antipodal points on its surface. Let a particle of mass mbe dropped at time t = 0 into this hole with initial speed zero, and let r(t) be its distance from the center of the earth at time t, where we take r < 0 when the mass is "below" the center of the earth. Conclude from Newton's second law and part (a) that $r''(t) = -k^2 r(t)$, where $k^{2} = GM/R^{3} = g/R$. (c) Take g = 32.2 ft/s², and conclude from part (b) that the particle undergoes simple harmonic motion back and forth between the ends of the hole, with a period of about 84 min. (d) Look up (or derive) the period of a satellite that just skims the surface of the earth; compare with the result in part (c). How do you explain the coincidence? Or is it a coincidence? (e) With what speed (in miles per hour) does the particle pass through the center of the earth? (f) Look up (or derive) the orbital velocity of a satellite that just skims the surface of the earth; compare with the result in part (e). How do you explain the coincidence? Or is it a coincidence?

- 13. Suppose that the mass in a mass-spring-dashpot system with m = 10, c = 9, and k = 2 is set in motion with x(0) = 0 and x'(0) = 5. (a) Find the position function x(t) and show that its graph looks as indicated in Fig. 5.4.14. (b) Find how far the mass moves to the right before starting back toward the origin.
- 14. Suppose that the mass in a mass-spring-dashpot system with m = 25, c = 10, and k = 226 is set in motion with x(0) = 20 and x'(0) = 41. (a) Find the position function x(t) and show that its graph looks as indicated in Fig. 5.4.15. (b) Find the pseudoperiod of the oscillations and the equations of the "envelope curves" that are dashed in the figure.

Free Damped Motion

The remaining problems in this section deal with free damped motion. In Problems 15 through 21, a mass m is attached to both a spring (with given spring constant k) and a dashpot (with given damping constant c). The mass is set in motion with initial position x_0 and initial velocity v_0 . Find the position function x(t) and determine whether the motion is



FIGURE 5.4.14. The position function x(t) of Problem 13.

overdamped, critically damped, or underdamped. If it is underdamped, write the position function in the form $x(t) = C_1 e^{-pt} \cos(\omega_1 t - \alpha_1)$. Also, find the undamped position function $u(t) = C_0 \cos(\omega_0 t - \alpha_0)$ that would result if the mass on the spring were set in motion with the same initial position and velocity, but with the dashpot disconnected (so c = 0). Finally, construct a figure that illustrates the effect of damping by comparing the graphs of x(t) and u(t).

- **15.** $m = \frac{1}{2}, c = 3, k = 4; x_0 = 2, v_0 = 0$ **16.** $m = 3, c = 30, k = 63; x_0 = 2, v_0 = 2$ **17.** $m = 1, c = 8, k = 16; x_0 = 5, v_0 = -10$ **18.** $m = 2, c = 12, k = 50; x_0 = 0, v_0 = -8$ **19.** $m = 4, c = 20, k = 169; x_0 = 4, v_0 = 16$ **20.** $m = 2, c = 16, k = 40; x_0 = 5, v_0 = 4$
- **21.** $m = 1, c = 10, k = 125; x_0 = 6, v_0 = 50$
- **22.** Vertical damped motion A 12-lb weight (mass m = 0.375 slugs in fps units) is attached both to a vertically suspended spring that it stretches 6 in. and to a dashpot that provides 3 lb of resistance for every foot per second of velocity. (a) If the weight is pulled down 1 ft below its static equilibrium position and then released from rest at time t = 0, find its position function x(t). (b) Find the frequency, time-varying amplitude, and phase angle of the motion.
- **23.** Car suspension This problem deals with a highly simplified model of a car of weight 3200 lb (mass m = 100 slugs in fps units). Assume that the suspension system acts like a single spring and its shock absorbers like a single dashpot, so that its vertical vibrations satisfy Eq. (4) with appropriate values of the coefficients. (a) Find the stiffness coefficient k of the spring if the car undergoes free vibrations at 80 cycles per minute (cycles/min) when its shock absorbers are disconnected. (b) With the shock absorbers connected, the car is set into vibration by driving it over a bump, and the resulting damped vibrations have a frequency of 78 cycles/min. After how long will the time-varying amplitude be 1% of its initial value?

Problems 24 through 34 deal with a mass-spring-dashpot system having position function x(t) satisfying Eq. (4). We write



FIGURE 5.4.15. The position function x(t) of Problem 14.

 $x_0 = x(0)$ and $v_0 = x'(0)$ and recall that p = c/(2m), $\omega_0^2 = k/m$, and $\omega_1^2 = \omega_0^2 - p^2$. The system is critically damped, overdamped, or underdamped, as specified in each problem.

24. (Critically damped) Show in this case that

$$x(t) = (x_0 + v_0t + px_0t)e^{-pt}$$

- **25.** (Critically damped) Deduce from Problem 24 that the mass passes through x = 0 at some instant t > 0 if and only if x_0 and $v_0 + px_0$ have opposite signs.
- **26.** (Critically damped) Deduce from Problem 24 that x(t) has a local maximum or minimum at some instant t > 0 if and only if v_0 and $v_0 + px_0$ have the same sign.
- 27. (Overdamped) Show in this case that

$$x(t) = \frac{1}{2\gamma} \left[(v_0 - r_2 x_0) e^{r_1 t} - (v_0 - r_1 x_0) e^{r_2 t} \right],$$

where $r_1, r_2 = -p \pm \sqrt{p^2 - \omega_0^2}$ and $\gamma = (r_1 - r_2)/2 > 0$. **28.** (Overdamped) If $x_0 = 0$, deduce from Problem 27 that

$$x(t) = \frac{v_0}{\gamma} e^{-pt} \sinh \gamma t.$$

- **29.** (Overdamped) Prove that in this case the mass can pass through its equilibrium position x = 0 at most once.
- 30. (Underdamped) Show that in this case

$$x(t) = e^{-pt} \left(x_0 \cos \omega_1 t + \frac{v_0 + px_0}{\omega_1} \sin \omega_1 t \right).$$

31. (Underdamped) If the damping constant c is small in comparison with $\sqrt{8mk}$, apply the binomial series to show that

$$\omega_1 \approx \omega_0 \left(1 - \frac{c^2}{8mk} \right).$$

32. (Underdamped) Show that the local maxima and minima of

$$x(t) = Ce^{-pt}\cos(\omega_1 t - \alpha)$$

occur where

$$\tan(\omega_1 t - \alpha) = -\frac{p}{\omega_1}.$$

Conclude that $t_2 - t_1 = 2\pi/\omega_1$ if two consecutive maxima occur at times t_1 and t_2 .

33. (Underdamped) Let x_1 and x_2 be two consecutive local maximum values of x(t). Deduce from the result of Problem 32 that

$$\ln \frac{x_1}{x_2} = \frac{2\pi p}{\omega_1}.$$

The constant $\Delta = 2\pi p/\omega_1$ is called the **logarithmic decrement** of the oscillation. Note also that $c = m\omega_1 \Delta/\pi$ because p = c/(2m).

Note: The result of Problem 33 provides an accurate method for measuring the viscosity of a fluid, which is an important parameter in fluid dynamics but is not easy to measure directly.

According to Stokes's drag law, a spherical body of radius a moving at a (relatively slow) speed through a fluid of viscosity μ experiences a resistive force $F_R = 6\pi\mu av$. Thus if a spherical mass on a spring is immersed in the fluid and set in motion, this drag resistance damps its oscillations with damping constant $c = 6\pi a\mu$. The frequency ω_1 and logarithmic decrement Δ of the oscillations can be measured by direct observation. The final formula in Problem 33 then gives c and hence the viscosity of the fluid.

34. (Underdamped) A body weighing 100 lb (mass m = 3.125 slugs in fps units) is oscillating attached to a spring and a dashpot. Its first two maximum displacements of 6.73 in. and 1.46 in. are observed to occur at times 0.34 s and 1.17 s, respectively. Compute the damping constant (in pound-seconds per foot) and spring constant (in pounds per foot).

Differential Equations and Determinism

Given a mass m, a dashpot constant c, and a spring constant k, Theorem 2 of Section 5.1 implies that the equation

$$mx'' + cx' + kx = 0 \tag{26}$$

has a unique solution for $t \ge 0$ satisfying given initial conditions $x(0) = x_0$, $x'(0) = v_0$. Thus the future motion of an ideal mass-spring-dashpot system is completely determined by the differential equation and the initial conditions. Of course in a real physical system it is impossible to measure the parameters m, c, and k precisely. Problems 35 through 38 explore the resulting uncertainty in predicting the future behavior of a physical system.

35. Suppose that m = 1, c = 2, and k = 1 in Eq. (26). Show that the solution with x(0) = 0 and x'(0) = 1 is

$$x_1(t) = te^{-t}$$

36. Suppose that m = 1 and c = 2 but $k = 1 - 10^{-2n}$. Show that the solution of Eq. (26) with x(0) = 0 and x'(0) = 1 is

$$x_2(t) = 10^n e^{-t} \sinh 10^{-n} t.$$

37. Suppose that m = 1 and c = 2 but that $k = 1 + 10^{-2n}$. Show that the solution of Eq. (26) with x(0) = 0 and x'(0) = 1 is

$$x_3(t) = 10^n e^{-t} \sin 10^{-n} t.$$

38. Whereas the graphs of $x_1(t)$ and $x_2(t)$ resemble those shown in Figs. 5.4.7 and 5.4.8, the graph of $x_3(t)$ exhibits damped oscillations like those illustrated in Fig. 5.4.9, but with a very long pseudoperiod. Nevertheless, show that for each fixed t > 0 it is true that

$$\lim_{n \to \infty} x_2(t) = \lim_{n \to \infty} x_3(t) = x_1(t).$$

Conclude that *on a given finite time interval* the three solutions are in "practical" agreement if *n* is sufficiently large.

5.5 Nonhomogeneous Equations and Undetermined Coefficients

We learned in Section 5.3 how to solve homogeneous linear equations with constant coefficients, but we saw in Section 5.4 that an external force in a simple mechanical system contributes a nonhomogeneous term to its differential equation. The general nonhomogeneous nth-order linear equation with constant coefficients has the form

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_1 y' + a_0 y = f(x).$$
 (1)

By Theorem 5 of Section 5.2, a general solution of Eq. (1) has the form

$$y = y_c + y_p \tag{2}$$

where the complementary function $y_c(x)$ is a general solution of the associated homogeneous equation

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_1 y' + a_0 y = 0,$$
(3)

and $y_p(x)$ is a particular solution of Eq. (1). Thus our remaining task is to find y_p .

The **method of undetermined coefficients** is a straightforward way of doing this when the given function f(x) in Eq. (1) is sufficiently simple that we can make an intelligent guess as to the general form of y_p . For example, suppose that f(x)is a polynomial of degree m. Then, because the derivatives of a polynomial are themselves polynomials of lower degree, it is reasonable to suspect a particular solution

$$y_p(x) = A_m x^m + A_{m-1} x^{m-1} + \dots + A_1 x + A_0$$

that is also a polynomial of degree m, but with as yet undetermined coefficients. We may, therefore, substitute this expression for y_p into Eq. (1), and then—by equating coefficients of like powers of x on the two sides of the resulting equation—attempt to determine the coefficients A_0, A_1, \ldots, A_m so that y_p will, indeed, be a particular solution of Eq. (1).

Similarly, suppose that

$$f(x) = a\cos kx + b\sin kx.$$

Then it is reasonable to expect a particular solution of the same form:

$$y_p(x) = A\cos kx + B\sin kx,$$

a linear combination with undetermined coefficients A and B. The reason is that any derivative of such a linear combination of $\cos kx$ and $\sin kx$ has the same form. We may therefore substitute this form of y_p in Eq. (1), and then—by equating coefficients of $\cos kx$ and $\sin kx$ on both sides of the resulting equation—attempt to determine the coefficients A and B so that y_p will, indeed, be a particular solution.

It turns out that this approach does succeed whenever all the derivatives of f(x) have the same form as f(x) itself. Before describing the method in full generality, we illustrate it with several preliminary examples.

Example 1 Solution	Find a particular solution of $y'' + 3y' + 4y = 3x + 2$. Here $f(x) = 3x + 2$ is a polynomial of degree 1, so our guess is that
	$y_p(x) = Ax + B.$
	Then $y'_p = A$ and $y''_p = 0$, so y_p will satisfy the differential equation provided that
	(0) + 3(A) + 4(Ax + B) = 3x + 2,
	that is,
	(4A)x + (3A + 4B) = 3x + 2
	for all x. This will be true if the x-terms and constant terms on the two sides of this equation agree. It therefore suffices for A and B to satisfy the two linear equations $4A = 3$ and $3A + 4B = 2$ that we readily solve for $A = \frac{3}{4}$ and $B = -\frac{1}{16}$. Thus we have found the particular solution
	$y_p(x) = \frac{3}{4}x - \frac{1}{16}.$
Example 2 Solution	Find a particular solution of $y'' - 4y = 2e^{3x}$. Any derivative of e^{3x} is a constant multiple of e^{3x} , so it is reasonable to try
	$y_p(x) = Ae^{3x}.$
	Then $y_p'' = 9Ae^{3x}$, so the given differential equation will be satisfied provided that
	$9Ae^{3x} - 4(Ae^{3x}) = 2e^{3x};$
	that is, $5A = 2$, so that $A = \frac{2}{5}$. Thus our particular solution is $y_p(x) = \frac{2}{5}e^{3x}$.
Example 3	Find a particular solution of $3y'' + y' - 2y = 2\cos x$.
Solution	A first guess might be $y_p(x) = A \cos x$, but the presence of y' on the left-hand side signals that we probably need a term involving $\sin x$ as well. So we try
	$y_p(x) = A\cos x + B\sin x,$
	$y'_p(x) = -A\sin x + B\cos x,$
	$y_p''(x) = -A\cos x - B\sin x.$
	Then substitution of y_p and its derivatives into the given differential equation yields
	$3(-A\cos x - B\sin x) + (-A\sin x + B\cos x) - 2(A\cos x + B\sin x) = 2\cos x,$
	that is (collecting coefficients on the left),
	$(-5A + B)\cos x + (-A - 5B)\sin x = 2\cos x.$
	This will be true for all x provided that the cosine and sine terms on the two sides of this equation agree. It therefore suffices for A and B to satisfy the two linear equations
	-5A + B = 2,

with readily found solution $A = -\frac{5}{13}$, $B = \frac{1}{13}$. Hence a particular solution is

$$y_p(x) = -\frac{5}{13}\cos x + \frac{1}{13}\sin x.$$

The following example, which superficially resembles Example 2, indicates that the method of undetermined coefficients is not always quite so simple as we have made it appear.

-A - 5B = 0

Example 4 Find a particular solution of $y'' - 4y = 2e^{2x}$. Solution If we try $y_p(x) = Ae^{2x}$, we find that

$$y_p'' - 4y_p = 4Ae^{2x} - 4Ae^{2x} = 0 \neq 2e^{2x}.$$

Thus, no matter how A is chosen, Ae^{2x} cannot satisfy the given nonhomogeneous equation. In fact, the preceding computation shows that Ae^{2x} satisfies instead the associated *homogeneous* equation. Therefore, we should begin with a trial function $y_p(x)$ whose derivative involves both e^{2x} and something else that can cancel upon substitution into the differential equation to leave the e^{2x} term that we need. A reasonable guess is

$$y_p(x) = Axe^{2x}$$

for which

$$y'_p(x) = Ae^{2x} + 2Axe^{2x}$$
 and $y''_p(x) = 4Ae^{2x} + 4Axe^{2x}$.

Substitution into the original differential equation yields

$$(4Ae^{2x} + 4Axe^{2x}) - 4(Axe^{2x}) = 2e^{2x}.$$

The terms involving xe^{2x} obligingly cancel, leaving only $4Ae^{2x} = 2e^{2x}$, so that $A = \frac{1}{2}$. Consequently, a particular solution is

$$y_p(x) = \frac{1}{2}xe^{2x}.$$

(4)

The General Approach

Our initial difficulty in Example 4 resulted from the fact that $f(x) = 2e^{2x}$ satisfies the associated homogeneous equation. Rule 1, given shortly, tells what to do when we do not have this difficulty, and Rule 2 tells what to do when we do have it.

The method of undetermined coefficients applies whenever the function f(x) in Eq. (1) is a linear combination of (finite) products of functions of the following three types:

- **1.** A polynomial in *x*;
- **2.** An exponential function e^{rx} ;
- 3. $\cos kx$ or $\sin kx$.

Any such function, for example,

$$f(x) = (3 - 4x^2)e^{5x} - 4x^3\cos 10x,$$

has the crucial property that only *finitely* many linearly independent functions appear as terms (summands) in f(x) and its derivatives of all orders. In Rules 1 and 2 we assume that Ly = f(x) is a nonhomogeneous linear equation with constant coefficients and that f(x) is a function of this kind.

RULE 1 Method of Undetermined Coefficients

Suppose that no term appearing either in f(x) or in any of its derivatives satisfies the associated homogeneous equation Ly = 0. Then take as a trial solution for y_p a linear combination of all linearly independent such terms and their derivatives. Then determine the coefficients by substitution of this trial solution into the nonhomogeneous equation Ly = f(x). Note that this rule is not a theorem requiring proof; it is merely a procedure to be followed in searching for a particular solution y_p . If we succeed in finding y_p , then nothing more need be said. (It can be proved, however, that this procedure will always succeed under the conditions specified here.)

In practice we check the supposition made in Rule 1 by first using the characteristic equation to find the complementary function y_c , and then write a list of all the terms appearing in f(x) and its successive derivatives. If none of the terms in this list duplicates a term in y_c , then we proceed with Rule 1.

Example 5 Find a particular solution of

$$y'' + 4y = 3x^3. (5)$$

Solution The (familiar) complementary solution of Eq. (5) is

$$y_c(x) = c_1 \cos 2x + c_2 \sin 2x.$$

The function $f(x) = 3x^3$ and its derivatives are constant multiples of the linearly independent functions x^3 , x^2 , x, and 1. Because none of these appears in y_c , we try

$$y_p = Ax^3 + Bx^2 + Cx + D,$$

 $y'_p = 3Ax^2 + 2Bx + C,$
 $y''_p = 6Ax + 2B.$

Substitution in Eq. (5) gives

$$y_p'' + 4y_p = (6Ax + 2B) + 4(Ax^3 + Bx^2 + Cx + D)$$

= $4Ax^3 + 4Bx^2 + (6A + 4C)x + (2B + 4D) = 3x^3.$

We equate coefficients of like powers of x in the last equation to get

$$4A = 3,$$
 $4B = 0,$
 $6A + 4C = 0,$ $2B + 4D = 0$

with solution $A = \frac{3}{4}$, B = 0, $C = -\frac{9}{8}$, and D = 0. Hence a particular solution of Eq. (5) is

$$y_p(x) = \frac{3}{4}x^3 - \frac{9}{8}x.$$

Example 6 Solve the initial value problem

$$y'' - 3y' + 2y = 3e^{-x} - 10\cos 3x;$$

y(0) = 1, y'(0) = 2. (6)

Solution The characteristic equation $r^2 - 3r + 2 = 0$ has roots r = 1 and r = 2, so the complementary function is

$$y_c(x) = c_1 e^x + c_2 e^{2x}.$$

The terms involved in $f(x) = 3e^{-x} - 10\cos 3x$ and its derivatives are e^{-x} , $\cos 3x$, and $\sin 3x$. Because none of these appears in y_c , we try

$$y_p = Ae^{-x} + B\cos 3x + C\sin 3x,$$

$$y'_p = -Ae^{-x} - 3B\sin 3x + 3C\cos 3x,$$

$$y''_p = Ae^{-x} - 9B\cos 3x - 9C\sin 3x.$$

After we substitute these expressions into the differential equation in (6) and collect coefficients, we get

$$y_p'' - 3y_p' + 2y_p = 6Ae^{-x} + (-7B - 9C)\cos 3x + (9B - 7C)\sin 3x$$
$$= 3e^{-x} - 10\cos 3x.$$

We equate the coefficients of the terms involving e^{-x} , those involving $\cos 3x$, and those involving $\sin 3x$. The result is the system

$$6A = 3,$$

$$-7B - 9C = -10,$$

$$9B - 7C = 0$$

with solution $A = \frac{1}{2}$, $B = \frac{7}{13}$, and $C = \frac{9}{13}$. This gives the particular solution

$$y_p(x) = \frac{1}{2}e^{-x} + \frac{7}{13}\cos 3x + \frac{9}{13}\sin 3x,$$

which, however, does not have the required initial values in (6).

To satisfy those initial conditions, we begin with the general solution

$$y(x) = y_c(x) + y_p(x)$$

= $c_1 e^x + c_2 e^{2x} + \frac{1}{2} e^{-x} + \frac{7}{13} \cos 3x + \frac{9}{13} \sin 3x$,

with derivative

 $y'(x) = c_1 e^x + 2c_2 e^{2x} - \frac{1}{2}e^{-x} - \frac{21}{13}\sin 3x + \frac{27}{13}\cos 3x.$

The initial conditions in (6) lead to the equations

$$y(0) = c_1 + c_2 + \frac{1}{2} + \frac{7}{13} = 1,$$

$$y'(0) = c_1 + 2c_2 - \frac{1}{2} + \frac{27}{13} = 2$$

with solution $c_1 = -\frac{1}{2}$, $c_2 = \frac{6}{13}$. The desired particular solution is therefore

$$y(x) = -\frac{1}{2}e^x + \frac{6}{13}e^{2x} + \frac{1}{2}e^{-x} + \frac{7}{13}\cos 3x + \frac{9}{13}\sin 3x.$$

Example 7 Find the general form of a particular solution of

$$y^{(3)} + 9y' = x\sin x + x^2 e^{2x}.$$
(7)

Solution The characteristic equation $r^3 + 9r = 0$ has roots r = 0, r = -3i, and r = 3i. So the complementary function is

$$y_c(x) = c_1 + c_2 \cos 3x + c_3 \sin 3x.$$

The derivatives of the right-hand side in Eq. (7) involve the terms

$$\cos x$$
, $\sin x$, $x \cos x$, $x \sin x$,
 e^{2x} , xe^{2x} , and x^2e^{2x} .

Because there is no duplication with the terms of the complementary function, the trial solution takes the form

$$y_{p}(x) = A\cos x + B\sin x + Cx\cos x + Dx\sin x + Ee^{2x} + Fxe^{2x} + Gx^{2}e^{2x}.$$

Upon substituting y_p in Eq. (7) and equating coefficients of like terms, we get seven equations determining the seven coefficients A, B, C, D, E, F, and G.

The Case of Duplication

Now we turn our attention to the situation in which Rule 1 does not apply: Some of the terms involved in f(x) and its derivatives satisfy the associated homogeneous equation. For instance, suppose that we want to find a particular solution of the differential equation

$$(D-r)^{3}y = (2x-3)e^{rx}.$$
(8)

Proceeding as in Rule 1, our first guess would be

$$y_p(x) = Ae^{rx} + Bxe^{rx}.$$
(9)

This form of $y_p(x)$ will not be adequate because the complementary function of Eq. (8) is

$$y_c(x) = c_1 e^{rx} + c_2 x e^{rx} + c_3 x^2 e^{rx},$$
(10)

so substitution of (9) in the left-hand side of (8) would yield zero rather than $(2x - 3)e^{rx}$.

To see how to amend our first guess, we observe that

$$(D-r)^{2}[(2x-3)e^{rx}] = [D^{2}(2x-3)]e^{rx} = 0$$

by Eq. (13) of Section 5.3. If y(x) is *any* solution of Eq. (8) and we apply the operator $(D - r)^2$ to both sides, we see that y(x) is also a solution of the equation $(D - r)^5 y = 0$. The general solution of this *homogeneous* equation can be written as

$$y(x) = \underbrace{c_1 e^{rx} + c_2 x e^{rx} + c_3 x^2 e^{rx}}_{y_c} + \underbrace{A x^3 e^{rx} + B x^4 e^{rx}}_{y_p}.$$

Thus *every* solution of our original equation in (8) is the sum of a complementary function and a *particular solution* of the form

$$y_p(x) = Ax^3 e^{rx} + Bx^4 e^{rx}.$$
 (11)

Note that the right-hand side in Eq. (11) can be obtained by multiplying each term of our first guess in (9) by the least positive integral power of x (in this case, x^3) that suffices to eliminate duplication between the terms of the resulting trial solution $y_p(x)$ and the complementary function $y_c(x)$ given in (10). This procedure succeeds in the general case.

To simplify the general statement of Rule 2, we observe that to find a particular solution of the nonhomogeneous linear differential equation

$$Ly = f_1(x) + f_2(x),$$
 (12)

it suffices to find *separately* particular solutions $Y_1(x)$ and $Y_2(x)$ of the two equations

$$Ly = f_1(x)$$
 and $Ly = f_2(x)$, (13)

respectively. For linearity then gives

$$L[Y_1 + Y_2] = LY_1 + LY_2 = f_1(x) + f_2(x),$$

Example 8

and therefore $y_p = Y_1 + Y_2$ is a particular solution of Eq. (12). (This is a type of "superposition principle" for nonhomogeneous linear equations.)

Now our problem is to find a particular solution of the equation Ly = f(x), where f(x) is a linear combination of products of the elementary functions listed in (4). Thus f(x) can be written as a sum of terms each of the form

$$P_m(x)e^{rx}\cos kx \quad \text{or} \quad P_m(x)e^{rx}\sin kx, \tag{14}$$

where $P_m(x)$ is a polynomial in x of degree m. Note that any derivative of such a term is of the same form but with *both* sines and cosines appearing. The procedure by which we arrived earlier at the particular solution in (11) of Eq. (8) can be generalized to show that the following procedure is always successful.

RULE 2 Method of Undetermined Coefficients

If the function f(x) is of either form in (14), take as the trial solution

$$y_p(x) = x^s [(A_0 + A_1 x + A_2 x^2 + \dots + A_m x^m) e^{rx} \cos kx + (B_0 + B_1 x + B_2 x^2 + \dots + B_m x^m) e^{rx} \sin kx],$$
(15)

where s is the smallest nonnegative integer such that no term in y_p duplicates a term in the complementary function y_c . Then determine the coefficients in Eq. (15) by substituting y_p into the nonhomogeneous equation.

In practice we seldom need to deal with a function f(x) exhibiting the full generality in (14). The table in Fig. 5.5.1 lists the form of y_p in various common cases, corresponding to the possibilities m = 0, r = 0, and k = 0.

On the other hand, it is common to have

$$f(x) = f_1(x) + f_2(x),$$

where $f_1(x)$ and $f_2(x)$ are different functions of the sort listed in the table in Fig. 5.5.1. In this event we take as y_p the sum of the trial solutions for $f_1(x)$ and $f_2(x)$, choosing *s separately* for each part to eliminate duplication with the complementary function. This procedure is illustrated in Examples 8 through 10.

Find a particular solution of		
	$y^{(3)} + y'' = 3e^x + 4x^2.$	

(16)

f(x)	Ур
$P_m(x) = b_0 + b_1 x + b_2 x^2 + \dots + b_m x^m$	$x^{s}(A_{0} + A_{1}x + A_{2}x^{2} + \dots + A_{m}x^{m})$
$a\cos kx + b\sin kx$	$x^{s}(A\cos kx + B\sin kx)$
$e^{rx}(a\cos kx + b\sin kx)$	$x^{s}e^{rx}(A\cos kx + B\sin kx)$
$P_m(x)e^{rx}$	$x^s(A_0 + A_1x + A_2x^2 + \dots + A_mx^m)e^{rx}$
$P_m(x)(a\cos kx + b\sin kx)$	$x^{s}[(A_{0}+A_{1}x+\cdots+A_{m}x^{m})\cos kx$
	$+ (B_0 + B_1 x + \dots + B_m x^m) \sin kx]$

FIGURE 5.5.1. Substitutions in the method of undetermined coefficients.

Solution The characteristic equation $r^3 + r^2 = 0$ has roots $r_1 = r_2 = 0$ and $r_3 = -1$, so the complementary function is

$$v_c(x) = c_1 + c_2 x + c_3 e^{-x}.$$

As a first step toward our particular solution, we form the sum

$$(Ae^x) + (B + Cx + Dx^2).$$

The part Ae^x corresponding to $3e^x$ does not duplicate any part of the complementary function, but the part $B + Cx + Dx^2$ must be multiplied by x^2 to eliminate duplication. Hence we take

$$y_p = Ae^x + Bx^2 + Cx^3 + Dx^4,$$

$$y'_p = Ae^x + 2Bx + 3Cx^2 + 4Dx^3,$$

$$y''_p = Ae^x + 2B + 6Cx + 12Dx^2, \text{ and}$$

$$y_p^{(3)} = Ae^x + 6C + 24Dx.$$

Substitution of these derivatives in Eq. (16) yields

$$2Ae^{x} + (2B + 6C) + (6C + 24D)x + 12Dx^{2} = 3e^{x} + 4x^{2}.$$

The system of equations

$$2A = 3,$$
 $2B + 6C = 0,$
 $6C + 24D = 0,$ $12D = 4$

has the solution $A = \frac{3}{2}$, B = 4, $C = -\frac{4}{3}$, and $D = \frac{1}{3}$. Hence the desired particular solution is

$$y_p(x) = \frac{3}{2}e^x + 4x^2 - \frac{4}{3}x^3 + \frac{1}{3}x^4.$$

Example 9 Determine the appropriate form for a particular solution of

 $y'' + 6y' + 13y = e^{-3x} \cos 2x.$

Solution The characteristic equation $r^2 + 6r + 13 = 0$ has roots $-3 \pm 2i$, so the complementary function is

$$y_c(x) = e^{-3x}(c_1 \cos 2x + c_2 \sin 2x).$$

This is the same form as a first attempt $e^{-3x}(A \cos 2x + B \sin 2x)$ at a particular solution, so we must multiply by x to eliminate duplication. Hence we would take

$$y_p(x) = e^{-3x} (Ax \cos 2x + Bx \sin 2x).$$

Example 10 Determine the appropriate form for a particular solution of the fifth-order equation

$$(D-2)^3(D^2+9)y = x^2e^{2x} + x\sin 3x.$$

Solution The characteristic equation $(r-2)^3(r^2+9) = 0$ has roots r = 2, 2, 2, 3i, and -3i, so the complementary function is

$$y_c(x) = c_1 e^{2x} + c_2 x e^{2x} + c_3 x^2 e^{2x} + c_4 \cos 3x + c_5 \sin 3x.$$

As a first step toward the form of a particular solution, we examine the sum

$$[(A + Bx + Cx^2)e^{2x}] + [(D + Ex)\cos 3x + (F + Gx)\sin 3x].$$

To eliminate duplication with terms of $y_c(x)$, the first part—corresponding to x^2e^{2x} —must be multiplied by x^3 , and the second part—corresponding to $x \sin 3x$ —must be multiplied by x. Hence we would take

$$y_p(x) = (Ax^3 + Bx^4 + Cx^5)e^{2x} + (Dx + Ex^2)\cos 3x + (Fx + Gx^2)\sin 3x.$$

Variation of Parameters

Finally, let us point out the kind of situation in which the method of undetermined coefficients cannot be used. Consider, for example, the equation

$$y'' + y = \tan x,\tag{17}$$

which at first glance may appear similar to those considered in the preceding examples. Not so; the function $f(x) = \tan x$ has *infinitely many* linearly independent derivatives

$$\sec^2 x$$
, $2 \sec^2 x \tan x$, $4 \sec^2 x \tan^2 x + 2 \sec^4 x$, ...

Therefore, we do not have available a *finite* linear combination to use as a trial solution.

We discuss here the method of **variation of parameters**, which—in principle (that is, if the integrals that appear can be evaluated)—can always be used to find a particular solution of the nonhomogeneous linear differential equation

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + \dots + p_1(x)y' + p_0(x)y = f(x),$$
(18)

provided that we already know the general solution

$$y_c = c_1 y_1 + c_2 y_2 + \dots + c_n y_n \tag{19}$$

of the associated homogeneous equation

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + \dots + p_1(x)y' + p_0(x)y = 0.$$
 (20)

Here, in brief, is the basic idea of the method of variation of parameters. Suppose that we replace the constants, or *parameters*, c_1, c_2, \ldots, c_n in the complementary function in Eq. (19) with *variables*: functions u_1, u_2, \ldots, u_n of x. We ask whether it is possible to choose these functions in such a way that the combination

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x) + \dots + u_n(x)y_n(x)$$
(21)

is a particular solution of the nonhomogeneous equation in (18). It turns out that this *is* always possible.

The method is essentially the same for all orders $n \ge 2$, but we will describe it in detail only for the case n = 2. So we begin with the second-order nonhomogeneous equation

$$L[y] = y'' + P(x)y' + Q(x)y = f(x)$$
(22)

with complementary function

>

$$y_c(x) = c_1 y_1(x) + c_2 y_2(x)$$
(23)

on some open interval I where the functions P and Q are continuous. We want to find functions u_1 and u_2 such that

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x)$$
(24)

is a particular solution of Eq. (22).

5.5 Nonhomogeneous Equations and Undetermined Coefficients 323

One condition on the two functions u_1 and u_2 is that $L[y_p] = f(x)$. Because two conditions are required to determine two functions, we are free to impose an additional condition of our choice. We will do so in a way that simplifies the computations as much as possible. But first, to impose the condition $L[y_p] = f(x)$, we must compute the derivatives y'_p and y''_p . The product rule gives

$$y'_p = (u_1y'_1 + u_2y'_2) + (u'_1y_1 + u'_2y_2).$$

To avoid the appearance of the second derivatives u_1'' and u_2'' , the additional condition that we now impose is that the second sum here must vanish:

$$u_1'y_1 + u_2'y_2 = 0. (25)$$

Then

$$y'_p = u_1 y'_1 + u_2 y'_2, (26)$$

and the product rule gives

$$y_p'' = (u_1 y_1'' + u_2 y_2'') + (u_1' y_1' + u_2' y_2').$$
⁽²⁷⁾

But both y_1 and y_2 satisfy the homogeneous equation

y'' + Py' + Qy = 0

associated with the nonhomogeneous equation in (22), so

$$y_i'' = -Py_i' - Qy_i$$
 (28)

for i = 1, 2. It therefore follows from Eq. (27) that

$$y_p'' = (u_1'y_1' + u_2'y_2') - P \cdot (u_1y_1' + u_2y_2') - Q \cdot (u_1y_1 + u_2y_2).$$

In view of Eqs. (24) and (26), this means that

$$y_p'' = (u_1'y_1' + u_2'y_2') - Py_p' - Qy_p;$$

hence

>

$$L[y_p] = u'_1 y'_1 + u'_2 y'_2.$$
⁽²⁹⁾

The requirement that y_p satisfy the nonhomogeneous equation in (22)—that is, that $L[y_p] = f(x)$ —therefore implies that

$$u_1'y_1' + u_2'y_2' = f(x). (30)$$

Finally, Eqs. (25) and (30) determine the functions u_1 and u_2 that we need. Collecting these equations, we obtain a system

$$u'_{1}y_{1} + u'_{2}y_{2} = 0,$$

$$u'_{1}y'_{1} + u'_{2}y'_{2} = f(x)$$
(31)

of two linear equations in the two *derivatives* u'_1 and u'_2 . Note that the determinant of coefficients in (31) is simply the Wronskian $W(y_1, y_2)$. Once we have solved

>

the equations in (31) for the derivatives u'_1 and u'_2 , we integrate each to obtain the functions u_1 and u_2 such that

$$y_p = u_1 y_1 + u_2 y_2 \tag{32}$$

is the desired particular solution of Eq. (22). In Problem 63 we ask you to carry out this process explicitly and thereby verify the formula for $y_p(x)$ in the following theorem.

THEOREM 1 Variation of Parameters

If the nonhomogeneous equation y'' + P(x)y' + Q(x)y = f(x) has complementary function $y_c(x) = c_1y_1(x) + c_2y_2(x)$, then a particular solution is given by

$$y_p(x) = -y_1(x) \int \frac{y_2(x)f(x)}{W(x)} dx + y_2(x) \int \frac{y_1(x)f(x)}{W(x)} dx, \qquad (33)$$

where $W = W(y_1, y_2)$ is the Wronskian of the two independent solutions y_1 and y_2 of the associated homogeneous equation.

Example 11 Solution Find a particular solution of the equation $y'' + y = \tan x$.

The complementary function is $y_c(x) = c_1 \cos x + c_2 \sin x$, and we could simply substitute directly in Eq. (33). But it is more instructive to set up the equations in (31) and solve for u'_1 and u'_2 , so we begin with

$$y_1 = \cos x,$$
 $y_2 = \sin x,$
 $y'_1 = -\sin x,$ $y'_2 = \cos x.$

Hence the equations in (31) are

$$(u'_1)(\cos x) + (u'_2)(\sin x) = 0,$$

$$(u'_1)(-\sin x) + (u'_2)(\cos x) = \tan x.$$

We easily solve these equations for

$$u'_1 = -\sin x \tan x = -\frac{\sin^2 x}{\cos x} = \cos x - \sec x,$$
$$u'_2 = \cos x \tan x = \sin x.$$

Hence we take

$$u_1 = \int (\cos x - \sec x) \, dx = \sin x - \ln |\sec x + \tan x]$$

and

$$u_2 = \int \sin x \, dx = -\cos x.$$

(Do you see why we choose the constants of integration to be zero?) Thus our particular solution is

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x)$$

= (sin x - ln |sec x + tan x|) cos x + (- cos x)(sin x):

that is,

$$y_p(x) = -(\cos x) \ln |\sec x + \tan x|.$$

5.5 Problems

In Problems 1 through 20, find a particular solution y_p of the given equation. In all these problems, primes denote derivatives with respect to x.

1. $y'' + 16y = e^{3x}$ 2. y'' - y' - 2y = 3x + 43. $y'' - y' - 6y = 2\sin 3x$ 4. $4v'' + 4v' + v = 3xe^x$ 5. $v'' + v' + v = \sin^2 x$ 6. $2v'' + 4v' + 7v = x^2$ 7. $y'' - 4y = \sinh x$ 8. $y'' - 4y = \cosh 2x$ 9. $y'' + 2y' - 3y = 1 + xe^x$ 10. $y'' + 9y = 2\cos 3x + 3\sin 3x$ 11. $v^{(3)} + 4v' = 3x - 1$ 12. $v^{(3)} + v' = 2 - \sin x$ **13.** $v'' + 2v' + 5v = e^x \sin x$ **14.** $v^{(4)} - 2v'' + v = xe^x$ 16. $v'' + 9v = 2x^2e^{3x} + 5$ **15.** $v^{(5)} + 5v^{(4)} - v = 17$ 17. $y'' + y = \sin x + x \cos x$ 18. $v^{(4)} - 5v'' + 4v = e^x - xe^{2x}$ **19.** $v^{(5)} + 2v^{(3)} + 2v'' = 3x^2 - 1$ **20.** $y^{(3)} - y = e^x + 7$

In Problems 21 through 30, set up the appropriate form of a particular solution y_p , but do not determine the values of the coefficients.

21. $v'' - 2v' + 2v = e^x \sin x$ **22.** $y^{(5)} - y^{(3)} = e^x + 2x^2 - 5$ **23.** $y'' + 4y = 3x \cos 2x$ **24.** $y^{(3)} - y'' - 12y' = x - 2xe^{-3x}$ **25.** $y'' + 3y' + 2y = x(e^{-x} - e^{-2x})$ **26.** $v'' - 6v' + 13v = xe^{3x} \sin 2x$ 27. $v^{(4)} + 5v'' + 4v = \sin x + \cos 2x$ **28.** $v^{(4)} + 9v'' = (x^2 + 1)\sin 3x$ **29.** $(D-1)^3(D^2-4)y = xe^x + e^{2x} + e^{-2x}$ **30.** $v^{(4)} - 2v'' + v = x^2 \cos x$

Solve the initial value problems in Problems 31 through 40.

31. v'' + 4v = 2x; v(0) = 1, v'(0) = 2**32.** $y'' + 3y' + 2y = e^x$; y(0) = 0, y'(0) = 3**33.** $v'' + 9v = \sin 2x$; v(0) = 1, v'(0) = 0**34.** $y'' + y = \cos x$; y(0) = 1, y'(0) = -1**35.** v'' - 2v' + 2v = x + 1; v(0) = 3, v'(0) = 0**36.** $y^{(4)} - 4y'' = x^2$; y(0) = y'(0) = 1, $y''(0) = y^{(3)}(0) = -1$ **37.** $y^{(3)} - 2y'' + y' = 1 + xe^x$; y(0) = y'(0) = 0, y''(0) = 1**38.** $v'' + 2v' + 2v = \sin 3x$; v(0) = 2, v'(0) = 0**39.** $y^{(3)} + y'' = x + e^{-x}$; y(0) = 1, y'(0) = 0, y''(0) = 1**40.** $v^{(4)} - v = 5$; $v(0) = v'(0) = v''(0) = v^{(3)}(0) = 0$

41. Find a particular solution of the equation

$$y^{(4)} - y^{(3)} - y'' - y' - 2y = 8x^5$$

42. Find the solution of the initial value problem consisting of the differential equation of Problem 41 and the initial conditions

$$y(0) = y'(0) = y''(0) = y^{(3)}(0) = 0.$$

43. (a) Write

 $\cos 3x + i \sin 3x = e^{3ix} = (\cos x + i \sin x)^3$

by Euler's formula, expand, and equate real and imaginary parts to derive the identities

$$\cos^{3} x = \frac{3}{4} \cos x + \frac{1}{4} \cos 3x,$$

$$\sin^{3} x = \frac{3}{4} \sin x - \frac{1}{4} \sin 3x.$$

(b) Use the result of part (a) to find a general solution of

$$y'' + 4y = \cos^3 x.$$

Use trigonometric identities to find general solutions of the equations in Problems 44 through 46.

44. $y'' + y' + y = \sin x \sin 3x$ **45.** $y'' + 9y = \sin^4 x$ **46.** $v'' + v = x \cos^3 x$

In Problems 47 through 56, use the method of variation of parameters to find a particular solution of the given differential equation.

- **48.** $v'' 2v' 8v = 3e^{-2x}$ **47.** $y'' + 3y' + 2y = 4e^x$ **49.** $y'' - 4y' + 4y = 2e^{2x}$ **50.** $v'' - 4v = \sinh 2x$ **51.** $v'' + 4v = \cos 3x$ 52. $v'' + 9v = \sin 3x$ 54. $y'' + y = \csc^2 x$ **53.** $v'' + 9v = 2 \sec 3x$ **55.** $y'' + 4y = \sin^2 x$ 56. $v'' - 4v = xe^x$
- 57. You can verify by substitution that $y_c = c_1 x + c_2 x^{-1}$ is a complementary function for the nonhomogeneous secondorder equation

$$x^2y'' + xy' - y = 72x^5.$$

But before applying the method of variation of parameters, you must first divide this equation by its leading coefficient x^2 to rewrite it in the standard form

$$y'' + \frac{1}{x}y' - \frac{1}{x^2}y = 72x^3.$$

Thus $f(x) = 72x^3$ in Eq. (22). Now proceed to solve the equations in (31) and thereby derive the particular solution $y_p = 3x^5$.

In Problems 58 through 62, a nonhomogeneous second-order linear equation and a complementary function y_c are given. Apply the method of Problem 57 to find a particular solution of the equation.

- **58.** $x^2y'' 4xy' + 6y = x^3$; $y_c = c_1x^2 + c_2x^3$
- 50. $x^{2}y'' 4xy + 6y x^{3}$, $y_{c} c_{1}x^{3} + c_{2}x^{3}$ 59. $x^{2}y'' 3xy' + 4y = x^{4}$; $y_{c} = x^{2}(c_{1} + c_{2}\ln x)$ 60. $4x^{2}y'' 4xy' + 3y = 8x^{4/3}$; $y_{c} = c_{1}x + c_{2}x^{3/4}$
- **61.** $x^2y'' + xy' + y = \ln x$; $y_c = c_1 \cos(\ln x) + c_2 \sin(\ln x)$ **62.** $(x^2 1)y'' 2xy' + 2y = x^2 1$; $y_c = c_1x + c_2(1 + x^2)$
- 63. Carry out the solution process indicated in the text to derive the variation of parameters formula in (33) from Eqs. (31) and (32).
- 64. Apply the variation of parameters formula in (33) to find the particular solution $y_p(x) = -x \cos x$ of the nonhomogeneous equation $y'' + y = 2 \sin x$.

5.5 Application Automated Variation of Parameters

Go to goo.gl/0hPCVi to download this application's computing resources including *Maple/Mathematica/*MATLAB. The variation of parameters formula in (33) is especially amenable to implementation in a computer algebra system when the indicated integrals would be too tedious or inconvenient for manual evaluation. For example, suppose that we want to find a particular solution of the nonhomogeneous equation

$$y'' + y = \tan x$$

of Example 11, with complementary function $y_c(x) = c_1 \cos x + c_2 \sin x$. Then the *Maple* commands

```
y1 := cos(x):
y2 := sin(x):
f := tan(x):
W := y1*diff(y2,x) - y2*diff(y1,x):
W := simplify(W):
yp := -y1*int(y2*f/W,x) + y2*int(y1*f/W,x):
simplify(yp);
```

implement (33) and produce the result

$$y_p(x) = -(\cos x) \ln\left(\frac{1+\sin x}{\cos x}\right)$$

equivalent to the result $y_p(x) = -(\cos x) \ln(\sec x + \tan x)$ found in Example 11. The analogous *Mathematica* commands

```
y1 = Cos[x];
y2 = Sin[x];
f = Tan[x];
W = y1*D[y2,x] - y2*D[y1,x] // Simplify
yp = -y1*Integrate[y2*f/W,x] + y2*Integrate[y1*f/W,x];
Simplify[yp]
```

produce the result

$$y_p(x) = -(\cos x) \ln \left(\frac{\cos(x/2) + \sin(x/2)}{\cos(x/2) - \sin(x/2)} \right)$$

which (by the usual difference-of-squares technique) also is equivalent to the result found in Example 11.

To solve similarly a second-order linear equation y'' + P(x)y' + Q(x)y = f(x) whose complementary function $y_c(x) = c_1y_1(x) + c_2y_2(x)$ is known, we need only insert the corresponding definitions of $y_1(x)$, $y_2(x)$, and f(x) in the initial lines shown here. Find in this way the indicated particular solution $y_p(x)$ of the nonhomogeneous equations in Problems 1 through 6.

1.	$y'' + y = 2\sin x$	$y_p(x) = -x \cos x$
2.	$y'' + y = 4x \sin x$	$y_p(x) = x \sin x - x^2 \cos x$
3.	$y'' + y = 12x^2 \sin x$	$y_p(x) = 3x^2 \sin x + (3x - 2x^3) \cos x$
4.	$y'' - 2y' + 2y = 2e^x \sin x$	$y_p(x) = -xe^x \cos x$
5.	$y'' - 2y' + 2y = 4xe^x \sin x$	$y_p(x) = e^x (x \sin x - x^2 \cos x)$
6.	$y'' - 2y' + 2y = 12x^2 e^x \sin x$	$y_p(x) = e^x \left[3x^2 \sin x + (3x - 2x^3) \cos x \right]$

5.6 Forced Oscillations and Resonance

In Section 5.4 we derived the differential equation

$$mx'' + cx' + kx = F(t)$$
 (1)

that governs the one-dimensional motion of a mass m that is attached to a spring (with constant k) and a dashpot (with constant c) and is also acted on by an external force F(t). Machines with rotating components commonly involve mass-spring systems (or their equivalents) in which the external force is simple harmonic:

$$F(t) = F_0 \cos \omega t$$
 or $F(t) = F_0 \sin \omega t$, (2)

where the constant F_0 is the amplitude of the periodic force and ω is its circular frequency.

For an example of how a rotating machine component can provide a simple harmonic force, consider the cart with a rotating vertical flywheel shown in Fig. 5.6.1. The cart has mass $m - m_0$, not including the flywheel of mass m_0 . The centroid of the flywheel is off center at a distance *a* from its center, and its angular speed is ω radians per second. The cart is attached to a spring (with constant *k*) as shown. Assume that the centroid of the cart itself is directly beneath the center of the flywheel, and denote by x(t) its displacement from its equilibrium position (where the spring is unstretched). Figure 5.6.1 helps us to see that the displacement \overline{x} of the centroid of the combined cart plus flywheel is given by

$$\overline{x} = \frac{(m - m_0)x + m_0(x + a\cos\omega t)}{m} = x + \frac{m_0 a}{m}\cos\omega t$$

Let us ignore friction and apply Newton's second law $m\overline{x}'' = -kx$, because the force exerted by the spring is -kx. We substitute for \overline{x} in the last equation to obtain

$$mx'' - m_0 a\omega^2 \cos \omega t = -kx;$$

that is,

>

$$mx'' + kx = m_0 a\omega^2 \cos \omega t. \tag{3}$$

Thus the cart with its rotating flywheel acts like a mass on a spring under the influence of a simple harmonic external force with amplitude $F_0 = m_0 a \omega^2$. Such a system is a reasonable model of a front-loading washing machine with the clothes being washed loaded off center. This illustrates the practical importance of analyzing solutions of Eq. (1) with external forces as in (2).

Undamped Forced Oscillations

To study undamped oscillations under the influence of the external force $F(t) = F_0 \cos \omega t$, we set c = 0 in Eq. (1), and thereby begin with the equation

$$mx'' + kx = F_0 \cos \omega t \tag{4}$$

whose complementary function is $x_c = c_1 \cos \omega_0 t + c_2 \sin \omega_0 t$. Here

 $\omega_0 = \sqrt{\frac{k}{m}}$



FIGURE 5.6.1. The cart-with-flywheel system.

(as in Eq. (9) of Section 5.4) is the (circular) **natural frequency** of the mass–spring system. The fact that the angle $\omega_0 t$ is measured in (dimensionless) radians reminds us that if t is measured in seconds (s), then ω_0 is measured in radians per second—that is, in inverse seconds (s⁻¹). Also recall from Eq. (14) in Section 5.4 that division of a circular frequency ω by the number 2π of radians in a cycle gives the corresponding (ordinary) **frequency** $\nu = \omega/2\pi$ in Hz (hertz = cycles per second).

Let us assume initially that the external and natural frequencies are *unequal*: $\omega \neq \omega_0$. We substitute $x_p = A \cos \omega t$ in Eq. (4) to find a particular solution. (No sine term is needed in x_p because there is no term involving x' on the left-hand side in Eq. (4).) This gives

$$-m\omega^2 A\cos\omega t + kA\cos\omega t = F_0\cos\omega t,$$

so

 $A = \frac{F_0}{k - m\omega^2} = \frac{F_0/m}{\omega_0^2 - \omega^2},$ (5)

and thus

$$x_p(t) = \frac{F_0/m}{\omega_0^2 - \omega^2} \cos \omega t.$$
(6)

Therefore, the general solution $x = x_c + x_p$ is given by

$$x(t) = c_1 \cos \omega_0 t + c_2 \sin \omega_0 t + \frac{F_0/m}{\omega_0^2 - \omega^2} \cos \omega t,$$
(7)

where the constants c_1 and c_2 are determined by the initial values x(0) and x'(0). Equivalently, as in Eq. (12) of Section 5.4, we can rewrite Eq. (7) as

$$x(t) = C\cos(\omega_0 t - \alpha) + \frac{F_0/m}{\omega_0^2 - \omega^2}\cos\omega t,$$
(8)

so we see that the resulting motion is a superposition of two oscillations, one with natural circular frequency ω_0 , the other with the frequency ω of the external force.

Example 1 Suppose that
$$m = 1, k = 9, F_0 = 80$$
, and $\omega = 5$, so the differential equation in (4) is

$$x'' + 9x = 80\cos 5t.$$

Solution

Here the natural frequency $\omega_0 = 3$ and the frequency $\omega = 5$ of the external force are unequal, as in the preceding discussion. First we substitute $x_p = A \cos 5t$ in the differential equation and find that -25A + 9A = 80, so that A = -5. Thus a particular solution is

$$x_p(t) = -5\cos 5t.$$

The complementary function is $x_c = c_1 \cos 3t + c_2 \sin 3t$, so the general solution of the given nonhomogeneous equation is

$$x(t) = c_1 \cos 3t + c_2 \sin 3t - 5 \cos 5t,$$

with derivative

Find x(t) if x(0) = x'(0) = 0.

$$x'(t) = -3c_1 \sin 3t + 3c_2 \cos 3t + 25 \sin 5t.$$

The initial conditions x(0) = 0 and x'(0) = 0 now yield $c_1 = 5$ and $c_2 = 0$, so the desired particular solution is

$$x(t) = 5\cos 3t - 5\cos 5t.$$

FIGURE 5.6.2. The response $x(t) = 5 \cos 3t - 5 \cos 5t$ in Example 1.

As indicated in Fig. 5.6.2, the period of x(t) is the least common multiple 2π of the periods $2\pi/3$ and $2\pi/5$ of the two cosine terms.



Beats

If we impose the initial conditions x(0) = x'(0) = 0 on the solution in (7), we find that

$$c_1 = -\frac{F_0}{m(\omega_0^2 - \omega^2)}$$
 and $c_2 = 0$,

so the particular solution is

$$x(t) = \frac{F_0}{m(\omega_0^2 - \omega^2)} (\cos \omega t - \cos \omega_0 t).$$
 (9)

The trigonometric identity $2 \sin A \sin B = \cos(A - B) - \cos(A + B)$, applied with $A = \frac{1}{2}(\omega_0 + \omega)t$ and $B = \frac{1}{2}(\omega_0 - \omega)t$, enables us to rewrite Eq. (9) in the form

$$x(t) = \frac{2F_0}{m(\omega_0^2 - \omega^2)} \sin \frac{1}{2}(\omega_0 - \omega)t \sin \frac{1}{2}(\omega_0 + \omega)t.$$
(10)

Suppose now that $\omega \approx \omega_0$, so that $\omega_0 + \omega$ is very large in comparison with $|\omega_0 - \omega|$. Then $\sin \frac{1}{2}(\omega_0 + \omega)t$ is a *rapidly* varying function, whereas $\sin \frac{1}{2}(\omega_0 - \omega)t$ is a *slowly* varying function. We may therefore interpret Eq. (10) as a rapid oscillation with circular frequency $\frac{1}{2}(\omega_0 + \omega)$,

$$x(t) = A(t)\sin\frac{1}{2}(\omega_0 + \omega)t,$$

but with a slowly varying amplitude

With m = 0.1, $F_0 = 50$, $\omega_0 = 55$, and $\omega = 45$, Eq. (10) gives

$$A(t) = \frac{2F_0}{m(\omega_0^2 - \omega^2)} \sin \frac{1}{2}(\omega_0 - \omega)t.$$

Example 2

$$x(t) = \sin 5t \sin 50t.$$



FIGURE 5.6.3. The phenomenon of beats.

Figure 5.6.3 shows the corresponding oscillation of frequency $\frac{1}{2}(\omega_0 + \omega) = 50$ that is "modulated" by the amplitude function $A(t) = \sin 5t$ of frequency $\frac{1}{2}(\omega_0 - \omega) = 5$.

A rapid oscillation with a (comparatively) slowly varying periodic amplitude exhibits the phenomenon of *beats*. For example, if two horns not exactly attuned to one another simultaneously play their middle C, one at $\omega_0/(2\pi) = 258$ Hz and the other at $\omega/(2\pi) = 254$ Hz, then one hears a beat—an audible variation in the *amplitude* of the combined sound—with a frequency of

$$\frac{(\omega_0 - \omega)/2}{2\pi} = \frac{258 - 254}{2} = 2 \text{ (Hz)}.$$

Resonance

Looking at Eq. (6), we see that the amplitude A of x_p is large when the natural and external frequencies ω_0 and ω are approximately equal. It is sometimes useful to rewrite Eq. (5) in the form

$$A = \frac{F_0}{k - m\omega^2} = \frac{F_0/k}{1 - (\omega/\omega_0)^2} = \pm \frac{\rho F_0}{k},$$
(11)

where F_0/k is the **static displacement** of a spring with constant k due to a *constant* force F_0 , and the **amplification factor** ρ is defined to be

$$\rho = \frac{1}{|1 - (\omega/\omega_0)^2|}.$$
(12)

It is clear that $\rho \to +\infty$ as $\omega \to \omega_0$. This is the phenomenon of **resonance**—the increase without bound (as $\omega \to \omega_0$) in the amplitude of oscillations of an undamped system with natural frequency ω_0 in response to an external force with frequency $\omega \approx \omega_0$.

We have been assuming that $\omega \neq \omega_0$. What sort of catastrophe should one expect if ω and ω_0 are precisely equal? Then Eq. (4), upon division of each term by m, becomes

$$x'' + \omega_0^2 x = \frac{F_0}{m} \cos \omega_0 t.$$
 (13)

Because $\cos \omega_0 t$ is a term of the complementary function, the method of undetermined coefficients calls for us to try

$$x_p(t) = t(A\cos\omega_0 t + B\sin\omega_0 t).$$

We substitute this in Eq. (13) and thereby find that A = 0 and $B = F_0/(2m\omega_0)$. Hence the particular solution is

$$x_p(t) = \frac{F_0}{2m\omega_0} t \sin \omega_0 t.$$
⁽¹⁴⁾

The graph of $x_p(t)$ in Fig. 5.6.4 (in which m = 1, $F_0 = 100$, and $\omega_0 = 50$) shows vividly how the amplitude of the oscillation theoretically would increase without bound in this case of *pure resonance*, $\omega = \omega_0$. We may interpret this phenomenon as reinforcement of the natural vibrations of the system by externally impressed vibrations at the same frequency.

Cart with rotating flywheel Suppose that m = 5 kg and k = 500 N/m in the cart with the flywheel of Fig. 5.6.1. Then the natural frequency is $\omega_0 = \sqrt{k/m} = 10$ rad/s; that is, $10/(2\pi) \approx 1.59$ Hz. We would therefore expect oscillations of very large amplitude to occur if the flywheel revolves at about $(1.59)(60) \approx 95$ revolutions per minute (rpm).

In practice, a mechanical system with very little damping can be destroyed by resonance vibrations. A spectacular example can occur when a column of soldiers marches in step over a bridge. Any complicated structure such as a bridge has many natural frequencies of vibration. If the frequency of the soldiers' cadence is approximately equal to one of the natural frequencies of the structure, then—just as in our simple example of a mass on a spring-resonance will occur. Indeed, the resulting resonance vibrations can be of such large amplitude that the bridge will collapse. This has actually happened—for example, the collapse of Broughton Bridge near Manchester, England, in 1831—and it is the reason for the now-standard practice of breaking cadence when crossing a bridge. Resonance may have been involved in the 1981 Kansas City disaster in which a hotel balcony (called a *skywalk*) collapsed with dancers on it. The collapse of a building in an earthquake is sometimes due to resonance vibrations caused by the ground oscillating at one of the natural frequencies of the structure; this happened to many buildings in the Mexico City earthquake of September 19, 1985. On occasion an airplane has crashed because of resonant wing oscillations caused by vibrations of the engines. It is reported that



FIGURE 5.6.4. The phenomenon of resonance.

```
Example 3
```

for some of the first commercial jet aircraft, the natural frequency of the vertical vibrations of the airplane during turbulence was almost exactly that of the mass–spring system consisting of the pilot's head (mass) and spine (spring). Resonance occurred, causing pilots to have difficulty in reading the instruments. Large modern commercial jets have different natural frequencies, so that this resonance problem no longer occurs.

Modeling Mechanical Systems

The avoidance of destructive resonance vibrations is an ever-present consideration in the design of mechanical structures and systems of all types. Often the most important step in determining the natural frequency of vibration of a system is the formulation of its differential equation. In addition to Newton's law F = ma, the principle of conservation of energy is sometimes useful for this purpose (as in the derivation of the pendulum equation in Section 5.4). The following kinetic and potential energy formulas are often useful.

- **1.** *Kinetic energy:* $T = \frac{1}{2}mv^2$ for translation of a mass *m* with velocity *v*;
- 2. *Kinetic energy:* $T = \frac{1}{2}I\omega^2$ for rotation of a body of a moment of inertia *I* with angular velocity ω ;
- **3.** *Potential energy:* $V = \frac{1}{2}kx^2$ for a spring with constant k stretched or compressed a distance x;
- 4. *Potential energy:* V = mgh for the gravitational potential energy of a mass m at height h above the reference level (the level at which V = 0), provided that g may be regarded as essentially constant.

Example 4

Rolling disk Find the natural frequency of a mass m on a spring with constant k if, instead of sliding without friction, it is a uniform disk of radius a that rolls without slipping, as shown in Fig 5.6.5.

Solution With the preceding notation, the principle of conservation of energy gives

$$\frac{1}{2}mv^2 + \frac{1}{2}I\omega^2 + \frac{1}{2}kx^2 = E$$





where E is a constant (the total mechanical energy of the system). We note that $v = a\omega$ and recall that $I = ma^2/2$ for a uniform circular disk. Then we may simplify the last equation to

$$\frac{3}{4}mv^2 + \frac{1}{2}kx^2 = E$$

Because the right-hand side of this equation is constant, differentiation with respect to t (with v = x' and v' = x'') now gives

$$\frac{3}{2}mx'x'' + kxx' = 0.$$

We divide each term by $\frac{3}{2}mx'$ to obtain

$$x'' + \frac{2k}{3m}x = 0.$$

Thus the natural frequency of horizontal back-and-forth oscillation of our rolling disk is $\sqrt{2k/3m}$, which is $\sqrt{2/3} \approx 0.8165$ times the familiar natural frequency $\sqrt{k/m}$ of a mass on a spring that is sliding without friction rather than rolling without sliding. It is interesting (and perhaps surprising) that this natural frequency does not depend on the *radius* of the disk. It could be either a dime or a large disk with a radius of one meter (but of the same mass).

Example 5 Car suspension Suppose that a car oscillates vertically as if it were a mass m = 800 kg on a single spring (with constant $k = 7 \times 10^4$ N/m), attached to a single dashpot (with constant c = 3000 N·s/m). Suppose that this car with the dashpot *disconnected* is driven along a washboard road surface with an amplitude of 5 cm and a wavelength of L = 10 m (Fig. 5.6.6). At what car speed will resonance vibrations occur?







FIGURE 5.6.7. The "unicycle model" of a car.

Solution We think of the car as a unicycle, as pictured in Fig. 5.6.7. Let x(t) denote the upward displacement of the mass *m* from its equilibrium position; we ignore the force of gravity, because it merely displaces the equilibrium position as in Problem 9 of Section 5.4. We write the equation of the road surface as

$$y = a \cos \frac{2\pi s}{L}$$
 (a = 0.05 m, L = 10 m). (15)

When the car is in motion, the spring is stretched by the amount x - y, so Newton's second law, F = ma, gives mx'' = -k(x - y);

that is,

>

$$mx'' + kx = ky. \tag{16}$$

If the velocity of the car is v, then s = vt in Eq. (15), so Eq. (16) takes the form

$$mx'' + kx = ka\cos\frac{2\pi vt}{L}.$$
(16')

This is the differential equation that governs the vertical oscillations of the car. In comparing it with Eq. (4), we see that we have forced oscillations with circular frequency $\omega = 2\pi v/L$. Resonance will occur when $\omega = \omega_0 = \sqrt{k/m}$. We use our numerical data to find the speed of the car at resonance:

$$v = \frac{L}{2\pi} \sqrt{\frac{k}{m}} = \frac{10}{2\pi} \sqrt{\frac{7 \times 10^4}{800}} \approx 14.89 \text{ (m/s)};$$

that is, about 33.3 mi/h (using the conversion factor of 2.237 mi/h per m/s).

Damped Forced Oscillations

In real physical systems there is always some damping, from frictional effects if nothing else. The complementary function x_c of the equation

$$mx'' + cx' + kx = F_0 \cos \omega t \tag{17}$$

is given by Eq. (19), (20), or (21) of Section 5.4, depending on whether $c > c_{cr} = \sqrt{4km}$, $c = c_{cr}$, or $c < c_{cr}$. The specific form is not important here. What is important is that, in any case, these formulas show that $x_c(t) \to 0$ as $t \to +\infty$. Thus x_c is a **transient solution** of Eq. (17)—one that dies out with the passage of time, leaving only the particular solution x_p .

The method of undetermined coefficients indicates that we should substitute

$$x(t) = A\cos\omega t + B\sin\omega t$$

in Eq. (17). When we do this, collect terms, and equate coefficients of $\cos \omega t$ and $\sin \omega t$, we obtain the two equations

$$(k - m\omega^2)A + c\omega B = F_0, \qquad -c\omega A + (k - m\omega^2)B = 0$$
(18)

that we solve without difficulty for

$$A = \frac{(k - m\omega^2)F_0}{(k - m\omega^2)^2 + (c\omega)^2}, \quad B = \frac{c\omega F_0}{(k - m\omega^2)^2 + (c\omega)^2}.$$
 (19)

If we write

$$A\cos\omega t + B\sin\omega t = C(\cos\omega t\cos\alpha + \sin\omega t\sin\alpha) = C\cos(\omega t - \alpha)$$

as usual, we see that the resulting steady periodic oscillation

$$x_p(t) = C\cos(\omega t - \alpha) \tag{20}$$

has amplitude

$$C = \sqrt{A^2 + B^2} = \frac{F_0}{\sqrt{(k - m\omega^2)^2 + (c\omega)^2}}.$$
 (21)

Now (19) implies that $\sin \alpha = B/C > 0$, so it follows that the phase angle α lies in the first or second quadrant. Thus

$$\tan \alpha = \frac{B}{A} = \frac{c\omega}{k - m\omega^2} \quad \text{with} \quad 0 < \alpha < \pi,$$
(22)

so

$$\alpha = \begin{cases} \tan^{-1} \frac{c\omega}{k - m\omega^2} & \text{if } k > m\omega^2, \\ \pi + \tan^{-1} \frac{c\omega}{k - m\omega^2} & \text{if } k < m\omega^2 \end{cases}$$

(whereas $\alpha = \pi/2$ if $k = m\omega^2$).

Note that if c > 0, then the "forced amplitude"—defined as a function $C(\omega)$ by (21)—always remains finite, in contrast with the case of resonance in the undamped case when the forcing frequency ω equals the critical frequency $\omega_0 = \sqrt{k/m}$. But the forced amplitude may attain a maximum for some value of ω , in which case we speak of *practical resonance*. To see if and when practical resonance occurs, we need only graph *C* as a function of ω and look for a global maximum. It can be shown (Problem 27) that *C* is a steadily decreasing function of ω if $c \ge \sqrt{2km}$. But if $c < \sqrt{2km}$, then the amplitude of *C* attains a maximum value—and so practical resonance occurs—at some value of ω less than ω_0 , and then approaches zero as $\omega \to +\infty$. It follows that an underdamped system typically will undergo forced oscillations whose amplitude is

- Large if ω is close to the critical resonance frequency;
- Close to F_0/k if ω is very small;
- Very small if ω is very large.

Example 6

Practical resonance Find the transient motion and steady periodic oscillations of a damped mass-and-spring system with m = 1, c = 2, and k = 26 under the influence of an external force $F(t) = 82 \cos 4t$ with x(0) = 6 and x'(0) = 0. Also investigate the possibility of practical resonance for this system.

Solution

The resulting motion
$$x(t) = x_{tr}(t) + x_{sp}(t)$$
 of the mass satisfies the initial value problem
 $x'' + 2x' + 26x = 82\cos 4t; \quad x(0) = 6, \quad x'(0) = 0.$ (23)

Instead of applying the general formulas derived earlier in this section, it is better in a concrete problem to work it directly. The roots of the characteristic equation

$$r^{2} + 2r + 26 = (r+1)^{2} + 25 = 0$$

are $r = -1 \pm 5i$, so the complementary function is

$$x_c(t) = e^{-t}(c_1\cos 5t + c_2\sin 5t).$$

When we substitute the trial solution

$$x(t) = A\cos 4t + B\sin 4t$$

in the given equation, collect like terms, and equate coefficients of $\cos 4t$ and $\sin 4t$, we get the equations

$$10A + 8B = 82$$
$$-8A + 10B = 0$$

with solution A = 5, B = 4. Hence the general solution of the equation in (23) is

$$x(t) = e^{-t}(c_1\cos 5t + c_2\sin 5t) + 5\cos 4t + 4\sin 4t.$$

At this point we impose the initial conditions x(0) = 6, x'(0) = 0 and find that $c_1 = 1$ and $c_2 = -3$. Therefore, the transient motion and the steady periodic oscillation of the mass are given by

$$x_{\rm tr}(t) = e^{-t}(\cos 5t - 3\sin 5t)$$

and

$$x_{\rm sp}(t) = 5\cos 4t + 4\sin 4t = \sqrt{41} \left(\frac{5}{\sqrt{41}}\cos 4t + \frac{4}{\sqrt{41}}\sin 4t\right)$$
$$= \sqrt{41}\cos(4t - \alpha)$$

where $\alpha = \tan^{-1}\left(\frac{4}{5}\right) \approx 0.6747.$

Figure 5.6.8 shows graphs of the solution $x(t) = x_{tr}(t) + x_{sp}(t)$ of the initial value problem

$$x'' + 2x' + 26x = 82\cos 4t, \quad x(0) = x_0, \quad x'(0) = 0$$
(24)

for the different values $x_0 = -20$, -10, 0, 10, and 20 of the initial position. Here we see clearly what it means for the transient solution $x_{tr}(t)$ to "die out with the passage of time," leaving only the steady periodic motion $x_{sp}(t)$. Indeed, because $x_{tr}(t) \rightarrow 0$ exponentially, within a very few cycles the full solution x(t) and the steady periodic solution $x_{sp}(t)$ are virtually indistinguishable (whatever the initial position x_0).

To investigate the possibility of practical resonance in the given system, we substitute the values m = 1, c = 2, and k = 26 in (21) and find that the forced amplitude at frequency ω is

$$C(\omega) = \frac{82}{\sqrt{676 - 48\omega^2 + \omega^4}}$$

The graph of $C(\omega)$ is shown in Fig. 5.6.9. The maximum amplitude occurs when

$$C'(\omega) = \frac{-41(4\omega^3 - 96\omega)}{(676 - 48\omega^2 + \omega^4)^{3/2}} = \frac{-164\omega(\omega^2 - 24)}{(676 - 48\omega^2 + \omega^4)^{3/2}} = 0$$

Thus practical resonance occurs when the external frequency is $\omega = \sqrt{24}$ (a bit less than the mass-and-spring's undamped critical frequency of $\omega_0 = \sqrt{k/m} = \sqrt{26}$).



FIGURE 5.6.8. Solutions of the initial value problem in (24) with $x_0 = -20, -10, 0, 10, \text{ and } 20.$



FIGURE 5.6.9. Plot of amplitude C versus external frequency ω .

5.6 Problems

In Problems 1 through 6, express the solution of the given initial value problem as a sum of two oscillations as in Eq. (8). Throughout, primes denote derivatives with respect to time t. In Problems 1–4, graph the solution function x(t) in such a way that you can identify and label (as in Fig. 5.6.2) its period.

1. $x'' + 9x = 10 \cos 2t$; x(0) = x'(0) = 02. $x'' + 4x = 5 \sin 3t$; x(0) = x'(0) = 03. $x'' + 100x = 225 \cos 5t + 300 \sin 5t$; x(0) = 375, x'(0) = 04. $x'' + 25x = 90 \cos 4t$; x(0) = 0, x'(0) = 905. $mx'' + kx = F_0 \cos \omega t$ with $\omega \neq \omega_0$; $x(0) = x_0$, x'(0) = 06. $mx'' + kx = F_0 \cos \omega t$ with $\omega = \omega_0$; x(0) = 0, $x'(0) = v_0$

In each of Problems 7 through 10, find the steady periodic solution $x_{sp}(t) = C \cos(\omega t - \alpha)$ of the given equation mx'' + cx' + kx = F(t) with periodic forcing function F(t) of frequency ω . Then graph $x_{sp}(t)$ together with (for comparison) the adjusted forcing function $F_1(t) = F(t)/m\omega$.

7. $x'' + 4x' + 4x = 10\cos 3t$ 8. $x'' + 3x' + 5x = -4\cos 5t$ 9. $2x'' + 2x' + x = 3\sin 10t$

10. $x'' + 3x' + 3x = 8\cos 10t + 6\sin 10t$

In each of Problems 11 through 14, find and plot both the steady periodic solution $x_{sp}(t) = C \cos(\omega t - \alpha)$ of the given differential equation and the actual solution $x(t) = x_{sp}(t) + x_{tr}(t)$ that satisfies the given initial conditions.

11. $x'' + 4x' + 5x = 10 \cos 3t$; x(0) = x'(0) = 0**12.** $x'' + 6x' + 13x = 10 \sin 5t$; x(0) = x'(0) = 0

- **13.** $x'' + 2x' + 26x = 600 \cos 10t$; x(0) = 10, x'(0) = 0
- **14.** $x'' + 8x' + 25x = 200 \cos t + 520 \sin t$; x(0) = -30, x'(0) = -10

Each of Problems 15 through 18 gives the parameters for a forced mass-spring-dashpot system with equation mx'' +

 $cx' + kx = F_0 \cos \omega t$. Investigate the possibility of practical resonance of this system. In particular, find the amplitude $C(\omega)$ of steady periodic forced oscillations with frequency ω . Sketch the graph of $C(\omega)$ and find the practical resonance frequency ω (if any).

15.
$$m = 1, c = 2, k = 2, F_0 = 2$$

16.
$$m = 1, c = 4, k = 5, F_0 = 10$$

- **17.** $m = 1, c = 6, k = 45, F_0 = 50$
- **18.** $m = 1, c = 10, k = 650, F_0 = 100$
- **19.** A mass weighing 100 lb (mass m = 3.125 slugs in fps units) is attached to the end of a spring that is stretched 1 in. by a force of 100 lb. A force $F_0 \cos \omega t$ acts on the mass. At what frequency (in hertz) will resonance oscillations occur? Neglect damping.
- **20.** A front-loading washing machine is mounted on a thick rubber pad that acts like a spring; the weight W = mg (with $g = 9.8 \text{ m/s}^2$) of the machine depresses the pad exactly 0.5 cm. When its rotor spins at ω radians per second, the rotor exerts a vertical force $F_0 \cos \omega t$ newtons on the machine. At what speed (in revolutions per minute) will resonance vibrations occur? Neglect friction.
- **21.** Pendulum-spring system Figure 5.6.10 shows a mass *m* on the end of a pendulum (of length *L*) also attached to a horizontal spring (with constant *k*). Assume small oscillations of *m* so that the spring remains essentially horizontal and neglect damping. Find the natural circular frequency ω_0 of motion of the mass in terms of *L*, *k*, *m*, and the gravitational constant *g*.
- **22.** Pulley-spring system A mass *m* hangs on the end of a cord around a pulley of radius *a* and moment of inertia *I*, as shown in Fig. 5.6.11. The rim of the pulley is attached to a spring (with constant k). Assume small oscillations so that the spring remains essentially horizontal and neglect friction. Find the natural circular frequency of the system in terms of *m*, *a*, *k*, *I*, and *g*.



FIGURE 5.6.10. The pendulum-and-spring system of Problem 21.

- **23.** Earthquake A building consists of two floors. The first floor is attached rigidly to the ground, and the second floor is of mass m = 1000 slugs (fps units) and weighs 16 tons (32,000 lb). The elastic frame of the building behaves as a spring that resists horizontal displacements of the second floor; it requires a horizontal force of 5 tons to displace the second floor a distance of 1 ft. Assume that in an earthquake the ground oscillates horizontally with amplitude A_0 and circular frequency ω , resulting in an external horizontal force $F(t) = mA_0\omega^2 \sin \omega t$ on the second floor. (a) What is the natural frequency (in hertz) of oscillations of the second floor? (b) If the ground undergoes one oscillation every 2.25 s with an amplitude of 3 in., what is the amplitude of the resulting forced oscillations of the second floor?
- **24.** A mass on a spring without damping is acted on by the external force $F(t) = F_0 \cos^3 \omega t$. Show that there are *two* values of ω for which resonance occurs, and find both.
- 25. Derive the steady periodic solution of

$$mx'' + cx' + kx = F_0 \sin \omega t.$$

In particular, show that it is what one would expect—the same as the formula in (20) with the same values of *C* and ω , except with $\sin(\omega t - \alpha)$ in place of $\cos(\omega t - \alpha)$.

26. Given the differential equation

$$mx'' + cx' + kx = E_0 \cos \omega t + F_0 \sin \omega t$$

$$x_{\rm sp}(t) = \frac{\sqrt{E_0^2 + F_0^2}}{\sqrt{(k - m\omega^2)^2 + (c\omega)^2}} \cos(\omega t - \alpha - \beta),$$

where α is defined in Eq. (22) and $\beta = \tan^{-1}(F_0/E_0)$. (*Suggestion:* Add the steady periodic solutions separately corresponding to $E_0 \cos \omega t$ and $F_0 \sin \omega t$ (see Problem 25).)

27. According to Eq. (21), the amplitude of forced steady periodic oscillations for the system mx'' + cx' + kx =



FIGURE 5.6.11. The mass–spring–pulley system of Problem 22.

 $F_0 \cos \omega t$ is given by

$$C(\omega) = \frac{F_0}{\sqrt{(k - m\omega^2)^2 + (c\omega)^2}}$$

(a) If $c \ge c_{\rm cr}/\sqrt{2}$, where $c_{\rm cr} = \sqrt{4km}$, show that C steadily decreases as ω increases. (b) If $c < c_{\rm cr}/\sqrt{2}$, show that C attains a maximum value (practical resonance) when

$$\omega = \omega_m = \sqrt{\frac{k}{m} - \frac{c^2}{2m^2}} < \omega_0 = \sqrt{\frac{k}{m}}.$$

28. As indicated by the cart-with-flywheel example discussed in this section, an unbalanced rotating machine part typically results in a force having amplitude proportional to the *square* of the frequency ω. (a) Show that the amplitude of the steady periodic solution of the differential equation

$$mx'' + cx' + kx = mA\omega^2 \cos \omega t$$

(with a forcing term similar to that in Eq. (17)) is given by

$$C(\omega) = \frac{mA\omega^2}{\sqrt{(k - m\omega^2)^2 + (c\omega)^2}}.$$

(b) Suppose that $c^2 < 2mk$. Show that the maximum amplitude occurs at the frequency ω_m given by

$$\omega_m = \sqrt{\frac{k}{m} \left(\frac{2mk}{2mk - c^2}\right)}.$$

Thus the resonance frequency in this case is *larger* (in contrast with the result of Problem 27) than the natural frequency $\omega_0 = \sqrt{k/m}$. (*Suggestion:* Maximize the *square* of *C*.)

Automobile Vibrations

Problems 29 and 30 deal further with the car of Example 5. Its upward displacement function satisfies the equation mx'' + cx' + kx = cy' + ky when the shock absorber is connected (so that c > 0). With $y = a \sin \omega t$ for the road surface, this differential equation becomes

$$mx'' + cx' + kx = E_0 \cos \omega t + F_0 \sin \omega t$$

where $E_0 = c\omega a$ and $F_0 = ka$.

29. Apply the result of Problem 26 to show that the amplitude *C* of the resulting steady periodic oscillation for the car is given by

$$C = \frac{a\sqrt{k^{2} + (c\omega)^{2}}}{\sqrt{(k - m\omega^{2})^{2} + (c\omega)^{2}}}$$

Because $\omega = 2\pi v/L$ when the car is moving with velocity v, this gives C as a function of v.

30. Figure 5.6.12 shows the graph of the amplitude function $C(\omega)$ using the numerical data given in Example 5 (including $c = 3000 \text{ N} \cdot \text{s/m}$). It indicates that, as the car accelerates gradually from rest, it initially oscillates with amplitude slightly over 5 cm. Maximum resonance vibrations with amplitude about 14 cm occur around 32 mi/h,

but then subside to more tolerable levels at high speeds. Verify these graphically based conclusions by analyzing the function $C(\omega)$. In particular, find the practical resonance frequency and the corresponding amplitude.



FIGURE 5.6.12. Amplitude of vibrations of the car on a washboard surface.

5.6 Application Forced Vibrations

Go to goo.gl/l2hK16 to

download this application's computing resources including *Maple/Mathematica/*MATLAB.

Here we investigate forced vibrations of the mass-spring-dashpot system with equation

$$mx'' + cx' + kx = F(t).$$
 (1)

To simplify the notation, let's take $m = p^2$, c = 2p, and $k = p^2q^2 + 1$, where p > 0 and q > 0. Then the complementary function of Eq. (1) is

$$x_c(t) = e^{-t/p} (c_1 \cos qt + c_2 \sin qt).$$
(2)

We will take p = 5, q = 3, and thus investigate the transient and steady periodic solutions corresponding to

$$25x'' + 10x' + 226x = F(t), \quad x(0) = 0, \quad x'(0) = 0$$
(3)

with several illustrative possibilities for the external force F(t). For your personal investigations to carry out similarly, you might select integers p and q with $6 \le p \le 9$ and $2 \le q \le 5$.

INVESTIGATION 1: With periodic external force $F(t) = 901 \cos 3t$, the MATLAB commands

produce the plot shown in Fig. 5.6.13. We see the (transient plus steady periodic) solution

$$x(t) = \cos 3t + 30\sin 3t + e^{-t/5} \left(-\cos 3t - \frac{451}{15}\sin 3t \right)$$

rapidly "building up" to the steady periodic oscillation $x_{sp}(t) = \cos 3t + 30 \sin 3t$.



FIGURE 5.6.13. The solution $x(t) = x_{tr}(t) + x_{sp}(t)$ and the steady periodic solution $x(t) = x_{sp}(t)$ with periodic external force $F(t) = 901 \cos 3t$.

INVESTIGATION 2: With damped oscillatory external force



FIGURE 5.6.14. The solution $x(t) = 6te^{-t/5} \sin 3t$ and the envelope curves $x(t) = \pm 6te^{-t/5}$ with damped oscillatory force $F(t) = 900e^{-t/5} \cos 3t$.



FIGURE 5.6.15. The solution $x(t) = e^{-t/5}[3t \cos t + (9t^2 - 1) \sin 3t]$ and the envelope curves $x(t) = \pm e^{-t/5} \sqrt{(3t)^2 + (9t^2 - 1)^2}$ with external force $F(t) = 2700te^{-t/5} \cos 3t.$

$$F(t) = 900e^{-t/5}\cos 3t,$$

we have duplication with the complementary function in (2). The Maple commands

produce the plot shown in Fig. 5.6.14. We see the solution

$$x(t) = 6t e^{-t/5} \sin 3t$$

oscillating up and down between the envelope curves $x = \pm 6te^{-t/5}$. (Note the factor of t that signals a resonance situation.)

INVESTIGATION 3: With damped oscillatory external force

$$F(t) = 2700t e^{-t/5} \cos 3t,$$

we have a still more complicated resonance situation. The Mathematica commands

produce the plot shown in Fig. 5.6.15. We see the solution

$$x(t) = e^{-t/5} \left[3t \cos 3t + (9t^2 - 1) \sin 3t \right]$$

oscillating up and down between the envelope curves

$$x = \pm e^{-t/5} \sqrt{(3t)^2 + (9t^2 - 1)^2}$$

6 Eigenvalues and Eigenvectors

6.1 Introduction to Eigenvalues

Given a square matrix **A**, let us pose the following question: Does there exist a nonzero vector **v** such that the result **Av** of multiplying **v** by the matrix **A** is simply a scalar multiple of **v**? Thus we ask whether or not there exist a nonzero vector **v** and a scalar λ such that

$$\mathbf{A}\mathbf{v} = \lambda \mathbf{v}.\tag{1}$$

Section 6.3 and subsequent chapters include interesting applications in which this question arises. The following definition provides the terminology that we use in discussing Eq. (1).

DEFINITION Eigenvalues and Eigenvectors

The number λ is said to be an **eigenvalue** of the $n \times n$ matrix **A** provided there exists a *nonzero* vector **v** such that

$$\mathbf{A}\mathbf{v} = \lambda \mathbf{v},\tag{1}$$

in which case the vector **v** is called an **eigenvector** of the matrix **A**. We also say that the eigenvector **v** is **associated** with the eigenvalue λ , or that the eigenvalue λ corresponds to the eigenvector **v**.

The prefix *eigen* is a German word that (in some contexts) may be translated as *proper*, so eigenvalues are sometimes called *proper values*. Eigenvalues and eigenvectors are also called *characteristic values* and *characteristic vectors* in some books.

Example 1

Consider the 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} 5 & -6 \\ 2 & -2 \end{bmatrix}.$$

If $\mathbf{v} = (2, 1) = \begin{bmatrix} 2 & 1 \end{bmatrix}^T$, then

$$\mathbf{A}\mathbf{v} = \begin{bmatrix} 5 & -6 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix} = 2\mathbf{v}.$$

Thus, $\mathbf{v} = \begin{bmatrix} 2 & 1 \end{bmatrix}^T$ is an eigenvector of **A** that is associated with the eigenvalue $\lambda = 2$. If $\mathbf{v} = \begin{bmatrix} 3 & 2 \end{bmatrix}^T$, then

$$\mathbf{A}\mathbf{v} = \begin{bmatrix} 5 & -6\\ 2 & -2 \end{bmatrix} \begin{bmatrix} 3\\ 2 \end{bmatrix} = \begin{bmatrix} 3\\ 2 \end{bmatrix} = 1\mathbf{v},$$

so $\mathbf{v} = \begin{bmatrix} 3 & 2 \end{bmatrix}^T$ is an eigenvector of **A** associated with the eigenvalue $\lambda = 1$. In summary, we see that the scalars $\lambda_1 = 2$ and $\lambda_2 = 1$ are both eigenvalues of the matrix **A**; they correspond to the eigenvectors $\mathbf{v}_1 = \begin{bmatrix} 2 & 1 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} 3 & 2 \end{bmatrix}^T$, respectively.

Eigenvalues and eigenvectors have a simple geometric interpretation. Suppose that λ is an eigenvalue of the matrix **A** and has the associated eigenvector **v**, so that $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$. Then the length $|\mathbf{A}\mathbf{v}|$ of the vector $\mathbf{A}\mathbf{v}$ is $\pm \lambda |\mathbf{v}|$, depending on the sign of λ . Thus, if $\lambda > 0$, then multiplication of **v** by the matrix **A** expands or contracts the vector **v** while preserving its direction; if $\lambda < 0$, then multiplication of **v** by **A** reverses the direction of **v** (see Fig. 6.1.1).

Remark 1 If $\mathbf{v} = \mathbf{0}$, then the equation $\mathbf{A}\mathbf{v} = \lambda \mathbf{v}$ holds for every scalar λ and hence is of no significance. This is why only *nonzero* vectors qualify as eigenvectors in the definition.

Remark 2 Let λ and **v** be an eigenvalue and associated eigenvector of the matrix **A**. If *k* is any nonzero scalar and $\mathbf{u} = k\mathbf{v}$, then

$$\mathbf{A}\mathbf{u} = \mathbf{A}(k\mathbf{v}) = k(\mathbf{A}\mathbf{v}) = k(\lambda\mathbf{v}) = \lambda(k\mathbf{v}) = \lambda\mathbf{u},$$

so $\mathbf{u} = k\mathbf{v}$ is also an eigenvector associated with λ . Thus, any nonzero scalar multiple of an eigenvector is also an eigenvector and is associated with the same eigenvalue. In Example 1, for instance, $\mathbf{u}_1 = -3\mathbf{v}_1 = \begin{bmatrix} -6 & -3 \end{bmatrix}^T$ is an eigenvector associated with $\lambda_1 = 2$, and $\mathbf{u}_2 = 4\mathbf{v}_2 = \begin{bmatrix} 12 & 8 \end{bmatrix}^T$ is an eigenvector associated with $\lambda_2 = 1$.

The Characteristic Equation

We now attack the problem of finding the eigenvalues and eigenvectors of a given $n \times n$ square matrix **A**. According to the definition, the nonzero vector **v** is an eigenvector of **A** associated with the eigenvalue λ exactly when

$$A\mathbf{v} = \lambda \mathbf{v} = \lambda \mathbf{I}\mathbf{v}$$

that is, when

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}.$$
 (2)

For a fixed value of λ , Eq. (2) is a homogeneous linear system of *n* equations in the *n* components of **v**. By Theorem 2 in Section 3.6 and Theorem 7 in Section 3.5, this system has a nontrivial solution $\mathbf{v} \neq \mathbf{0}$ if and only if the determinant

$$\det(\mathbf{A} - \lambda \mathbf{I}) = |\mathbf{A} - \lambda \mathbf{I}|$$

of its coefficient matrix is *zero*. The equation $|\mathbf{A} - \lambda \mathbf{I}| = 0$ is called the **characteristic equation** of the square matrix \mathbf{A} , and we have proved that there exists an eigenvector \mathbf{v} associated with λ if and only if λ satisfies this equation.



FIGURE 6.1.1. (a) A positive eigenvalue; (b) a negative eigenvalue.

THEOREM 1 The Characteristic Equation

The number λ is an eigenvalue of the $n \times n$ matrix **A** if and only if λ satisfies the characteristic equation

$$\mathbf{A} - \lambda \mathbf{I} = 0. \tag{3}$$

Now let us see just what sort of equation the characteristic equation in (3) is. Note that the matrix

$$\mathbf{A} - \lambda \mathbf{I} = \begin{bmatrix} a_{11} - \lambda & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} - \lambda & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} - \lambda \end{bmatrix}$$
(4)

is obtained simply by subtracting λ from each diagonal element of **A**. If we think of expanding the determinant by cofactors, we see that $|\mathbf{A} - \lambda \mathbf{I}|$ is a *polynomial* in the variable λ , and that the highest power of λ comes from the product of the diagonal elements of the matrix in (4). Therefore, the characteristic equation of the $n \times n$ matrix **A** takes the form

$$(-1)^{n}\lambda^{n} + c_{n-1}\lambda^{n-1} + \dots + c_{1}\lambda + c_{0} = 0,$$
(5)

an *n*th-degree polynomial equation in λ .

According to the fundamental theorem of algebra, every *n*th-degree polynomial equation in one variable has *n* solutions (counting multiple solutions), but some of them can be complex. Hence, we can say that an $n \times n$ matrix **A** always has *n* eigenvalues, though they might not be distinct and might not all be real. In this chapter we confine our attention mainly to real eigenvalues, but in Chapter 7 we will see important applications of complex eigenvalues to the solution of linear systems of differential equations.

ALGORITHM Eigenvalues and Eigenvectors

To find the eigenvalues and associated eigenvectors of the $n \times n$ matrix **A**:

1. First solve the characteristic equation

$$\mathbf{A} - \lambda \mathbf{I} = 0.$$

2. Then, for *each* eigenvalue λ thereby found, solve the linear system

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$$

to find the eigenvectors associated with λ .

Solving the characteristic equation is almost always easier said than done. In the examples that follow and in the problems, we have chosen matrices for which the characteristic polynomial $|\mathbf{A} - \lambda \mathbf{I}|$ readily factors to reveal the eigenvalues.

Example 2

Find the eigenvalues and associated eigenvectors of the matrix

$$\mathbf{A} = \begin{bmatrix} 5 & 7 \\ -2 & -4 \end{bmatrix}.$$

Solution Here

$$\mathbf{A} - \lambda \mathbf{I} = \begin{bmatrix} 5 - \lambda & 7\\ -2 & -4 - \lambda \end{bmatrix},\tag{6}$$

so the characteristic equation of A is

$$0 = |\mathbf{A} - \lambda \mathbf{I}|$$

= $\begin{vmatrix} 5 - \lambda & 7 \\ -2 & -4 - \lambda \end{vmatrix}$
= $(5 - \lambda)(-4 - \lambda) - (-2)(7)$
= $(\lambda - 5)(\lambda + 4) + 14 = \lambda^2 - \lambda - 6;$

that is, $(\lambda + 2)(\lambda - 3) = 0$. Thus, the matrix **A** has the two eigenvalues -2 and 3. To distinguish them, we write $\lambda_1 = -2$ and $\lambda_2 = 3$. To find the associated eigenvectors, we must *separately* substitute each eigenvalue in (6) and then solve the resulting system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$.

CASE 1:
$$\lambda_1 = -2$$
. With $\mathbf{v} = \begin{bmatrix} x & y \end{bmatrix}^T$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is $\begin{bmatrix} 7 & 7 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$.

Each of the two scalar equations here is a multiple of the equation x + y = 0, and any nontrivial solution $\mathbf{v} = \begin{bmatrix} x & y \end{bmatrix}^T$ of this equation is a nonzero multiple of $\begin{bmatrix} 1 & -1 \end{bmatrix}^T$. Hence, to within a constant multiple, the only eigenvector associated with $\lambda_1 = -2$ is $\mathbf{v}_1 = \begin{bmatrix} 1 & -1 \end{bmatrix}^T$.

CASE 2:
$$\lambda_2 = 3$$
. With $\mathbf{v} = \begin{bmatrix} x & y \end{bmatrix}^T$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is $\begin{bmatrix} 2 & 7 \\ -2 & -7 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$.

Again, we have only a single equation, 2x + 7y = 0, and any nontrivial solution of this equation will suffice. The choice y = -2 yields x = 7, so (to within a constant multiple) the only eigenvector associated with $\lambda_2 = 3$ is $\mathbf{v}_2 = \begin{bmatrix} 7 & -2 \end{bmatrix}^T$.

Finally, note that it is not enough to say simply that the given matrix **A** has eigenvalues -2 and 3 and has eigenvectors $\begin{bmatrix} 1 & -1 \end{bmatrix}^T$ and $\begin{bmatrix} 7 & -2 \end{bmatrix}^T$. To give complete information, we must say which eigenvector is associated with each eigenvalue.

If the elements of the matrix **A** are all real, then so are the coefficients of its characteristic equation $|\mathbf{A} - \lambda \mathbf{I}| = 0$. In this event, it follows that the complex eigenvalues (if any) of **A** occur only in conjugate pairs. The next example illustrates the possibility of imaginary or complex eigenvalues and eigenvectors.

Example 3 The characteristic equation of the matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 8 \\ -2 & 0 \end{bmatrix}$$

is

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{bmatrix} -\lambda & 8\\ -2 & -\lambda \end{bmatrix} = \lambda^2 + 16 = 0.$$

Hence the matrix **A** has the complex conjugate eigenvalues $\lambda = \pm 4i$.

CASE 1:
$$\lambda_1 = -4i$$
. With $\mathbf{v} = \begin{bmatrix} x & y \end{bmatrix}^T$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is
$$\begin{bmatrix} 4i & 8\\ -2 & 4i \end{bmatrix} \begin{bmatrix} x\\ y \end{bmatrix} = \begin{bmatrix} 0\\ 0 \end{bmatrix}.$$

The first equation here, 4ix + 8y = 0, is -2i times the second one and, obviously, is satisfied by x = 2i, y = 1. Thus $\mathbf{v}_1 = \begin{bmatrix} 2i & 1 \end{bmatrix}^T$ is a complex eigenvector associated with the complex eigenvalue $\lambda_1 = -4i$. **CASE 2:** $\lambda_2 = +4i$. In this case, you may verify similarly that the conjugate $\mathbf{v}_2 = \begin{bmatrix} -2i & 1 \end{bmatrix}^T$ of \mathbf{v}_1 is an eigenvector associated with the conjugate λ_2 of the eigenvalue λ_1 . Thus the complex conjugate eigenvalues $\pm 4i$ of the matrix **A** correspond to the complex conjugate eigenvectors $\begin{bmatrix} \pm 2i & 1 \end{bmatrix}^T$ (taking either both upper signs or both lower signs).

Example 4

The 2×2 identity matrix **I** has characteristic equation

$$\begin{vmatrix} 1-\lambda & 0\\ 0 & 1-\lambda \end{vmatrix} = (1-\lambda)^2 = 0,$$

so **I** has the single eigenvalue $\lambda = 1$. The equation $(\mathbf{I} - 1\lambda)\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

so *every* nonzero vector $\mathbf{v} = \begin{bmatrix} x & y \end{bmatrix}^T$ is an eigenvector of **I**. In particular, the single eigenvalue $\lambda = 1$ corresponds to the *two* linearly independent eigenvectors $\mathbf{v}_1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$.

Example 5

The characteristic equation of the matrix

$$= \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$$

is $(2 - \lambda)^2 = 0$, so **A** has the single eigenvalue $\lambda = 2$. The equation $(\mathbf{A} - 2\mathbf{I})\mathbf{v} = \mathbf{0}$ is

А

$$\begin{bmatrix} 0 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Thus x is arbitrary, but y = 0, so the eigenvalue $\lambda = 2$ corresponds (to within a constant multiple) to the single eigenvector $\mathbf{v} = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$.

Examples 2–5 illustrate the four possibilities for a 2×2 matrix **A**. It can have either

- two distinct real eigenvalues, each corresponding to a single eigenvector;
- a single real eigenvalue corresponding to a single eigenvector;
- a single real eigenvalue corresponding to two linearly independent eigenvectors; or
- two complex conjugate eigenvalues corresponding to complex conjugate eigenvectors.

The characteristic equation of a 3×3 matrix is, by Eq. (5), of the form

$$-\lambda^3 + c_2 \lambda^2 + c_1 \lambda + c_0 = 0. \tag{7}$$

Now the value of the continuous function $-\lambda^3 + \cdots + c_0$ on the left-hand side here approaches $+\infty$ as $\lambda \to -\infty$ and $-\infty$ as $\lambda \to +\infty$. Hence every such cubic equation has at least one real solution, so every 3×3 matrix has (in contrast with 2×2 matrices) at least one real eigenvalue. A 3×3 matrix may have one, two, or three distinct eigenvalues, and a single eigenvalue of a 3×3 matrix may correspond to one, two, or three linearly independent eigenvectors. The remaining two examples of this section illustrate some of the possibilities. The next example shows also that, whereas the zero vector **0** cannot be an eigenvector of a matrix, there is nothing to prevent $\lambda = 0$ from being an eigenvalue. **Example 6** Find the eigenvalues and associated eigenvectors of the matrix

$$\mathbf{A} = \begin{bmatrix} 3 & 0 & 0 \\ -4 & 6 & 2 \\ 16 & -15 & -5 \end{bmatrix}.$$

Solution The matrix $\mathbf{A} - \lambda \mathbf{I}$ is

$$\mathbf{A} - \lambda \mathbf{I} = \begin{bmatrix} 3 - \lambda & 0 & 0 \\ -4 & 6 - \lambda & 2 \\ 16 & -15 & -5 - \lambda \end{bmatrix}.$$
 (8)

Upon expansion of the determinant along its first row, we find that

$$|\mathbf{A} - \lambda \mathbf{I}| = (3 - \lambda)[(6 - \lambda)(-5 - \lambda) + 30]$$
$$= (3 - \lambda)(\lambda^2 - \lambda) = \lambda(\lambda - 1)(3 - \lambda)$$

Hence the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}| = 0$ yields the three eigenvalues $\lambda_1 = 0$, $\lambda_2 = 1$, and $\lambda_3 = 3$. To find the associated eigenvectors, we must solve the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ separately for each of these three eigenvalues.

CASE 1: $\lambda_1 = 0$. We write $\mathbf{v} = \begin{bmatrix} x & y & z \end{bmatrix}^T$ and substitute $\lambda = 0$ in the coefficient matrix in (8) to obtain the system

3	0	0	$\begin{bmatrix} x \end{bmatrix}$		0	
-4	6	2	y	=	0	.
16	-15	-5	_ <i>z</i>		0	

From the first of the three equations here, 3x = 0, we see that x = 0. Then each of the remaining two equations is a multiple of the equation 3y + z = 0. The choice y = 1 yields z = -3. Thus the eigenvector $\mathbf{v}_1 = \begin{bmatrix} 0 & 1 & -3 \end{bmatrix}^T$ is associated with $\lambda_1 = 0$.

CASE 2: $\lambda_2 = 1$. Substitution of $\lambda = 1$ in the coefficient matrix in (8) yields the system

2	0	0]	$\begin{bmatrix} x \end{bmatrix}$		[0]
-4	5	2	y	=	0
16	-15	-6			0

for $\mathbf{v} = \begin{bmatrix} x & y & z \end{bmatrix}^T$. The first equation 2x = 0 implies that x = 0. Then the third equation is a multiple of the second equation, 5y + 2z = 0. The choice of y = 2 yields z = -5, so the eigenvector $\mathbf{v}_2 = \begin{bmatrix} 0 & 2 & -5 \end{bmatrix}^T$ is associated with $\lambda_2 = 1$.

CASE 3: $\lambda_3 = 3$. Substitution of $\lambda = 3$ in the coefficient matrix in (8) yields the system

0	0	0]	$\begin{bmatrix} x \end{bmatrix}$]	0]	
-4	3	2	y y	=	0	
16	-15	-8	_ <i>z</i> _		0	

In this case, the first equation yields no information, but the result of adding 4 times the second equation to the third equation is -3y = 0, so y = 0. Consequently, the second and third equations are both multiples of the equation 2x - z = 0. The choice x = 1 yields z = 2, so the eigenvector $\mathbf{v}_3 = \begin{bmatrix} 1 & 0 & 2 \end{bmatrix}^T$ is associated with $\lambda_3 = 3$.

In summary, we have found the eigenvectors $\mathbf{v}_1 = \begin{bmatrix} 0 & 1 & -3 \end{bmatrix}^T$, $\mathbf{v}_2 = \begin{bmatrix} 0 & 2 & -5 \end{bmatrix}^T$, and $\mathbf{v}_3 = \begin{bmatrix} 1 & 0 & 2 \end{bmatrix}^T$ associated with the distinct eigenvalues $\lambda_1 = 0$, $\lambda_2 = 1$, and $\lambda_3 = 3$, respectively.

Remark Substitution of $\lambda = 0$ in the characteristic equation $|\mathbf{A} - \lambda \mathbf{I}| = 0$ yields $|\mathbf{A}| = 0$. Therefore, $\lambda = 0$ is an eigenvalue of the matrix \mathbf{A} if and only if \mathbf{A} is singular: $|\mathbf{A}| = 0$.

Eigenspaces

Let λ be a fixed eigenvalue of the $n \times n$ matrix **A**. Then the set of all eigenvectors associated with **A** is the set of all nonzero solution vectors of the system

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}.\tag{9}$$

The solution space of this system is called the **eigenspace** of **A** associated with the eigenvalue λ . This subspace of **R**^{*n*} consists of all eigenvectors associated with λ together with the zero vector. In Example 6 we found (to within a constant multiple) only a single eigenvector associated with each eigenvalue λ ; in this case, the eigenspace of λ is 1-dimensional. In the case of an eigenspace of higher dimension, we generally want to find a basis for the solution space of Eq. (9).

xample 7	Find bases for the eigenspaces of the matrix				
	$\mathbf{A} =$	4 2 2	$-2 \\ 0 \\ -2$	1 1 3	

Solution Here we have

$$\mathbf{A} - \lambda \mathbf{I} = \begin{bmatrix} 4 - \lambda & -2 & 1\\ 2 & -\lambda & 1\\ 2 & -2 & 3 - \lambda \end{bmatrix}.$$
 (10)

We expand along the first row to obtain

$$|\mathbf{A} - \lambda \mathbf{I}| = (4 - \lambda)(\lambda^2 - 3\lambda + 2) - (-2)(4 - 2\lambda) + (1)(-4 + 2\lambda)$$
$$= -\lambda^3 + 7\lambda^2 - 16\lambda + 12.$$

Thus, to find the eigenvalues we need to solve the cubic equation

$$\lambda^3 - 7\lambda^2 + 16\lambda - 12 = 0. \tag{11}$$

We look (hopefully) for integer solutions. The factor theorem of algebra implies that if the polynomial equation

$$\lambda^n + c_{n-1}\lambda^{n-1} + \dots + c_1\lambda + c_0 = 0$$

with integer coefficients and leading coefficient 1 has an integer solution, then that integer is a divisor of the constant c_0 . In the case of the cubic equation in (11), the possibilities for such a solution are ± 1 , ± 2 , ± 3 , ± 4 , ± 6 , and ± 12 . We substitute these numbers successively in (11) and thereby find that +1 and -1 are not solutions but that $\lambda = +2$ is a solution. Hence $\lambda - 2$ is a factor of the cubic polynomial in (11). Next, the long division

$$\lambda - 2 \boxed{\begin{array}{c} \lambda^2 - 5\lambda + 6\\ \lambda^3 - 7\lambda^2 + 16\lambda - 12\\ \hline \lambda^3 - 2\lambda^2\\ \hline -5\lambda^2 + 16\lambda - 12\\ \hline -5\lambda^2 + 10\lambda\\ \hline 6\lambda - 12\\ \hline 6\lambda - 12\\ \hline 0 \end{array}}$$

shows that

$$\lambda^3 - 7\lambda^2 + 16\lambda - 12 = (\lambda - 2)(\lambda^2 - 5\lambda + 6)$$
$$= (\lambda - 2)(\lambda - 2)(\lambda - 3)$$
$$= (\lambda - 2)^2(\lambda - 3).$$

Thus we see finally that the given matrix **A** has the repeated eigenvalue $\lambda = 2$ and the eigenvalue $\lambda = 3$.
CASE 1: $\lambda = 2$. The system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

2	-2	1]	$\int x$		0	
2	-2	1	y y	=	0	,
2	-2	1			0	

which reduces to the single equation 2x - 2y + z = 0. This equation obviously has a 2dimensional solution space. With y = 1 and z = 0, we get x = 1 and, hence, obtain the basis eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 & 0 \end{bmatrix}^T$. With y = 0 and z = 2, we get x = -1 and, hence, the basis eigenvector $\mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 2 \end{bmatrix}^T$. The 2-dimensional eigenspace of **A** associated with the repeated eigenvalue $\lambda = 2$ has basis $\{\mathbf{v}_1, \mathbf{v}_2\}$.

CASE 2: $\lambda = 3$. The system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

1	-2	1]	$\int x$]	0]	
2	-3	1	y y	=	0	
2	-2	0			0	

The last equation here implies that x = y, and then each of the first two equations yields x = y = z. It follows that the eigenspace of **A** associated with $\lambda = 3$ is 1-dimensional and has $\mathbf{v}_3 = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^T$ as a basis eigenvector.

Remark The typical higher-degree polynomial is not so easy to factor as the one in Example 7. Hence a numerical technique such as Newton's method is often needed to solve the characteristic equation. Moreover, for an $n \times n$ matrix with n greater than about 4, the amount of labor required to find the characteristic equation by expanding the determinant $|\mathbf{A} - \lambda \mathbf{I}|$ is generally prohibitive; because of the presence of the variable λ , row and column elimination methods do not work as they do with numerical determinants. Consequently, specialized techniques, beyond the scope of the present discussion, are often required to find the eigenvalues and eigenvectors of the large matrices that occur in many applications. Problems 40 and 41 at the end of this section outline a numerical technique that sometimes is useful with matrices of moderate size.

6.1 **Problems**

In Problems 1 through 26, find the (real) eigenvalues and associated eigenvectors of the given matrix **A**. Find a basis for each eigenspace of dimension 2 or larger.

1.	$\left[\begin{array}{rr} 4 & -2 \\ 1 & 1 \end{array}\right]$	2. $\begin{bmatrix} 5 & -3 \\ 3 & -3 \end{bmatrix}$	-6 -4	3. $\begin{bmatrix} 8\\3 \end{bmatrix}$	$\begin{pmatrix} -6 \\ -1 \end{bmatrix}$
4.	$\left[\begin{array}{rrr} 4 & -3 \\ 2 & -1 \end{array}\right]$	5. $\begin{bmatrix} 10 \\ 6 \end{bmatrix}$	$\begin{bmatrix} -9 \\ -5 \end{bmatrix}$	6. $\begin{bmatrix} 6 \\ 3 \end{bmatrix}$	$\begin{pmatrix} -4 \\ -1 \end{bmatrix}$
7.	$\left[\begin{array}{rrr} 10 & -8 \\ 6 & -4 \end{array}\right]$	8. $\begin{bmatrix} 7\\12 \end{bmatrix}$	$\begin{pmatrix} -6 \\ -10 \end{bmatrix}$	9. $\begin{bmatrix} 8 \\ 2 \end{bmatrix}$	$\begin{pmatrix} -10 \\ -1 \end{bmatrix}$
10.	$\begin{bmatrix} 9 & -10 \\ 2 & 0 \end{bmatrix}$	11. $\begin{bmatrix} 19\\21 \end{bmatrix}$	$\begin{bmatrix} -10 \\ -10 \end{bmatrix}$	12. $\begin{bmatrix} 13 \\ 6 \end{bmatrix}$	$\begin{bmatrix} -15\\ -6 \end{bmatrix}$
13.	$\begin{bmatrix} 2 & 0 \\ 2 & -2 & -2 \\ -2 & 6 \end{bmatrix}$	$\begin{bmatrix} 0 \\ -1 \\ 3 \end{bmatrix} $ 14.	$\begin{bmatrix} 5\\4\\-2 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 \\ -4 & -2 \\ 12 & 6 \end{bmatrix}$	
15.	$\begin{bmatrix} 2 & -2 \\ 2 & -2 \\ -2 & 2 \end{bmatrix} -2$	$\begin{bmatrix} 0\\-1\\3 \end{bmatrix}$ 16.	$\begin{bmatrix} 1\\ -2\\ -6 \end{bmatrix}$	$\begin{bmatrix} 0 & -1 \\ 3 & -1 \\ 6 & 0 \end{bmatrix}$	
17.	$\begin{bmatrix} 3 & 5 & -2 \\ 0 & 2 & 0 \\ 0 & 2 & 1 \end{bmatrix}$] 18.	$\begin{bmatrix} 1\\ -6\\ 12 \end{bmatrix}$	$ \begin{array}{ccc} 0 & 0 \\ 8 & 2 \\ -15 & -3 \end{array} $	

$19. \begin{bmatrix} 3 & 6 & -2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$	20. $\begin{bmatrix} 1 & 0 & 0 \\ -4 & 7 & 2 \\ 10 & -15 & -4 \end{bmatrix}$
21. $\begin{bmatrix} 4 & -3 & 1 \\ 2 & -1 & 1 \\ 0 & 0 & 2 \end{bmatrix}$	22. $\begin{bmatrix} 5 & -6 & 3 \\ 6 & -7 & 3 \\ 6 & -6 & 2 \end{bmatrix}$
$23. \begin{bmatrix} 1 & 2 & 2 & 2 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & 3 & 2 \\ 0 & 0 & 0 & 4 \end{bmatrix}$	$24. \begin{bmatrix} 1 & 0 & 4 & 0 \\ 0 & 1 & 4 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$
$25. \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$	$26. \begin{bmatrix} 4 & 0 & 0 & -3 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 6 & 0 & 0 & -5 \end{bmatrix}$

Find the complex conjugate eigenvalues and corresponding eigenvectors of the matrices given in Problems 27 through 32.

27.
$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$
 28. $\mathbf{A} = \begin{bmatrix} 0 & -6 \\ 6 & 0 \end{bmatrix}$
29. $\mathbf{A} = \begin{bmatrix} 0 & -3 \\ 12 & 0 \end{bmatrix}$ **30.** $\mathbf{A} = \begin{bmatrix} 0 & -12 \\ 12 & 0 \end{bmatrix}$

31.
$$\mathbf{A} = \begin{bmatrix} 0 & 24 \\ -6 & 0 \end{bmatrix}$$
 32. $\mathbf{A} = \begin{bmatrix} 0 & -4 \\ 36 & 0 \end{bmatrix}$

- **33.** Suppose that λ is an eigenvalue of the matrix **A** with associated eigenvector **v** and that *n* is a positive integer. Show that λ^n is an eigenvalue of **A**^{*n*} with associated eigenvector **v**.
- **34.** Show that λ is an eigenvalue of the invertible matrix **A** if and only if λ^{-1} is an eigenvalue of \mathbf{A}^{-1} . Are the associated eigenvectors the same?
- **35.** (a) Suppose that A is a square matrix. Use the characteristic equation to show that A and A^T have the same eigenvalues.
 - (b) Give an example of a 2×2 matrix **A** such that **A** and \mathbf{A}^T do not have the same eigenvectors.
- **36.** Show that the eigenvalues of a triangular $n \times n$ matrix are its diagonal elements.
- **37.** Suppose that the characteristic equation $|\mathbf{A} \lambda \mathbf{I}| = 0$ is written as a polynomial equation [Eq. (5)]. Show that the constant term is $c_0 = \det \mathbf{A}$. Suggestion: Substitute an appropriate value for λ .

Problems 38 through 42 introduce the trace of a square matrix and explore its connections with the determinant and the characteristic polynomial of the matrix.

38. If $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ is an $n \times n$ matrix, then the **trace** Tr A of A is defined to be

$$\operatorname{Tr} \mathbf{A} = a_{11} + a_{22} + \dots + a_{nn},$$

the sum of the diagonal elements of **A**. It can be proved that the coefficient of λ^{n-1} in Eq. (5) is $c_{n-1} = (-1)^{n-1}(\text{Tr }\mathbf{A})$. Show explicitly that this is true in the case of a 2×2 matrix.

39. Suppose that the $n \times n$ matrix **A** has *n* (real) eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$. Assuming the general result stated in Problem 38, prove that

$$\lambda_1 + \lambda_2 + \dots + \lambda_n = \operatorname{Tr} \mathbf{A}$$
$$= a_{11} + a_{22} + \dots + a_{nn}$$

40. According to the results stated in Problems 37 and 38, the characteristic polynomial

$$p(\lambda) = |\mathbf{A} - \lambda \mathbf{I}|$$

6.2 **Diagonalization of Matrices**

Given an $n \times n$ matrix **A**, we may ask *how many* linearly independent eigenvectors the matrix **A** has. In Section 6.1, we saw several examples (with n = 2 and n = 3) in which the $n \times n$ matrix **A** has *n* linearly independent eigenvectors—the largest possible number. By contrast, in Example 5 of Section 6.1, we saw that the 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$$

has the single eigenvalue $\lambda = 2$ corresponding to the single eigenvector $\mathbf{v} = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$. Something very nice happens when the $n \times n$ matrix **A** does have *n* linearly independent eigenvectors. Suppose that the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (not necessar-

of a 3×3 matrix **A** is given by

$$p(\lambda) = -\lambda^3 + (\operatorname{Tr} \mathbf{A})\lambda^2 + c_1\lambda + (\det \mathbf{A}).$$

The remaining coefficient c_1 can be found by substituting $\lambda = 1$ and then calculating the two determinants $|\mathbf{A}|$ and $p(1) = |\mathbf{A} - \mathbf{I}|$. Use this method to find the characteristic equation, eigenvalues, and associated eigenvectors of the matrix

$$\mathbf{A} = \begin{bmatrix} 32 & -67 & 47 \\ 7 & -14 & 13 \\ -7 & 15 & -6 \end{bmatrix}$$

41. According to the results stated in Problems 37 and 38, the characteristic polynomial

$$p(\lambda) = |\mathbf{A} - \lambda \mathbf{I}|$$

of a 4×4 matrix **A** is given by

$$p(\lambda) = \lambda^4 - (\operatorname{Tr} \mathbf{A})\lambda^3 + c_2\lambda^2 + c_1\lambda + (\det \mathbf{A}).$$

The remaining coefficients c_1 and c_2 can be found by substituting $\lambda = \pm 1$ and calculating the three determinants $|\mathbf{A}|$, $p(1) = |\mathbf{A} - \mathbf{I}|$, and $p(-1) = |\mathbf{A} + \mathbf{I}|$. Use this method to find the characteristic equation, eigenvalues, and associated eigenvectors of the matrix

$$\mathbf{A} = \begin{bmatrix} 22 & -9 & -8 & -8\\ 10 & -7 & -14 & 2\\ 10 & 0 & 8 & -10\\ 29 & -9 & -3 & -15 \end{bmatrix}.$$

- **42.** Combine ideas from Problems 37–40 to show explicitly that
 - (a) If A is a 2×2 matrix then $det(\mathbf{A} \lambda \mathbf{I}) = (-\lambda)^2 + Tr(\mathbf{A})(-\lambda) + det(\mathbf{A})$.
 - (b) If **A** is a 3×3 matrix then det $(\mathbf{A} \lambda \mathbf{I}) = (-\lambda)^3 + \text{Tr}(\mathbf{A})(-\lambda)^2 + c_1(-\lambda) + \text{det}(\mathbf{A})$, where c_1 is the sum of the minors of the diagonal elements of **A**. (Recall from Section 3.6 the distinction between minors and cofactors.)

348 CHAPTER 6 EIGENVALUES AND EIGENVECTORS

ily distinct) of **A** correspond to the *n* linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$, respectively. Let

 $\mathbf{P} = \begin{bmatrix} | & | & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & | \end{bmatrix}$ (1)

be the $n \times n$ matrix having these eigenvectors as its *column* vectors. Then

г

$$\mathbf{AP} = \mathbf{A} \begin{bmatrix} | & | & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | \\ \mathbf{Av}_1 & \mathbf{Av}_2 & \cdots & \mathbf{Av}_n \\ | & | & | \end{bmatrix}$$

and hence

$$\mathbf{AP} = \begin{bmatrix} | & | & | \\ \lambda_1 \mathbf{v}_1 & \lambda_2 \mathbf{v}_2 & \cdots & \lambda_n \mathbf{v}_n \\ | & | & | & | \end{bmatrix},$$
(2)

-

because $\mathbf{A}\mathbf{v}_j = \lambda_j \mathbf{v}_j$ for each j = 1, 2, ..., n. Thus the product matrix **AP** has column vectors $\lambda_1 \mathbf{v}_1, \lambda_2 \mathbf{v}_2, ..., \lambda_n \mathbf{v}_n$.

Now consider the diagonal matrix

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0\\ 0 & \lambda_2 & \cdots & 0\\ \vdots & \vdots & \ddots & \vdots\\ 0 & 0 & \cdots & \lambda_n \end{bmatrix},$$
(3)

whose diagonal elements are the eigenvalues corresponding (in the same order) to the eigenvectors forming the columns of **P**. Then

$$\mathbf{PD} = \begin{bmatrix} | & | & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & | \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}$$
$$= \begin{bmatrix} | & | & | \\ \lambda_1 \mathbf{v}_1 & \lambda_2 \mathbf{v}_2 & \cdots & \lambda_n \mathbf{v}_n \\ | & | & | & | \end{bmatrix}, \qquad (4)$$

because the product of the *i*th row of **P** and the *j*th column of **D** is simply the product of λ_j and the *i*th component of \mathbf{v}_j .

Finally, upon comparing the results in (2) and (4), we see that

$$\mathbf{AP} = \mathbf{PD}.\tag{5}$$

But the matrix **P** is invertible, because its *n* column vectors are linearly independent. So we may multiply on the right by \mathbf{P}^{-1} to obtain

$$\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}.$$
 (6)

Equation (6) expresses the $n \times n$ matrix **A** having *n* linearly independent eigenvectors in terms of the *eigenvector matrix* **P** and the diagonal *eigenvalue matrix* **D**. It can be rewritten as $\mathbf{D} = \mathbf{P}^{-1}\mathbf{AP}$, but the form in (6) is the one that should be memorized.

In Example 1 of Section 6.1 we saw that the matrix

$$\mathbf{A} = \begin{bmatrix} 5 & -6 \\ 2 & -2 \end{bmatrix}$$

has eigenvalues $\lambda_1 = 2$ and $\lambda_2 = 1$ corresponding to the linearly independent eigenvectors $\mathbf{v}_1 = \begin{bmatrix} 2 & 1 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} 3 & 2 \end{bmatrix}^T$, respectively. Then

$$\mathbf{P} = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}, \quad \mathbf{D} = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}, \quad \text{and} \quad \mathbf{P}^{-1} = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}.$$

So

Example 1

$$\mathbf{PDP}^{-1} = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}$$
$$= \begin{bmatrix} 4 & 3 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & -6 \\ 2 & -2 \end{bmatrix} = \mathbf{A}$$

in accord with Eq. (6).

Similarity and Diagonalization

The following definition embodies the precise relationship in (6) between the original matrix \mathbf{A} and the diagonal matrix \mathbf{D} .

DEFINITION Similar Matrices

The $n \times n$ matrices **A** and **B** are called **similar** provided that there exists an invertible matrix **P** such that

$$\mathbf{B} = \mathbf{P}^{-1} \mathbf{A} \mathbf{P}.$$
 (7)

Note that this relationship between **A** and **B** is symmetric, for if $\mathbf{B} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$, then $\mathbf{A} = \mathbf{Q}^{-1}\mathbf{B}\mathbf{Q}$ for some invertible matrix \mathbf{Q} —just take $\mathbf{Q} = \mathbf{P}^{-1}$.

An $n \times n$ matrix **A** is called **diagonalizable** if it is similar to a diagonal matrix **D**; that is, there exist a diagonal matrix **D** and an invertible matrix **P** such that $\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$, and so

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}.$$
 (8)

The process of finding the diagonalizing matrix \mathbf{P} and the diagonal matrix \mathbf{D} in (8) is called **diagonalization** of the matrix \mathbf{A} . In Example 1 we showed that the matrices

$$\mathbf{A} = \begin{bmatrix} 5 & -6 \\ 2 & -2 \end{bmatrix} \quad \text{and} \quad \mathbf{D} = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$$

are similar, and hence that the 2×2 matrix **A** is diagonalizable.

Now we ask under what conditions a given square matrix is diagonalizable. In deriving Eq. (6), we showed that *if* the $n \times n$ matrix **A** has *n* linearly independent eigenvectors, *then* **A** is diagonalizable. The converse of this statement is also true.

THEOREM 1 Criterion for Diagonalizability

The $n \times n$ matrix **A** is diagonalizable if and only if it has *n* linearly independent eigenvectors.

Proof: It remains only to show that, if the $n \times n$ matrix **A** is diagonalizable, then it has *n* linearly independent eigenvectors. Suppose that **A** is similar to the diagonal matrix **D** with diagonal elements d_1, d_2, \ldots, d_n , and let

$$\mathbf{P} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix}$$

be an invertible matrix such that $\mathbf{D} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$. Then

$$\mathbf{AP} = \mathbf{A} \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix} = \begin{bmatrix} \mathbf{Av}_1 & \mathbf{Av}_2 & \cdots & \mathbf{Av}_n \end{bmatrix}$$

and

$$\mathbf{PD} = \begin{bmatrix} d_1 \mathbf{v}_1 & d_2 \mathbf{v}_2 & \cdots & d_n \mathbf{v}_n \end{bmatrix},$$

by essentially the same computation as in Eq. (4). But $\mathbf{AP} = \mathbf{PD}$ because $\mathbf{D} = \mathbf{P}^{-1}\mathbf{AP}$, so it follows that

$$\mathbf{A}\mathbf{v}_j = d_j\mathbf{v}_j$$

for each j = 1, 2, ..., n. Thus the vectors $\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_n$ are eigenvectors of **A** associated with the eigenvalues $d_1, d_2, ..., d_n$, respectively. And it follows from Theorem 2 in Section 3.6 and Theorem 2 in Section 4.3 that these *n* eigenvectors of the matrix **A** are linearly independent, because they are the column vectors of the invertible matrix **P**.

Remark It is important to remember not only the fact that an $n \times n$ matrix **A** having *n* linearly independent eigenvectors is diagonalizable, but also the specific diagonalization $\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$ in Eq. (6), where the matrix **P** has the *n* eigenvectors as its columns, and the corresponding eigenvalues are the diagonal elements of the diagonal matrix **D**.

Example 2 In Example 5 of Section 6.1 we saw that the matrix

$$\mathbf{A} = \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$$

has only one eigenvalue, $\lambda = 2$, and that (to within a constant multiple) only the single eigenvector $\mathbf{v} = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ is associated with this eigenvalue. Thus the 2 × 2 matrix **A** does *not* have n = 2 linearly independent eigenvectors. Hence Theorem 1 implies that **A** is not diagonalizable.

Example 3 In Example 6 of Section 6.1 we saw that the matrix

$$\mathbf{A} = \begin{bmatrix} 3 & 0 & 0 \\ -4 & 6 & 2 \\ 16 & -15 & -5 \end{bmatrix}$$

has the following eigenvalues and associated eigenvectors:

$$\lambda_1 = 3: \quad \mathbf{v}_1 = \begin{bmatrix} 1 & 0 & 2 \end{bmatrix}^T$$
$$\lambda_2 = 1: \quad \mathbf{v}_2 = \begin{bmatrix} 0 & 2 & -5 \end{bmatrix}^T$$
$$\lambda_3 = 0: \quad \mathbf{v}_3 = \begin{bmatrix} 0 & 1 & -3 \end{bmatrix}^T.$$

It is obvious (why?) that the three eigenvectors v_1 , v_2 , v_3 are linearly independent, so Theorem 1 implies that the 3×3 matrix **A** is diagonalizable. In particular, the inverse of the eigenvector matrix

$$\mathbf{P} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 2 & -5 & -3 \end{bmatrix}$$
$$\mathbf{P}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 3 & 1 \\ 4 & -5 & -2 \end{bmatrix},$$

is

and the diagonal eigenvalue matrix is

$$\mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Therefore, Eq. (6) in the form $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}$ yields the diagonalization

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 3 & 1 \\ 4 & -5 & -2 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0 \\ -4 & 6 & 2 \\ 16 & -15 & -5 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 2 & -5 & -3 \end{bmatrix}$$
$$= \begin{bmatrix} 3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

of the matrix A.

The following theorem tells us that any set of eigenvectors associated with *distinct* eigenvalues (as in Example 3) is automatically linearly independent.

THEOREM 2 Eigenvectors Associated with Distinct Eigenvalues

Suppose that the eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are associated with the distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$ of the matrix **A**. Then these k eigenvectors are linearly independent.

Proof: Our proof will be by induction on k. The theorem is certainly true in the case k = 1, because any single (nonzero) eigenvector constitutes a linearly independent set. Now assume inductively that any set of k - 1 eigenvectors associated with distinct eigenvalues is linearly independent. Supposing that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0},\tag{9}$$

we need to show that $c_1 = c_2 = \cdots = c_k = 0$. To do this, we will multiply in Eq. (9) by the matrix $\mathbf{A} - \lambda_1 \mathbf{I}$. First note that

$$(\mathbf{A} - \lambda_1 \mathbf{I})\mathbf{v}_j = \mathbf{A}\mathbf{v}_j - \lambda_1 \mathbf{v}_j = \begin{cases} \mathbf{0} & \text{if } j = 1, \\ (\lambda_j - \lambda_1)\mathbf{v}_j & \text{if } j > 1, \end{cases}$$

because $\mathbf{A}\mathbf{v}_j = \lambda_j \mathbf{v}_j$ for each *j*. Therefore, the result of multiplying Eq. (9) by $\mathbf{A} - \lambda_1 \mathbf{I}$ is

$$c_2(\lambda_2 - \lambda_1)\mathbf{v}_2 + \dots + c_k(\lambda_k - \lambda_1)\mathbf{v}_k = \mathbf{0}.$$
 (10)

But the k - 1 eigenvectors $\mathbf{v}_2, \mathbf{v}_3, \dots, \mathbf{v}_k$ are linearly independent by the inductive assumption, so each of the scalar coefficients $c_j(\lambda_j - \lambda_1)$ here must be zero. Now our hypothesis that the eigenvalues of **A** are distinct implies that $\lambda_j - \lambda_1 \neq 0$ for each j > 1. It therefore follows from Eq. (10) that $c_2 = c_3 = \dots = c_k = 0$. But then Eq. (9) reduces to $c_1\mathbf{v}_1 = \mathbf{0}$, so it now follows (because $\mathbf{v}_1 \neq \mathbf{0}$) that $c_1 = 0$ as well. Thus we have shown that all the coefficients in Eq. (9) must vanish, and hence that the k eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent. Theorem 2 now follows by induction.

If the $n \times n$ matrix **A** has *n* distinct eigenvalues, then by Theorem 2 the *n* associated eigenvectors are linearly independent, so Theorem 1 implies that the matrix **A** is diagonalizable. Thus we have the sufficient condition for diagonalizability given in Theorem 3.

THEOREM 3 An n × n Matrix with n Distinct Eigenvalues

If the $n \times n$ matrix **A** has *n* distinct eigenvalues, then it is diagonalizable.

In general, however, an $n \times n$ matrix **A** can be expected to have fewer than n distinct eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_k$. If k < n, then we may *attempt* to diagonalize **A** by carrying out the following procedure.

Step 1. Find a basis S_i for the eigenspace associated with each eigenvalue λ_i .

Step 2. Form the union *S* of the bases $S_1, S_2, ..., S_k$. According to Theorem 4 in this section, the set *S* of eigenvectors of **A** is linearly independent.

Step 3. If *S* contains *n* eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$, then the matrix

$$\mathbf{P} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix}$$

diagonalizes A: that is, $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}$, where the diagonal elements of \mathbf{D} are the eigenvalues (repeated as necessary) corresponding to the *n* eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$.

If the set *S*—obtained by "merging" the bases for all the eigenspaces of A— contains fewer than *n* eigenvectors, then it can be proved that the matrix A is not diagonalizable.

Example 4 In Example 7 of Section 6.1, we saw that the matrix

	4	-2	1
$\mathbf{A} =$	2	0	1
	2	-2	3

has only two distinct eigenvalues, $\lambda_1 = 2$ and $\lambda_2 = 3$. We found that the eigenvalue $\lambda_1 = 2$ corresponds to a 2-dimensional eigenspace with basis vectors $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 & 0 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 2 \end{bmatrix}^T$, and that $\lambda_2 = 3$ corresponds to a 1-dimensional eigenspace with basis vector $\mathbf{v}_3 = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^T$. By Theorem 4 (or by explicit verification), these three eigenvectors are linearly independent, so Theorem 1 implies that the 3×3 matrix **A** is diagonalizable. The diagonalizing matrix

$$\mathbf{P} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 \end{bmatrix} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}$$

has inverse matrix

$$\mathbf{P}^{-1} = \begin{bmatrix} -2 & 3 & -1 \\ -1 & 1 & 0 \\ 2 & -2 & 1 \end{bmatrix},$$

so we obtain the diagonalization

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{bmatrix} -2 & 3 & -1 \\ -1 & 1 & 0 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 4 & -2 & 1 \\ 2 & 0 & 1 \\ 2 & -2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}$$
$$= \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix} = \mathbf{D}$$

of the matrix A.

THEOREM 4 Complete Independence of Eigenvectors

Let $\lambda_1, \lambda_2, \ldots, \lambda_k$ be the distinct eigenvalues of the $n \times n$ matrix **A**. For each $i = 1, 2, \ldots, k$, let S_i be a basis for the eigenspace associated with λ_i . Then the union *S* of the bases S_1, S_2, \ldots, S_k is a linearly independent set of eigenvectors of **A**.

Proof: To simplify the notation, we will illustrate the proof with the typical case k = 3, with A having three distinct eigenvalues λ_1 , λ_2 , and λ_3 . Let

$$S_1 = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p\},$$

$$S_2 = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_q\}, \text{ and }$$

$$S_3 = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_r\}$$

be bases for the eigenspaces associated with the eigenvalues λ_1 , λ_2 , and λ_3 . Assuming that a linear combination of the vectors in $S = S_1 \cup S_2 \cup S_3$ vanishes—

$$a_{1}\mathbf{u}_{1} + a_{2}\mathbf{u}_{2} + \dots + a_{p}\mathbf{u}_{p}$$

+ $b_{1}\mathbf{v}_{1} + b_{2}\mathbf{v}_{2} + \dots + b_{q}\mathbf{v}_{q}$
+ $c_{1}\mathbf{w}_{1} + c_{2}\mathbf{w}_{2} + \dots + c_{r}\mathbf{w}_{r} = \mathbf{0}$ (11)

-we need to show that the coefficients are all zero. If we write

$$\mathbf{u} = a_1 \mathbf{u}_1 + a_2 \mathbf{u}_2 + \dots + a_p \mathbf{u}_p,$$

$$\mathbf{v} = b_1 \mathbf{v}_1 + b_2 \mathbf{v}_2 + \dots + b_q \mathbf{v}_q, \text{ and}$$

$$\mathbf{w} = c_1 \mathbf{w}_1 + c_2 \mathbf{w}_2 + \dots + c_r \mathbf{w}_r,$$

then Eq. (11) takes the simple form

$$\mathbf{u} + \mathbf{v} + \mathbf{w} = \mathbf{0}.\tag{12}$$

But the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} either are zero vectors or are eigenvectors associated with the distinct eigenvalues λ_1 , λ_2 , and λ_3 . In the latter event, Theorem 2 would imply that \mathbf{u} , \mathbf{v} , and \mathbf{w} are linearly independent. Therefore, Eq. (12) implies that $\mathbf{u} = \mathbf{v} = \mathbf{w} = \mathbf{0}$. Finally, the fact that the $\{\mathbf{u}_i\}$ are linearly independent implies that $a_1 = a_2 = \cdots = a_p = 0$; the fact that the $\{\mathbf{v}_i\}$ are linearly independent implies that $b_1 = b_2 = \cdots = b_q = 0$; similarly, $c_1 = c_2 = \cdots = c_r = 0$. Thus we have shown that the coefficients in (11) all vanish, and hence that the vectors in $S = S_1 \cup S_2 \cup S_3$ are linearly independent.

6.2 **Problems**

In Problems 1 throu matrix A is diagond	ugh 28, determine whe alizable. If it is, find a	ther or not the given diagonalizing matrix	7. $\begin{bmatrix} 6 & -10 \\ 2 & -3 \end{bmatrix}$	8. $\begin{bmatrix} 11 & -15 \\ 6 & -8 \end{bmatrix}$	9. $\begin{bmatrix} -1 & 4 \\ -1 & 3 \end{bmatrix}$
P and a diagonal m	natrix D such that \mathbf{P}^{-1}	$\mathbf{AP}=\mathbf{D}.$	Го 1 Л	с _с 17	Г 11 0 Л
1. $\begin{bmatrix} 5 & -4 \\ 2 & -1 \end{bmatrix}$	2. $\begin{bmatrix} 6 & -6 \\ 4 & -4 \end{bmatrix}$	3. $\begin{bmatrix} 5 & -3 \\ 2 & 0 \end{bmatrix}$	10. $\begin{bmatrix} 5 & -1 \\ 1 & 1 \end{bmatrix}$	11. $\begin{bmatrix} 5 & 1 \\ -9 & -1 \end{bmatrix}$	12. $\begin{bmatrix} 11 & 9 \\ -16 & -13 \end{bmatrix}$
$4. \begin{bmatrix} 5 & -4 \\ 3 & -2 \end{bmatrix}$	5. $\begin{bmatrix} 9 & -8 \\ 6 & -5 \end{bmatrix}$	6. $\begin{bmatrix} 10 & -6 \\ 12 & -7 \end{bmatrix}$	13. $\begin{bmatrix} 1 & 3 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$	14. $\begin{bmatrix} 2\\ 2\\ 2\\ 2 \end{bmatrix}$	$ \begin{array}{ccc} -2 & 1 \\ -2 & 1 \\ -2 & 1 \end{array} $



- **29.** Prove: If the matrices **A** and **B** are similar and the matrices **B** and **C** are similar, then the matrices **A** and **C** are similar.
- **30.** Suppose that the matrices **A** and **B** are similar and that *n* is a positive integer. Prove that the matrices \mathbf{A}^n and \mathbf{B}^n are similar.
- **31.** Suppose that the invertible matrices **A** and **B** are similar. Prove that their inverses A^{-1} and B^{-1} are also similar.
- **32.** Show that if the $n \times n$ matrices **A** and **B** are similar, then they have the same characteristic equation and therefore have the same eigenvalues.
- **33.** Suppose that the $n \times n$ matrices **A** and **B** are similar and that each has *n* real eigenvalues. Show that det **A** = det **B**

and that $\operatorname{Tr} \mathbf{A} = \operatorname{Tr} \mathbf{B}$. See Problems 38 and 39 in Section 6.1.

34. Consider the 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

and let $\Delta = (a - d)^2 + 4bc$. Then show that

- (a) A is diagonalizable if $\Delta > 0$;
- (b) A is not diagonalizable if $\Delta < 0$;
- (c) If $\Delta = 0$, then A may be diagonalizable or it may not be.
- **35.** Let **A** be a 3×3 matrix with three distinct eigenvalues. Tell how to construct six different invertible matrices $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_6$ and six different diagonal matrices $\mathbf{D}_1, \mathbf{D}_2, \dots, \mathbf{D}_6$ such that $\mathbf{P}_i \mathbf{D}_i (\mathbf{P}_i)^{-1} = \mathbf{A}$ for each $i = 1, 2, \dots, 6$.
- **36.** Prove: If the diagonalizable matrices **A** and **B** have the same eigenvalues (with the same multiplicities), then **A** and **B** are similar.
- **37.** Given: The diagonalizable matrix **A**. Show that the eigenvalues of \mathbf{A}^2 are the squares of the eigenvalues of **A** but that **A** and \mathbf{A}^2 have the same eigenvectors.
- **38.** Suppose that the $n \times n$ matrix **A** has *n* linearly independent eigenvectors associated with a single eigenvalue λ . Show that **A** is a diagonal matrix.
- **39.** Let λ_i be an eigenvalue of the $n \times n$ matrix **A**, and assume that the characteristic equation of **A** has only real solutions. The **algebraic multiplicity** of λ_i is the largest positive integer p(i) such that $(\lambda \lambda_i)^{p(i)}$ is a factor of the characteristic polynomial $|\mathbf{A} \lambda \mathbf{I}|$. The **geometric multiplicity** of λ_i is the dimension q(i) of the eigenspace associated with λ_i . It can be shown that $p(i) \ge q(i)$ for every eigenvalue λ_i . Taking this as already established, prove that the given matrix **A** is diagonalizable if and only if the geometric multiplicity of each eigenvalue is equal to its algebraic multiplicity.

6.3 Applications Involving Powers of Matrices

In this section we discuss applications that depend on an ability to compute the matrix \mathbf{A}^k for large values of k, given the $n \times n$ matrix \mathbf{A} . If \mathbf{A} is diagonalizable, then \mathbf{A}^k can be found directly by a method that avoids the labor of calculating the powers $\mathbf{A}^2, \mathbf{A}^3, \mathbf{A}^4, \ldots$ by successive matrix multiplications.

Recall from Section 6.2 that, if the $n \times n$ matrix **A** has *n* linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ associated with the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (not necessarily distinct), then

$$\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1},\tag{1}$$

where

$$\mathbf{P} = \begin{bmatrix} | & | & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & | \end{bmatrix} \text{ and } \mathbf{D} = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix}.$$

Note that (1) yields

$$\mathbf{A}^2 = (\mathbf{P}\mathbf{D}\mathbf{P}^{-1})(\mathbf{P}\mathbf{D}\mathbf{P}^{-1}) = \mathbf{P}\mathbf{D}(\mathbf{P}^{-1}\mathbf{P})\mathbf{D}\mathbf{P}^{-1} = \mathbf{P}\mathbf{D}^2\mathbf{P}^{-1}$$

because $\mathbf{P}^{-1}\mathbf{P} = \mathbf{I}$. More generally, for each positive integer k,

$$\mathbf{A}^{k} = (\mathbf{P}\mathbf{D}\mathbf{P}^{-1})^{k}$$

= (\mathbf{P}\mathbf{D}\mathbf{P}^{-1})(\mathbf{P}\mathbf{D}\mathbf{P}^{-1})(\mathbf{P}\mathbf{D}\mathbf{P}^{-1})
= \mathbf{P}\mathbf{O}(\mathbf{P}^{-1}\mathbf{P})\mathbf{D}\mathbf{V}^{-1}\mathbf{P})\mathbf{D}\mathbf{P}^{-1};
$$\mathbf{A}^{k} = \mathbf{P}\mathbf{D}^{k}\mathbf{P}^{-1}.$$
 (2)

But the *k*th power \mathbf{D}^k of the diagonal matrix \mathbf{D} is easily computed:

$$\mathbf{D}^{k} = \begin{bmatrix} \lambda_{1}^{k} & 0 & \cdots & 0 \\ 0 & \lambda_{2}^{k} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_{n}^{k} \end{bmatrix}.$$
 (3)

Consequently the formula in (2) gives a quick and effective way to calculate any desired power of the diagonalizable matrix **A**, once its eigenvalues and associated eigenvectors—and hence the matrices **P** and **D**—have been found.

Example 1 Find \mathbf{A}^5 if $\mathbf{A} = \begin{bmatrix} 4 & -2 & 1 \\ 2 & 0 & 1 \\ 2 & -2 & 3 \end{bmatrix}.$ Solution In Example 7 of Section 6.1, we found that the 3 × 3 matrix \mathbf{A} has the eigenvalue $\lambda_1 = 3$ with associated eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^T$, and the repeated eigenvalue $\lambda_2 = 2$ with associated eigenvectors $\mathbf{v}_2 = \begin{bmatrix} 1 & 1 & 0 \end{bmatrix}^T$ and $\mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 2 \end{bmatrix}^T$. Therefore, $\mathbf{A} = \mathbf{PDP}^{-1}$ with $\mathbf{P} = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 2 \end{bmatrix} \text{ and } \mathbf{D} = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$ We first calculate $\mathbf{P}^{-1} = \begin{bmatrix} 2 & -2 & 1 \\ -2 & 3 & -1 \\ -1 & 1 & 0 \end{bmatrix} \text{ and } \mathbf{D}^5 = \begin{bmatrix} 243 & 0 & 0 \\ 0 & 322 & 0 \\ 0 & 0 & 32 \end{bmatrix}.$ Then the formula in (3) yields $\mathbf{A}^5 = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} 243 & 0 & 0 \\ 0 & 322 & 0 \\ 0 & 0 & 32 \end{bmatrix} \begin{bmatrix} 2 & -2 & 1 \\ -2 & 3 & -1 \\ -1 & 1 & 0 \end{bmatrix}$ $= \begin{bmatrix} 243 & 32 & -32 \\ 243 & 32 & -32 \\ 243 & 0 & 64 \end{bmatrix} \begin{bmatrix} 2 & -2 & 1 \\ -2 & 3 & -1 \\ -1 & 1 & 0 \end{bmatrix}$ $= \begin{bmatrix} 454 & -422 & 211 \\ 422 & -390 & 211 \\ 422 & -422 & 243 \end{bmatrix}.$

Transition Matrices

We want to apply this method of computing \mathbf{A}^k to the analysis of a certain type of physical system that can be described by means of the following kind of mathematical model. Suppose that the sequence

$$\mathbf{x}_0, \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k, \dots \tag{4}$$

of *n*-vectors is defined by its **initial vector** \mathbf{x}_0 and an $n \times n$ **transition matrix** \mathbf{A} in the following manner:

$$\mathbf{x}_{k+1} = \mathbf{A}\mathbf{x}_k \qquad \text{for each } k \ge 0. \tag{5}$$

We envision a physical system—such as a population with *n* specified subpopulations—that evolve through a sequence of successive *states* described by the vectors in (4). Then our goal is to calculate the *k*th *state vector* \mathbf{x}_k . But using (5) repeatedly, we find that

$$\mathbf{x}_1 = \mathbf{A}\mathbf{x}_0, \quad \mathbf{x}_2 = \mathbf{A}\mathbf{x}_1 = \mathbf{A}^2\mathbf{x}_0, \quad \mathbf{x}_3 = \mathbf{A}\mathbf{x}_2 = \mathbf{A}^3\mathbf{x}_0,$$

and in general that

$$\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0. \tag{6}$$

Thus our task amounts to calculating the *k*th power \mathbf{A}^k of the transition matrix \mathbf{A} .

Example 2 Urban/suburban population Consider a metropolitan area with a *constant* total population of 1 million individuals. This area consists of a city and its suburbs, and we want to analyze the changing urban and suburban populations. Let C_k denote the city population and S_k the suburban population after k years. Suppose that each year 15% of the people in the city move to the suburbs, whereas 10% of the people in the suburbs move to the city. Then it follows that

$$C_{k+1} = 0.85C_k + 0.10S_k$$

$$S_{k+1} = 0.15C_k + 0.90S_k$$
(7)

for each $k \ge 0$. (For instance, next year's city population C_{k+1} will equal 85% of this year's city population C_k plus 10% of this year's suburban population S_k .) Thus the metropolitan area's *population vector* $\mathbf{x}_k = \begin{bmatrix} C_k & S_k \end{bmatrix}^T$ satisfies

$$\mathbf{x}_{k+1} = \mathbf{A}\mathbf{x}_k$$
 and hence $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0$ (8)

(for each $k \ge 0$) with *transition matrix*

$$\mathbf{A} = \begin{bmatrix} 0.85 & 0.10\\ 0.15 & 0.90 \end{bmatrix}.$$

The characteristic equation of the transition matrix A is

$$\binom{17}{20} - \lambda \binom{9}{10} - \lambda - \binom{3}{20} \binom{1}{10} = 0;$$

$$(17 - 20\lambda)(9 - 10\lambda) - 3 = 0;$$

$$200\lambda^2 - 350\lambda + 150 = 0;$$

$$4\lambda^2 - 7\lambda + 3 = 0;$$

$$(\lambda - 1)(4\lambda - 3) = 0.$$

Thus the eigenvalues of **A** are $\lambda_1 = 1$ and $\lambda_2 = 0.75$. For $\lambda_1 = 1$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.15 & 0.10 \\ 0.15 & -0.10 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

6.3 APPLICATIONS INVOLVING POWERS OF MATRICES 357

so an associated eigenvector is $\mathbf{v}_1 = \begin{bmatrix} 2 & 3 \end{bmatrix}^T$. For $\lambda_2 = 0.75$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} 0.10 & 0.10 \\ 0.15 & 0.15 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

so an associated eigenvector is $\mathbf{v}_2 = \begin{bmatrix} -1 & 1 \end{bmatrix}^T$. It follows that $\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$, where

$$\mathbf{P} = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix}, \quad \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & \frac{3}{4} \end{bmatrix}, \quad \text{and} \quad \mathbf{P}^{-1} = \frac{1}{5} \begin{bmatrix} 1 & 1 \\ -3 & 2 \end{bmatrix}.$$

Now suppose that our goal is to determine the long-term distribution of population between the city and its suburbs. Note first that $(\frac{3}{4})^k$ is "negligible" when k is sufficiently large; for instance, $(\frac{3}{4})^{40} \approx 0.00001$. It follows that if $k \ge 40$, then the formula $\mathbf{A}^k = \mathbf{P}\mathbf{D}^k\mathbf{P}^{-1}$ yields

$$\mathbf{A}^{k} = \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & (\frac{3}{4})^{k} \end{bmatrix} \begin{pmatrix} \frac{1}{5} \\ \frac{1}{5} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -3 & 2 \end{bmatrix}$$
$$\approx \frac{1}{5} \begin{bmatrix} 2 & -1 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -3 & 2 \end{bmatrix}$$
$$= \frac{1}{5} \begin{bmatrix} 2 & 0 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -3 & 2 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix}.$$
(9)

Hence it follows that, if k is sufficiently large, then

$$\mathbf{x}_{k} = \mathbf{A}^{k} \mathbf{x}_{0} \approx \frac{1}{5} \begin{bmatrix} 2 & 2\\ 3 & 3 \end{bmatrix} \begin{bmatrix} C_{0}\\ S_{0} \end{bmatrix} = (C_{0} + S_{0}) \begin{bmatrix} 0.4\\ 0.6 \end{bmatrix} = \begin{bmatrix} 0.4\\ 0.6 \end{bmatrix}$$

because $C_0 + S_0 = 1$ (million), the constant total population of the metropolitan area. Thus, our analysis shows that, irrespective of the initial distribution of population between the city and its suburbs, the long-term distribution consists of 40% in the city and 60% in the suburbs.

Remark The result in Example 2—that the long-term situation is independent of the initial situation—is characteristic of a general class of common problems. Note that the transition matrix **A** in (8) has the property that *the sum of the elements in each column is* 1. An $n \times n$ matrix with nonnegative entries having this property is called a **stochastic matrix**. It can be proved that, if **A** is a stochastic matrix having only positive entries, then $\lambda_1 = 1$ is one eigenvalue of **A** and $|\lambda_i| < 1$ for the others. (See Problems 39 and 40.) Moreover, as $k \to \infty$, the matrix **A**^k approaches the constant matrix

$$\begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_1 & \cdots & \mathbf{v}_1 \end{bmatrix}$$
,

each of whose identical columns is the eigenvector of **A** associated with $\lambda_1 = 1$ that has the sum of its elements equal to 1. The 2 × 2 stochastic matrix **A** of Example 1 illustrates this general result, with $\lambda_1 = 1$, $\lambda_2 = \frac{3}{4}$, and $\mathbf{v}_1 = (\frac{2}{5}, \frac{3}{5})$.

Predator-Prey Models

Next, we consider a predator-prey population consisting of the foxes and rabbits living in a certain forest. Initially, there are F_0 foxes and R_0 rabbits; after k months, there are F_k foxes and R_k rabbits. We assume that the transition from each month to the next is described by the equations

$$F_{k+1} = 0.4F_k + 0.3R_k$$

$$R_{k+1} = -rF_k + 1.2R_k,$$
(10)

where the constant r > 0 is the "capture rate" representing the average number of rabbits consumed monthly by each fox. Thus

$$\mathbf{x}_{k+1} = \mathbf{A}\mathbf{x}_k$$
 and hence $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0$, (11)

where

$$\mathbf{x}_{k} = \begin{bmatrix} F_{k} \\ R_{k} \end{bmatrix} \quad \text{and} \quad \mathbf{A} = \begin{bmatrix} 0.4 & 0.3 \\ -r & 1.2 \end{bmatrix}. \tag{12}$$

The term $0.4F_k$ in the first equation in (10) indicates that, without rabbits to eat, only 40% of the foxes would survive each month, while the term $0.3R_k$ represents the growth in the fox population due to the available food supply of rabbits. The term $1.2R_k$ in the second equation indicates that, in the absence of any foxes, the rabbit population would increase by 20% each month. We want to investigate the long-term behavior of the fox and rabbit populations for different values of the capture rate r of rabbits by foxes.

The characteristic equation of the transition matrix \mathbf{A} in (12) is

$$(0.4 - \lambda)(1.2 - \lambda) + (0.3)r = 0;$$

$$(4 - 10\lambda)(12 - 10\lambda) + 30r = 0;$$

$$100\lambda^2 - 160\lambda + (48 + 30r) = 0.$$

The quadratic formula then yields the equation

$$\lambda = \frac{1}{200} \left[160 \pm \sqrt{(160)^2 - (400)(48 + 30r)} \right] = \frac{1}{10} \left(8 \pm \sqrt{16 - 30r} \right), \quad (13)$$

which gives the eigenvalues of **A** in terms of the capture rate r. Examples 3, 4, and 5 illustrate three possibilities (for different values of r) for what may happen to the fox and rabbit populations as k increases:

- F_k and R_k may approach constant nonzero values. This is the case of *stable limiting populations* that coexist in equilibrium with one another.
- F_k and R_k may both approach zero. This is the case of *mutual extinction* of the two species.
- F_k and R_k may both increase without bound. This is the case of a *population* explosion.

Example 3 Stable limiting population If r = 0.4, then Eq. (13) gives the eigenvalues $\lambda_1 = 1$ and $\lambda_2 = 0.6$. For $\lambda_1 = 1$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.6 & 0.3 \\ -0.4 & 0.2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

so an associated eigenvector is $\mathbf{v}_1 = \begin{bmatrix} 1 & 2 \end{bmatrix}^T$. For $\lambda_2 = 0.6$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.2 & 0.3 \\ -0.4 & 0.6 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

so an associated eigenvector is $\mathbf{v}_2 = \begin{bmatrix} 3 & 2 \end{bmatrix}^T$. It follows that $\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$, where

$$\mathbf{P} = \begin{bmatrix} 1 & 3\\ 2 & 2 \end{bmatrix}, \quad \mathbf{D} = \begin{bmatrix} 1 & 0\\ 0 & 0.6 \end{bmatrix}, \quad \text{and} \quad \mathbf{P}^{-1} = -\frac{1}{4} \begin{bmatrix} 2 & -3\\ -2 & 1 \end{bmatrix}.$$

We are now ready to calculate \mathbf{A}^k . If k is sufficiently large that $(0.6)^k \approx 0$ —for

instance, $(0.6)^{25} \approx 0.000003$ —then the formula $\mathbf{A}^k = \mathbf{P}\mathbf{D}^k\mathbf{P}^{-1}$ yields

$$\mathbf{A}^{k} = \begin{bmatrix} 1 & 3\\ 2 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0\\ 0 & (0.6)^{k} \end{bmatrix} \begin{pmatrix} -\frac{1}{4} \end{pmatrix} \begin{bmatrix} 2 & -3\\ -2 & 1 \end{bmatrix}$$
$$\approx -\frac{1}{4} \begin{bmatrix} 1 & 3\\ 2 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0\\ 0 & 0 \end{bmatrix} \begin{bmatrix} 2 & -3\\ -2 & 1 \end{bmatrix}$$
$$= -\frac{1}{4} \begin{bmatrix} 1 & 0\\ 2 & 0 \end{bmatrix} \begin{bmatrix} 2 & -3\\ -2 & 1 \end{bmatrix}$$
$$= \frac{1}{4} \begin{bmatrix} -2 & 3\\ -4 & 6 \end{bmatrix}.$$

Hence it follows that if k is sufficiently large, then

$$\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \frac{1}{4} \begin{bmatrix} -2 & 3\\ -4 & 6 \end{bmatrix} \begin{bmatrix} F_0\\ R_0 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 3R_0 - 2F_0\\ 6R_0 - 4F_0 \end{bmatrix}$$

-that is,

$$\begin{bmatrix} F_k \\ R_k \end{bmatrix} = \alpha \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \text{ where } \alpha = \frac{1}{4}(3R_0 - 2F_0). \tag{14}$$

Assuming that the initial populations are such that $\alpha > 0$ (that is, $3R_0 > 2F_0$), (14) implies that, as *k* increases, the fox and rabbit populations approach a stable situation in which there are twice as many rabbits as foxes. For instance, if $F_0 = R_0 = 100$, then when *k* is sufficiently large, there will be 25 foxes and 50 rabbits.

Example 4 Mutual extinction If r = 0.5, then Eq. (13) gives the eigenvalues $\lambda_1 = 0.9$ and $\lambda_2 = 0.7$. For $\lambda_1 = 0.9$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.5 & 0.3 \\ -0.5 & 0.3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

so an associated eigenvector is $\mathbf{v}_1 = \begin{bmatrix} 3 & 5 \end{bmatrix}^T$. For $\lambda_2 = 0.7$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.3 & 0.3 \\ -0.5 & 0.5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

so an associated eigenvector is $\mathbf{v}_2 = \begin{bmatrix} 1 & 1 \end{bmatrix}^T$. It follows that $\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$, with

$$\mathbf{P} = \begin{bmatrix} 3 & 1 \\ 5 & 1 \end{bmatrix}, \quad \mathbf{D} = \begin{bmatrix} 0.9 & 0 \\ 0 & 0.7 \end{bmatrix}, \quad \text{and} \quad \mathbf{P}^{-1} = -\frac{1}{2} \begin{bmatrix} 1 & -1 \\ -5 & 3 \end{bmatrix}$$

Now both $(0.9)^k$ and $(0.7)^k$ approach 0 as k increases without bound $(k \to +\infty)$. Hence if k is sufficiently large, then the formula $\mathbf{A}^k = \mathbf{P}\mathbf{D}^k\mathbf{P}^{-1}$ yields

$$\mathbf{A}^{k} = \begin{bmatrix} 3 & 1 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} (0.9)^{k} & 0 \\ 0 & (0.7)^{k} \end{bmatrix} (-\frac{1}{2}) \begin{bmatrix} 1 & -1 \\ -5 & 3 \end{bmatrix}$$
$$\approx -\frac{1}{2} \begin{bmatrix} 3 & 1 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -5 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$$

so

 $\begin{bmatrix} F_k \\ R_k \end{bmatrix} = \mathbf{A}^k \begin{bmatrix} F_0 \\ R_0 \end{bmatrix} \approx \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} F_0 \\ R_0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$ (15) Thus F_k and R_k both approach zero as $k \to +\infty$, so both the foxes and the rabbits die out—mutual extinction occurs.

Example 5 Population explosion If r = 0.325, then Eq. (13) gives the eigenvalues $\lambda_1 = 1.05$ and $\lambda_2 = 0.55$. For $\lambda_1 = 1.05$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.650 & 0.30 \\ -0.325 & 0.15 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Each equation is a multiple of -13x + 6y = 0, so an associated eigenvector is $\mathbf{v}_1 = \begin{bmatrix} 6 & 13 \end{bmatrix}^T$. For $\lambda_2 = 0.55$, the system $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$\begin{bmatrix} -0.150 & 0.30 \\ -0.325 & 0.65 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

so an associated eigenvector is $\mathbf{v}_2 = \begin{bmatrix} 2 & 1 \end{bmatrix}^T$. It follows that $\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$ with

$$\mathbf{P} = \begin{bmatrix} 6 & 2 \\ 13 & 1 \end{bmatrix}, \quad \mathbf{D} = \begin{bmatrix} 1.05 & 0 \\ 0 & 0.55 \end{bmatrix}, \text{ and } \mathbf{P}^{-1} = -\frac{1}{20} \begin{bmatrix} 1 & -2 \\ -13 & 6 \end{bmatrix}.$$

Note that $(0.55)^k$ approaches zero but that $(1.05)^k$ increases without bound as $k \to +\infty$. It follows that if k is sufficiently large, then the formula $\mathbf{A}^k = \mathbf{P}\mathbf{D}^k\mathbf{P}^{-1}$ yields

$$\mathbf{A}^{k} = \begin{bmatrix} 6 & 2\\ 13 & 1 \end{bmatrix} \begin{bmatrix} (1.05)^{k} & 0\\ 0 & (0.55)^{k} \end{bmatrix} \begin{pmatrix} -\frac{1}{20} \\ -13 & 6 \end{bmatrix}$$
$$\approx -\frac{1}{20} \begin{bmatrix} 6 & 2\\ 13 & 1 \end{bmatrix} \begin{bmatrix} (1.05)^{k} & 0\\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2\\ -13 & 6 \end{bmatrix}$$
$$= -\frac{1}{20} \begin{bmatrix} (6)(1.05)^{k} & 0\\ (13)(1.05)^{k} & 0 \end{bmatrix} \begin{bmatrix} 1 & -2\\ -13 & 6 \end{bmatrix},$$

and therefore

$$\mathbf{A}^{k} \approx -\frac{1}{20} (1.05)^{k} \begin{bmatrix} 6 & -12\\ 13 & -26 \end{bmatrix}.$$
 (16)

Hence, if k is sufficiently large, then

$$\mathbf{x}_{k} = \mathbf{A}^{k} \mathbf{x}_{0} \approx -\frac{1}{20} (1.05)^{k} \begin{bmatrix} 6 & -12\\ 13 & -26 \end{bmatrix} \begin{bmatrix} F_{0}\\ R_{0} \end{bmatrix}$$
$$= -\frac{1}{20} (1.05)^{k} \begin{bmatrix} 6F_{0} - 12R_{0}\\ 13F_{0} - 26R_{0} \end{bmatrix},$$

and so

$$\begin{bmatrix} F_k \\ R_k \end{bmatrix} \approx (1.05)^k \gamma \begin{bmatrix} 6 \\ 13 \end{bmatrix}, \text{ where } \gamma = \frac{1}{20}(2R_0 - F_0). \tag{17}$$

If $\gamma > 0$ (that is, if $2R_0 > F_0$), then the factor $(1.05)^k$ in (17) implies that the fox and rabbit populations both increase at a rate of 5% per month, and thus each increases without bound as $k \to +\infty$. Moreover, when k is sufficiently large, the two populations maintain a constant ratio of 6 foxes for every 13 rabbits. It is also of interest to note that the monthly "population multiplier" is the larger eigenvalue $\lambda_1 = 1.05$ and that the limiting *ratio* of populations is determined by the associated eigenvector $\mathbf{v}_1 = \begin{bmatrix} 6 & 13 \end{bmatrix}^T$.

In summary, let us compare the results in Examples 3, 4, and 5. The critical capture rate r = 0.4 of Example 3 represents a monthly consumption of 0.4 rabbits per fox, resulting in stable limiting populations of both species. But if the foxes are greedier and consume more than 0.4 rabbits per fox monthly, then the result is extinction of both species (as in Example 4). If the rabbits become more skilled at evading foxes, so that less than 0.4 rabbits per fox are consumed each month, then both populations grow without bound, as in Example 5.

The Cayley-Hamilton Theorem

One of the more notable and important theorems in advanced linear algebra says that *every matrix satisfies its own characteristic equation* (as proved for the 2×2 case in Problem 29 of Section 3.4).

THEOREM 1 Cayley-Hamilton

If the $n \times n$ matrix **A** has the characteristic polynomial

$$p(\lambda) = (-1)^n \lambda^n + c_{n-1} \lambda^{n-1} + \dots + c_2 \lambda^2 + c_1 \lambda + c_0,$$

then

$$p(\mathbf{A}) = (-1)^{n} \mathbf{A}^{n} + c_{n-1} \mathbf{A}^{n-1} + \dots + c_{2} \mathbf{A}^{2} + c_{1} \mathbf{A} + c_{0} \mathbf{I} = \mathbf{0}.$$
 (18)

Proof: We establish (18) only for the special case in which the matrix **A** is *diagonalizable*; the general case will be discussed in Section 7.6. If $\lambda_1, \lambda_2, \ldots, \lambda_n$ are the eigenvalues of **A**, then

$$\mathbf{A}^{k} = \mathbf{P}\mathbf{D}^{k}\mathbf{P}^{-1}, \text{ where } \mathbf{D}^{k} = \begin{bmatrix} \lambda_{1}^{k} & 0 & \cdots & 0\\ 0 & \lambda_{2}^{k} & \cdots & 0\\ \vdots & \vdots & \ddots & \vdots\\ 0 & 0 & \cdots & \lambda_{n}^{k} \end{bmatrix}$$

as in Eqs. (2) and (3) at the beginning of this section. First, we note that

$$p(\mathbf{D}) = (-1)^{n} \mathbf{D}^{n} + c_{n-1} \mathbf{D}^{n-1} + \dots + c_{2} \mathbf{D}^{2} + c_{1} \mathbf{D} + c_{0} \mathbf{I}$$

$$= (-1)^{n} \begin{bmatrix} \lambda_{1}^{n} & 0 & \cdots & 0 \\ 0 & \lambda_{2}^{n} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_{n}^{n} \end{bmatrix} + c_{n-1} \begin{bmatrix} \lambda_{1}^{n-1} & 0 & \cdots & 0 \\ 0 & \lambda_{2}^{n-1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_{n}^{n-1} \end{bmatrix}$$

$$+ \dots + c_{2} \begin{bmatrix} \lambda_{1}^{2} & 0 & \cdots & 0 \\ 0 & \lambda_{2}^{2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_{n}^{2} \end{bmatrix} + c_{1} \begin{bmatrix} \lambda_{1} & 0 & \cdots & 0 \\ 0 & \lambda_{2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_{n} \end{bmatrix}$$

$$+ c_{0} \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix} = \begin{bmatrix} p(\lambda_{1}) & 0 & \cdots & 0 \\ 0 & p(\lambda_{2}) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & p(\lambda_{n}) \end{bmatrix}$$

$$= \mathbf{0},$$

because $p(\lambda_k) = 0$ for k = 1, 2, ..., n. Thus the diagonal form **D** of the matrix **A** satisfies the characteristic equation of **A**. It therefore follows that

$$p(\mathbf{A}) = (-1)^n \mathbf{P} \mathbf{D}^n \mathbf{P}^{-1} + c_{n-1} \mathbf{P} \mathbf{D}^{n-1} \mathbf{P}^{-1} + \cdots$$
$$+ c_2 \mathbf{P} \mathbf{D}^2 \mathbf{P}^{-1} + c_1 \mathbf{P} \mathbf{D} \mathbf{P}^{-1} + c_0 \mathbf{P} \mathbf{I} \mathbf{P}^{-1}$$
$$= \mathbf{P} p(\mathbf{D}) \mathbf{P}^{-1}$$
$$= \mathbf{P} \mathbf{0} \mathbf{P}^{-1} = \mathbf{0},$$

as desired.

Example 6

In Example 7 of Section 6.1, we found that the matrix

$$\mathbf{A} = \begin{bmatrix} 4 & -2 & 1 \\ 2 & 0 & 1 \\ 2 & -2 & 3 \end{bmatrix}$$

has the characteristic polynomial

$$p(\lambda) = -\lambda^3 + 7\lambda^2 - 16\lambda + 12,$$

so

$$-\mathbf{A}^3 + 7\mathbf{A}^2 - 16\mathbf{A} + 12\mathbf{I} = \mathbf{0},$$
 (19)

by the Cayley-Hamilton theorem. Because

$$\mathbf{A}^2 = \begin{bmatrix} 14 & -10 & 5\\ 10 & -6 & 5\\ 10 & -10 & 9 \end{bmatrix},$$

it follows from (19) that

$$\mathbf{A}^{3} = 7\mathbf{A}^{2} - 16\mathbf{A} + 12\mathbf{I}$$

$$= 7\begin{bmatrix} 14 & -10 & 5\\ 10 & -6 & 5\\ 10 & -10 & 9 \end{bmatrix} - 16\begin{bmatrix} 4 & -2 & 1\\ 2 & 0 & 1\\ 2 & -2 & 3 \end{bmatrix} + 12\begin{bmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 46 & -38 & 19\\ 38 & -30 & 19\\ 38 & -38 & 27 \end{bmatrix}.$$

Multiplication by A now gives

$$\begin{aligned} \mathbf{A}^{4} &= 7\mathbf{A}^{3} - 16\mathbf{A}^{2} + 12\mathbf{A} \\ &= 7(7\mathbf{A}^{2} - 16\mathbf{A} + 12\mathbf{I}) - 16\mathbf{A}^{2} + 12\mathbf{A} \\ &= 33\mathbf{A}^{2} - 100\mathbf{A} + 84\mathbf{I} \\ &= 33\begin{bmatrix} 14 & -10 & 5\\ 10 & -6 & 5\\ 10 & -10 & 9 \end{bmatrix} - 100\begin{bmatrix} 4 & -2 & 1\\ 2 & 0 & 1\\ 2 & -2 & 3 \end{bmatrix} + 84\begin{bmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 146 & -130 & 65\\ 130 & -114 & 65\\ 130 & -130 & 81 \end{bmatrix}. \end{aligned}$$

Thus we can use the Cayley-Hamilton theorem to express (positive integral) powers of the 3×3 matrix **A** in terms of **A** and \mathbf{A}^2 . We also can use the Cayley-Hamilton theorem to find the inverse matrix \mathbf{A}^{-1} . If we multiply Eq. (19) by \mathbf{A}^{-1} , we can solve the resulting equation $-\mathbf{A}^2 + 7\mathbf{A} - 16\mathbf{I} + 12\mathbf{A}^{-1} = \mathbf{0}$:

$$\mathbf{A}^{-1} = \frac{1}{12} (\mathbf{A}^2 - 7\mathbf{A} + 16\mathbf{I})$$

= $\frac{1}{12} \begin{bmatrix} 14 & -10 & 5\\ 10 & -6 & 5\\ 10 & -10 & 9 \end{bmatrix} - \frac{7}{12} \begin{bmatrix} 4 & -2 & 1\\ 2 & 0 & 1\\ 2 & -2 & 3 \end{bmatrix} + \frac{16}{12} \begin{bmatrix} 1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1 \end{bmatrix}$
= $\frac{1}{6} \begin{bmatrix} 1 & 2 & -1\\ -2 & 5 & -1\\ -2 & 2 & 2 \end{bmatrix}$.

6.3 **Problems**

In Problems 1 through 10, a matrix \mathbf{A} is given. Use the method of Example 1 to compute \mathbf{A}^5 .

1.	$\begin{bmatrix} 3\\1 \end{bmatrix}$	$\begin{bmatrix} -2\\0 \end{bmatrix}$	2.	5 3	$\begin{pmatrix} -6 \\ -4 \end{bmatrix}$	
3.	$\begin{bmatrix} 6\\ 4 \end{bmatrix}$	$\begin{bmatrix} -6 \\ -4 \end{bmatrix}$	4.	$\begin{bmatrix} 4\\2 \end{bmatrix}$	$\begin{bmatrix} -3 \\ -1 \end{bmatrix}$	
5.	5 3	$\begin{bmatrix} -4 \\ -2 \end{bmatrix}$	6.	$\begin{bmatrix} 6\\2 \end{bmatrix}$	$\begin{bmatrix} -10 \\ -3 \end{bmatrix}$	
7.	$\begin{bmatrix} 1\\0\\0 \end{bmatrix}$	$\begin{bmatrix} 3 & 0 \\ 2 & 0 \\ 0 & 2 \end{bmatrix}$	8.	$\left[\begin{array}{c}1\\0\\0\end{array}\right]$	$-2 \\ 1 \\ -2$	$\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$
9.	$\begin{bmatrix} 1\\0\\0 \end{bmatrix}$	$\begin{bmatrix} -3 & 1 \\ 2 & 0 \\ 0 & 2 \end{bmatrix}$	10.	$\begin{bmatrix} 4\\2\\0 \end{bmatrix}$	$-3 \\ -1 \\ 0$	$\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$

Find A^{10} for each matrix A given in Problems 11 through 14.

11.	$\begin{bmatrix} 1\\6\\21&-1 \end{bmatrix}$	$\begin{bmatrix} 0 & 0 \\ 5 & 2 \\ 5 & -6 \end{bmatrix}$	12. $\begin{bmatrix} 11 & -6 & -2 \\ 20 & -11 & -4 \\ 0 & 0 & 1 \end{bmatrix}$
13.	$\begin{bmatrix} 1 & -1 \\ 2 & -2 \\ 4 & -4 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$	14. $\begin{bmatrix} 5 & -5 & -3 \\ 2 & -2 & -1 \\ 4 & -4 & -3 \end{bmatrix}$

15-24. Use the Cayley-Hamilton theorem (as in Example 6) to find A⁻¹, A³, and A⁴ for each matrix A given in Problems 5-14 (respectively).

In Problems 25 through 30, a city-suburban population transition matrix \mathbf{A} (as in Example 2) is given. Find the resulting long-term distribution of a constant total population between the city and its suburbs.

25.
$$\mathbf{A} = \begin{bmatrix} 0.9 & 0.1 \\ 0.1 & 0.9 \end{bmatrix}$$
26. $\mathbf{A} = \begin{bmatrix} 0.85 & 0.05 \\ 0.15 & 0.95 \end{bmatrix}$ **27.** $\mathbf{A} = \begin{bmatrix} 0.75 & 0.15 \\ 0.25 & 0.85 \end{bmatrix}$ **28.** $\mathbf{A} = \begin{bmatrix} 0.8 & 0.1 \\ 0.2 & 0.9 \end{bmatrix}$ **29.** $\mathbf{A} = \begin{bmatrix} 0.9 & 0.05 \\ 0.1 & 0.95 \end{bmatrix}$ **30.** $\mathbf{A} = \begin{bmatrix} 0.8 & 0.15 \\ 0.2 & 0.85 \end{bmatrix}$

Predator-Prey

Problems 31 through 33 deal with a fox-rabbit population as in Examples 3 through 5, except with the transition matrix

$$\mathbf{A} = \begin{bmatrix} 0.6 & 0.5\\ -r & 1.2 \end{bmatrix}$$

in place of the one used in the text.

31. If r = 0.16, show that in the long term the populations of foxes and rabbits are stable, with 5 foxes for each 4 rabbits.

- **32.** If r = 0.175, show that in the long term the populations of foxes and rabbits both die out.
- **33.** If r = 0.135, show that in the long term the fox and rabbit populations both increase at the rate of 5% per month, maintaining a constant ratio of 10 foxes for each 9 rabbits.
- **34.** Suppose that the 2 × 2 matrix **A** has eigenvalues $\lambda_1 = 1$ and $\lambda_2 = -1$ with eigenvectors $\mathbf{v}_1 = (3, 4)$ and $\mathbf{v}_2 = (5, 7)$, respectively. Find the matrix **A** and the powers \mathbf{A}^{99} and \mathbf{A}^{100} .
- **35.** Suppose that $|\lambda| = 1$ for each eigenvalue λ of the diagonalizable matrix **A**. Show that $\mathbf{A}^n = \mathbf{I}$ for every even positive integer *n*.
- **36.** Suppose that

 $\mathbf{A} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$

Show that $\mathbf{A}^{2n} = \mathbf{I}$ and that $\mathbf{A}^{2n+1} = \mathbf{A}$ for every positive integer *n*.

37. Suppose that

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Show that $\mathbf{A}^{4n} = \mathbf{I}$, $\mathbf{A}^{4n+1} = \mathbf{A}$, $\mathbf{A}^{4n+2} = -\mathbf{I}$, and $\mathbf{A}^{4n+3} = -\mathbf{A}$ for every positive integer *n*.

38. The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

is not diagonalizable. (Why not?) Write $\mathbf{A} = \mathbf{I} + \mathbf{B}$. Show that $\mathbf{B}^2 = \mathbf{0}$ and thence that

$$\mathbf{A}^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}.$$

39. Consider the stochastic matrix

$$\mathbf{A} = \begin{bmatrix} p & 1-q \\ 1-p & q \end{bmatrix},$$

where 0 and <math>0 < q < 1. Show that the eigenvalues of **A** are $\lambda_1 = 1$ and $\lambda_2 = p + q - 1$, so that $|\lambda_2| < 1$.

- **40.** Suppose that **A** is an $n \times n$ stochastic matrix—the sum of the elements of each column vector is 1. If $\mathbf{v} = (1, 1, ..., 1)$, show that $\mathbf{A}^T \mathbf{v} = \mathbf{v}$. Why does it follow that $\lambda = 1$ is an eigenvalue of **A**?
- **41.** In his *Liber abaci* (Book of Calculation) published in 1202, Leonardo Fibonacci^{*} asked the following question: How many pairs of rabbits are produced from a single original pair in one year, if every month each pair begets a new pair, which is similarly productive beginning in the second succeeding month? The answer is provided by the **Fibonacci sequence**

in which each term is the sum of its two immediate predecessors. That is, the sequence is defined recursively as follows:

$$s_0 = 1 = s_1$$
, $s_{n+1} = s_n + s_{n-1}$ for $n \ge 1$.

* "Leonardo, son of Bonacci," 1175–1250?, one of the outstanding mathematicians of the middle ages. He is also known as Leonardo of Pisa (Leonardo Pisano).

364 CHAPTER 6 EIGENVALUES AND EIGENVECTORS

Then s_n is the number of rabbit pairs present after n months. Note that if

$$\mathbf{x}_n = \begin{bmatrix} s_{n+1} \\ s_n \end{bmatrix}$$
 and $\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

then

$$\mathbf{x}_n = \mathbf{A}\mathbf{x}_{n-1}$$
 and so $\mathbf{x}_n = \mathbf{A}^n\mathbf{x}_0$,

where $x_0 = (1, 1)$.

- (a) Show that A has eigenvalues $\lambda_1 = \frac{1}{2}(1 + \sqrt{5})$ and $\lambda_2 = \frac{1}{2}(1 \sqrt{5})$ with associated eigenvectors $\mathbf{v}_1 = (1 + \sqrt{5}, 2)$ and $\mathbf{v}_2 = (1 \sqrt{5}, 2)$, respectively.
- (b) Compute

$$\begin{bmatrix} s_{n+1} \\ s_n \end{bmatrix} = \mathbf{A}^n \mathbf{x}_0 = \mathbf{P} \mathbf{D}^n \mathbf{P}^{-1} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

to derive the amazing formula

$$s_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^{n+1} - \left(\frac{1-\sqrt{5}}{2} \right)^{n+1} \right].$$

Thus after 1 year the number of rabbit pairs is

$$s_{12} = \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^{13} - \left(\frac{1-\sqrt{5}}{2} \right)^{13} \right] = 233.$$

Show that there are 75,025 rabbit pairs after 2 years and over 2.5 trillion rabbit pairs after 5 years.

Linear Systems of Differential Equations

7.1 First-Order Systems and Applications

In Chapters 1 and 5 we discussed methods for solving an ordinary differential equation that involves only one dependent variable. Many applications, however, require the use of two or more dependent variables, each a function of a single independent variable (typically time). Such a problem leads naturally to a *system* of simultaneous ordinary differential equations. We will usually denote the independent variable by t and the dependent variables (the unknown functions of t) by x_1 , x_2 , x_3 , ..., or by x, y, z, Primes will indicate derivatives with respect to t.

We will restrict our attention to systems in which the number of equations is the same as the number of dependent variables (unknown functions). For instance, a system of two first-order equations in the dependent variables x and y has the general form

$$f(t, x, y, x', y') = 0,$$

$$g(t, x, y, x', y') = 0,$$
(1)

where the functions f and g are given. A **solution** of this system is a pair x(t), y(t) of functions of t that satisfy both equations identically over some interval of values of t.

For an example of a second-order system, consider a particle of mass *m* that moves in space under the influence of a force field **F** that depends on time *t*, the position (x(t), y(t), z(t)) of the particle, and its velocity (x'(t), y'(t), z'(t)). Writing Newton's law $m\mathbf{a} = \mathbf{F}$ componentwise, we get the system

$$mx'' = F_1(t, x, y, z, x', y', z'),$$

$$my'' = F_2(t, x, y, z, x', y', z'),$$

$$mz'' = F_3(t, x, y, z, x', y', z')$$
(2)

of three second-order equations with independent variable *t* and dependent variables *x*, *y*, *z*; the three right-hand side functions F_1 , F_2 , F_3 are the components of the vector-valued function **F**.

Initial Applications

Examples 1 and 2 further illustrate how systems of differential equations arise naturally in scientific problems.

Dual mass-spring system Consider the system of two masses and two springs shown in Fig. 7.1.1, with a given external force f(t) acting on the right-hand mass m_2 . We denote by x(t) the displacement (to the right) of the mass m_1 from its static equilibrium position [when the system is motionless and in equilibrium and f(t) = 0] and by y(t) the displacement of the mass m_2 from its static position. Thus the two springs are neither stretched nor compressed when x and y are zero.

In the configuration in Fig. 7.1.1, the first spring is stretched x units and the second by y - x units. We apply Newton's law of motion to the two "free body diagrams" shown in Fig. 7.1.2; we thereby obtain the system

$$m_1 x'' = -k_1 x + k_2 (y - x),$$

$$m_2 y'' = -k_2 (y - x) + f(t)$$
(3)

of differential equations that the position functions x(t) and y(t) must satisfy. For instance, if $m_1 = 2$, $m_2 = 1$, $k_1 = 4$, $k_2 = 2$, and $f(t) = 40 \sin 3t$ in appropriate physical units, then the system in (3) reduces to

$$2x'' = -6x + 2y,$$

$$y'' = 2x - 2y + 40 \sin 3t.$$
(4)

Dual brine tanks Consider two brine tanks connected as shown in Fig. 7.1.3. Tank 1 contains x(t) pounds of salt in 100 gal of brine and tank 2 contains y(t) pounds of salt in 200 gal of brine. The brine in each tank is kept uniform by stirring, and brine is pumped from each tank to the other at the rates indicated in Fig. 7.1.3. In addition, fresh water flows into tank 1 at 20 gal/min, and the brine in tank 2 flows out at 20 gal/min (so the total volume of brine in the two tanks remains constant). The salt concentrations in the two tanks are x/100 pounds per gallon and y/200 pounds per gallon, respectively. When we compute the rates of change of the amount of salt in the two tanks, we therefore get the system of differential equations that x(t) and y(t) must satisfy:

$$x' = -30 \cdot \frac{x}{100} + 10 \cdot \frac{y}{200} = -\frac{3}{10}x + \frac{1}{20}y,$$

$$y' = 30 \cdot \frac{x}{100} - 10 \cdot \frac{y}{200} - 20 \cdot \frac{y}{200} = \frac{3}{10}x - \frac{3}{20}y$$

$$20x' = -6x + y,$$

$$20y' = 6x - 3y.$$
(5)

First-Order Systems

-that is,

Consider a system of differential equations that can be solved for the highest-order derivatives of the dependent variables that appear, as explicit functions of t and lower-order derivatives of the dependent variables. For instance, in the case of a system of two second-order equations, our assumption is that it can be written in the form

$$\begin{aligned} x_1'' &= f_1(t, x_1, x_2, x_1', x_2'), \\ x_2'' &= f_2(t, x_1, x_2, x_1', x_2'). \end{aligned}$$
(6)

It is of both practical and theoretical importance that any such higher-order system can be transformed into an equivalent system of *first-order* equations.

Equilibrium positions

x(t)

 k_1

 m_1

FIGURE 7.1.1. The mass-and-spring system of Example 1.

 k_2

 m_2

y(t)



FIGURE 7.1.2. The free body diagrams for the system of Example 1.

Example 2



FIGURE 7.1.3. The two brine tanks of Example 2.

7.1 FIRST-ORDER SYSTEMS AND APPLICATION \$67

To describe how such a transformation is accomplished, we consider first the "system" consisting of the single nth-order equation

$$x^{(n)} = f(t, x, x', \dots, x^{(n-1)}).$$
⁽⁷⁾

We introduce the dependent variables x_1, x_2, \ldots, x_n defined as follows:

$$x_1 = x, \quad x_2 = x', \quad x_3 = x'', \quad \dots, \quad x_n = x^{(n-1)}.$$
 (8)

Note that $x'_1 = x' = x_2$, $x'_2 = x'' = x_3$, and so on. Hence the substitution of (8) in Eq. (7) yields the system

$$x'_{1} = x_{2},$$

$$x'_{2} = x_{3},$$

$$\vdots$$

$$x'_{n-1} = x_{n},$$

$$x'_{n} = f(t, x_{1}, x_{2}, \dots, x_{n})$$
(9)

of *n* first-order equations. Evidently, this system is equivalent to the original *n*thorder equation in (7), in the sense that x(t) is a solution of Eq. (7) if and only if the functions $x_1(t), x_2(t), \ldots, x_n(t)$ defined in (8) satisfy the system of equations in (9).

Example 3	The third-order equation	
	$x^{(3)} + 3x'' + 2x' - 5x = \sin 2t$	
	is of the form in (7) with	
	$f(t, x, x', x'') = 5x - 2x' - 3x'' + \sin 2t.$	

Hence the substitutions

$$x_1 = x$$
, $x_2 = x' = x'_1$, $x_3 = x'' = x'_2$

yield the system

$$x'_1 = x_2,$$

 $x'_2 = x_3,$
 $x'_3 = 5x_1 - 2x_2 - 3x_3 + \sin 2t$

of three first-order equations.

It may appear that the first-order system obtained in Example 3 offers little advantage, because we could use the methods of Chapter 5 to solve the original (linear) third-order equation. But suppose that we were confronted with the nonlinear equation

$$x'' = x^3 + (x')^3,$$

to which none of our earlier methods can be applied. The corresponding first-order system is

$$x'_1 = x_2,$$

 $x'_2 = (x_1)^3 + (x_2)^3,$
(10)

and we will see in Section 7.7 that there exist effective numerical techniques for approximating the solution of essentially any first-order system. So, in this case, the transformation to a first-order system *is* advantageous. From a practical viewpoint, large systems of higher-order differential equations typically are solved numerically with the aid of the computer, and the first step is to transform such a system into a first-order system for which a standard computer program is available.

Example 4

$$2x'' = -6x + 2y,$$

$$y'' = 2x - 2y + 40\sin 3t$$
(4)

of second-order equations was derived in Example 1. Transform this system into an equivalent first-order system.

Solution

$$x_1 = x$$
, $x_2 = x' = x'_1$, $y_1 = y$, $y_2 = y' = y'_1$.

Then the system in (4) yields the system

Motivated by the equations in (8), we define

$$x'_{1} = x_{2},$$

$$2x'_{2} = -6x_{1} + 2y_{1},$$

$$y'_{1} = y_{2},$$

$$y'_{2} = 2x_{1} - 2y_{1} + 40\sin 3t$$
(11)

of four first-order equations in the dependent variables x_1 , x_2 , y_1 , and y_2 .

Simple Two-Dimensional Systems

The linear second-order differential equation

$$x'' + px' + qx = 0 (12)$$

(with constant coefficients and independent variable *t*) transforms via the substitutions x' = y, x'' = y' into the two-dimensional linear system

$$\begin{aligned} x' &= y, \\ y' &= -qx - py. \end{aligned} \tag{13}$$

Conversely, we can solve this system in (13) by solving the familiar single equation in (12).

Example 5 To solve the two-dimensional system

x' = -2y, $y' = \frac{1}{2}x,$ (14)

we begin with the observation that

$$x'' = -2y' = -2\left(\frac{1}{2}x\right) = -x$$

This gives the single second-order equation x'' + x = 0 having general solution

$$x(t) = A\cos t + B\sin t = C\cos(t - \alpha),$$

where $A = C \cos \alpha$ and $B = C \sin \alpha$. Then

$$y(t) = -\frac{1}{2}x'(t) = -\frac{1}{2}(-A\sin t + B\cos t)$$

= $\frac{1}{2}C\sin(t-\alpha)$.

The identity $\cos^2 \theta + \sin^2 \theta = 1$ therefore implies that, for each value of t, the point (x(t), y(t)) lies on the ellipse

$$\frac{x^2}{C^2} + \frac{y^2}{(C/2)^2} = 1$$

with semiaxes C and C/2. Figure 7.1.4 shows several such ellipses in the xy-plane.



FIGURE 7.1.4. Direction field and solution curves for the system x' = -2y, $y' = \frac{1}{2}x$ of Example 5.

A solution (x(t), y(t)) of a two-dimensional system

$$x' = f(t, x, y),$$
$$y' = g(t, x, y)$$

can be regarded as a parametrization of a **solution curve** or **trajectory** of the system in the xy-plane. Thus the trajectories of the system in (14) are the ellipses of Fig. 7.1.4. The choice of an initial point (x(0), y(0)) determines which one of these trajectories a particular solution parametrizes.

The picture showing a system's trajectories in the *xy*-plane—its so-called *phase plane portrait*—fails to reveal precisely how the point (x(t), y(t)) moves along its trajectory. If the functions f and g do not involve the independent variable t, then a direction field—showing typical arrows representing vectors with components (proportional to) the derivatives x' = f(x, y) and y' = g(x, y)—can be plotted. Because the moving point (x(t), y(t)) has velocity vector (x'(t), y'(t)), this direction field indicates the point's direction of motion along its trajectory. For instance, the direction field plotted in Fig. 7.1.4 indicates that each such point moves counterclockwise around its elliptical trajectory. Additional information can be shown in the separate graphs of x(t) and y(t) as functions of t.

Continued With initial values x(0) = 2, y(0) = 0, the general solution in Example 5 yields

$$x(0) = A = 2, \quad y(0) = -\frac{1}{2}B = 0.$$

The resulting particular solution is given by

$$x(t) = 2\cos t, \quad y(t) = \sin t.$$

The graphs of the two functions are shown in Fig. 7.1.5. We see that x(t) initially decreases while y(t) increases. It follows that, as t increases, the solution point (x(t), y(t)) traverses the trajectory $\frac{1}{4}x^2 + y^2 = 1$ in the counterclockwise direction, as indicated by the direction field vectors in Fig. 7.1.4.

Example 6 To find a general solution of the system

$$\begin{aligned} x' &= y, \\ y' &= 2x + y, \end{aligned} \tag{15}$$

we begin with the observation that

$$x'' = y' = 2x + y = x' + 2x.$$

This gives the single linear second-order equation

$$x'' - x' - 2x = 0$$

having the characteristic equation

$$r^2 - r - 2 = (r+1)(r-2) = 0$$

and the general solution

$$x(t) = Ae^{-t} + Be^{2t}.$$
 (16)

Therefore,

FIGURE 7.1.6. Direction field and solution curves for the system x' = y, y' = 2x + y of Example 6.

$$y(t) = x^{*}(t) = -Ae^{-t} + 2Be^{2t}.$$
 (1/

 $t = t \cdot c \cdot p \cdot 2t$

Typical phase plane trajectories of the system in (15) parametrized by Eqs. (16) and (17) are shown in Fig. 7.1.6. These trajectories may resemble hyperbolas sharing common asymptotes, but Problem 23 shows that their actual form is somewhat more complicated.



 $v = \sin t$

4

3

2

1

- 1

-2

-3

0

 $x = 2 \cos t$



Example 7

To solve the initial value problem

$$x' = -y,$$

$$y' = (1.01)x - (0.2)y,$$

$$x(0) = 0, \quad y(0) = 1,$$

(18)

we begin with the observation that

$$x'' = -y' = -[(1.01)x - (0.2)y] = (-1.01)x - (0.2)x'.$$

This gives the single linear second-order equation

$$x'' + (0.2)x' + (1.01)x = 0$$

having the characteristic equation

$$r^{2} + (0.2)r + 1.01 = (r + 0.1)^{2} + 1 = 0$$

characteristic roots $-0.1 \pm i$, and the general solution

$$x(t) = e^{-t/10} (A \cos t + B \sin t).$$

Then x(0) = A = 0, so

$$x(t) = Be^{-t/10} \sin t,$$

$$y(t) = -x'(t) = \frac{1}{10}Be^{-t/10} \sin t - Be^{-t/10} \cos t.$$

Finally, y(0) = -B = 1, so the desired solution of the system in (18) is

$$x(t) = e^{-t/10} \sin t,$$

$$y(t) = \frac{1}{10}e^{-t/10}(\sin t + 10\cos t).$$
(19)

These equations parametrize the spiral trajectory in Fig. 7.1.7; the trajectory approaches the origin as $t \to +\infty$. Figure 7.1.8 shows the *x*- and *y*-solution curves given in (19).

When we study linear systems in subsequent sections, we will learn why the superficially similar systems in Examples 5 through 7 have the markedly different trajectories shown in Figs. 7.1.4, 7.1.6, and 7.1.7.

Linear Systems

In addition to practical advantages for numerical computation, the general theory of systems and systematic solution techniques are more easily and more concisely described for first-order systems than for higher-order systems. For instance, consider a *linear* first-order system of the form

$$\begin{aligned} x_1' &= p_{11}(t)x_1 + p_{12}(t)x_2 + \dots + p_{1n}(t)x_n + f_1(t), \\ x_2' &= p_{21}(t)x_1 + p_{22}(t)x_2 + \dots + p_{2n}(t)x_n + f_2(t), \\ \vdots \\ x_n' &= p_{n1}(t)x_1 + p_{n2}(t)x_2 + \dots + p_{nn}(t)x_n + f_n(t). \end{aligned}$$
(20)

We say that this system is **homogeneous** if the functions f_1, f_2, \ldots, f_n are all identically zero; otherwise, it is **nonhomogeneous**. Thus the linear system in (5) is homogeneous, whereas the linear system in (11) is nonhomogeneous. The system in (10) is nonlinear because the right-hand side of the second equation is not a linear function of the dependent variables x_1 and x_2 .



FIGURE 7.1.8. *x*- and *y*-solution curves for the initial value problem of Example 7.



FIGURE 7.1.7. Direction field and solution curve for the system x' = -y,

y' = (1.01)x - (0.2)y of Example 7.

7.1 FIRST-ORDER SYSTEMS AND APPLICATION \$371

A solution of the system in (20) is an *n*-tuple of functions $x_1(t)$, $x_2(t)$, ..., $x_n(t)$ that (on some interval) identically satisfy each of the equations in (20). We will see that the general theory of a system of *n* linear first-order equations shares many similarities with the general theory of a single *n*th-order linear differential equation. Theorem 1 (proved in Appendix A) is analogous to Theorem 2 of Section 5.2. It tells us that if the coefficient functions p_{ij} and f_j in (20) are continuous, then the system has a unique solution satisfying given initial conditions.

THEOREM 1 Existence and Uniqueness for Linear Systems

Suppose that the functions p_{11} , p_{12} , ..., p_{nn} and the functions f_1 , f_2 , ..., f_n are continuous on the open interval I containing the point a. Then, given the n numbers b_1 , b_2 , ..., b_n , the system in (20) has a unique solution on the entire interval I that satisfies the n initial conditions

$$x_1(a) = b_1, \quad x_2(a) = b_2, \quad \dots, \quad x_n(a) = b_n.$$
 (21)

Thus n initial conditions are needed to determine a solution of a system of n linear first-order equations, and we therefore expect a general solution of such a system to involve n arbitrary constants. For instance, we saw in Example 4 that the second-order linear system

$$2x'' = -6x + 2y, y'' = 2x - 2y + 40 \sin 3t,$$

which describes the position functions x(t) and y(t) of Example 1, is equivalent to the system of *four* first-order linear equations in (11). Hence four initial conditions would be needed to determine the subsequent motions of the two masses in Example 1. Typical initial values would be the initial positions x(0) and y(0) and the initial velocities x'(0) and y'(0). On the other hand, we found that the amounts x(t) and y(t) of salt in the two tanks of Example 2 are described by the system

$$20x' = -6x + y,$$

$$20y' = 6x - 3y$$

of *two* first-order linear equations. Hence the two initial values x(0) and y(0) should suffice to determine the solution. Given a higher-order system, we often must transform it into an equivalent first-order system to discover how many initial conditions are needed to determine a unique solution. Theorem 1 tells us that the number of such conditions is precisely the same as the number of equations in the equivalent first-order system.

7.1 **Problems**

In Problems 1 through 10, transform the given differential equation or system into an equivalent system of first-order differential equations.

1. $x'' + 3x' + 7x = t^2$ 2. $x^{(4)} + 6x'' - 3x' + x = \cos 3t$ 3. $t^2x'' + tx' + (t^2 - 1)x = 0$ 4. $t^3x^{(3)} - 2t^2x'' + 3tx' + 5x = \ln t$ 5. $x^{(3)} = (x')^2 + \cos x$ 6. x'' - 5x + 4y = 0, y'' + 4x - 5y = 07. $x'' = -\frac{kx}{(x^2 + y^2)^{3/2}}, y'' = -\frac{ky}{(x^2 + y^2)^{3/2}}$ 8. $x'' + 3x' + 4x - 2y = 0, y'' + 2y' - 3x + y = \cos t$ 9. x'' = 3x - y + 2z, y'' = x + y - 4z, z'' = 5x - y - z10. x'' = (1 - y)x, y'' = (1 - x)y Use the method of Examples 5, 6, and 7 to find general solutions of the systems in Problems 11 through 20. If initial conditions are given, find the corresponding particular solution. For each problem, use a computer system or graphing calculator to construct a direction field and typical solution curves for the given system.

- 11. x' = y, y' = -x
- **12.** x' = y, y' = x
- **13.** x' = -2y, y' = 2x; x(0) = 1, y(0) = 0
- **14.** x' = 10y, y' = -10x; x(0) = 3, y(0) = 4
- **15.** $x' = \frac{1}{2}y, y' = -8x$
- **16.** x' = 8y, y' = -2x
- **17.** x' = y, y' = 6x y; x(0) = 1, y(0) = 2
- **18.** x' = -y, y' = 10x 7y; x(0) = 2, y(0) = -7
- **19.** x' = -y, y' = 13x + 4y; x(0) = 0, y(0) = 3
- **20.** x' = y, y' = -9x + 6y
- **21.** (a) Calculate $[x(t)]^2 + [y(t)]^2$ to show that the trajectories of the system x' = y, y' = -x of Problem 11 are circles. (b) Calculate $[x(t)]^2 [y(t)]^2$ to show that the trajectories of the system x' = y, y' = x of Problem 12 are hyperbolas.
- 22. (a) Beginning with the general solution of the system x' = -2y, y' = 2x of Problem 13, calculate $x^2 + y^2$ to show that the trajectories are circles. (b) Show similarly that the trajectories of the system $x' = \frac{1}{2}y$, y' = -8x of Problem 15 are ellipses with equations of the form $16x^2 + y^2 = C^2$.
- **23.** First solve Eqs. (16) and (17) for e^{-t} and e^{2t} in terms of x(t), y(t), and the constants *A* and *B*. Then substitute the results in $(e^{2t})(e^{-t})^2 = 1$ to show that the trajectories of the system x' = y, y' = 2x + y in Example 6 satisfy an equation of the form

$$4x^3 - 3xy^2 + y^3 = C$$
 (constant).

Then show that C = 0 yields the straight lines y = -x and y = 2x that are visible in Fig. 7.1.6.

24. Dual mass-spring system Derive the equations

$$m_1 x_1'' = -(k_1 + k_2)x_1 + k_2 x_2,$$

$$m_2 x_2'' = k_2 x_1 - (k_2 + k_3)x_2$$

for the displacements (from equilibrium) of the two masses shown in Fig. 7.1.9.



FIGURE 7.1.10. The mechanical system of Problem 25.



FIGURE 7.1.11. The fermentation tanks of Problem 26.



FIGURE 7.1.9. The system of Problem 24.

25. Oscillating particles Two particles each of mass *m* are attached to a string under (constant) tension *T*, as indicated in Fig. 7.1.10. Assume that the particles oscillate vertically (that is, parallel to the *y*-axis) with amplitudes so small that the sines of the angles shown are accurately approximated by their tangents. Show that the displacements y_1 and y_2 satisfy the equations

$$ky_1'' = -2y_1 + y_2, \quad ky_2'' = y_1 - 2y_2,$$

where k = mL/T.

26. Fermentation vats Three 100-gal fermentation vats are connected as indicated in Fig. 7.1.11, and the mixtures in each tank are kept uniform by stirring. Denote by $x_i(t)$ the amount (in pounds) of alcohol in tank T_i at time t (i = 1, 2, 3). Suppose that the mixture circulates between the tanks at the rate of 10 gal/min. Derive the equations

$$10x'_1 = -x_1 + x_3$$

$$10x'_2 = x_1 - x_2$$

$$10x'_3 = x_2 - x_3.$$

27. A particle of mass *m* moves in the plane with coordinates (x(t), y(t)) under the influence of a force that is directed toward the origin and has magnitude $k/(x^2 + y^2)$ —an inverse-square central force field. Show that

$$mx'' = -\frac{kx}{r^3}$$
 and $my'' = -\frac{ky}{r^3}$,

where $r = \sqrt{x^2 + y^2}$.

28. Suppose that a projectile of mass *m* moves in a vertical plane in the atmosphere near the surface of the earth under the influence of two forces: a downward gravitational force \mathbf{F}_{G} of magnitude mg, and a resistive force \mathbf{F}_{R} that is directed opposite to the velocity vector \mathbf{v} and has magnitude kv^{2} (where $v = |\mathbf{v}|$ is the speed of the projectile; see Fig. 7.1.12). Show that the equations of motion of the projectile are

$$mx'' = -kvx', \quad my'' = -kvy' - mg,$$

where $v = \sqrt{(x')^2 + (y')^2}.$



FIGURE 7.1.12. The trajectory of the projectile of Problem 28.

7.1 FIRST-ORDER SYSTEMS AND APPLICATION \$373

29. Suppose that a particle with mass *m* and electrical charge *q* moves in the *xy*-plane under the influence of the magnetic field $\mathbf{B} = B\mathbf{k}$ (thus a uniform field parallel to the *z*-axis),

so the force on the particle is $\mathbf{F} = q\mathbf{v} \times \mathbf{B}$ if its velocity is **v**. Show that the equations of motion of the particle are

$$mx'' = +qBy', \quad my'' = -qBx'.$$

7.1 Application Gravitation and Kepler's Laws of Planetary Motion



Around the turn of the seventeenth century, Johannes Kepler analyzed a lifetime of planetary observations by the astronomer Tycho Brahe. Kepler concluded that the motion of the planets around the sun is described by the following three propositions, now known as **Kepler's laws of planetary motion:**

- 1. The orbit of each planet is an ellipse with the sun at one focus.
- **2.** The radius vector from the sun to each planet sweeps out area at a constant rate.
- **3.** The square of the planet's period of revolution is proportional to the cube of the major semiaxis of its elliptical orbit.

In his *Principia Mathematica* (1687) Isaac Newton deduced the inverse square law of gravitation from Kepler's laws. In this application we lead you (in the opposite direction) through a derivation of Kepler's first two laws from Newton's law of gravitation. Newton's explanation of planetary motion in terms of gravitation is a landmark of scientific and intellectual history.

Assume that the sun is located at the origin in the plane of motion of a planet, and write the position vector of the planet in the form

$$\mathbf{r}(t) = (x(t), y(t)) = x\mathbf{i} + y\mathbf{j},\tag{1}$$

where $\mathbf{i} = (1, 0)$ and $\mathbf{j} = (0, 1)$ denote the unit vectors in the positive *x*- and *y*-directions. Then the inverse-square law of gravitation implies (Problem 27) that the acceleration vector $\mathbf{r}''(t)$ of the planet is given by

$$\mathbf{r}'' = -\frac{k\mathbf{r}}{r^3},\tag{2}$$

where $r = \sqrt{x^2 + y^2}$ is the distance from the sun to the planet. If the polar coordinates of the planet at time t are $(r(t), \theta(t))$, then the radial and transverse unit vectors shown in Fig. 7.1.13 are given by

$$\mathbf{u}_r = \mathbf{i}\cos\theta + \mathbf{j}\sin\theta$$
 and $\mathbf{u}_{\theta} = -\mathbf{i}\sin\theta + \mathbf{j}\cos\theta$. (3)

The radial unit vector \mathbf{u}_r (when located at the planet's position) always points directly away from the origin, so $\mathbf{u}_r = \mathbf{r}/r$, and the transverse unit vector \mathbf{u}_{θ} is obtained from \mathbf{u}_r by a 90° counterclockwise rotation.

STEP 1: Differentiate the equations in (3) componentwise to show that

$$\frac{d\mathbf{u}_r}{dt} = \mathbf{u}_\theta \frac{d\theta}{dt} \quad \text{and} \quad \frac{d\mathbf{u}_\theta}{dt} = -\mathbf{u}_r \frac{d\theta}{dt}.$$
(4)

STEP 2: Use the equations in (4) to differentiate the planet's position vector $\mathbf{r} = r\mathbf{u}_r$ and thereby show that its velocity vector is given by

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} = \mathbf{u}_r \frac{dr}{dt} + r \frac{d\theta}{dt} \mathbf{u}_{\theta}.$$
 (5)



FIGURE 7.1.13. The radial and transverse unit vectors \mathbf{u}_r and \mathbf{u}_{θ} .

STEP 3: Differentiate again to show that the planet's acceleration vector $\mathbf{a} = d\mathbf{v}/dt$ is given by

$$\mathbf{a} = \left[\frac{d^2r}{dt^2} - r\left(\frac{d\theta}{dt}\right)^2\right]\mathbf{u}_r + \left[\frac{1}{r}\frac{d}{dt}\left(r^2\frac{d\theta}{dt}\right)\right]\mathbf{u}_\theta.$$
 (6)

STEP 4: The radial and transverse components on the right-hand sides in Eqs. (2) and (6) must agree. Equating the transverse components—that is, the coefficients of \mathbf{u}_{θ} —we get

$$\frac{1}{r}\frac{d}{dt}\left(r^2\frac{d\theta}{dt}\right) = 0,\tag{7}$$

so it follows that

$$r^2 \frac{d\theta}{dt} = h,\tag{8}$$

where *h* is a constant. Because the polar-coordinate area element—for computation of the area A(t) in Fig. 7.1.14—is given by $dA = \frac{1}{2}r^2d\theta$, Eq. (8) implies that the derivative A'(t) is *constant*, which is a statement of Kepler's second law.

STEP 5: Equate radial components in (2) and (6), and then use the result in (8) to show that the planet's radial coordinate function r(t) satisfies the second-order differential equation

$$\frac{d^2r}{dt^2} - \frac{h^2}{r^3} = -\frac{k}{r^2}.$$
(9)

STEP 6: Although the differential equation in (9) is nonlinear, it can be transformed to a linear equation by means of a simple substitution. For this purpose, assume that the orbit can be written in the polar-coordinate form $r = r(\theta)$, and first use the chain rule and Eq. (8) to show that if r = 1/z, then

$$\frac{dr}{dt} = -h\frac{dz}{d\theta}.$$

Differentiate again, to deduce from Eq. (9) that the function $z(\theta) = 1/r(\theta)$ satisfies the second-order equation

$$\frac{d^2z}{d\theta^2} + z = \frac{k}{h^2}.$$
(10)

STEP 7: Show that the general solution of Eq. (10) is

$$z(\theta) = A\cos\theta + B\sin\theta + \frac{k}{h^2}.$$
 (11)

STEP 8: Finally, deduce from Eq. (11) that $r(\theta) = 1/z(\theta)$ is given by

$$r(\theta) = \frac{L}{1 + e\cos(\theta - \alpha)},\tag{12}$$

with $e = Ch^2/k$, $C \cos \alpha = A$, $C \sin \alpha = B$, and $L = h^2/k$. The polar-coordinate graph of Eq. (12) is a conic section of eccentricity *e*—an ellipse if $0 \le e < 1$, a parabola if e = 1, and a hyperbola if e > 1—with focus at the origin. Planetary orbits are bounded and therefore are ellipses, with eccentricity e < 1. As indicated in Fig. 7.1.15, the major axis of the ellipse lies along the radial line $\theta = \alpha$.



FIGURE 7.1.14. Area swept out by the radius vector.



FIGURE 7.1.15. The elliptical orbit

$$r = \frac{L}{1 + e\cos(\theta - \alpha)}$$

with perihelion distance L/(1 + z) and a shall

 $r_1 = L/(1 + e)$ and aphelion distance $r_2 = L/(1 - e)$.

STEP 9: Plot some typical elliptical orbits—as described by (12)—having different eccentricities, sizes, and orientations. In rectangular coordinates you can write

$$x(t) = r(t)\cos t$$
, $y(t) = r(t)\sin t$, $0 \le t \le 2\pi$

to plot an elliptical orbit with eccentricity e, semilatus rectum L (Fig. 7.1.15), and rotation angle α . The eccentricity of the earth's orbit is $e \approx 0.0167$, so close to zero that the orbit looks nearly circular (though with the sun off center), and the eccentricities of the other planetary orbits range from 0.0068 for Venus and 0.0933 for Mars to 0.2056 for Mercury (and 0.2486 for Pluto, no longer classified a planet). But many comets have highly eccentric orbits—Halley's comet has $e \approx 0.97$ (Fig.7.1.16).

7.2 Matrices and Linear Systems

A system of differential equations often can be simplified by expressing it as a single differential equation involving a matrix-valued function. A **matrix-valued function**, or simply **matrix function**, is a matrix such as

$$\mathbf{x}(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_n(t) \end{bmatrix}$$
(1)

or

If

then

$$\mathbf{A}(t) = \begin{bmatrix} a_{11}(t) & a_{12}(t) & \cdots & a_{1n}(t) \\ a_{21}(t) & a_{22}(t) & \cdots & a_{2n}(t) \\ \vdots & \vdots & & \vdots \\ a_{m1}(t) & a_{m2}(t) & \cdots & a_{mn}(t) \end{bmatrix},$$
(2)

in which each entry is a function of t. We say that the matrix function $\mathbf{A}(t)$ is **continuous** (or **differentiable**) at a point (or on an interval) if each of its elements has the same property. The **derivative** of a differentiable matrix function is defined by elementwise differentiation; that is,

$$\mathbf{A}'(t) = \frac{d\mathbf{A}}{dt} = \left[\frac{da_{ij}}{dt}\right].$$
(3)

Example 1

$$\mathbf{x}(t) = \begin{bmatrix} t \\ t^2 \\ e^{-t} \end{bmatrix} \text{ and } \mathbf{A}(t) = \begin{bmatrix} \sin t & 1 \\ t & \cos t \end{bmatrix},$$
$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} 1 \\ 2t \\ -e^{-t} \end{bmatrix} \text{ and } \mathbf{A}'(t) = \begin{bmatrix} \cos t & 0 \\ 1 & -\sin t \end{bmatrix}.$$





The differentiation rules

$$\frac{d}{dt}(\mathbf{A} + \mathbf{B}) = \frac{d\mathbf{A}}{dt} + \frac{d\mathbf{B}}{dt}$$
(4)

and

$$\frac{d}{dt}(\mathbf{AB}) = \mathbf{A}\frac{d\mathbf{B}}{dt} + \frac{d\mathbf{A}}{dt}\mathbf{B}$$
(5)

follow readily by elementwise application of the analogous differentiation rules of elementary calculus for real-valued functions. If c is a (constant) real number and **C** is a constant matrix, then

$$\frac{d}{dt}(c\mathbf{A}) = c\frac{d\mathbf{A}}{dt}, \quad \frac{d}{dt}(\mathbf{C}\mathbf{A}) = \mathbf{C}\frac{dA}{dt}, \quad \text{and} \quad \frac{d}{dt}(\mathbf{A}\mathbf{C}) = \frac{d\mathbf{A}}{dt}\mathbf{C}.$$
 (6)

Because of the noncommutativity of matrix multiplication, it is important not to reverse the order of the factors in Eqs. (5) and (6).

First-Order Linear Systems

The notation and terminology of matrices and vectors may seem rather elaborate when first encountered, but it is readily assimilated with practice. Our main use for matrix notation will be the simplification of computations with systems of differential equations, especially those computations that would be burdensome in scalar notation.

We discuss here the general system of *n* first-order linear equations

$$\begin{aligned} x_1' &= p_{11}(t)x_1 + p_{12}(t)x_2 + \dots + p_{1n}(t)x_n + f_1(t), \\ x_2' &= p_{21}(t)x_1 + p_{22}(t)x_2 + \dots + p_{2n}(t)x_n + f_2(t), \\ x_3' &= p_{31}(t)x_1 + p_{32}(t)x_2 + \dots + p_{3n}(t)x_n + f_3(t), \\ \vdots \end{aligned}$$
(7)

 $x'_n = p_{n1}(t)x_1 + p_{n2}(t)x_2 + \dots + p_{nn}(t)x_n + f_n(t).$

If we introduce the *coefficient matrix*

$$\mathbf{P}(t) = \left[p_{ij}(t) \right]$$

and the column vectors

 \succ

$$\mathbf{x} = \begin{bmatrix} x_i \end{bmatrix}$$
 and $\mathbf{f}(t) = \begin{bmatrix} f_i(t) \end{bmatrix}$,

then—as illustrated in Example 2 below—the system in (7) takes the form of a single matrix equation

$$\frac{d\mathbf{x}}{dt} = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t). \tag{8}$$

We will see that the general theory of the linear system in (7) closely parallels that of a single *n*th-order equation. The matrix notation used in Eq. (8) not only emphasizes this analogy, but also saves a great deal of space.

A solution of Eq. (8) on the open interval *I* is a column vector function $\mathbf{x}(t) = [x_i(t)]$ such that the component functions of \mathbf{x} satisfy the system in (7) identically on *I*. If the functions $p_{ij}(t)$ and $f_i(t)$ are all continuous on *I*, then Theorem 1 of Section 7.1 guarantees the existence on *I* of a unique solution $\mathbf{x}(t)$ satisfying preassigned initial conditions $\mathbf{x}(a) = \mathbf{b}$.

Example 2 The first-order system

$$\begin{aligned} x_1' &= 4x_1 - 3x_2, \\ x_2' &= 6x_1 - 7x_2 \end{aligned}$$

can be recast as a matrix equation by writing

$$\mathbf{x}' = \begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 4x_1 - 3x_2 \\ 6x_1 - 7x_2 \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 6 & -7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 6 & -7 \end{bmatrix} \mathbf{x}$$

Thus we obtain a matrix equation of the form in (8),

$$\frac{d\mathbf{x}}{dt} = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t) \quad \text{with} \quad \mathbf{P}(t) = \begin{bmatrix} 4 & -3\\ 6 & -7 \end{bmatrix} \quad \text{and} \quad \mathbf{f}(t) = \begin{bmatrix} 0\\ 0 \end{bmatrix} = \mathbf{0}$$

To verify that the vector functions

$$\mathbf{x}_1(t) = \begin{bmatrix} 3e^{2t} \\ 2e^{2t} \end{bmatrix}$$
 and $\mathbf{x}_2(t) = \begin{bmatrix} e^{-5t} \\ 3e^{-5t} \end{bmatrix}$

are both solutions of the matrix differential equation with coefficient matrix **P**, we need only calculate

$$\mathbf{P}\mathbf{x}_1 = \begin{bmatrix} 4 & -3 \\ 6 & -7 \end{bmatrix} \begin{bmatrix} 3e^{2t} \\ 2e^{2t} \end{bmatrix} = \begin{bmatrix} 6e^{2t} \\ 4e^{2t} \end{bmatrix} = \mathbf{x}_1'$$

and

>

 $\mathbf{P}\mathbf{x}_2 = \begin{bmatrix} 4 & -3\\ 6 & -7 \end{bmatrix} \begin{bmatrix} e^{-5t}\\ 3e^{-5t} \end{bmatrix} = \begin{bmatrix} -5e^{-5t}\\ -15e^{-5t} \end{bmatrix} = \mathbf{x}'_2.$

To investigate the general nature of the solutions of Eq. (8), we consider first the **associated homogeneous equation**

 \overline{d}

$$\frac{\mathbf{x}}{t} = \mathbf{P}(t)\mathbf{x},\tag{9}$$

which has the form shown in Eq. (8), but with $\mathbf{f}(t) \equiv \mathbf{0}$. We expect it to have *n* solutions $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ that are independent in some appropriate sense, and that are such that every solution of Eq. (9) is a linear combination of these *n* particular solutions. Given *n* solutions $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ of Eq. (9), let us write

$$\mathbf{x}_{j}(t) = \begin{bmatrix} x_{1j}(t) \\ \vdots \\ x_{ij}(t) \\ \vdots \\ x_{ni}(t) \end{bmatrix}.$$
 (10)

Thus $x_{ij}(t)$ denotes the *i*th component of the vector $\mathbf{x}_j(t)$, so the second subscript refers to the vector function $\mathbf{x}_j(t)$, whereas the first subscript refers to a component of this function. Theorem 1 is analogous to Theorem 1 of Section 5.2.

THEOREM 1 Principle of Superposition

Let $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ be *n* solutions of the homogeneous linear equation in (9) on the open interval *I*. If c_1, c_2, \ldots, c_n are constants, then the linear combination

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$$
(11)

is also a solution of Eq. (9) on *I*.

Proof: We know that $\mathbf{x}'_i = \mathbf{P}(t)\mathbf{x}_i$ for each $i \ (1 \le i \le n)$, so it follows immediately that

$$\mathbf{x}' = c_1 \mathbf{x}'_1 + c_2 \mathbf{x}'_2 + \dots + c_n \mathbf{x}'_n$$

= $c_1 \mathbf{P}(t) \mathbf{x}_1 + c_2 \mathbf{P}(t) \mathbf{x}_2 + \dots + c_n \mathbf{P}(t) \mathbf{x}_n$
= $\mathbf{P}(t) (c_1 \mathbf{x}_1 + c_2 \mathbf{x}_2 + \dots + c_n \mathbf{x}_n).$

That is, $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$, as desired. The remarkable simplicity of this proof demonstrates clearly one advantage of matrix notation.

Example 2 Continued If \mathbf{x}_1 and \mathbf{x}_2 are the two solutions of

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} 4 & -3\\ 6 & -7 \end{bmatrix} \mathbf{x}$$

discussed in Example 2, then the linear combination

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) = c_1 \begin{bmatrix} 3e^{2t} \\ 2e^{2t} \end{bmatrix} + c_2 \begin{bmatrix} e^{-5t} \\ 3e^{-5t} \end{bmatrix}$$

is also a solution. In scalar form with $\mathbf{x} = \begin{bmatrix} x_1 & x_2 \end{bmatrix}^T$, this gives the solution

$$x_1(t) = 3c_1e^{2t} + c_2e^{-5t},$$

$$x_2(t) = 2c_1e^{2t} + 3c_2e^{-5t}.$$

Independence and General Solutions

Linear independence is defined in the same way for vector-valued functions as for real-valued functions (Section 5.2). The vector-valued functions $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ are **linearly dependent** on the interval *I* provided that there exist constants c_1, c_2, \ldots, c_n , not all zero, such that

$$c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t) = \mathbf{0}$$

$$\tag{12}$$

for all *t* in *I*. Otherwise, they are **linearly independent**. Equivalently, they are linearly independent provided that no one of them is a linear combination of the others. For instance, the two solutions \mathbf{x}_1 and \mathbf{x}_2 of Example 2 are linearly independent because, clearly, neither is a scalar multiple of the other.

Just as in the case of a single *n*th-order equation, there is a Wronskian determinant that tells us whether or not *n* given solutions of the homogeneous equation in (9) are linearly dependent. If $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ are such solutions, then their **Wronskian** is the $n \times n$ determinant

$$W(t) = \begin{vmatrix} x_{11}(t) & x_{12}(t) & \cdots & x_{1n}(t) \\ x_{21}(t) & x_{22}(t) & \cdots & x_{2n}(t) \\ \vdots & \vdots & & \vdots \\ x_{n1}(t) & x_{n2}(t) & \cdots & x_{nn}(t) \end{vmatrix},$$
(13)

using the notation in (10) for the components of the solutions. We may write either W(t) or $W(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_n)$. Note that W is the determinant of the matrix that has as its *column* vectors the solutions $\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_n$. Theorem 2 is analogous to Theorem 3 of Section 5.2. Moreover, its proof is essentially the same, with the definition of $W(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_n)$ in Eq. (13) substituted for the definition of the Wronskian of *n* solutions of a single *n*th-order equation. (See Problems 34 through 36.)

THEOREM 2 Wronskians of Solutions

Suppose that $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are *n* solutions of the homogeneous linear equation $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ on an open interval *I*. Suppose also that $\mathbf{P}(t)$ is continuous on *I*. Let

$$W = W(\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n).$$

Then

- If $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are linearly dependent on *I*, then W = 0 at every point of *I*.
- If x₁, x₂,..., x_n are linearly independent on *I*, then W ≠ 0 at each point of *I*.

Thus there are only two possibilities for solutions of homogeneous systems: Either W = 0 at *every* point of *I*, or W = 0 at *no* point of *I*.

Example 3

It is readily verified (as in Example 2) that

$$\mathbf{x}_{1}(t) = \begin{bmatrix} 2e^{t} \\ 2e^{t} \\ e^{t} \end{bmatrix}, \quad \mathbf{x}_{2}(t) = \begin{bmatrix} 2e^{3t} \\ 0 \\ -e^{3t} \end{bmatrix}, \quad \text{and} \quad \mathbf{x}_{3}(t) = \begin{bmatrix} 2e^{5t} \\ -2e^{5t} \\ e^{5t} \end{bmatrix}$$

are solutions of the equation

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} 3 & -2 & 0\\ -1 & 3 & -2\\ 0 & -1 & 3 \end{bmatrix} \mathbf{x}.$$
 (14)

The Wronskian of these solutions is

$$W = \begin{vmatrix} 2e^t & 2e^{3t} & 2e^{5t} \\ 2e^t & 0 & -2e^{5t} \\ e^t & -e^{3t} & e^{5t} \end{vmatrix} = e^{9t} \begin{vmatrix} 2 & 2 & 2 \\ 2 & 0 & -2 \\ 1 & -1 & 1 \end{vmatrix} = -16e^{9t},$$

which is never zero. Hence Theorem 2 implies that the solutions \mathbf{x}_1 , \mathbf{x}_2 , and \mathbf{x}_3 are linearly independent (on any open interval).

Theorem 3 is analogous to Theorem 4 of Section 5.2. It says that the **general** solution of the *homogeneous* $n \times n$ system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ is a linear combination

$$\mathbf{x} = c_1 \mathbf{x}_1 + c_2 \mathbf{x}_2 + \dots + c_n \mathbf{x}_n \tag{15}$$

of any *n* given linearly independent solutions $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$.

THEOREM 3 General Solutions of Homogeneous Systems

Let $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ be *n* linearly independent solutions of the homogeneous linear equation $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ on an open interval *I* where $\mathbf{P}(t)$ is continuous. If $\mathbf{x}(t)$ is any solution whatsoever of the equation $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ on *I*, then there exist numbers c_1 , c_2, \dots, c_n such that

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$$
(15)

for all t in I.

Proof: Let *a* be a fixed point of *I*. We first show that there exist numbers c_1 , c_2 , ..., c_n such that the solution

$$\mathbf{y}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$$
(16)

has the same initial values at t = a as does the given solution $\mathbf{x}(t)$; that is, such that

$$c_1\mathbf{x}_1(a) + c_2\mathbf{x}_2(a) + \dots + c_n\mathbf{x}_n(a) = \mathbf{x}(a).$$
(17)

Let $\mathbf{X}(t)$ be the $n \times n$ matrix with column vectors $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$, and let \mathbf{c} be the column vector with components c_1, c_2, \ldots, c_n . Then Eq. (17) may be written in the form

$$\mathbf{X}(a)\mathbf{c} = \mathbf{x}(a). \tag{18}$$

The Wronskian determinant $W(a) = |\mathbf{X}(a)|$ is nonzero because the solutions \mathbf{x}_1 , $\mathbf{x}_2, \ldots, \mathbf{x}_n$ are linearly independent. Hence the matrix $\mathbf{X}(a)$ has an inverse matrix $\mathbf{X}(a)^{-1}$. Therefore the vector $\mathbf{c} = \mathbf{X}(a)^{-1}\mathbf{x}(a)$ satisfies Eq. (18), as desired.

Finally, note that the given solution $\mathbf{x}(t)$ and the solution $\mathbf{y}(t)$ of Eq. (16) with the values of c_i determined by the equation $\mathbf{c} = \mathbf{X}(a)^{-1}\mathbf{x}(a)$ —have the same initial values (at t = a). It follows from the existence-uniqueness theorem of Section 7.1 that $\mathbf{x}(t) = \mathbf{y}(t)$ for all t in I. This establishes Eq. (15).

Remark Every $n \times n$ system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ with continuous coefficient matrix does have a set of *n* linearly independent solutions $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ as in the hypotheses of Theorem 3. It suffices to choose for $\mathbf{x}_i(t)$ the unique solution such that

$$\mathbf{x}_{j}(a) = \begin{bmatrix} 0\\0\\0\\\vdots\\0\\1\\0\\\vdots\\0 \end{bmatrix} \leftarrow \text{position } j$$

—that is, the column vector with all elements zero except for a 1 in row j. (In other words, $\mathbf{x}_j(a)$ is merely the *j*th column of the identity matrix.) Then

$$W(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)\Big|_{t=a} = |\mathbf{I}| \neq 0,$$

so the solutions $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are linearly independent by Theorem 2. How to actually find these solutions explicitly is another matter—one that we address in Section 7.3 (for the case of constant-coefficient matrices).

Initial Value Problems and Elementary Row Operations

The general solution in Eq. (15) of the homogeneous linear system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ can be written in the form

$$\mathbf{x}(t) = \mathbf{X}(t)\mathbf{c},\tag{19}$$

where

$$\mathbf{X}(t) = \begin{bmatrix} \mathbf{x}_1(t) & \mathbf{x}_2(t) & \cdots & \mathbf{x}_n(t) \end{bmatrix}$$
(20)

is the $n \times n$ matrix whose *column vectors* are the linearly independent solutions \mathbf{x}_1 , $\mathbf{x}_2, \ldots, \mathbf{x}_n$, and where $\mathbf{c} = \begin{bmatrix} c_1 & c_2 & \cdots & c_n \end{bmatrix}^T$ is the vector of coefficients in the linear combination

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t).$$
(15)

7.2 MATRICES AND LINEAR SYSTEMS 381

Suppose now that we wish to solve the *initial value problem*

$$\frac{d\mathbf{x}}{dt} = \mathbf{P}\mathbf{x}, \quad \mathbf{x}(a) = \mathbf{b}, \tag{21}$$

where the initial vector $\mathbf{b} = \begin{bmatrix} b_1 & b_2 & \cdots & b_n \end{bmatrix}^T$ is given. Then, according to Eq. (19), it suffices to solve the system

$$\mathbf{X}(a)\mathbf{c} = \mathbf{b} \tag{22}$$

to find the coefficients c_1, c_2, \ldots, c_n in Eq. (15). For instance, we can use the row-reduction techniques of Sections 3.2 and 3.3.

Example 4 Use the solution vectors given in Example 3 to solve the initial value problem

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} 3 & -2 & 0\\ -1 & 3 & -2\\ 0 & -1 & 3 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 0\\ 2\\ 6 \end{bmatrix}.$$
(23)

Solution It follows from Theorem 3 that the linear combination

>

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + c_3 \mathbf{x}_3(t)$$

$$= c_1 \begin{bmatrix} 2e^t \\ 2e^t \\ e^t \end{bmatrix} + c_2 \begin{bmatrix} 2e^{3t} \\ 0 \\ -e^{3t} \end{bmatrix} + c_3 \begin{bmatrix} 2e^{5t} \\ -2e^{5t} \\ e^{5t} \end{bmatrix}$$

is a general solution of the 3×3 linear system in (23). In scalar form, this gives the general solution

$$\begin{aligned} x_1(t) &= 2c_1e^t + 2c_2e^{3t} + 2c_3e^{5t}, \\ x_2(t) &= 2c_1e^t - 2c_3e^{5t}, \\ x_3(t) &= c_1e^t - c_2e^{3t} + c_3e^{5t}. \end{aligned}$$

We seek the particular solution satisfying the initial conditions

$$x_1(0) = 0$$
, $x_2(0) = 2$, $x_3(0) = 6$.

When we substitute these values in the preceding three scalar equations, we get the algebraic linear system

$$2c_1 + 2c_2 + 2c_3 = 0,$$

$$2c_1 - 2c_3 = 2,$$

$$c_1 - c_2 + c_3 = 6$$

having the augmented coefficient matrix

$$\begin{bmatrix} 2 & 2 & 2 & 0 \\ 2 & 0 & -2 & 2 \\ 1 & -1 & 1 & 6 \end{bmatrix}.$$

Multiplication of each of the first two rows by $\frac{1}{2}$ gives

1	1	1	0	
1	0	-1	1	;
1	-1	1	6	

then subtraction of the first row both from the second row and from the third row gives the matrix

1	1	1	0	
0	-1	-2	1	
0	-2	0	6	

The first column of this matrix now has the desired form.
>

 $\mathbf{>}$

Now, we multiply the second row by -1, then add twice the result to the third row. Thereby, we get the upper triangular augmented coefficient matrix

$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 2 & -1 \\ 0 & 0 & 4 & 4 \end{bmatrix}$$

that corresponds to the transformed system

$$c_1 + c_2 + c_3 = 0,$$

 $c_2 + 2c_3 = -1,$
 $4c_3 = 4.$

We finally solve in turn for $c_3 = 1$, $c_2 = -3$, and $c_1 = 2$. Thus the desired particular solution is given by

$$\mathbf{x}(t) = 2\mathbf{x}_1(t) - 3\mathbf{x}_2(t) + \mathbf{x}_3(t) = \begin{bmatrix} 4e^t - 6e^{3t} + 2e^{5t} \\ 4e^t & -2e^{5t} \\ 2e^t + 3e^{3t} + e^{5t} \end{bmatrix}.$$

Nonhomogeneous Solutions

We finally turn our attention to a *nonhomogeneous* linear system of the form

$$\frac{d\mathbf{x}}{dt} = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t).$$
(24)

The following theorem is analogous to Theorem 5 of Section 5.2 and is proved in precisely the same way, substituting the preceding theorems in this section for the analogous theorems of Section 5.2. In brief, Theorem 4 means that the general solution of Eq. (24) has the form

$$\mathbf{x}(t) = \mathbf{x}_c(t) + \mathbf{x}_p(t), \tag{25}$$

where $\mathbf{x}_p(t)$ is a single particular solution of Eq. (24) and the **complementary func**tion $\mathbf{x}_c(t)$ is a general solution of the associated homogeneous equation $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$.

THEOREM 4 Solutions of Nonhomogeneous Systems

Let \mathbf{x}_p be a particular solution of the nonhomogeneous linear equation in (24) on an open interval *I* on which the functions $\mathbf{P}(t)$ and $\mathbf{f}(t)$ are continuous. Let $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$ be linearly independent solutions of the associated homogeneous equation on *I*. If $\mathbf{x}(t)$ is any solution whatsoever of Eq. (24) on *I*, then there exist numbers c_1, c_2, \ldots, c_n such that

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t) + \mathbf{x}_p(t)$$
(26)

for all t in I.

Thus, finding a general solution of a homogeneous linear system involves two separate steps:

- 1. Finding the general solution $\mathbf{x}_{c}(t)$ of the associated homogeneous system;
- 2. Finding a single particular solution $\mathbf{x}_p(t)$ of the nonhomogeneous system.

The sum $\mathbf{x}(t) = \mathbf{x}_c(t) + \mathbf{x}_p(t)$ will then be a general solution of the nonhomogeneous system.

Example 5 The nonhomogeneous linear system

$$\begin{aligned} x_1' &= 3x_1 - 2x_2 &- 9t + 13, \\ x_2' &= -x_1 + 3x_2 - 2x_3 + 7t - 15, \\ x_3' &= -x_2 + 3x_3 - 6t + 7 \end{aligned}$$

can be recast as a matrix equation by writing

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} x_1' \\ x_2' \\ x_3' \end{bmatrix} = \begin{bmatrix} 3x_1 - 2x_2 & -9t + 13 \\ -x_1 + 3x_2 - 2x_3 + 7t - 15 \\ -x_2 + 3x_3 - 6t + 7 \end{bmatrix}$$
$$= \begin{bmatrix} 3x_1 - 2x_2 \\ -x_1 + 3x_2 - 2x_3 \\ -x_2 + 3x_3 \end{bmatrix} + \begin{bmatrix} -9t + 13 \\ 7t - 15 \\ -6t + 7 \end{bmatrix}$$
$$= \begin{bmatrix} 3 & -2 & 0 \\ -1 & 3 & -2 \\ 0 & -1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + \begin{bmatrix} -9t + 13 \\ 7t - 15 \\ -6t + 7 \end{bmatrix}.$$

Thus we get the form in (24) with

$$\mathbf{P}(t) = \begin{bmatrix} 3 & -2 & 0 \\ -1 & 3 & -2 \\ 0 & -1 & 3 \end{bmatrix}, \quad \mathbf{f}(t) = \begin{bmatrix} -9t + 13 \\ 7t - 15 \\ -6t + 7 \end{bmatrix}.$$

In Example 3 we saw that a general solution of the associated homogeneous linear system

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} 3 & -2 & 0\\ -1 & 3 & -2\\ 0 & -1 & 3 \end{bmatrix} \mathbf{x}$$

is given by

$$\mathbf{x}_{c}(t) = \begin{bmatrix} 2c_{1}e^{t} + 2c_{2}e^{3t} + 2c_{3}e^{5t} \\ 2c_{1}e^{t} & -2c_{3}e^{5t} \\ c_{1}e^{t} - c_{2}e^{3t} + c_{3}e^{5t} \end{bmatrix},$$

and we can verify, by substitution, that the function

$$\mathbf{x}_p(t) = \begin{bmatrix} 3t\\5\\2t \end{bmatrix}$$

(found by using a computer algebra system, or perhaps by a human being using a method discussed in Section 8.2) is a particular solution of the original nonhomogeneous system. Consequently, Theorem 4 implies that a general solution of the nonhomogeneous system is given by

$$\mathbf{x}(t) = \mathbf{x}_{c}(t) + \mathbf{x}_{p}(t)$$

-that is, by

$$x_1(t) = 2c_1e^t + 2c_2e^{3t} + 2c_3e^{5t} + 3t,$$

$$x_2(t) = 2c_1e^t - 2c_3e^{5t} + 5,$$

$$x_3(t) = c_1e^t - c_2e^{3t} + c_3e^{5t} + 2t.$$

7.2 Problems

In Problems 1 and 2, verify the product law for differentiation, (AB)' = A'B + AB'.

1.
$$\mathbf{A}(t) = \begin{bmatrix} t & 2t-1 \\ t^3 & \frac{1}{t} \end{bmatrix} \text{ and } \mathbf{B}(t) = \begin{bmatrix} 1-t & 1+t \\ 3t^2 & 4t^3 \end{bmatrix}.$$

2.
$$\mathbf{A}(t) = \begin{bmatrix} e^t & t & t^2 \\ -t & 0 & 2 \\ 8t & -1 & t^3 \end{bmatrix} \text{ and } \mathbf{B}(t) = \begin{bmatrix} 3 \\ 2e^{-t} \\ 3t \end{bmatrix}.$$

In Problems 3 through 12, write the given system in the form $\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t)$.

3. x' = -3y, y' = 3x4. x' = 3x - 2y, y' = 2x + y5. $x' = 2x + 4y + 3e^{t}, y' = 5x - y - t^{2}$ 6. $x' = tx - e^{t}y + \cos t, y' = e^{-t}x + t^{2}y - \sin t$ 7. x' = y + z, y' = z + x, z' = x + y8. x' = 2x - 3y, y' = x + y + 2z, z' = 5y - 7z9. $x' = 3x - 4y + z + t, y' = x - 3z + t^{2}, z' = 6y - 7z + t^{3}$ 10. $x' = tx - y + e^{t}z, y' = 2x + t^{2}y - z, z' = e^{-t}x + 3ty + t^{3}z$ 11. $x'_{1} = x_{2}, x'_{2} = 2x_{3}, x'_{3} = 3x_{4}, x'_{4} = 4x_{1}$ 12. $x'_{1} = x_{2} + x_{3} + 1, x'_{2} = x_{3} + x_{4} + t$.

$$x'_{1} = x_{2} + x_{3} + t, x'_{2} = x_{3} + x_{4} + t, x'_{3} = x_{1} + x_{4} + t^{2}, x'_{4} = x_{1} + x_{2} + t^{3}$$

In Problems 13 through 22, first verify that the given vectors are solutions of the given system. Then use the Wronskian to show that they are linearly independent. Finally, write the general solution of the system.

13.
$$\mathbf{x}' = \begin{bmatrix} 4 & 2 \\ -3 & -1 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = \begin{bmatrix} 2e^t \\ -3e^t \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} e^{2t} \\ -e^{2t} \end{bmatrix}$$

14. $\mathbf{x}' = \begin{bmatrix} -3 & 2 \\ -3 & 4 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = \begin{bmatrix} e^{3t} \\ 3e^{3t} \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} 2e^{-2t} \\ e^{-2t} \end{bmatrix}$
15. $\mathbf{x}' = \begin{bmatrix} 3 & -1 \\ 5 & -3 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = e^{2t} \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{x}_2 = e^{-2t} \begin{bmatrix} 1 \\ 5 \end{bmatrix}$
16. $\mathbf{x}' = \begin{bmatrix} 4 & 1 \\ -2 & 1 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = e^{3t} \begin{bmatrix} -1 \\ -1 \end{bmatrix}, \mathbf{x}_2 = e^{2t} \begin{bmatrix} -1 \\ -2 \end{bmatrix}$
17. $\mathbf{x}' = \begin{bmatrix} 4 & -3 \\ 6 & -7 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = \begin{bmatrix} 3e^{2t} \\ 2e^{2t} \end{bmatrix}, \mathbf{x}_2 = \begin{bmatrix} e^{-5t} \\ 3e^{-5t} \end{bmatrix}$
18. $\mathbf{x}' = \begin{bmatrix} 3 & -2 & 0 \\ -1 & 3 & -2 \\ 0 & -1 & 3 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = e^t \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$
19. $\mathbf{x}' = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \mathbf{x}; \mathbf{x}_1 = e^{2t} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{x}_2 = e^{-t} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \mathbf{x}_3 = e^{-t} \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$



In Problems 23 through 32, find a particular solution of the indicated linear system that satisfies the given initial conditions.

- **23.** The system of Problem 14: $x_1(0) = 0, x_2(0) = 5$
- **24.** The system of Problem 15: $x_1(0) = 5$, $x_2(0) = -3$
- **25.** The system of Problem 16: $x_1(0) = 11, x_2(0) = -7$
- **26.** The system of Problem 17: $x_1(0) = 8$, $x_2(0) = 0$
- **27.** The system of Problem 18: $x_1(0) = 0$, $x_2(0) = 0$, $x_3(0) = 4$
- **28.** The system of Problem 19: $x_1(0) = 10, x_2(0) = 12, x_3(0) = -1$
- **29.** The system of Problem 21: $x_1(0) = 1$, $x_2(0) = 2$, $x_3(0) = 3$
- **30.** The system of Problem 21: $x_1(0) = 5$, $x_2(0) = -7$, $x_3(0) = 11$
- **31.** The system of Problem 22: $x_1(0) = x_2(0) = x_3(0) = x_4(0) = 1$
- **32.** The system of Problem 22: $x_1(0) = 1$, $x_2(0) = 3$, $x_3(0) = 4$, $x_4(0) = 7$
- 33. (a) Show that the vector functions

$$\mathbf{x}_1(t) = \begin{bmatrix} t \\ t^2 \end{bmatrix}$$
 and $\mathbf{x}_2 = \begin{bmatrix} t^2 \\ t^3 \end{bmatrix}$

are linearly independent on the real line. (b) Why does it follow from Theorem 2 that there is *no* continuous matrix P(t) such that x₁ and x₂ are both solutions of x' = P(t)x?
34. Suppose that one of the vector functions

$$\mathbf{x}_1(t) = \begin{bmatrix} x_{11}(t) \\ x_{21}(t) \end{bmatrix} \text{ and } \mathbf{x}_2(t) = \begin{bmatrix} x_{12}(t) \\ x_{22}(t) \end{bmatrix}$$

is a constant multiple of the other on the open interval *I*. Show that their Wronskian $W(t) = |[x_{ij}(t)]|$ must vanish identically on *I*. This proves part (a) of Theorem 2 in the case n = 2.

7.3 THE EIGENVALUE METHOD FOR LINEAR SYSTEMS 385

35. Suppose that the vectors $\mathbf{x}_1(t)$ and $\mathbf{x}_2(t)$ of Problem 34 are solutions of the equation $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$, where the 2 × 2 matrix $\mathbf{P}(t)$ is continuous on the open interval *I*. Show that if there exists a point *a* of *I* at which their Wronskian W(a) is zero, then there exist numbers c_1 and c_2 not both zero such that $c_1\mathbf{x}_1(a) + c_2\mathbf{x}_2(a) = \mathbf{0}$. Then conclude from the uniqueness of solutions of the equation $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ that

$$c_1\mathbf{x}_1(t) + c_2\mathbf{x}_2(t) = \mathbf{0}$$

for all *t* in *I*; that is, that \mathbf{x}_1 and \mathbf{x}_2 are linearly dependent. This proves part (b) of Theorem 2 in the case n = 2.

- **36.** Generalize Problems 34 and 35 to prove Theorem 2 for *n* an arbitrary positive integer.
- **37.** Let $\mathbf{x}_1(t)$, $\mathbf{x}_2(t)$, ..., $\mathbf{x}_n(t)$ be vector functions whose *i*th components (for some fixed *i*) $x_{i1}(t)$, $x_{i2}(t)$, ..., $x_{in}(t)$ are linearly independent real-valued functions. Conclude that the vector functions are themselves linearly independent.

7.3 The Eigenvalue Method for Linear Systems

We now introduce a powerful method for constructing the general solution of a *homogeneous* first-order system with *constant* coefficients,

$$\begin{aligned} x_1' &= a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n, \\ x_2' &= a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n, \\ \vdots \\ x_n' &= a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n. \end{aligned}$$
(1)

By Theorem 3 of Section 7.2, we know that it suffices to find *n* linearly independent solution vectors $\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_n$; the linear combination

 $\mathbf{x}(t) = c_1 \mathbf{x}_1 + c_2 \mathbf{x}_2 + \dots + c_n \mathbf{x}_n \tag{2}$

with arbitrary coefficients will then be a general solution of the system in (1).

To search for the n needed linearly independent solution vectors, we proceed by analogy with the characteristic root method for solving a single homogeneous equation with constant coefficients (Section 5.3). It is reasonable to anticipate solution vectors of the form

$$\mathbf{x}(t) = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} v_1 e^{\lambda t} \\ v_2 e^{\lambda t} \\ v_3 e^{\lambda t} \\ \vdots \\ v_n e^{\lambda t} \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ \vdots \\ v_n \end{bmatrix} e^{\lambda t} = \mathbf{v} e^{\lambda t},$$
(3)

where λ , v_1 , v_2 , v_3 , ..., v_n are appropriate scalar constants. For if we substitute

$$x_i = v_i e^{\lambda t}, \qquad x'_i = \lambda v_i e^{\lambda t} \qquad (i = 1, 2, \dots, n)$$

in (1), then each term in the resulting equations will have the factor $e^{\lambda t}$, so we can cancel it throughout. This will leave us with *n* linear equations that—for appropriate values of λ —we can hope to solve for values of the coefficients v_1, v_2, \ldots, v_n in Eq. (3), so that $\mathbf{x}(t) = \mathbf{v}e^{\lambda t}$ is, indeed, a solution of the system in (1).

To investigate this possibility, it is more efficient to write the system in (1) in the matrix form

$$\mathbf{x}' = \mathbf{A}\mathbf{x},\tag{4}$$

where $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$. When we substitute the trial solution $\mathbf{x} = \mathbf{v}e^{\lambda t}$ (having derivative $\mathbf{x}' = \lambda \mathbf{v}e^{\lambda t}$) in Eq. (4), the result is

$$\lambda \mathbf{v} e^{\lambda t} = \mathbf{A} \mathbf{v} e^{\lambda t}$$

>

We cancel the nonzero scalar factor $e^{\lambda t}$ to get

$$\mathbf{A}\mathbf{v} = \lambda \mathbf{v}.\tag{5}$$

Recall, from Section 6.1, that Eq. (5) means that $\mathbf{v} \neq \mathbf{0}$ is an *eigenvector* of the matrix **A** associated with the *eigenvalue* λ . Our discussion of Eqs. (3)–(5) therefore provides a proof of the following theorem, which is the basis for the **eigenvalue method** of solving a first-order system with constant coefficients.

THEOREM 1 Eigenvalue Solutions of x' = Ax

Let λ be an eigenvalue of the [constant] coefficient matrix **A** of the first-order linear system

 $\frac{d\mathbf{x}}{dt} = \mathbf{A}\mathbf{x}.$

If **v** is an eigenvector associated with λ , then

 $\mathbf{x}(t) = \mathbf{v}e^{\lambda t}$

is a nontrivial solution of the system.

Recall that an eigenvalue λ of the matrix **A** is a solution of the *characteristic* equation

$$\mathbf{A} - \lambda \mathbf{I} = \det(\mathbf{A} - \lambda \mathbf{I}) = 0 \tag{6}$$

and that an eigenvector **v** associated with λ is then a solution of the *eigenvector* equation

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}.\tag{7}$$

The Eigenvalue Method

In outline, this method for solving the $n \times n$ homogeneous constant-coefficient system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ proceeds as follows:

- 1. First, we solve the characteristic equation in (6) for the eigenvalues λ_1 , λ_2 , ..., λ_n of the matrix **A**.
- **2.** Next, we attempt to find *n* linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ associated with these eigenvalues.
- **3.** Step 2 is not always possible, but, when it is, we get *n* linearly independent solutions

$$\mathbf{x}_1(t) = \mathbf{v}_1 e^{\lambda_1 t}, \quad \mathbf{x}_2(t) = \mathbf{v}_2 e^{\lambda_2 t}, \quad \dots, \quad \mathbf{x}_n(t) = \mathbf{v}_n e^{\lambda_n t}.$$
(8)

In this case the general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is a linear combination

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$$

of these n solutions.

We will discuss separately the various cases that can occur, depending on whether the eigenvalues are distinct or repeated, real or complex. The case of repeated eigenvalues—multiple roots of the characteristic equation—will be deferred to Section 7.6.

Distinct Real Eigenvalues

If the eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$ are real and distinct, then we substitute each of them in turn in Eq. (7) and solve for the associated eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$. It then follows from Theorem 2 in Section 6.2 that the particular solutions given in (8) are linearly independent. (In any particular example, such linear independence can always be verified by using the Wronskian determinant of Section 7.2.) The following example illustrates the eigenvalue method.

Example 1 Find a general solution of the system

$$\begin{aligned} x_1' &= 4x_1 + 2x_2, \\ x_2' &= 3x_1 - x_2. \end{aligned}$$
 (9)

Solution The matrix form of the system in (9) is

$$\mathbf{x}' = \begin{bmatrix} 4 & 2\\ 3 & -1 \end{bmatrix} \mathbf{x}.$$
 (10)

The characteristic equation of the coefficient matrix is

$$\begin{vmatrix} 4-\lambda & 2\\ 3 & -1-\lambda \end{vmatrix} = (4-\lambda)(-1-\lambda) - 6\\ = \lambda^2 - 3\lambda - 10 = (\lambda+2)(\lambda-5) = 0,$$

so we have the distinct real eigenvalues $\lambda_1 = -2$ and $\lambda_2 = 5$.

For the coefficient matrix **A** in Eq. (10), the eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ takes the form

$$\begin{bmatrix} 4-\lambda & 2\\ 3 & -1-\lambda \end{bmatrix} \begin{bmatrix} a\\ b \end{bmatrix} = \begin{bmatrix} 0\\ 0 \end{bmatrix}$$
(11)

for the associated eigenvector $\mathbf{v} = \begin{bmatrix} a & b \end{bmatrix}^T$.

CASE 1: $\lambda_1 = -2$. Substitution of the first eigenvalue $\lambda_1 = -2$ in Eq. (11) yields the system

$$\begin{bmatrix} 6 & 2 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

—that is, the two scalar equations

$$6a + 2b = 0,$$

 $3a + b = 0.$ (12)

In contrast with the typical nonsingular (algebraic) linear system that has a unique solution, the homogeneous linear system in (12) is *singular*—the two scalar equations obviously are equivalent (each being a multiple of the other). Therefore, Eq. (12) has infinitely many nonzero solutions—we can choose a arbitrarily (but nonzero) and then solve for b.

Substitution of an eigenvalue λ in the eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ always yields a singular homogeneous linear system, and among its infinity of solutions we generally seek a "simple" solution with small integer values (if possible). Looking at the second equation in (12), the choice a = 1 yields b = -3, and thus

$$\mathbf{v}_1 = \left[\begin{array}{c} 1\\ -3 \end{array} \right]$$

is an eigenvector associated with $\lambda_1 = -2$ (as is any nonzero constant multiple of \mathbf{v}_1).

Remark If, instead of the "simplest" choice a = 1, b = -3, we had made another choice a = c, b = -3c, we would have obtained the eigenvector

$$\mathbf{v}_1 = \begin{bmatrix} c \\ -3c \end{bmatrix} = c \begin{bmatrix} 1 \\ -3 \end{bmatrix}.$$

Because this is a constant multiple of our previous result, any choice we make leads to (a constant multiple of) the same solution

$$\mathbf{x}_1(t) = \begin{bmatrix} 1\\ -3 \end{bmatrix} e^{-2t}.$$

CASE 2: $\lambda_2 = 5$. Substitution of the second eigenvalue $\lambda = 5$ in (11) yields the pair

$$-a + 2b = 0,$$

3a - 6b = 0 (13)

of equivalent scalar equations. With b = 1 in the first equation we get a = 2, so

$$\mathbf{v}_2 = \left[\begin{array}{c} 2\\1 \end{array} \right]$$

is an eigenvector associated with $\lambda_2 = 5$. A different choice—a = 2c, b = c—would merely give a [constant] multiple of \mathbf{v}_2 .

These two eigenvalues and their associated eigenvectors yield the two solutions

$$\mathbf{x}_1(t) = \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t}$$
 and $\mathbf{x}_2(t) = \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{5t}$.

They are linearly independent, because their Wronskian

$$\begin{vmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{vmatrix} = 7e^{3t}$$

is nonzero. Hence a general solution of the system in (10) is

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) = c_1 \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t} + c_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{5t};$$

in scalar form,

$$x_1(t) = c_1 e^{-2t} + 2c_2 e^{5t},$$

$$x_2(t) = -3c_1 e^{-2t} + c_2 e^{5t}.$$

Figure 7.3.1 shows some typical solution curves of the system in (10). We see two families of hyperbolas sharing the same pair of asymptotes: the line $x_1 = 2x_2$ obtained from the general solution with $c_1 = 0$, and the line $x_2 = -3x_1$ obtained with $c_2 = 0$. Given initial values $x_1(0) = b_1, x_2(0) = b_2$, it is apparent from the figure that

- If (b_1, b_2) lies to the right of the line $x_2 = -3x_1$, then $x_1(t)$ and $x_2(t)$ both tend to $+\infty$ as $t \to +\infty$;
- If (b_1, b_2) lies to the left of the line $x_2 = -3x_1$, then $x_1(t)$ and $x_2(t)$ both tend to $-\infty$ as $t \to +\infty$.

Remark As in Example 1, it is convenient when discussing a linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ to use vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ to denote different vector-valued solutions of the system, whereas the *scalars* x_1, x_2, \dots, x_n denote the components of a single vector-valued solution \mathbf{x} .

Compartmental Analysis

Frequently, a complex process or system can be broken down into simpler subsystems or "compartments" that can be analyzed separately. The whole system can then be modeled by describing the interactions between the various compartments. Thus a chemical plant may consist of a succession of separate stages (or even physical compartments) in which various reactants and products combine or are mixed. It may happen that a single differential equation describes each compartment of the system; then the whole physical system is modeled by a system of differential equations.



FIGURE 7.3.1. Direction field and solution curves for the linear system $x'_1 = 4x_1 + 2x_2, x'_2 = 3x_1 - x_2$ of Example 1.

7.3 THE EIGENVALUE METHOD FOR LINEAR SYSTEMS 389

As a simple example of a three-stage system, Fig. 7.3.2 shows three brine tanks containing V_1 , V_2 , and V_3 gallons of brine, respectively. Fresh water flows into tank 1, while mixed brine flows from tank 1 into tank 2, from tank 2 into tank 3, and out of tank 3. Let $x_i(t)$ denote the amount (in pounds) of salt in tank *i* at time *t*, for i = 1, 2, and 3. If each flow rate is *r* gallons per minute, then a simple accounting of salt concentrations, as in Example 2 of Section 7.1, yields the first-order system

where

$$k_i = \frac{r}{V_i}, \quad i = 1, 2, 3.$$
 (15)

Example 2 Three brine tanks If $V_1 = 20$, $V_2 = 40$, $V_3 = 50$, r = 10 (gal/min), and the initial amounts of salt in the three brine tanks, in pounds, are

 $x_1(0) = 15, \quad x_2(0) = x_3(0) = 0,$

Substituting the given numerical values in (14) and (15), we get the initial value problem

find the amount of salt in each tank at time $t \ge 0$.

Solution

$$\mathbf{x}'(t) = \begin{bmatrix} -0.5 & 0.0 & 0.0\\ 0.5 & -0.25 & 0.0\\ 0.0 & 0.25 & -0.2 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 15\\ 0\\ 0 \end{bmatrix}$$
(16)

for the vector $\mathbf{x}(t) = \begin{bmatrix} x_1(t) & x_2(t) & x_3(t) \end{bmatrix}^T$. The simple form of the matrix

$$\mathbf{A} - \lambda \mathbf{I} = \begin{bmatrix} -0.5 - \lambda & 0.0 & 0.0 \\ 0.5 & -0.25 - \lambda & 0.0 \\ 0.0 & 0.25 & -0.2 - \lambda \end{bmatrix}$$
(17)

leads readily to the characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = (-0.5 - \lambda)(-0.25 - \lambda)(-0.2 - \lambda) = 0.$$

Thus, the coefficient matrix **A** in (16) has the distinct eigenvalues $\lambda_1 = -0.5$, $\lambda_2 = -0.25$, and $\lambda_3 = -0.2$ and therefore has three linearly independent eigenvectors.

CASE 1: $\lambda_1 = -0.5$. Substituting $\lambda = -0.5$ in (17), we get the equation

$$\begin{bmatrix} \mathbf{A} + (0.5) \cdot \mathbf{I} \end{bmatrix} \mathbf{v} = \begin{bmatrix} 0.0 & 0.0 & 0.0 \\ 0.5 & 0.25 & 0.0 \\ 0.0 & 0.25 & 0.3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

for the associated eigenvector $\mathbf{v} = \begin{bmatrix} a & b & c \end{bmatrix}^T$. The last two rows, after division by 0.25 and 0.05, respectively, yield the scalar equations

$$2a + b = 0,$$

$$5b + 6c = 0.$$

The second equation is satisfied by b = -6 and c = 5, and then the first equation gives a = 3. Thus the eigenvector

$$\mathbf{v}_1 = \begin{bmatrix} 3 & -6 & 5 \end{bmatrix}^T$$

is associated with the eigenvalue $\lambda_1 = -0.5$.



FIGURE 7.3.2. The three brine tanks of Example 2.

CASE 2:
$$\lambda_2 = -0.25$$
. Substituting $\lambda = -0.25$ in (17), we get the equation

$$\begin{bmatrix} \mathbf{A} + (0.25) \cdot \mathbf{I} \end{bmatrix} \mathbf{v} = \begin{bmatrix} -0.25 & 0 & 0\\ 0.5 & 0 & 0\\ 0 & 0.25 & 0.05 \end{bmatrix} \begin{bmatrix} a\\ b\\ c \end{bmatrix} = \begin{bmatrix} 0\\ 0\\ 0 \end{bmatrix}$$

for the associated eigenvector $\mathbf{v} = \begin{bmatrix} a & b & c \end{bmatrix}^T$. Each of the first two rows implies that a = 0, and division of the third row by 0.05 gives the equation

$$5b + c = 0$$

which is satisfied by b = 1, c = -5. Thus the eigenvector

$$\mathbf{v}_2 = \begin{bmatrix} 0 & 1 & -5 \end{bmatrix}^T$$

is associated with the eigenvalue $\lambda_2 = -0.25$.

CASE 3:
$$\lambda_3 = -0.2$$
. Substituting $\lambda = -0.2$ in (17), we get the equation

	-0.3	0.0	0.0]	$\begin{bmatrix} a \end{bmatrix}$		0	
$\begin{bmatrix} \mathbf{A} + (0.2) \cdot \mathbf{I} \end{bmatrix} \mathbf{v} =$	0.5	-0.05	0.0	b	=	0	
	0.0	0.25	0.0	_ c		0	

for the eigenvector **v**. The first and third rows imply that a = 0, and b = 0, respectively, but the all-zero third column leaves *c* arbitrary (but nonzero). Thus

$$\mathbf{v}_3 = \begin{bmatrix} 0 & 0 & 1 \end{bmatrix}^T$$

is an eigenvector associated with $\lambda_3 = -0.2$.

The general solution

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t} + c_3 \mathbf{v}_3 e^{\lambda_3 t}$$

therefore takes the form

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 3\\-6\\5 \end{bmatrix} e^{(-0.5)t} + c_2 \begin{bmatrix} 0\\1\\-5 \end{bmatrix} e^{(-0.25)t} + c_3 \begin{bmatrix} 0\\0\\1 \end{bmatrix} e^{(-0.2)t}.$$

The resulting scalar equations are

$$\begin{aligned} x_1(t) &= 3c_1 e^{(-0.5)t}, \\ x_2(t) &= -6c_1 e^{(-0.5)t} + c_2 e^{(-0.25)t}, \\ x_3(t) &= 5c_1 e^{(-0.5)t} - 5c_2 e^{(-0.25)t} + c_3 e^{(-0.2)t} \end{aligned}$$

When we impose the initial conditions $x_1(0) = 15$, $x_2(0) = x_3(0) = 0$, we get the equations

$$3c_1 = 15, -6c_1 + c_2 = 0, 5c_1 - 5c_2 + c_3 = 0$$

that are readily solved (in turn) for $c_1 = 5$, $c_2 = 30$, and $c_3 = 125$. Thus, finally, the amounts of salt at time *t* in the three brine tanks are given by

$$\begin{aligned} x_1(t) &= 15e^{(-0.5)t}, \\ x_2(t) &= -30e^{(-0.5)t} + 30e^{(-0.25)t}, \\ x_3(t) &= 25e^{(-0.5)t} - 150e^{(-0.25)t} + 125e^{(-0.2)t}. \end{aligned}$$

Figure 7.3.3 shows the graphs of $x_1(t)$, $x_2(t)$, and $x_3(t)$. As we would expect, tank 1 is rapidly "flushed" by the incoming fresh water, and $x_1(t) \rightarrow 0$ as $t \rightarrow +\infty$. The amounts $x_2(t)$ and $x_3(t)$ of salt in tanks 2 and 3 peak in turn and then approach zero as the whole three-tank system is purged of salt as $t \rightarrow +\infty$.



FIGURE 7.3.3. The salt content functions of Example 2.

Complex Eigenvalues

Even if some of the eigenvalues are complex, so long as they are distinct the method described previously still yields n linearly independent solutions. The only complication is that the eigenvectors associated with complex eigenvalues are ordinarily complex-valued, so we will have complex-valued solutions.

To obtain real-valued solutions, we note that—because we are assuming that the matrix **A** has only real entries—the coefficients in the characteristic equation will all be real. Consequently any complex eigenvalues must appear in complexconjugate pairs. Suppose, then, that $\lambda = p + qi$ and $\overline{\lambda} = p - qi$ are such a pair of eigenvalues. If **v** is an eigenvector associated with λ , so that

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0},$$

then taking complex conjugates in this equation yields

$$(\mathbf{A} - \overline{\lambda}\mathbf{I})\overline{\mathbf{v}} = \mathbf{0}$$

since $\overline{\mathbf{A}} = \mathbf{A}$ and $\overline{\mathbf{I}} = \mathbf{I}$ (these matrices being real) and the conjugate of a complex product is the product of the conjugates of the factors. Thus the conjugate $\overline{\mathbf{v}}$ of \mathbf{v} is an eigenvector associated with $\overline{\lambda}$. Of course, the conjugate of a vector is defined componentwise; if

$$\mathbf{v} = \begin{bmatrix} a_1 + b_1 i \\ a_2 + b_2 i \\ \vdots \\ a_n + b_n i \end{bmatrix} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} i = \mathbf{a} + \mathbf{b}i,$$
(18)

then $\overline{\mathbf{v}} = \mathbf{a} - \mathbf{b}i$. The complex-valued solution associated with λ and \mathbf{v} is then

$$\mathbf{x}(t) = \mathbf{v}e^{\lambda t} = \mathbf{v}e^{(p+qi)t} = (\mathbf{a} + \mathbf{b}i)e^{pt}(\cos qt + i\sin qt)$$

-that is,

$$\mathbf{x}(t) = e^{pt} (\mathbf{a} \cos qt - \mathbf{b} \sin qt) + i e^{pt} (\mathbf{b} \cos qt + \mathbf{a} \sin qt).$$
(19)

Because the real and imaginary parts of a complex-valued solution are also solutions, we thus get the two *real-valued* solutions

$$\mathbf{x}_{1}(t) = \operatorname{Re}[\mathbf{x}(t)] = e^{pt}(\mathbf{a}\cos qt - \mathbf{b}\sin qt),$$

$$\mathbf{x}_{2}(t) = \operatorname{Im}[\mathbf{x}(t)] = e^{pt}(\mathbf{b}\cos qt + \mathbf{a}\sin qt)$$
(20)

associated with the complex conjugate eigenvalues $p \pm qi$. It is easy to check that the same two real-valued solutions result from taking real and imaginary parts of $\overline{\mathbf{v}}e^{\overline{\lambda}t}$. Rather than memorizing the formulas in (20), it is preferable in a specific example to proceed as follows:

- First, find explicitly a single complex-valued solution **x**(*t*) associated with the complex eigenvalue λ;
- Then, find the real and imaginary parts $\mathbf{x}_1(t)$ and $\mathbf{x}_2(t)$, to get two independent real-valued solutions corresponding to the two complex conjugate eigenvalues λ and $\overline{\lambda}$.

Example 3 Find a general solution of the system

$$\frac{dx_1}{dt} = 4x_1 - 3x_2,$$

$$\frac{dx_2}{dt} = 3x_1 + 4x_2.$$
(21)

Solution The coefficient matrix

$$\mathbf{A} = \begin{bmatrix} 4 & -3 \\ 3 & 4 \end{bmatrix}$$

has the characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{vmatrix} 4 - \lambda & -3 \\ 3 & 4 - \lambda \end{vmatrix} = (4 - \lambda)^2 + 9 = 0$$

and hence has the complex conjugate eigenvalues $\lambda = 4 - 3i$ and $\overline{\lambda} = 4 + 3i$. Substituting $\lambda = 4 - 3i$ in the eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$, we get the equation

$$[\mathbf{A} - (4 - 3i) \cdot \mathbf{I}]\mathbf{v} = \begin{bmatrix} 3i & -3 \\ 3 & 3i \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

for an associated eigenvalue $\mathbf{v} = \begin{bmatrix} a & b \end{bmatrix}^T$. Division of each row by 3 yields the two scalar equations

$$ia - b = 0,$$

$$a + ib = 0,$$

each of which is satisfied by a = 1 and b = i. Thus, $\mathbf{v} = \begin{bmatrix} 1 & i \end{bmatrix}^T$ is a complex eigenvector associated with the complex eigenvalue $\lambda = 4 - 3i$.

The corresponding complex-valued solution $\mathbf{x}(t) = \mathbf{v}e^{\lambda t}$ of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is then

$$\mathbf{x}(t) = \begin{bmatrix} 1\\i \end{bmatrix} e^{(4-3i)t} = \begin{bmatrix} 1\\i \end{bmatrix} e^{4t} (\cos 3t - i\sin 3t) = e^{4t} \begin{bmatrix} \cos 3t - i\sin 3t\\i\cos 3t + \sin 3t \end{bmatrix}$$

The real and imaginary parts of $\mathbf{x}(t)$ are the real-valued solutions

$$\mathbf{x}_1(t) = e^{4t} \begin{bmatrix} \cos 3t \\ \sin 3t \end{bmatrix}$$
 and $\mathbf{x}_2(t) = e^{4t} \begin{bmatrix} -\sin 3t \\ \cos 3t \end{bmatrix}$

A real-valued general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is then given by

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) = e^{4t} \begin{bmatrix} c_1 \cos 3t - c_2 \sin 3t \\ c_1 \sin 3t + c_2 \cos 3t \end{bmatrix}.$$

Finally, a general solution of the system in (21) in scalar form is

$$x_1(t) = e^{4t} (c_1 \cos 3t - c_2 \sin 3t),$$

$$x_2(t) = e^{4t} (c_1 \sin 3t + c_2 \cos 3t).$$

Figure 7.3.4 shows some typical solution curves of the system in (21). Each appears to spiral counterclockwise as it emanates from the origin in the x_1x_2 -plane. Actually, because of the factor e^{4t} in the general solution, we see that

• Along each solution curve, the point $(x_1(t), x_2(t))$ approaches the origin as $t \to -\infty$, whereas

• The absolute values of $x_1(t)$ and $x_2(t)$ both increase without bound as $t \to +\infty$.



FIGURE 7.3.4. Direction field and solution curves for the linear system $x'_1 = 4x_1 - 3x_2$, $x'_2 = 3x_1 + 4x_2$ of Example 3.

7.3 THE EIGENVALUE METHOD FOR LINEAR SYSTEMS 393



FIGURE 7.3.5. The three brine tanks of Example 4.

Figure 7.3.5 shows a "closed" system of three brine tanks with volumes V_1 , V_2 , and V_3 . The difference between this system and the "open" system of Fig. 7.3.2 is that now the inflow to tank 1 is the outflow from tank 3. With the same notation as in Example 2, the appropriate modification of Eq. (14) is

$$\frac{dx_1}{dt} = -k_1 x_1 + k_3 x_3,
\frac{dx_2}{dt} = k_1 x_1 - k_2 x_2,
\frac{dx_3}{dt} = k_2 x_2 - k_3 x_3,$$
(22)

where $k_i = r/V_i$ as in (15).

Example 4

Find the amounts $x_1(t)$, $x_2(t)$, and $x_3(t)$ of salt at time t in the three brine tanks of Fig. 7.3.5 if $V_1 = 50$ gal, $V_2 = 25$ gal, $V_3 = 50$ gal, and r = 10 gal/min.

Solution With the given numerical values, (22) takes the form

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} -0.2 & 0 & 0.2\\ 0.2 & -0.4 & 0\\ 0 & 0.4 & -0.2 \end{bmatrix} \mathbf{x},$$
(23)

where $\mathbf{x} = \begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix}^T$ as usual. When we expand the determinant of the matrix

$$\mathbf{A} - \lambda \cdot \mathbf{I} = \begin{bmatrix} -0.2 - \lambda & 0.0 & 0.2 \\ 0.2 & -0.4 - \lambda & 0.0 \\ 0.0 & 0.4 & -0.2 - \lambda \end{bmatrix}$$
(24)

along its first row, we find that the characteristic equation of A is

$$(-0.2 - \lambda)(-0.4 - \lambda)(-0.2 - \lambda) + (0.2)(0.2)(0.4)$$

= $-\lambda^3 - (0.8) \cdot \lambda^2 - (0.2) \cdot \lambda$
= $-\lambda \left[(\lambda + 0.4)^2 + (0.2)^2 \right] = 0.$

Thus **A** has the zero eigenvalue $\lambda_0 = 0$ and the complex conjugate eigenvalues λ , $\overline{\lambda} = -0.4 \pm (0.2)i$. We anticipate one solution corresponding to the zero eigenvalue and two additional linearly independent solutions corresponding to the complex conjugate eigenvalues.

CASE 1: $\lambda_0 = 0$. Substitution of $\lambda = 0$ in Eq. (24) gives the eigenvector equation

$$(\mathbf{A} - 0 \cdot \mathbf{I})\mathbf{v} = \begin{bmatrix} -0.2 & 0.0 & 0.2 \\ 0.2 & -0.4 & 0.0 \\ 0.0 & 0.4 & -0.2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$
(25)

for $\mathbf{v} = \begin{bmatrix} a & b & c \end{bmatrix}^T$. The first row gives a = c and the second row gives a = 2b, so $\mathbf{v}_0 = \begin{bmatrix} 2 & 1 & 2 \end{bmatrix}^T$ is an eigenvector associated with the eigenvalue $\lambda_0 = 0$. The corresponding solution $\mathbf{x}_0(t) = \mathbf{v}_0 e^{\lambda_0 t}$ of Eq. (23) is the constant solution

$$\mathbf{x}_{\mathbf{0}}(t) = \begin{bmatrix} 2\\1\\2 \end{bmatrix}.$$
 (26)

394 CHAPTER 7 LINEAR SYSTEMS OF DIFFERENTIAL EQUATIONS

CASE 2: $\lambda = -0.4 - (0.2)i$. Substitution of $\lambda = -0.4 - (0.2)i$ in Eq. (24) gives the eigenvector equation

$$[\mathbf{A} - (-0.4 - (0.2)i)\mathbf{I}]\mathbf{v} = \begin{bmatrix} 0.2 + (0.2)i & 0.0 & 0.2 \\ 0.2 & (0.2)i & 0.0 \\ 0.0 & 0.4 & 0.2 + (0.2)i \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

The second equation (0.2)a + (0.2)ib = 0 is satisfied by a = 1 and b = i. Then the first equation

$$[0.2 + (0.2)i]a + (0.2)c = 0$$

gives c = -1 - i. Thus, $\mathbf{v} = \begin{bmatrix} 1 & i & (-1 - i) \end{bmatrix}^T$ is a complex eigenvector associated with the complex eigenvalue $\lambda = -0.4 - (0.2)i$.

The corresponding complex-valued solution $\mathbf{x}(t) = \mathbf{v}e^{\lambda t}$ of (23) is

$$\mathbf{x}(t) = \begin{bmatrix} 1 & i & -1-i \end{bmatrix}^T e^{(-0.4-0.2i)t}$$

= $\begin{bmatrix} 1 & i & -1-i \end{bmatrix}^T e^{(-0.4)t} (\cos 0.2t - i \sin 0.2t)$
= $e^{(-0.4)t} \begin{bmatrix} \cos 0.2t - i \sin 0.2t \\ \sin 0.2t + i \cos 0.2t \\ -\cos 0.2t - \sin 0.2t - i \cos 0.2t + i \sin 0.2t \end{bmatrix}$.

The real and imaginary parts of $\mathbf{x}(t)$ are the real-valued solutions

$$\mathbf{x}_{1}(t) = e^{(-0.4)t} \begin{bmatrix} \cos 0.2t \\ \sin 0.2t \\ -\cos 0.2t - \sin 0.2t \end{bmatrix},$$

$$\mathbf{x}_{2}(t) = e^{(-0.4)t} \begin{bmatrix} -\sin 0.2t \\ \cos 0.2t \\ -\cos 0.2t + \sin 0.2t \end{bmatrix}.$$
(27)

The general solution

$$\mathbf{x}(t) = c_0 \mathbf{x}_0(t) + c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t)$$

has scalar components

$$x_{1}(t) = 2c_{0} + e^{(-0.4)t} (c_{1} \cos 0.2t - c_{2} \sin 0.2t),$$

$$x_{2}(t) = c_{0} + e^{(-0.4)t} (c_{1} \sin 0.2t + c_{2} \cos 0.2t),$$

$$x_{3}(t) = 2c_{0} + e^{(-0.4)t} [(-c_{1} - c_{2}) \cos 0.2t + (-c_{1} + c_{2}) \sin 0.2t]$$
(28)

giving the amounts of salt in the three tanks at time t. Observe that

$$x_1(t) + x_2(t) + x_3(t) \equiv 5c_0.$$
⁽²⁹⁾

Of course, the total amount of salt in the closed system is constant; the constant c_0 in (29) is one-fifth the total amount of salt. Because of the factors of $e^{(-0.4)t}$ in (28), we see that

$$\lim_{t \to \infty} x_1(t) = 2c_0, \quad \lim_{t \to \infty} x_2(t) = c_0, \quad \text{and} \quad \lim_{t \to \infty} x_3(t) = 2c_0$$

Thus, as $t \to +\infty$ the salt in the system approaches a *steady-state* distribution with 40% of the salt in each of the two 50-gallon tanks and 20% in the 25-gallon tank. So whatever the initial distribution of salt among the three tanks, the limiting distribution is one of uniform concentration throughout the system. Figure 7.3.6 shows the graphs of the three solution functions with $c_0 = 10$, $c_1 = 30$, and $c_2 = -10$, in which case

$$x_1(0) = 50$$
 and $x_2(0) = x_3(0) = 0$.



FIGURE 7.3.6. The salt content functions of Example 4.

7.3 Problems

In Problems 1 through 16, apply the eigenvalue method of this section to find a general solution of the given system. If initial values are given, find also the corresponding particular solution. For each problem, use a computer system or graphing calculator to construct a direction field and typical solution curves for the given system.

1.
$$x'_1 = x_1 + 2x_2$$
, $x'_2 = 2x_1 + x_2$
2. $x'_1 = 2x_1 + 3x_2$, $x'_2 = 2x_1 + x_2$
3. $x'_1 = 3x_1 + 4x_2$, $x'_2 = 3x_1 + 2x_2$; $x_1(0) = x_2(0) = 1$
4. $x'_1 = 4x_1 + x_2$, $x'_2 = 6x_1 - x_2$
5. $x'_1 = 6x_1 - 7x_2$, $x'_2 = x_1 - 2x_2$
6. $x'_1 = 9x_1 + 5x_2$, $x'_2 = -6x_1 - 2x_2$; $x_1(0) = 1$, $x_2(0) = 0$
7. $x'_1 = -3x_1 + 4x_2$, $x'_2 = 6x_1 - 5x_2$
8. $x'_1 = x_1 - 5x_2$, $x'_2 = x_1 - x_2$
9. $x'_1 = 2x_1 - 5x_2$, $x'_2 = 4x_1 - 2x_2$; $x_1(0) = 2$, $x_2(0) = 3$
10. $x'_1 = -3x_1 - 2x_2$, $x'_2 = 9x_1 + 3x_2$
11. $x'_1 = x_1 - 2x_2$, $x'_2 = 2x_1 + x_2$; $x_1(0) = 0$, $x_2(0) = 4$
12. $x'_1 = x_1 - 5x_2$, $x'_2 = 2x_1 - x_2$
13. $x'_1 = 5x_1 - 9x_2$, $x'_2 = 4x_1 + 3x_2$
15. $x'_1 = 7x_1 - 5x_2$, $x'_2 = 4x_1 + 3x_2$
16. $x'_1 = -50x_1 + 20x_2$, $x'_2 = 100x_1 - 60x_2$

In Problems 17 through 25, the eigenvalues of the coefficient matrix can be found by inspection and factoring. Apply the eigenvalue method to find a general solution of each system.

- **17.** $x'_1 = 4x_1 + x_2 + 4x_3$, $x'_2 = x_1 + 7x_2 + x_3$, $x'_3 = 4x_1 + x_2 + 4x_3$
- **18.** $x'_1 = x_1 + 2x_2 + 2x_3, \ x'_2 = 2x_1 + 7x_2 + x_3,$ $x'_3 = 2x_1 + x_2 + 7x_3$
- **19.** $x_1^{\overline{i}} = 4x_1 + x_2 + x_3$, $x_2' = x_1 + 4x_2 + x_3$, $x_3' = x_1 + 4x_2 + x_3$ $x_1 + x_2 + 4x_3$
- **20.** $x'_1 = 5x_1 + x_2 + 3x_3, \ x'_2 = x_1 + 7x_2 + x_3,$
- $x'_{3} = 3x_{1} + x_{2} + 5x_{3}$ **21.** $x'_{1} = 5x_{1} 6x_{3}, x'_{2} = 2x_{1} x_{2} 2x_{3}, x'_{3} = 4x_{1} 2x_{2} 2x_{3}$ $4x_3$
- **22.** $x'_1 = 3x_1 + 2x_2 + 2x_3, x'_2 = -5x_1 4x_2 2x_3, x'_3 =$ $5x_1 + 5x_2 + 3x_3$
- **23.** $x'_1 = 3x_1 + x_2 + x_3$, $x'_2 = -5x_1 3x_2 x_3$, $x'_3 = 5x_1 + 5x_2 + 3x_3$
- **24.** $x'_1 = 2x_1 + x_2 x_3, \ x'_2 = -4x_1 3x_2 x_3,$ $x_3' = 4x_1 + 4x_2 + 2x_3$
- **25.** $x'_1 = 5x_1 + 5x_2 + 2x_3$, $x'_2 = -6x_1 6x_2 5x_3$, $x'_3 =$ $6x_1 + 6x_2 + 5x_3$
- **26.** Find the particular solution of the system

$$\frac{dx_1}{dt} = 3x_1 + x_3,
\frac{dx_2}{dt} = 9x_1 - x_2 + 2x_3,
\frac{dx_3}{dt} = -9x_1 + 4x_2 - x_3$$

that satisfies the initial conditions $x_1(0) = 0$, $x_2(0) = 0$, $x_3(0) = 17.$

Cascading Brine Tanks

The amounts $x_1(t)$ and $x_2(t)$ of salt in the two brine tanks of Fig. 7.3.7 satisfy the differential equations

$$\frac{dx_1}{dt} = -k_1 x_1, \quad \frac{dx_2}{dt} = k_1 x_1 - k_2 x_2,$$

where $k_i = r/V_i$ for i = 1, 2. In Problems 27 and 28 the volumes V_1 and V_2 are given. First solve for $x_1(t)$ and $x_2(t)$, assuming that r = 10 (gal/min), $x_1(0) = 15$ (lb), and $x_2(0) = 0$. Then find the maximum amount of salt ever in tank 2. Finally, construct a figure showing the graphs of $x_1(t)$ and $x_2(t)$.



FIGURE 7.3.7. The two brine tanks of Problems 27 and 28.

27. $V_1 = 50$ (gal), $V_2 = 25$ (gal) **28.** $V_1 = 25$ (gal), $V_2 = 40$ (gal)

Interconnected Brine Tanks

The amounts $x_1(t)$ and $x_2(t)$ of salt in the two brine tanks of Fig. 7.3.8 satisfy the differential equations

$$\frac{dx_1}{dt} = -k_1 x_1 + k_2 x_2, \quad \frac{dx_2}{dt} = k_1 x_1 - k_2 x_2,$$

where $k_i = r/V_i$ as usual. In Problems 29 and 30, solve for $x_1(t)$ and $x_2(t)$, assuming that r = 10 (gal/min), $x_1(0) = 15$ (lb), and $x_2(0) = 0$. Then construct a figure showing the graphs of $x_1(t)$ and $x_2(t)$.



FIGURE 7.3.8. The two brine tanks of Problems 29 and 30.

29. $V_1 = 50$ (gal), $V_2 = 25$ (gal) **30.** $V_1 = 25$ (gal), $V_2 = 40$ (gal)

Open Three-Tank System

Problems 31 through 34 deal with the open three-tank system of Fig. 7.3.2. Fresh water flows into tank 1; mixed brine flows from tank 1 into tank 2, from tank 2 into tank 3, and out of tank 3; all at the given flow rate r gallons per minute. The initial amounts $x_1(0) = x_0$ (lb), $x_2(0) = 0$, and $x_3(0) = 0$ of salt in the three tanks are given, as are their volumes V_1 , V_2 , and V_3 (in gallons). First solve for the amounts of salt in the three tanks at time t, then determine the maximal amount of salt that tank 3 ever contains. Finally, construct a figure showing the graphs of $x_1(t)$, $x_2(t)$, and $x_3(t)$.

31. r = 30, $x_0 = 27$, $V_1 = 30$, $V_2 = 15$, $V_3 = 10$ **32.** r = 60, $x_0 = 45$, $V_1 = 20$, $V_2 = 30$, $V_3 = 60$ **33.** r = 60, $x_0 = 45$, $V_1 = 15$, $V_2 = 10$, $V_3 = 30$ **34.** r = 60, $x_0 = 40$, $V_1 = 20$, $V_2 = 12$, $V_3 = 60$

Closed Three-Tank System

Problems 35 through 37 deal with the closed three-tank system of Fig. 7.3.5, which is described by the equations in (24). Mixed brine flows from tank 1 into tank 2, from tank 2 into tank 3, and from tank 3 into tank 1, all at the given flow rate r gallons per minute. The initial amounts $x_1(0) = x_0$ (pounds), $x_2(0) = 0$, and $x_3(0) = 0$ of salt in the three tanks are given, as are their volumes V_1 , V_2 , and V_3 (in gallons). First solve for the amounts of salt in the three tanks at time t, then determine the limiting amount (as $t \to +\infty$) of salt in each tank. Finally, construct a figure showing the graphs of $x_1(t)$, $x_2(t)$, and $x_3(t)$.

35. $r = 120, x_0 = 33, V_1 = 20, V_2 = 6, V_3 = 40$ **36.** $r = 10, x_0 = 18, V_1 = 20, V_2 = 50, V_3 = 20$ **37.** $r = 60, x_0 = 55, V_1 = 60, V_2 = 20, V_3 = 30$

For each matrix **A** given in Problems 38 through 40, the zeros in the matrix make its characteristic polynomial easy to calculate. Find the general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

38.
$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 2 & 0 & 0 \\ 0 & 3 & 3 & 0 \\ 0 & 0 & 4 & 4 \end{bmatrix}$$

39.
$$\mathbf{A} = \begin{bmatrix} -2 & 0 & 0 & 9 \\ 4 & 2 & 0 & -10 \\ 0 & 0 & -1 & 8 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

40.
$$\mathbf{A} = \begin{bmatrix} 2 & 0 & 0 & 0 \\ -21 & -5 & -27 & -9 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & -21 & -2 \end{bmatrix}$$

41. The coefficient matrix **A** of the 4×4 system

$x'_1 =$	$4x_1$	+	x_2	+	<i>x</i> ₃	+	$7x_4$,
$x'_2 =$	x_1	+	$4x_2$	+	$10x_{3}$	+	<i>x</i> ₄ ,
$x'_3 =$	x_1	+	$10x_2$	+	$4x_{3}$	+	<i>x</i> 4,
$x'_4 =$	$7x_1$	+	<i>x</i> ₂	+	х3	+	4 <i>x</i> ₄

has eigenvalues $\lambda_1 = -3$, $\lambda_2 = -6$, $\lambda_3 = 10$, and $\lambda_4 = 15$. Find the particular solution of this system that satisfies the initial conditions

$$x_1(0) = 3$$
, $x_2(0) = x_3(0) = 1$, $x_4(0) = 3$.

In Problems 42 through 50, use a calculator or computer system to calculate the eigenvalues and eigenvectors (as illustrated in Application 7.3) in order to find a general solution of the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with the given coefficient matrix \mathbf{A} .

42.	$\mathbf{A} =$	$\begin{bmatrix} -40 & -12 & 54 \\ 35 & 13 & -46 \\ -25 & -7 & 34 \end{bmatrix}$
43.	$\mathbf{A} =$	$\begin{bmatrix} -20 & 11 & 13 \\ 12 & -1 & -7 \\ -48 & 21 & 31 \end{bmatrix}$
44.	$\mathbf{A} =$	$\begin{bmatrix} 147 & 23 & -202 \\ -90 & -9 & 129 \\ 90 & 15 & -123 \end{bmatrix}$
45.	$\mathbf{A} =$	$\begin{bmatrix} 9 & -7 & -5 & 0 \\ -12 & 7 & 11 & 9 \\ 24 & -17 & -19 & -9 \\ -18 & 13 & 17 & 9 \end{bmatrix}$
46.	$\mathbf{A} =$	$\begin{bmatrix} 13 & -42 & 106 & 139 \\ 2 & -16 & 52 & 70 \\ 1 & 6 & -20 & -31 \\ -1 & -6 & 22 & 33 \end{bmatrix}$
47.	$\mathbf{A} =$	$\begin{bmatrix} 23 & -18 & -16 & 0 \\ -8 & 6 & 7 & 9 \\ 34 & -27 & -26 & -9 \\ -26 & 21 & 25 & 12 \end{bmatrix}$
48.	$\mathbf{A} =$	$\begin{bmatrix} 47 & -8 & 5 & -5 \\ -10 & 32 & 18 & -2 \\ 139 & -40 & -167 & -121 \\ -232 & 64 & 360 & 248 \end{bmatrix}$
49.	$\mathbf{A} =$	$\begin{bmatrix} 139 & -14 & -52 & -14 & 28 \\ -22 & 5 & 7 & 8 & -7 \\ 370 & -38 & -139 & -38 & 76 \\ 152 & -16 & -59 & -13 & 35 \\ 95 & -10 & -38 & -7 & 23 \end{bmatrix}$
50.	$\mathbf{A} =$	$\begin{bmatrix} 9 & 13 & 0 & 0 & -13 \\ -14 & 19 & -10 & -20 & 10 & 4 \\ -30 & 12 & -7 & -30 & 12 & 18 \\ -12 & 10 & -10 & -9 & 10 & 2 \\ 6 & 9 & 0 & 6 & 5 & -15 \\ -14 & 23 & -10 & -20 & 10 & 0 \end{bmatrix}$

7.3 Application Automatic Calculation of Eigenvalues and Eigenvectors



computing resources including *Maple/Mathematica/*MATLAB.

1.	1 Þ		D	ocument 1 🤜	7	<10
	-0.5	0	0	-0.5	0	0]
a:=	0.5	-0.25	0	0.5	-0.25	0
L	0	0.25	-0.2	0	0.25	-0.2
eig	л(a)			{-0.	2,-0.25	,-0.5}
eigVc(a)		0.	0.	0.35	8569	
			0.	0.196116	-0.71	7137
			1.	-0.980581	0.59	7614
I.						
-						3/

FIGURE 7.3.9. TI-Nspire CX CAS calculation of the eigenvalues and eigenvectors of the matrix **A**.

Most computational systems offer the capability to find eigenvalues and eigenvectors readily. For instance, Fig. 7.3.9 shows a graphing calculator computation of the eigenvalues and eigenvectors of the matrix

	-0.5	0.0	0.0
$\mathbf{A} = $	0.5	-0.25	0.0
	0.0	0.25	-0.2

of Example 2. We see the three eigenvectors displayed as column vectors, appearing in the same order as their corresponding eigenvalues. In this display the eigenvectors are *normalized*, that is, multiplied by an appropriate scalar so as to have length 1. You can verify, for example, that the displayed eigenvector corresponding to the third eigenvalue $\lambda = -\frac{1}{2}$ is a scalar multiple of $\mathbf{v} = \begin{bmatrix} 1 & -2 & \frac{5}{3} \end{bmatrix}^T$. The *Maple* commands

```
with(linalg)
A := matrix(3,3,[-0.5,0,0,0.5,-0.25,0,0,0.25,-0.2]);
eigenvects(A);
```

the Mathematica commands

the Wolfram Alpha query

```
((-0.5, 0, 0), (0.5, -0.25, 0), (0, 0.25, -0.2))
```

and the MATLAB commands

$$A = [-0.5,0,0; 0.5,-0.25,0; 0,0.25,-0.2]$$

[V,D] = eig(A)

(where D will be a diagonal matrix displaying the eigenvalues of A and the column vectors of v are the corresponding eigenvectors) produce similar results. You can use these commands to find the eigenvalues and eigenvectors needed for any of the problems in this section.

For a more substantial investigation, choose a positive integer n < 10 (n = 5, for instance) and let q_1, q_2, \ldots, q_n denote the first *n* nonzero digits in your student ID number. Now consider an open system of brine tanks as in Fig. 7.3.2, except with *n* rather than three successive tanks having volumes $V_i = 10q_i$ ($i = 1, 2, \ldots, n$) in gallons. If each flow rate is r = 10 gallons per minute, then the salt amounts $x_1(t), x_2(t), \ldots, x_n(t)$ satisfy the linear system

$$\begin{aligned} x'_1 &= -k_1 x_1, \\ x'_i &= k_{i-1} x_{i-1} - k_i x_i \quad (i = 2, 3, \dots, n), \end{aligned}$$

where $k_i = r/V_i$. Apply the eigenvalue method to solve this system with initial conditions

$$x_1(0) = 10, \quad x_2(0) = x_3(0) = \dots = x_n(0) = 0.$$

Graph the solution functions and estimate graphically the maximum amount of salt that each tank ever contains.

For an alternative investigation, suppose that the system of n tanks is closed as in Fig. 7.3.5, so that tank 1 receives as inflow the outflow from tank n (rather than

fresh water). Then the first equation should be replaced with $x'_1 = k_n x_n - k_1 x_1$. Now show that, in this *closed* system, as $t \to +\infty$ the salt originally in tank 1 distributes itself with constant density throughout the various tanks. A plot like Fig. 7.3.6 should make this fairly obvious.

7.4 A Gallery of Solution Curves of Linear Systems

In the preceding section we saw that the eigenvalues and eigenvectors of the $n \times n$ matrix **A** are of central importance to the solutions of the homogeneous linear constant-coefficient system

$$\mathbf{x}' = \mathbf{A}\mathbf{x}.\tag{1}$$

Indeed, according to Theorem 1 from Section 7.3, if λ is an eigenvalue of **A** and **v** is an eigenvector of **A** associated with λ , then

$$\mathbf{x}(t) = \mathbf{v}e^{\lambda t} \tag{2}$$

is a nontrivial solution of the system (1). Moreover, if **A** has *n* linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ associated with its *n* eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$, then in fact all solutions of the system (1) are given by linear combinations

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t} + \dots + c_n \mathbf{v}_n e^{\lambda_n t},$$
(3)

where c_1, c_2, \ldots, c_n are arbitrary constants. If the eigenvalues include complex conjugate pairs, then we can obtain a real-valued general solution from Eq. (3) by taking real and imaginary parts of the terms in (3) corresponding to the complex eigenvalues.

Our goal in this section is to gain a *geometric* understanding of the role that the eigenvalues and eigenvectors of the matrix **A** play in the solutions of the system (1). We will see, illustrating primarily with the case n = 2, that particular arrangements of eigenvalues and eigenvectors correspond to identifiable patterns—"fingerprints," so to speak—in the phase plane portrait of the system (1). Just as in algebra we learn to recognize when an equation in x and y corresponds to a line or parabola, we can predict the general appearance of the solution curves of the system (1) from the eigenvalues and eigenvectors we will create a "gallery"— Figure 7.4.16 appearing at the end of this section—of typical phase plane portraits that gives, in essence, a complete catalog of the geometric behaviors that the solutions of a 2×2 homogeneous linear constant-coefficient system can exhibit. This will help us analyze not only systems of the form (1), but also more complicated systems that can be approximated by linear systems, a topic we explore in Section 9.2.

Systems of Dimension n = 2

3

Until stated otherwise, we henceforth assume that n = 2, so that the eigenvalues of the matrix **A** are λ_1 and λ_2 . According to Theorem 2 of Section 6.2, if λ_1 and λ_2 are distinct, then the associated eigenvectors \mathbf{v}_1 and \mathbf{v}_2 of **A** are linearly independent. In this event, the general solution of the system (1) is given by

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t} \tag{4}$$

if λ_1 and λ_2 are real, and by

$$\mathbf{x}(t) = c_1 e^{pt} (\mathbf{a} \cos qt - \mathbf{b} \sin qt) + c_2 e^{pt} (\mathbf{b} \cos qt + \mathbf{a} \sin qt)$$
(5)

7.4 A GALLERY OF SOLUTION CURVES OF LINEAR SYSTEM 99

if λ_1 and λ_2 are the complex conjugate numbers $p \pm iq$; here the vectors **a** and **b** are the real and imaginary parts, respectively, of a (complex-valued) eigenvector of **A** associated with the eigenvalue $p \pm iq$. If instead λ_1 and λ_2 are equal (to a common value λ , say), then as we will see in Section 7.6, the matrix **A** may or may not have two linearly independent eigenvectors \mathbf{v}_1 and \mathbf{v}_2 . If it does, then the eigenvalue method of Section 7.3 applies once again, and the general solution of the system (1) is given by the linear combination

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 \mathbf{v}_2 e^{\lambda t}$$
(6)

as before. If A does not have two linearly independent eigenvectors, then—as we will see—we can find a vector v_2 such that the general solution of the system (1) is given by

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t}, \tag{7}$$

where \mathbf{v}_1 is an eigenvector of **A** associated with the lone eigenvalue λ . The nature of the vector \mathbf{v}_2 and other details of the general solution in (7) will be discussed in Section 7.6, but we include this case here in order to make our gallery complete.

With this algebraic background in place, we begin our analysis of the solution curves of the system (1). First we assume that the eigenvalues λ_1 and λ_2 of the matrix **A** are real, and subsequently we take up the case where λ_1 and λ_2 are complex conjugates.

Real Eigenvalues

We will divide the case where λ_1 and λ_2 are real into the following possibilities:

Distinct eigenvalues

- Nonzero and of opposite sign ($\lambda_1 < 0 < \lambda_2$)
- Both negative $(\lambda_1 < \lambda_2 < 0)$
- Both positive $(0 < \lambda_2 < \lambda_1)$
- One zero and one negative $(\lambda_1 < \lambda_2 = 0)$
- One zero and one positive $(0 = \lambda_2 < \lambda_1)$

Repeated eigenvalue

- Positive $(\lambda_1 = \lambda_2 > 0)$
- Negative $(\lambda_1 = \lambda_2 < 0)$
- Zero $(\lambda_1 = \lambda_2 = 0)$

Saddle Points

NONZERO DISTINCT EIGENVALUES OF OPPOSITE SIGN: The key observation when $\lambda_1 < 0 < \lambda_2$ is that the positive scalar factors $e^{\lambda_1 t}$ and $e^{\lambda_2 t}$ in the general solution

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t} \tag{4}$$

of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ move in opposite directions (on the real line) as t varies. For example, as t grows large and positive, $e^{\lambda_2 t}$ grows large, because $\lambda_2 > 0$, whereas $e^{\lambda_1 t}$ approaches zero, because $\lambda_1 < 0$; thus the term $c_1\mathbf{v}_1e^{\lambda_1 t}$ in the solution $\mathbf{x}(t)$ in (4) vanishes and $\mathbf{x}(t)$ approaches $c_2\mathbf{v}_2e^{\lambda_2 t}$. If instead t grows large and negative, then the opposite occurs: The factor $e^{\lambda_1 t}$ grows large whereas $e^{\lambda_2 t}$ becomes small, and the solution $\mathbf{x}(t)$ approaches $c_1\mathbf{v}_1e^{\lambda_1 t}$. If we assume for the moment that both c_1 and c_2 are nonzero, then loosely speaking, as t ranges from $-\infty$ to $+\infty$, the solution $\mathbf{x}(t)$ shifts from being "mostly" a multiple of the eigenvector \mathbf{v}_1 to being "mostly" a multiple of \mathbf{v}_2 .



FIGURE 7.4.1. Solution curves $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t}$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when the eigenvalues λ_1, λ_2 of \mathbf{A} are real with $\lambda_1 < 0 < \lambda_2$.

Geometrically, this means that all solution curves given by (4) with both c_1 and c_2 nonzero have two asymptotes, namely the lines l_1 and l_2 passing through the origin and parallel to the eigenvectors \mathbf{v}_1 and \mathbf{v}_2 , respectively; the solution curves approach l_1 as $t \to -\infty$ and l_2 as $t \to +\infty$. Indeed, as Fig. 7.4.1 illustrates, the lines l_1 and l_2 effectively divide the plane into four "quadrants" within which all solution curves flow from the asymptote l_1 to the asymptote l_2 as t increases. (The eigenvectors shown in Fig. 7.4.1—and in other figures—are scaled so as to have equal length.) The particular quadrant in which a solution curve lies is determined by the signs of the coefficients c_1 and c_2 . If c_1 and c_2 are both positive, for example, then the corresponding solution curve extends asymptotically in the direction of the eigenvector \mathbf{v}_1 as $t \to -\infty$, and asymptotically in the direction of \mathbf{v}_2 as $t \to \infty$. If instead $c_1 > 0$ but $c_2 < 0$, then the corresponding solution curve still extends asymptotically in the direction of \mathbf{v}_1 as $t \to -\infty$, but extends asymptotically in the direction *opposite* \mathbf{v}_2 as $t \to +\infty$ (because the negative coefficient c_2 causes the vector $c_2\mathbf{v}_2$ to point "backwards" from \mathbf{v}_2).

If c_1 or c_2 equals zero, then the solution curve remains confined to one of the lines l_1 and l_2 . For example, if $c_1 \neq 0$ but $c_2 = 0$, then the solution (4) becomes $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t}$, which means that the corresponding solution curve lies along the line l_1 . It approaches the origin as $t \to +\infty$, because $\lambda_1 < 0$, and recedes farther and farther from the origin as $t \to -\infty$, either in the direction of \mathbf{v}_1 (if $c_1 > 0$) or the direction opposite \mathbf{v}_1 (if $c_1 < 0$). Similarly, if $c_1 = 0$ and $c_2 \neq 0$, then because $\lambda_2 > 0$, the solution curve flows along the line l_2 away from the origin as $t \to +\infty$ and toward the origin as $t \to -\infty$.

Figure 7.4.1 illustrates typical solution curves corresponding to nonzero values of the coefficients c_1 and c_2 . Because the overall picture of the solution curves is suggestive of the level curves of a saddle-shaped surface (like z = xy), we call the origin a **saddle point** for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

Example 1

The solution curves in Fig. 7.4.1 correspond to the choice

$$\mathbf{A} = \begin{bmatrix} 4 & 1\\ 6 & -1 \end{bmatrix} \tag{8}$$

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$; as you can verify, the eigenvalues of \mathbf{A} are $\lambda_1 = -2$ and $\lambda_2 = 5$ (thus $\lambda_1 < 0 < \lambda_2$), with associated eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} -1 \\ 6 \end{bmatrix}$$
 and $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

According to Eq. (4), the resulting general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -1\\ 6 \end{bmatrix} e^{-2t} + c_2 \begin{bmatrix} 1\\ 1 \end{bmatrix} e^{5t},$$
(9)

or, in scalar form,

$$x_1(t) = -c_1 e^{-2t} + c_2 e^{5t},$$

$$x_2(t) = 6c_1 e^{-2t} + c_2 e^{5t}.$$
(10)

Our gallery Fig. 7.4.16 at the end of this section shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (8). (In Problem 29 we explore "Cartesian" equations for the solution curves (10) relative to the "axes" defined by the lines l_1 and l_2 , which form a natural frame of reference for the solution curves.)

Nodes: Sinks and Sources

DISTINCT NEGATIVE EIGENVALUES: When $\lambda_1 < \lambda_2 < 0$, the factors $e^{\lambda_1 t}$ and $e^{\lambda_2 t}$ both *decrease* as *t* increases. Indeed, as $t \to +\infty$, both $e^{\lambda_1 t}$ and $e^{\lambda_2 t}$ approach zero, which means that the solution curve

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t}$$
(4)

approaches the origin; likewise, as $t \to -\infty$, both $e^{\lambda_1 t}$ and $e^{\lambda_2 t}$ grow without bound, and so the solution curve "goes off to infinity." Moreover, differentiation of the solution in (4) gives

$$\mathbf{x}'(t) = c_1 \lambda_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \lambda_2 \mathbf{v}_2 e^{\lambda_2 t} = e^{\lambda_2 t} \left[c_1 \lambda_1 \mathbf{v}_1 e^{(\lambda_1 - \lambda_2)t} + c_2 \lambda_2 \mathbf{v}_2 \right].$$
(11)

This shows that the tangent vector $\mathbf{x}'(t)$ to the solution curve $\mathbf{x}(t)$ is a scalar multiple of the vector $c_1\lambda_1\mathbf{v}_1e^{(\lambda_1-\lambda_2)t} + c_2\lambda_2\mathbf{v}_2$, which approaches the fixed nonzero multiple $c_2\lambda_2\mathbf{v}_2$ of the vector \mathbf{v}_2 as $t \to +\infty$ (because $e^{(\lambda_1-\lambda_2)t}$ approaches zero). It follows that if $c_2 \neq 0$, then as $t \to +\infty$, the solution curve $\mathbf{x}(t)$ becomes more and more nearly parallel to the eigenvector \mathbf{v}_2 . (More specifically, note that if $c_2 > 0$, for example, then $\mathbf{x}(t)$ approaches the origin in the direction *opposite to* \mathbf{v}_2 , because the scalar $c_2\lambda_2$ is negative.) Thus, if $c_2 \neq 0$, then with increasing t the solution curve approaches the origin and is tangent there to the line l_2 passing through the origin and parallel to \mathbf{v}_2 .

If $c_2 = 0$, on the other hand, then the solution curve $\mathbf{x}(t)$ flows similarly along the line l_1 passing through the origin and parallel to the eigenvector \mathbf{v}_1 . Once again, the net effect is that the lines l_1 and l_2 divide the plane into four "quadrants" as shown in Figure 7.4.2, which illustrates typical solution curves corresponding to nonzero values of the coefficients c_1 and c_2 .

To describe the appearance of phase portraits like Fig. 7.4.2, we introduce some new terminology, which will be useful both now and in Chapter 9, when we study nonlinear systems. In general, we call the origin a **node** of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ provided that *both* of the following conditions are satisfied:

- *Either* every trajectory approaches the origin as $t \to +\infty$ or every trajectory recedes from the origin as $t \to +\infty$;
- Every trajectory is tangent at the origin to some straight line through the origin.

Moreover, we say that the origin is a **proper** node provided that no two different pairs of "opposite" trajectories are tangent to the same straight line through the origin. This is the situation in Fig. 7.4.6, in which the trajectories *are* straight lines, not merely tangent to straight lines; indeed, a proper node might be called a "star point." However, in Fig. 7.4.2, all trajectories—apart from those that flow along the line l_1 —are tangent to the line l_2 ; as a result we call the node **improper**.

Further, if every trajectory for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ approaches the origin as $t \to +\infty$ (as in Fig. 7.4.2), then the origin is called a **sink**; if instead every trajectory recedes from the origin, then the origin is a **source**. Thus we describe the characteristic pattern of the trajectories in Fig. 7.4.2 as an **improper nodal sink**.

Example 2 The solution curves in Fig. 7.4.2 correspond to the choice

$$\mathbf{A} = \begin{bmatrix} -8 & 3\\ 2 & -13 \end{bmatrix} \tag{12}$$



FIGURE 7.4.2. Solution curves $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t}$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when the eigenvalues λ_1, λ_2 of \mathbf{A} are real with $\lambda_1 < \lambda_2 < 0$.

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. The eigenvalues of \mathbf{A} are $\lambda_1 = -14$ and $\lambda_2 = -7$ (and thus $\lambda_1 < \lambda_2 < 0$), with associated eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} -1\\2 \end{bmatrix}$$
 and $\mathbf{v}_2 = \begin{bmatrix} 3\\1 \end{bmatrix}$.

Equation (4) then gives the general solution

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -1\\2 \end{bmatrix} e^{-14t} + c_2 \begin{bmatrix} 3\\1 \end{bmatrix} e^{-7t},$$
(13)

or, in scalar form,

$$x_1(t) = -c_1 e^{-14t} + 3c_2 e^{-7t},$$

$$x_2(t) = 2c_1 e^{-14t} + c_2 e^{-7t}.$$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (12).

The case of distinct positive eigenvalues mirrors that of distinct negative eigenvalues. But instead of analyzing it independently, we can rely on the following principle, whose verification is a routine matter of checking signs (Problem 30).

PRINCIPLE Time Reversal in Linear Systems

Let $\mathbf{x}(t)$ be a solution of the 2-dimensional linear system

$$\mathbf{x}' = \mathbf{A}\mathbf{x} \tag{1}$$

Then the function $\tilde{\mathbf{x}}(t) = \mathbf{x}(-t)$ is a solution of the system

$$\tilde{\mathbf{x}}' = -\mathbf{A}\tilde{\mathbf{x}}.\tag{14}$$

We note furthermore that the two vector-valued functions $\mathbf{x}(t)$ and $\tilde{\mathbf{x}}(t)$ for $-\infty < t < \infty$ have the same solution curve (or image) in the plane. However, the chain rule gives $\tilde{\mathbf{x}}'(t) = -\mathbf{x}'(t)$; since $\tilde{\mathbf{x}}(t)$ and $\mathbf{x}(-t)$ represent the same point, it follows that at each point of their common solution curve the velocity vectors of the two functions $\mathbf{x}(t)$ and $\tilde{\mathbf{x}}(t)$ are negatives of each other. Therefore the two solutions traverse their common solution curve in opposite directions as *t* increases—or, alternatively, in the same direction as *t* increases for one solution and decreases for the other. In short, we may say that the solutions of the systems (1) and (14) correspond to each other under "time reversal," since we get the solutions of one system by letting time "run backwards" in the solutions of the other.

DISTINCT POSITIVE EIGENVALUES: If the matrix **A** has positive eigenvalues with $0 < \lambda_2 < \lambda_1$, then as you can verify (Problem 31), the matrix $-\mathbf{A}$ has negative eigenvalues $-\lambda_1 < -\lambda_2 < 0$ but the same eigenvectors \mathbf{v}_1 and \mathbf{v}_2 . The preceding case then shows that the system $\mathbf{x}' = -\mathbf{A}\mathbf{x}$ has an improper nodal sink at the origin. But the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ has the same trajectories, except with the direction of motion (as *t* increases) along each solution curve reversed. Thus the origin is now a *source*, rather than a sink, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$, and we call the origin an **improper nodal source**. Figure 7.4.3 illustrates typical solution curves given by $\mathbf{x}(t) = c_1\mathbf{v}_1e^{\lambda_1t} + c_2\mathbf{v}_2e^{\lambda_2t}$ corresponding to nonzero values of the coefficients c_1 and c_2 .

Example 3

3 The solution curves in Fig. 7.4.3 correspond to the choice

$$\mathbf{A} = -\begin{bmatrix} -8 & 3\\ 2 & -13 \end{bmatrix} = \begin{bmatrix} 8 & -3\\ -2 & 13 \end{bmatrix}$$
(15)



FIGURE 7.4.3. Solution curves $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t}$ for the

system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when the eigenvalues

 λ_1, λ_2 of **A** are real with

 $0 < \lambda_2 < \lambda_1$.

7.4 A GALLERY OF SOLUTION CURVES OF LINEAR SYSTEM 63

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$; thus \mathbf{A} is the negative of the matrix in Example 2. Therefore we can solve the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ by applying the principle of time reversal to the solution in Eq. (13): Replacing *t* with -t in the righthand side of (13) leads to

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -1\\2 \end{bmatrix} e^{14t} + c_2 \begin{bmatrix} 3\\1 \end{bmatrix} e^{7t}.$$
(16)

Of course, we could also have "started from scratch" by finding the eigenvalues λ_1 , λ_2 and eigenvectors \mathbf{v}_1 , \mathbf{v}_2 of **A**. These can be found from the definition of eigenvalue, but it is easier to note (see Problem 31 again) that because **A** is the negative of the matrix in Eq. (12), λ_1 and λ_2 are likewise the negatives of their values in Example 2, whereas we can take \mathbf{v}_1 and \mathbf{v}_2 to be the same as in Example 2. By either means we find that $\lambda_1 = 14$ and $\lambda_2 = 7$ (so that $0 < \lambda_2 < \lambda_1$), with associated eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$
 and $\mathbf{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$.

From Eq. (4), then, the general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -1\\2 \end{bmatrix} e^{14t} + c_2 \begin{bmatrix} 3\\1 \end{bmatrix} e^{7t}$$

(in agreement with Eq. (16)), or, in scalar form,

$$x_1(t) = -c_1 e^{14t} + 3c_2 e^{7t},$$

$$x_2(t) = 2c_1 e^{14t} + c_2 e^{7t}.$$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (15).

Zero Eigenvalues and Straight-Line Solutions

ONE ZERO AND ONE NEGATIVE EIGENVALUE: When $\lambda_1 < \lambda_2 = 0$, the general solution (4) becomes

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2. \tag{17}$$

For any fixed nonzero value of the coefficient c_1 , the term $c_1\mathbf{v}_1e^{\lambda_1 t}$ in Eq. (17) is a scalar multiple of the eigenvector \mathbf{v}_1 , and thus (as t varies) travels along the line l_1 passing through the origin and parallel to \mathbf{v}_1 ; the direction of travel is toward the origin as $t \to +\infty$ because $\lambda_1 < 0$. If $c_1 > 0$, for example, then $c_1\mathbf{v}_1e^{\lambda_1 t}$ extends in the direction of \mathbf{v}_1 , approaching the origin as t increases, and receding from the origin as t decreases. If instead $c_1 < 0$, then $c_1\mathbf{v}_1e^{\lambda_1 t}$ extends in the direction opposite \mathbf{v}_1 while still approaching the origin as t increases. Loosely speaking, we can visualize the flow of the term $c_1\mathbf{v}_1e^{\lambda_1 t}$ taken alone as a pair of arrows opposing each other head-to-head at the origin. The solution curve $\mathbf{x}(t)$ in Eq. (17) is simply this same trajectory $c_1\mathbf{v}_1e^{\lambda_1 t}$, then shifted (or offset) by the constant vector $c_2\mathbf{v}_2$. Thus in this case the phase portrait of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ consists of all lines parallel to the eigenvector \mathbf{v}_1 , where along each such line the solution flows (from both directions) toward the line l_2 passing through the origin and parallel to \mathbf{v}_1 . Figure 7.4.4 illustrates typical solution curves corresponding to nonzero values of the coefficients c_1 and c_2 .

It is noteworthy that each single point represented by a constant vector **b** lying on the line l_2 represents a *constant* solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Indeed, if **b** lies on l_2 , then **b** is a scalar multiple $k \cdot \mathbf{v}_2$ of the eigenvector \mathbf{v}_2 of **A** associated with the eigenvalue $\lambda_2 = 0$. In this case, the constant-valued solution $\mathbf{x}(t) \equiv \mathbf{b}$ is given by Eq. (17) with $c_1 = 0$ and $c_2 = k$. This constant solution, with its "trajectory" being



FIGURE 7.4.4. Solution curves $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when the eigenvalues λ_1, λ_2 of \mathbf{A} are real with $\lambda_1 < \lambda_2 = 0$.

a single point lying on the line l_2 , is then the unique solution of the initial value problem

$$\mathbf{x}' = \mathbf{A}\mathbf{x}, \quad \mathbf{x}(0) = \mathbf{b}$$

guaranteed by Theorem 1 of Section 7.1. Note that this situation is in marked contrast with the other eigenvalue cases we have considered so far, in which $\mathbf{x}(t) \equiv \mathbf{0}$ is the only constant solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. (In Problem 32 we explore the general circumstances under which the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ has constant solutions other than $\mathbf{x}(t) \equiv \mathbf{0}$.)

Example 4 The solution curves in Fig. 7.4.4 correspond to the choice

$$\mathbf{A} = \begin{bmatrix} -36 & -6\\ 6 & 1 \end{bmatrix} \tag{18}$$

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. The eigenvalues of \mathbf{A} are $\lambda_1 = -35$ and $\lambda_2 = 0$, with associated eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 6\\-1 \end{bmatrix}$$
 and $\mathbf{v}_2 = \begin{bmatrix} 1\\-6 \end{bmatrix}$

Based on Eq. (17), the general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 6\\-1 \end{bmatrix} e^{-35t} + c_2 \begin{bmatrix} 1\\-6 \end{bmatrix},$$
(19)

or, in scalar form,

$$x_1(t) = 6c_1 e^{-35t} + c_2,$$

$$x_2(t) = -c_1 e^{-35t} - 6c_2.$$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (18).

ONE ZERO AND ONE POSITIVE EIGENVALUE: When $0 = \lambda_2 < \lambda_1$, the solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is again given by

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2. \tag{17}$$

By the principle of time reversal, the trajectories of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ are identical to those of the system $\mathbf{x}' = -\mathbf{A}\mathbf{x}$, except that they flow in the opposite direction. Since the eigenvalues $-\lambda_1$ and $-\lambda_2$ of the matrix $-\mathbf{A}$ satisfy $-\lambda_1 < -\lambda_2 = 0$, by the preceding case the trajectories of $\mathbf{x}' = -\mathbf{A}\mathbf{x}$ are lines parallel to the eigenvector \mathbf{v}_1 and flowing toward the line l_2 from both directions. Therefore the trajectories of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ are lines parallel to \mathbf{v}_1 and flowing *away* from the line l_2 . Figure 7.4.5 illustrates typical solution curves given by $\mathbf{x}(t) = c_1\mathbf{v}_1e^{\lambda_1t} + c_2\mathbf{v}_2$ corresponding to nonzero values of the coefficients c_1 and c_2 .

Ie 5 The solution curves in Fig. 7.4.5 correspond to the choice

$$\mathbf{A} = -\begin{bmatrix} -36 & -6\\ 6 & 1 \end{bmatrix} = \begin{bmatrix} 36 & 6\\ -6 & -1 \end{bmatrix}$$
(20)

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$; thus \mathbf{A} is the negative of the matrix in Example 4. Once again we can solve the system using the principle of time reversal: Replacing *t* with -t in the right-hand side of the solution in Eq. (19) of Example 4 leads to

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 6\\-1 \end{bmatrix} e^{35t} + c_2 \begin{bmatrix} 1\\-6 \end{bmatrix}.$$
 (21)



FIGURE 7.4.5. Solution curves $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when the eigenvalues λ_1, λ_2 of \mathbf{A} are real with $0 = \lambda_2 < \lambda_1$.

Example 5

Alternatively, directly finding the eigenvalues and eigenvectors of A leads to $\lambda_1 = 35$ and $\lambda_2 = 0$, with associated eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 6\\ -1 \end{bmatrix}$$
 and $\mathbf{v}_2 = \begin{bmatrix} 1\\ -6 \end{bmatrix}$.

Equation (17) gives the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ as

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 6\\-1 \end{bmatrix} e^{35t} + c_2 \begin{bmatrix} 1\\-6 \end{bmatrix}$$

(in agreement with Eq. (21)), or, in scalar form,

$$x_1(t) = 6c_1 e^{35t} + c_2,$$

$$x_2(t) = -c_1 e^{35t} - 6c_2.$$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (20).

Repeated Eigenvalues; Proper and Improper Nodes

REPEATED POSITIVE EIGENVALUE: As we noted earlier, if the matrix **A** has one repeated eigenvalue, then **A** may or may not have two associated linearly independent eigenvectors. Because these two possibilities lead to quite different phase portraits, we will consider them separately. We let λ denote the repeated eigenvalue of **A** with $\lambda > 0$.

With two independent eigenvectors: First, if A does have two linearly independent eigenvectors, then it is easy to show (Problem 33) that in fact *every* nonzero vector is an eigenvector of A, from which it follows that A must be equal to the scalar λ times the identity matrix of order two, that is,

$$\mathbf{A} = \lambda \begin{bmatrix} 1 & 0\\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \lambda & 0\\ 0 & \lambda \end{bmatrix}.$$
(22)

Therefore the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ becomes (in scalar form)

$$x'_{1}(t) = \lambda x_{1}(t),$$

 $x'_{2}(t) = \lambda x_{2}(t).$
(23)

The general solution of Eq. (23) is

$$x_1(t) = c_1 e^{\lambda t}$$

$$x_2(t) = c_2 e^{\lambda t},$$
(24)

or in vector format,

$$\mathbf{x}(t) = e^{\lambda t} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}.$$
 (25)

We could also have arrived at Eq. (25) by starting, as in previous cases, from our general solution (4): Because all nonzero vectors are eigenvectors of **A**, we are free to take $\mathbf{v}_1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$ as a representative pair of linearly independent eigenvectors, each associated with the eigenvalue λ . Then Eq. (4) leads to the same result as Eq. (25):

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 \mathbf{v}_2 e^{\lambda t} = e^{\lambda t} (c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2) = e^{\lambda t} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}.$$

Either way, our solution in Eq. (25) shows that $\mathbf{x}(t)$ is always a positive scalar multiple of the fixed vector $\begin{bmatrix} c_1 & c_2 \end{bmatrix}^T$. Thus apart from the case $c_1 = c_2 = 0$, the trajectories of the system (1) are half-lines, or rays, emanating from the origin and (because $\lambda > 0$) flowing away from it. As noted above, the origin in this case represents a *proper* node, because no two pairs of "opposite" solution curves are tangent to the same straight line through the origin. Moreover the origin is also a source (rather than a sink), and so in this case we call the origin a **proper nodal source**. Figure 7.4.6 shows the "exploding star" pattern characteristic of such points.

Example 6

The solution curves in Fig. 7.4.6 correspond to the case where the matrix **A** is given by Eq. (22) with $\lambda = 2$:

$$\mathbf{A} = \begin{bmatrix} 2 & 0\\ 0 & 2 \end{bmatrix}. \tag{26}$$



FIGURE 7.4.6. Solution curves $\mathbf{x}(t) = e^{\lambda t} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when **A** has one repeated

positive eigenvalue and two linearly independent eigenvectors.

Equation (25) then gives the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ as

$$\mathbf{x}(t) = e^{2t} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix},\tag{27}$$

or, in scalar form,

$$x_1(t) = c_1 e^{2t},$$

 $x_2(t) = c_2 e^{2t}.$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (26).

Without two independent eigenvectors: The remaining possibility is that the matrix **A** has a repeated positive eigenvalue yet fails to have two linearly independent eigenvectors. In this event the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is given by Eq. (7) above:

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t}.$$
(7)

Here \mathbf{v}_1 is an eigenvector of the matrix **A** associated with the repeated eigenvalue λ and \mathbf{v}_2 is a (nonzero) "generalized eigenvector" that will be described more fully in Section 7.6. To analyze this trajectory, we first distribute the factor $e^{\lambda t}$ in Eq. (7), leading to

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 (\mathbf{v}_1 t e^{\lambda t} + \mathbf{v}_2 e^{\lambda t}).$$
(28)

Our assumption that $\lambda > 0$ implies that both $e^{\lambda t}$ and $te^{\lambda t}$ approach zero as $t \to -\infty$, and so by Eq. (28) the solution $\mathbf{x}(t)$ approaches the origin as $t \to -\infty$. Except for the trivial solution given by $c_1 = c_2 = 0$, all trajectories given by Eq. (7) "emanate" from the origin as t increases.

The direction of flow of these curves can be understood from the tangent vector $\mathbf{x}'(t)$. Rewriting Eq. (28) as

$$\mathbf{x}(t) = e^{\lambda t} \left[c_1 \mathbf{v}_1 + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) \right]$$

and applying the product rule for vector-valued functions gives

$$\mathbf{x}'(t) = e^{\lambda t} c_2 \mathbf{v}_1 + \lambda e^{\lambda t} [c_1 \mathbf{v}_1 + c_2 (\mathbf{v}_1 t + \mathbf{v}_2)]$$

= $e^{\lambda t} (c_2 \mathbf{v}_1 + \lambda c_1 \mathbf{v}_1 + \lambda c_2 \mathbf{v}_1 t + \lambda c_2 \mathbf{v}_2).$ (29)

For $t \neq 0$, we can factor out t in Eq. (29) and rearrange terms to get

$$\mathbf{x}'(t) = t e^{\lambda t} \left[\lambda c_2 \mathbf{v}_1 + \frac{1}{t} (\lambda c_1 \mathbf{v}_1 + \lambda c_2 \mathbf{v}_2 + c_2 \mathbf{v}_1) \right].$$
(30)

Equation (30) shows that for $t \neq 0$, the tangent vector $\mathbf{x}'(t)$ is a nonzero scalar multiple of the vector $\lambda c_2 \mathbf{v}_1 + \frac{1}{t}(\lambda c_1 \mathbf{v}_1 + \lambda c_2 \mathbf{v}_2 + c_2 \mathbf{v}_1)$, which, if $c_2 \neq 0$, approaches the fixed nonzero multiple $\lambda c_2 \mathbf{v}_1$ of the eigenvector \mathbf{v}_1 as $t \to +\infty$ or as $t \to -\infty$. In this case it follows that as t gets larger and larger numerically (in either direction), the tangent line to the solution curve at the point $\mathbf{x}(t)$ —since it is parallel to the tangent vector $\mathbf{x}'(t)$ which approaches $\lambda c_2 \mathbf{v}_1$ —becomes more and more nearly parallel to the eigenvector \mathbf{v}_1 . In short, we might say that as t increases numerically, the point $\mathbf{x}(t)$ on the solution curve moves in a direction that is more and more nearly parallel to the vector \mathbf{v}_1 , or still more briefly, that near $\mathbf{x}(t)$ the solution curve itself is virtually parallel to \mathbf{v}_1 .

We conclude that if $c_2 \neq 0$, then as $t \to -\infty$ the point $\mathbf{x}(t)$ approaches the origin along the solution curve which is tangent there to the vector \mathbf{v}_1 . But as $t \to +\infty$ and the point $\mathbf{x}(t)$ recedes farther and farther from the origin, the tangent line to the trajectory at this point tends to differ (in direction) less and less from the (moving) line through $\mathbf{x}(t)$ that is parallel to the (fixed) vector \mathbf{v}_1 . Speaking loosely but suggestively, we might therefore say that at points sufficiently far from the origin, all trajectories are essentially parallel to the single vector \mathbf{v}_1 .

If instead $c_2 = 0$, then our solution (7) becomes

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t},\tag{31}$$

and thus runs along the line l_1 passing through the origin and parallel to the eigenvector \mathbf{v}_1 . Because $\lambda > 0$, $\mathbf{x}(t)$ flows away from the origin as t increases; the flow is in the direction of \mathbf{v}_1 if $c_1 > 0$, and opposite \mathbf{v}_1 if $c_1 < 0$.

We can further see the influence of the coefficient c_2 by writing Eq. (7) in yet a different way:

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t} = (c_1 + c_2 t) \mathbf{v}_1 e^{\lambda t} + c_2 \mathbf{v}_2 e^{\lambda t}.$$
 (32)

It follows from Eq. (32) that if $c_2 \neq 0$, then the solution curve $\mathbf{x}(t)$ does not cross the line l_1 . Indeed, if $c_2 > 0$, then Eq. (32) shows that for all t, the solution curve $\mathbf{x}(t)$ lies on the same side of l_1 as \mathbf{v}_2 , whereas if $c_2 < 0$, then $\mathbf{x}(t)$ lies on the opposite side of l_1 .

To see the overall picture, then, suppose for example that the coefficient $c_2 > 0$. Starting from a large *negative* value of t, Eq. (30) shows that as t increases, the direction in which the solution curve $\mathbf{x}(t)$ initially proceeds from the origin is roughly that of the vector $te^{\lambda t}\lambda c_2\mathbf{v}_1$. Since the scalar $te^{\lambda t}\lambda c_2$ is negative (because t < 0 and $\lambda c_2 > 0$), the direction of the trajectory is *opposite* that of \mathbf{v}_1 . For large *positive* values of t, on the other hand, the scalar $te^{\lambda t}\lambda c_2$ is positive, and so $\mathbf{x}(t)$ flows in nearly the *same* direction as \mathbf{v}_1 . Thus, as t increases from $-\infty$ to $+\infty$, the solution curve leaves the origin flowing in the direction opposite \mathbf{v}_1 , makes a "U-turn" as it moves away from the origin, and ultimately flows in the direction of \mathbf{v}_1 .

Because all nonzero trajectories are tangent at the origin to the line l_1 , the origin represents an **improper nodal source**. Figure 7.4.7 illustrates typical solution curves given by $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t}$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when **A** has a repeated eigenvalue but does not have two linearly independent eigenvectors.

Example 7 The solution curves in Fig. 7.4.7 correspond to the choice

$$\mathbf{A} = \begin{bmatrix} 1 & -3\\ 3 & 7 \end{bmatrix} \tag{33}$$



FIGURE 7.4.7. Solution curves $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda t} + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t}$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when **A** has one repeated positive eigenvalue λ with associated eigenvector \mathbf{v}_1 and "generalized eigenvector" \mathbf{v}_2 .

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. In Examples 2 and 3 of Section 7.6 we will see that \mathbf{A} has the repeated eigenvalue $\lambda = 4$ with associated eigenvector and generalized eigenvector given by

$$\mathbf{v}_1 = \begin{bmatrix} -3\\ 3 \end{bmatrix}$$
 and $\mathbf{v}_2 = \begin{bmatrix} 1\\ 0 \end{bmatrix}$, (34)

respectively. According to Eq. (7) the resulting general solution is

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -3\\3 \end{bmatrix} e^{4t} + c_2 \begin{bmatrix} -3t+1\\3t \end{bmatrix} e^{4t}, \qquad (35)$$

· 1 f

or, in scalar form,

$$x_1(t) = (-3c_2t - 3c_1 + c_2)e^{4t},$$

$$x_2(t) = (3c_2t + 3c_1)e^{4t}.$$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (33).

REPEATED NEGATIVE EIGENVALUE: Once again the principle of time reversal shows that the solutions $\mathbf{x}(t)$ of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ are identical to those of $\mathbf{x}' = -\mathbf{A}\mathbf{x}$ with *t* replaced by -t; hence these two systems share the same trajectories while flowing in opposite directions. Further, if the matrix \mathbf{A} has the repeated negative eigenvalue λ , then the matrix $-\mathbf{A}$ has the repeated positive eigenvalue $-\lambda$ (Problem 31 again). Therefore, to construct phase portraits for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when \mathbf{A} has a repeated negative eigenvalue, we simply reverse the directions of the trajectories in the phase portraits corresponding to a repeated positive eigenvalue. These portraits are illustrated in Figs. 7.4.8 and 7.4.9. In Fig. 7.4.8 the origin represents a **proper nodal sink**, whereas in Fig. 7.4.9 it represents an **improper nodal sink**.



The solution curves in Fig. 7.4.8 correspond to the choice

$$\mathbf{A} = -\begin{bmatrix} 2 & 0\\ 0 & 2 \end{bmatrix} = \begin{bmatrix} -2 & 0\\ 0 & -2 \end{bmatrix}$$
(36)

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$; thus \mathbf{A} is the negative of the matrix in Example 6. We can solve this system by applying the principle of time reversal to the solution found in Eq. (27): Replacing *t* with -t in the right-hand side of Eq. (27) leads to

$$\mathbf{x}(t) = e^{-2t} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix},\tag{37}$$

or, in scalar form,

$$x_1(t) = c_1 e^{-2t},$$

$$x_2(t) = c_2 e^{-2t}.$$

Alternatively, because **A** is given by Eq. (22) with $\lambda = -2$, Eq. (25) leads directly to the solution in Eq. (37). Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with **A** given by Eq. (36).

Example 9 The solution curves in Fig. 7.4.9 correspond to the choice

$$\mathbf{A} = -\begin{bmatrix} 1 & -3\\ 3 & 7 \end{bmatrix} = \begin{bmatrix} -1 & 3\\ -3 & -7 \end{bmatrix}$$
(38)

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Thus \mathbf{A} is the negative of the matrix in Example 7, and once again we can apply the principle of time reversal to the solution found in Eq. (35): Replacing *t* with -t in the right-hand side of Eq. (35) yields

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -3\\ 3 \end{bmatrix} e^{-4t} + c_2 \begin{bmatrix} 3t+1\\ -3t \end{bmatrix} e^{-4t}.$$
(39)

We could also arrive at an equivalent form of the solution in Eq. (39) in the following way. You can verify that A has the repeated eigenvalue $\lambda = -2$ with eigenvector \mathbf{v}_1 given by Eq. (34), that is,

$$\mathbf{v}_1 = \begin{bmatrix} -3\\ 3 \end{bmatrix}.$$

However, as the methods of Section 7.6 will show, a generalized eigenvector v_2 associated with v_1 is now given by

$$\mathbf{v}_2 = -\begin{bmatrix} 1\\0 \end{bmatrix} = \begin{bmatrix} -1\\0 \end{bmatrix};$$

that is, v_2 is the negative of the generalized eigenvector in Eq. (34). Equation (7) then gives the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ as

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -3\\3 \end{bmatrix} e^{-4t} + c_2 \begin{bmatrix} -3t-1\\3t \end{bmatrix} e^{-4t}, \tag{40}$$

or, in scalar form,

$$x_1(t) = (-3c_2t - 3c_1 - c_2)e^{-4t},$$

$$x_2(t) = (3c_2t + 3c_1)e^{-4t}.$$

Note that replacing c_2 with $-c_2$ in the solution (39) yields the solution (40), thus confirming that the two solutions are indeed equivalent. Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (38).

The Special Case of a Repeated Zero Eigenvalue

REPEATED ZERO EIGENVALUE: Once again the matrix **A** may or may not have two linearly independent eigenvectors associated with the repeated eigenvalue $\lambda = 0$. If it does, then (using Problem 33 once more) we conclude that every nonzero vector is an eigenvector of **A**, that is, that $\mathbf{A}\mathbf{v} = 0 \cdot \mathbf{v} = \mathbf{0}$ for all two-dimensional vectors **v**. It follows that **A** is the zero matrix of order two, that is,

$$\mathbf{A} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Therefore the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ reduces to $x'_1(t) = x'_2(t) = 0$, which is to say that $x_1(t)$ and $x_2(t)$ are each constant functions. Thus the general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is simply

$$\mathbf{x}(t) = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix},\tag{41}$$

where c_1 and c_2 are arbitrary constants, and the "trajectories" given by Eq. (41) are simply the fixed points (c_1, c_2) in the phase plane.

If instead **A** does not have two linearly independent eigenvectors associated with $\lambda = 0$, then the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is given by Eq. (7) with $\lambda = 0$:

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 + c_2 (\mathbf{v}_1 t + \mathbf{v}_2) = (c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2) + c_2 \mathbf{v}_1 t.$$
(42)

Once again \mathbf{v}_1 denotes an eigenvector of the matrix **A** associated with the repeated eigenvalue $\lambda = 0$ and \mathbf{v}_2 denotes a corresponding nonzero "generalized eigenvector." If $c_2 \neq 0$, then the trajectories given by Eq. (42) are lines parallel to the eigenvector \mathbf{v}_1 and "starting" at the point $c_1\mathbf{v}_1 + c_2\mathbf{v}_2$ (when t = 0). When $c_2 > 0$ the trajectory proceeds in the same direction as \mathbf{v}_1 , whereas when $c_2 < 0$ the solution curve flows in the direction opposite \mathbf{v}_1 . Once again the lines l_1 and l_2 passing through the origin and parallel to the vectors \mathbf{v}_1 and \mathbf{v}_2 , respectively, divide the plane into "quadrants" corresponding to the signs of the coefficients c_1 and c_2 . The particular quadrant in which the "starting point" $c_1\mathbf{v}_1 + c_2\mathbf{v}_2$ of the trajectory falls is determined by the signs of c_1 and c_2 . Finally, if $c_2 = 0$, then Eq. (42) gives $\mathbf{x}(t) \equiv c_1\mathbf{v}_1$ for all t, which means that each fixed point $c_1\mathbf{v}_1$ along the line l_1 corresponds to a solution curve. (Thus the line l_1 could be thought of as a median strip dividing two opposing lanes of traffic.) Figure 7.4.10 illustrates typical solution curves corresponding to nonzero values of the coefficients c_1 and c_2 .

Example 10 The solution curves in Fig. 7.4.10 correspond to the choice

$$\mathbf{A} = \begin{bmatrix} 2 & 4\\ -1 & -2 \end{bmatrix} \tag{43}$$

 $\begin{array}{c|c} c_1 > 0, c_2 > 0 & c_1 < 0, c_2 > 0 \\ \hline l_1 & v_2 & v_1 \\ \hline l_2 & v_1 & c_1 < 0, c_2 < 0 \\ \hline c_1 > 0, c_2 < 0 & c_1 < 0, c_2 < 0 \\ \hline x_1 & v_1 & c_1 < 0, c_2 < 0 \\ \hline \end{array}$

FIGURE 7.4.10. Solution curves $\mathbf{x}(t) = (c_1\mathbf{v}_1 + c_2\mathbf{v}_2) + c_2\mathbf{v}_1t$ for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when **A** has a repeated zero eigenvalue with associated eigenvector \mathbf{v}_1 and "generalized eigenvector" \mathbf{v}_2 . The emphasized point on each solution curve corresponds to t = 0.

in the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. You can verify that $\mathbf{v}_1 = \begin{bmatrix} 2 & -1 \end{bmatrix}^T$ is an eigenvector of \mathbf{A} associated with the repeated eigenvalue $\lambda = 0$. Further, using the methods of Section 7.6 we can show that $\mathbf{v}_2 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ is a corresponding "generalized eigenvector" of \mathbf{A} . According to Eq. (42) the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is therefore

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 2\\-1 \end{bmatrix} + c_2 \left(\begin{bmatrix} 2\\-1 \end{bmatrix} t + \begin{bmatrix} 1\\0 \end{bmatrix} \right), \tag{44}$$

or, in scalar form,

$$x_1(t) = 2c_1 + (2t+1)c_2,$$

$$x_2(t) = -c_1 - tc_2.$$

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (43).

Complex Conjugate Eigenvalues and Eigenvectors

We turn now to the situation in which the eigenvalues λ_1 and λ_2 of the matrix **A** are complex conjugate. As we noted at the beginning of this section, the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is given by Eq. (5):

$$\mathbf{x}(t) = c_1 e^{pt} (\mathbf{a} \cos qt - \mathbf{b} \sin qt) + c_2 e^{pt} (\mathbf{b} \cos qt + \mathbf{a} \sin qt).$$
(5)

Here the vectors **a** and **b** are the real and imaginary parts, respectively, of a (complexvalued) eigenvector of **A** associated with the eigenvalue $\lambda_1 = p + iq$. We will divide the case of complex conjugate eigenvalues according to whether the real part *p* of λ_1 and λ_2 is zero, positive, or negative:

- Pure imaginary $(\lambda_1, \lambda_2 = \pm iq \text{ with } q \neq 0)$
- Complex with negative real part $(\lambda_1, \lambda_2 = p \pm iq \text{ with } p < 0 \text{ and } q \neq 0)$
- Complex with positive real part $(\lambda_1, \lambda_2 = p \pm iq \text{ with } p > 0 \text{ and } q \neq 0)$

Pure Imaginary Eigenvalues: Centers and Elliptical Orbits

PURE IMAGINARY EIGENVALUES: Here we assume that the eigenvalues of the matrix **A** are given by $\lambda_1 \lambda_2 = \pm iq$ with $q \neq 0$. Taking p = 0 in Eq. (5) gives the general solution

$$\mathbf{x}(t) = c_1(\mathbf{a}\cos qt - \mathbf{b}\sin qt) + c_2(\mathbf{b}\cos qt + \mathbf{a}\sin qt)$$
(45)

for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Rather than directly analyze the trajectories given by Eq. (45), as we have done in the previous cases, we begin instead with an example that will shed light on the nature of these solution curves.

Example 11 Solve the initial value problem

$$\mathbf{x}' = \begin{bmatrix} 6 & -17\\ 8 & -6 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 4\\ 2 \end{bmatrix}.$$
(46)

Solution The coefficient matrix

$$\mathbf{A} = \begin{bmatrix} 6 & -17\\ 8 & -6 \end{bmatrix} \tag{47}$$

has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{bmatrix} 6 - \lambda & -17\\ 8 & -6 - \lambda \end{bmatrix} = \lambda^2 + 100 = 0,$$

and hence has the complex conjugate eigenvalues λ_1 , $\lambda_2 = \pm 10i$. If $\mathbf{v} = \begin{bmatrix} a & b \end{bmatrix}^T$ is an eigenvector associated with $\lambda = 10i$, then the eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ yields

$$\begin{bmatrix} \mathbf{A} - 10i \cdot \mathbf{I} \end{bmatrix} \mathbf{v} = \begin{bmatrix} 6 - 10i & -17 \\ 8 & -6 - 10i \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Upon division of the second row by 2, this gives the two scalar equations

$$(6-10i)a - 17b = 0,$$

$$4a - (3+5i)b = 0.$$
(48)

each of which is satisfied by a = 3 + 5i and b = 4. Thus the desired eigenvector is $\mathbf{v} = \begin{bmatrix} 3+5i & 4 \end{bmatrix}^T$, with real and imaginary parts

$$\mathbf{a} = \begin{bmatrix} 3\\4 \end{bmatrix}$$
 and $\mathbf{b} = \begin{bmatrix} 5\\0 \end{bmatrix}$, (49)

respectively. Taking q = 10 in Eq. (45) therefore gives the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$:

$$\mathbf{x}(t) = c_1 \left(\begin{bmatrix} 3\\4 \end{bmatrix} \cos 10t - \begin{bmatrix} 5\\0 \end{bmatrix} \sin 10t \right) + c_2 \left(\begin{bmatrix} 5\\0 \end{bmatrix} \cos 10t + \begin{bmatrix} 3\\4 \end{bmatrix} \sin 10t \right)$$
$$= \begin{bmatrix} c_1(3\cos 10t - 5\sin 10t) + c_2(5\cos 10t + 3\sin 10t) \\ 4c_1\cos 10t + 4c_2\sin 10t \end{bmatrix}.$$
(50)

To solve the given initial value problem it remains only to determine values of the coefficients c_1 and c_2 . The initial condition $\mathbf{x}(0) = \begin{bmatrix} 4 & 2 \end{bmatrix}^T$ readily yields $c_1 = c_2 = \frac{1}{2}$, and with these values Eq. (50) becomes (in scalar form)

$$x_1(t) = 4\cos 10t - \sin 10t,$$

$$x_2(t) = 2\cos 10t + 2\sin 10t.$$
(51)

Figure 7.4.11 shows the trajectory given by Eq. (51) together with the initial point (4, 2).



FIGURE 7.4.11. Solution curve $x_1(t) = 4 \cos 10t - \sin 10t$, $x_2(t) = 2 \cos 10t + 2 \sin 10t$ for the initial value problem in Eq. (46).

This solution curve appears to be an ellipse rotated counterclockwise by the angle $\theta = \arctan \frac{2}{4} \approx 0.4636$. We can verify this by finding the equations of the solution curve relative to the rotated *u*- and *v*-axes shown in Fig. 7.4.11. By a standard formula from analytic geometry, these new equations are given by

$$u = x_1 \cos \theta + x_2 \sin \theta = \frac{2}{\sqrt{5}} x_1 + \frac{1}{\sqrt{5}} x_2,$$

$$v = -x_1 \sin \theta + x_2 \cos \theta = -\frac{1}{\sqrt{5}} x_1 + \frac{2}{\sqrt{5}} x_2.$$
(52)

In Problem 34 we ask you to substitute the expressions for x_1 and x_2 from Eq. (51) into Eq. (52), leading (after simplification) to

$$u = 2\sqrt{5}\cos 10t, \quad v = \sqrt{5}\sin 10t.$$
 (53)

Equation (53) not only confirms that the solution curve in Eq. (51) is indeed an ellipse rotated by the angle θ , but it also shows that the lengths of the semi-major and semi-minor axes of the ellipse are $2\sqrt{5}$ and $\sqrt{5}$, respectively.

Furthermore, we can demonstrate that any choice of initial point (apart from the origin) leads to a solution curve that is an ellipse rotated by the same angle θ and "concentric" (in an obvious sense) with the trajectory in Fig. 7.4.11 (see Problems 35–37). All these concentric rotated ellipses are centered at the origin (0,0), which is therefore called a **center** for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ whose coefficient matrix \mathbf{A} has pure imaginary eigenvalues. Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (47).

Further investigation: Geometric significance of the eigenvector. Our general solution in Eq. (50) was based upon the vectors **a** and **b** in Eq. (49), that is, the real and imaginary parts of the complex eigenvector $\mathbf{v} = \begin{bmatrix} 3+5i & 4 \end{bmatrix}^T$ of the matrix **A**. We might therefore expect **a** and **b** to have some clear geometric connection to the solution curve in Fig. 7.4.11. For example, we might guess that **a** and **b** would be parallel to the major and minor axes of the elliptical trajectory. However, it is clear from Fig. 7.4.12—which shows the vectors **a** and **b** together with the solution curve given by Eq. (51)—that this is not the case. Do the eigenvectors of **A**, then, play any *geometric* role in the phase portrait of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$?

The (affirmative) answer lies in the fact that any nonzero *real or complex* multiple of a complex eigenvector of the matrix **A** is still an eigenvector of **A** associated with that eigenvalue. Perhaps, then, if we multiply the eigenvector $\mathbf{v} = \begin{bmatrix} 3+5i & 4 \end{bmatrix}^T$ by a suitable nonzero complex constant *z*, the resulting eigenvector $\tilde{\mathbf{v}}$ will have real and imaginary parts $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ that can be readily identified with geometric features of the ellipse. To this end, let us multiply **v** by the complex scalar $z = \frac{1}{2}(1+i)$. (The reason for this particular choice will become clear shortly.) The resulting new complex eigenvector $\tilde{\mathbf{v}}$ of the matrix **A** is

$$\tilde{\mathbf{v}} = z \cdot \mathbf{v} = \frac{1}{2}(1+i) \cdot \begin{bmatrix} 3+5i\\4 \end{bmatrix} = \begin{bmatrix} -1+4i\\2+2i \end{bmatrix},$$

and has real and imaginary parts

$$\tilde{\mathbf{a}} = \begin{bmatrix} -1\\2 \end{bmatrix}$$
 and $\tilde{\mathbf{b}} = \begin{bmatrix} 4\\2 \end{bmatrix}$.



FIGURE 7.4.12. Solution curve for the initial value problem in Eq. (46) showing the vectors **a**, **b**, $\tilde{\mathbf{a}}$, and $\tilde{\mathbf{b}}$.

It is clear that the vector **b** is parallel to the major axis of our elliptical trajectory. Further, you can easily check that $\tilde{\mathbf{a}} \cdot \tilde{\mathbf{b}} = 0$, which means that $\tilde{\mathbf{a}}$ is perpendicular to $\tilde{\mathbf{b}}$, and hence is parallel to the minor axis of the ellipse, as Fig. 7.4.12 illustrates. Moreover, the length of $\tilde{\mathbf{b}}$ is twice that of $\tilde{\mathbf{a}}$, reflecting the fact that the lengths of the major and minor axes of the ellipse are in this same ratio. Thus for a matrix \mathbf{A} with pure imaginary eigenvalues, the complex eigenvector of \mathbf{A} used in the general solution (45)—if suitably chosen—is indeed of great significance to the geometry of the elliptical solution curves of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

How was the value $\frac{1}{2}(1 + i)$ chosen for the scalar *z*? In order that the real and imaginary parts $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ of $\tilde{\mathbf{v}} = z \cdot \mathbf{v}$ be parallel to the axes of the ellipse, at a minimum $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ must be perpendicular to each other. In Problem 38 we ask you to show that this condition is satisfied if and only if *z* is of the form $r(1 \pm i)$, where *r* is a nonzero real number, and that if *z* is chosen in this way, then $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ are in fact parallel to the axes of the ellipse. The value $r = \frac{1}{2}$ then aligns the lengths of $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ with those of the semi-minor and -major axes of the elliptical trajectory. More generally, we can show that given any eigenvector \mathbf{v} of a matrix \mathbf{A} with pure imaginary eigenvalues, there exists a constant *z* such that the real and imaginary parts $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ of the eigenvector $\tilde{\mathbf{v}} = z \cdot \mathbf{v}$ are parallel to the axes of the (elliptical) trajectories of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

Further investigation: Direction of flow. Figs. 7.4.11 and 7.4.12 suggest that the solution curve in Eq. (51) flows in a counterclockwise direction with increasing *t*. However, you can check that the matrix

$$-\mathbf{A} = \begin{bmatrix} -6 & 17\\ -8 & 6 \end{bmatrix}$$

has the same eigenvalues and eigenvectors as the matrix **A** in Eq. (47) itself, and yet (by the principle of time reversal) the trajectories of the system $\mathbf{x}' = -\mathbf{A}\mathbf{x}$ are identical to those of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ while flowing in the *opposite* direction, that is, clockwise. Clearly, mere knowledge of the eigenvalues and eigenvectors of the matrix **A** is not sufficient to predict the direction of flow of the elliptical trajectories of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ as *t* increases. How then can we determine this direction of flow?

One simple approach is to use the tangent vector \mathbf{x}' to monitor the direction in which the solution curves flow as they cross the positive x_1 -axis. If *s* is any positive number (so that the point (*s*, 0) lies on the positive x_1 -axis), and if the matrix \mathbf{A} is given by

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

then any trajectory for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ passing through (s, 0) satisfies

$$\mathbf{x}' = \mathbf{A}\mathbf{x} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} s \\ 0 \end{bmatrix} = \begin{bmatrix} as \\ cs \end{bmatrix} = s \begin{bmatrix} a \\ c \end{bmatrix}$$

at the point (s, 0). Therefore, at this point the direction of flow of the solution curve is a positive scalar multiple of the vector $\begin{bmatrix} a & c \end{bmatrix}^T$. Since *c* cannot be zero (see Problem 39), this vector either points "upward" into the first quadrant of the phase plane (if c > 0), or "downward" into the fourth quadrant (if c < 0). If upward, then the flow of the solution curve is counterclockwise; if downward, then clockwise. For the matrix **A** in Eq. (47), the vector $\begin{bmatrix} a & c \end{bmatrix}^T = \begin{bmatrix} 6 & 8 \end{bmatrix}^T$ points into the first quadrant because c = 8 > 0, thus indicating a counterclockwise direction of flow (as Figs. 7.4.11 and 7.4.12 suggest).

Complex Eigenvalues: Spiral Sinks and Sources

COMPLEX EIGENVALUES WITH NEGATIVE REAL PART: Now we assume that the eigenvalues of the matrix **A** are given by λ_1 , $\lambda_2 = p \pm iq$ with $q \neq 0$ and p < 0. In this case the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is given directly by Eq. (5):

$$\mathbf{x}(t) = c_1 e^{pt} (\mathbf{a} \cos qt - \mathbf{b} \sin qt) + c_2 e^{pt} (\mathbf{b} \cos qt + \mathbf{a} \sin qt),$$
(5)

where the vectors **a** and **b** have their usual meaning. Once again we begin with an example to gain an understanding of these solution curves.

Example 12 Solve the initial value problem

$$\mathbf{x}' = \begin{bmatrix} 5 & -17\\ 8 & -7 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 4\\ 2 \end{bmatrix}.$$
(54)

Solution The coefficient matrix

$$\mathbf{A} = \begin{bmatrix} 5 & -17\\ 8 & -7 \end{bmatrix} \tag{55}$$

has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{vmatrix} 5 - \lambda & -17 \\ 8 & -7 - \lambda \end{vmatrix} = (\lambda + 1)^2 + 100 = 0,$$

and hence has the complex conjugate eigenvalues λ_1 , $\lambda_2 = -1 \pm 10i$. If $\mathbf{v} = \begin{bmatrix} a & b \end{bmatrix}^T$ is an eigenvector associated with $\lambda = -1 + 10i$, then the eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ yields the same system (48) of equations found in Example 11:

$$(6-10i)a - 17b = 0,$$

$$4a - (3+5i)b = 0.$$
(48)

As in Example 11, each of these equations is satisfied by a = 3 + 5i and b = 4. Thus the desired eigenvector, associated with $\lambda_1 = -1 + 10i$, is once again $\mathbf{v} = \begin{bmatrix} 3+5i & 4 \end{bmatrix}^T$, with real and imaginary parts

$$\mathbf{a} = \begin{bmatrix} 3\\4 \end{bmatrix}$$
 and $\mathbf{b} = \begin{bmatrix} 5\\0 \end{bmatrix}$, (56)

respectively. Taking p = -1 and q = 10 in Eq. (5) therefore gives the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$:

$$\mathbf{x}(t) = c_1 e^{-t} \left(\begin{bmatrix} 3\\4 \end{bmatrix} \cos 10t - \begin{bmatrix} 5\\0 \end{bmatrix} \sin 10t \right) + c_2 e^{-t} \left(\begin{bmatrix} 5\\0 \end{bmatrix} \cos 10t + \begin{bmatrix} 3\\4 \end{bmatrix} \sin 10t \right)$$

=
$$\begin{bmatrix} c_1 e^{-t} (3\cos 10t - 5\sin 10t) + c_2 e^{-t} (5\cos 10t + 3\sin 10t) \\ 4c_1 e^{-t} \cos 10t + 4c_2 e^{-t} \sin 10t \end{bmatrix}.$$
 (57)

The initial condition $\mathbf{x}(0) = \begin{bmatrix} 4 & 2 \end{bmatrix}^T$ gives $c_1 = c_2 = \frac{1}{2}$ once again, and with these values Eq. (57) becomes (in scalar form)

$$x_1(t) = e^{-t} (4\cos 10t - \sin 10t),$$

$$x_2(t) = e^{-t} (2\cos 10t + 2\sin 10t).$$
(58)

Figure 7.4.13 shows the trajectory given by Eq. (58) together with the initial point (4, 2). It is noteworthy to compare this spiral trajectory with the elliptical trajectory in Eq. (51). The equations for $x_1(t)$ and $x_2(t)$ in (58) are obtained by multiplying their counterparts in (51) by the common factor e^{-t} , which is positive and decreasing with increasing t. Thus for positive values of t, the spiral trajectory is generated, so to speak, by standing at the origin and "reeling in" the point on the elliptical trajectory (51) as it is traced out. When t is negative, the picture is rather



FIGURE 7.4.13. Solution curve $x_1(t) = e^{-t} (4 \cos 10t - \sin 10t), x_2(t) = e^{-t} (2 \cos 10t + 2 \sin 10t)$ for the initial value problem in Eq. (54). The dashed and solid portions of the curve correspond to negative and positive values of *t*, respectively.

one of "casting away" the point on the ellipse farther out from the origin to create the corresponding point on the spiral.

Our gallery Fig. 7.4.16 shows a more complete set of solution curves, together with a direction field, for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (55). Because the solution curves all "spiral into" the origin, we call the origin in this a case a **spiral sink**.

COMPLEX EIGENVALUES WITH POSITIVE REAL PART: We conclude with the case where the eigenvalues of the matrix **A** are given by λ_1 , $\lambda_2 = p \pm iq$ with $q \neq 0$ and p > 0. Just as in the preceding case, the general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is given by Eq. (5):

$$\mathbf{x}(t) = c_1 e^{pt} (\mathbf{a} \cos qt - \mathbf{b} \sin qt) + c_2 e^{pt} (\mathbf{b} \cos qt + \mathbf{a} \sin qt).$$
(5)

An example will illustrate the close relation between the cases p > 0 and p < 0.

Example 13 Solve the initial value problem

 $\mathbf{x}' = \begin{bmatrix} -5 & 17\\ -8 & 7 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 4\\ 2 \end{bmatrix}.$ (59)

Solution Although we could directly apply the eigenvalue/eigenvector method as in previous cases (see Problem 40), here it is more convenient to notice that the coefficient matrix

$$\mathbf{A} = \begin{bmatrix} -5 & 17\\ -8 & 7 \end{bmatrix} \tag{60}$$

FIGURE 7.4.14. Solution curve $x_1(t) = e^t (4 \cos 10t + \sin 10t), x_2(t) = e^t (2 \cos 10t - 2 \sin 10t)$ for the initial value problem in Eq. (59). The dashed and solid portions of the curve correspond to negative and positive values of *t*, respectively.

is the negative of the matrix in Eq. (55) used in Example 12. By the principle of time reversal, therefore, the solution of the initial value problem (59) is given by simply replacing t with -t in the right-hand sides of the solution (58) of the initial value problem in that example:

$$x_1(t) = e^t (4\cos 10t + \sin 10t),$$

$$x_2(t) = e^t (2\cos 10t - 2\sin 10t).$$
(61)

Figure 7.4.14 shows the trajectory given by Eq. (61) together with the initial point (4, 2). Our gallery Fig. 7.4.16 shows this solution curve together with a direction field for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (60). Because the solution curve "spirals away from" the origin, we call the origin in this case a **spiral source**.

A 3-Dimensional Example

Figure 7.4.15 illustrates the space trajectories of solutions of the 3-dimensional system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with constant coefficient matrix

$$\mathbf{A} = \begin{bmatrix} 4 & 10 & 0 \\ -5 & -6 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$
 (62)

To portray the motion in space of a point $\mathbf{x}(t)$ moving on a trajectory of this system, we can regard this trajectory as a necklace string on which colored beads are placed to mark its successive positions at fixed increments of time (so the point is moving fastest where the spacing between beads is greatest). In order to aid the eye in following the moving point's progress, the size of the beads decreases continuously with the passage of time and motion along the trajectory.



FIGURE 7.4.15. Three-dimensional trajectories for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with the matrix **A** given by Eq. (62).

The matrix **A** has the single real eigenvalue -1 with the single (real) eigenvector $\begin{bmatrix} 0 & 0 & 1 \end{bmatrix}^T$ and the complex conjugate eigenvalues $-1 \pm 5i$. The negative real eigenvalue corresponds to trajectories that lie on the x_3 -axis and approach the origin as $t \to 0$ (as illustrated by the beads on the vertical axis of the figure). Thus the origin (0, 0, 0) is a sink that "attracts" all the trajectories of the system.

The complex conjugate eigenvalues with negative real part correspond to trajectories in the horizontal x_1x_2 -plane that spiral around the origin while approaching it. Any other trajectory—one which starts at a point lying neither on the *z*-axis nor in the x_1x_2 -plane—combines the preceding behaviors by spiraling around the surface of a cone while approaching the origin at its vertex.

Gallery of Typical Phase Portraits for the System x' = Ax: Nodes



Proper Nodal Source: A repeated positive real eigenvalue with two linearly independent eigenvectors.





Proper Nodal Sink: A repeated negative real eigenvalue with two linearly independent eigenvectors.



Improper Nodal Source: Distinct positive real eigenvalues (left) or a repeated positive real eigenvalue without two linearly independent eigenvectors (right).





Improper Nodal Sink: Distinct negative real eigenvalues (left) or a repeated negative real eigenvalue without two linearly independent eigenvectors (right).

FIGURE 7.4.16. Gallery of typical phase plane portraits for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.
Gallery of Typical Phase Portraits for the System x' = Ax: Saddles, Centers, Spirals, and Parallel Lines



Saddle Point: Real eigenvalues of opposite sign.



Center: Pure imaginary eigenvalues.



Spiral Source: Complex conjugate eigenvalues with positive real part.



Parallel Lines: One zero and one negative real eigenvalue. (If the nonzero eigenvalue is positive, then the trajectories flow *away* from the dotted line.)

FIGURE 7.4.16. (Continued)



Spiral Sink: Complex conjugate eigenvalues with negative real part.



Parallel Lines: A repeated zero eigenvalue without two linearly independent eigenvectors.

7.4 **Problems**

For each of the systems in Problems 1 through 16 in Section 7.3, categorize the eigenvalues and eigenvectors of the coefficient matrix \mathbf{A} according to Fig. 7.4.16 and sketch the phase portrait of the system by hand. Then use a computer system or graphing calculator to check your answer.

The phase portraits in Problems 17 through 28 correspond to linear systems of the form $\mathbf{x}' = \mathbf{A}\mathbf{x}$ in which the matrix \mathbf{A} has two linearly independent eigenvectors. Determine the nature of the eigenvalues and eigenvectors of each system. For example, you may discern that the system has pure imaginary eigenvalues, or that it has real eigenvalues of opposite sign; that an eigenvector associated with the positive eigenvalue is roughly $\begin{bmatrix} 2 & -1 \end{bmatrix}^T$, etc.





420 CHAPTER 7 LINEAR SYSTEMS OF DIFFERENTIAL EQUATIONS





29. We can give a simpler description of the general solution

$$\mathbf{x}(t) = c_1 \begin{bmatrix} -1\\ 6 \end{bmatrix} e^{-2t} + c_2 \begin{bmatrix} 1\\ 1 \end{bmatrix} e^{5t}$$
(9)

of the system

$$\mathbf{x}' = \begin{bmatrix} 4 & 1\\ 6 & -1 \end{bmatrix} \mathbf{x}$$

in Example 1 by introducing the oblique *uv*-coordinate system indicated in Fig. 7.4.17, in which the *u*- and *v*- axes are determined by the eigenvectors $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 6 \end{bmatrix}$ and $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, respectively.



FIGURE 7.4.17. The oblique uv-coordinate system determined by the eigenvectors v_1 and v_2 .

The *uv*-coordinate functions u(t) and v(t) of the moving point $\mathbf{x}(t)$ are simply its distances from the origin measured in the directions parallel to \mathbf{v}_1 and \mathbf{v}_2 . It follows from (9) that a trajectory of the system is described by

$$u(t) = u_0 e^{-2t}, \quad v(t) = v_0 e^{5t}$$
 (63)

where $u_0 = u(0)$ and $v_0 = v(0)$. (a) Show that if $v_0 = 0$, then this trajectory lies on the *u*-axis, whereas if $u_0 = 0$, then it lies on the *v*-axis. (b) Show that if u_0 and v_0 are both nonzero, then a "Cartesian" equation of the parametric curve in Eq. (63) is given by $v = Cu^{-5/2}$.

30. Use the chain rule for vector-valued functions to verify the principle of time reversal.

In Problems 31–33 A represents a 2×2 matrix.

- 31. Use the definitions of eigenvalue and eigenvector (Section 7.3) to prove that if λ is an eigenvalue of A with associated eigenvector v, then -λ is an eigenvalue of the matrix -A with associated eigenvector v. Conclude that if A has positive eigenvalues 0 < λ₂ < λ₁ with associated eigenvectors v₁ and v₂, then -A has negative eigenvalues -λ₁ < -λ₂ < 0 with the same associated eigenvectors.
- 32. Show that the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ has constant solutions other than $\mathbf{x}(t) \equiv \mathbf{0}$ if and only if there exists a (constant) vector $\mathbf{x} \neq \mathbf{0}$ with $\mathbf{A}\mathbf{x} = \mathbf{0}$. (It is shown in linear algebra that such a vector \mathbf{x} exists exactly when det(\mathbf{A}) = 0.)
- **33.** (a) Show that if **A** has the repeated eigenvalue λ with two linearly independent associated eigenvectors, then every nonzero vector **v** is an eigenvector of **A**. (*Hint:* Express **v** as a linear combination of the linearly independent eigenvectors and multiply both sides by **A**.) (b) Conclude that **A** must be given by Eq. (22). (*Suggestion:* In the equation $A\mathbf{v} = \lambda \mathbf{v}$ take $\mathbf{v} = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ and $\mathbf{v} = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$.)
- **34.** Verify Eq. (53) by substituting the expressions for $x_1(t)$ and $x_2(t)$ from Eq. (51) into Eq. (52) and simplifying.

Problems 35–37 show that all nontrivial solution curves of the system in Example 11 are ellipses rotated by the same angle as the trajectory in Fig. 7.4.11.

35. The system in Example 11 can be rewritten in scalar form as

$$\begin{aligned} x_1' &= 6x_1 - 17x_2, \\ x_2' &= 8x_1 - 6x_2, \end{aligned}$$

leading to the first-order differential equation

$$\frac{dx_2}{dx_1} = \frac{dx_2/dt}{dx_1/dt} = \frac{8x_1 - 6x_2}{6x_1 - 17x_2}$$

or, in differential form,

$$(6x_2 - 8x_1) \, dx_1 + (6x_1 - 17x_2) \, dx_2 = 0$$

Verify that this equation is exact with general solution

$$-4x_1^2 + 6x_1x_2 - \frac{17}{2}x_2^2 = k, (64)$$

where k is a constant.

36. In analytic geometry it is shown that the general quadratic equation

$$Ax_1^2 + Bx_1x_2 + Cx_2^2 = k (65)$$

represents an ellipse centered at the origin if and only if Ak > 0 and the *discriminant* $B^2 - 4AC < 0$. Show that Eq. (64) satisfies these conditions if k < 0, and thus conclude that all nondegenerate solution curves of the system in Example 11 are elliptical.

37. It can be further shown that Eq. (65) represents in general a conic section rotated by the angle θ given by

$$\tan 2\theta = \frac{B}{A-C}$$

Show that this formula applied to Eq. (64) leads to the angle θ = arctan $\frac{2}{4}$ found in Example 11, and thus conclude that all elliptical solution curves of the system in Example 11 are rotated by the same angle θ . (*Suggestion:* You may find useful the double-angle formula for the tangent function.)

- **38.** Let $\mathbf{v} = \begin{bmatrix} 3+5i & 4 \end{bmatrix}^T$ be the complex eigenvector found in Example 11 and let z be a complex number. (a) Show that the real and imaginary parts $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$, respectively, of the vector $\tilde{\mathbf{v}} = z \cdot \mathbf{v}$ are perpendicular if and only if $z = r(1 \pm i)$ for some nonzero real number r. (b) Show that if this is the case, then $\tilde{\mathbf{a}}$ and $\tilde{\mathbf{b}}$ are parallel to the axes of the elliptical trajectory found in Example 11 (as Fig. 7.4.12 indicates).
- **39.** Let **A** denote the 2×2 matrix

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

(a) Show that the characteristic equation of A (Eq. (3), Section 6.1) is given by

$$\lambda^2 - (a+d)\lambda + (ad-bc) = 0.$$

- (b) Suppose that the eigenvalues of **A** are pure imaginary. Show that the *trace* $T(\mathbf{A}) = a + d$ of **A** must be zero and that the *determinant* $D(\mathbf{A}) = ad - bc$ must be positive. Conclude that $c \neq 0$.
- **40.** Use the eigenvalue/eigenvector method to confirm the solution in Eq. (61) of the initial value problem in Eq. (59).

7.4 Application **Dynamic Phase Plane Graphics**

Using computer systems we can "bring to life" the static gallery of phase portraits in Fig. 7.4.16 by allowing initial conditions, eigenvalues, and even eigenvectors to vary in "real time." Such dynamic phase plane graphics afford additional insight into the relationship between the algebraic properties of the 2×2 matrix **A** and the phase plane portrait of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

For example, the basic linear system

$$\frac{dx_1}{dt} = -x_1,$$

$$\frac{dx_2}{dt} = -kx_2 \quad (k \text{ a nonzero constant}),$$

has general solution

$$x_1(t) = ae^{-t}, \quad x_2(t) = be^{-kt}$$

where (a, b) is the initial point. If $a \neq 0$, then we can write

$$x_2 = be^{-kt} = \frac{b}{a^k}(ae^{-t}) = cx_1^k,$$
(1)

where $c = b/a^k$. A version of the *Maple* commands

```
with(plots):
createPlot := proc(k)
    soln := plot([exp(-t), exp(-k*t),
        t = -10..10], x = -5..5, y = -5..5):
return display(soln):
end proc:
Explore(createPlot(k),
    parameters = [k = -2.0..2.0])
```

produces Fig. 7.4.18, which allows the user to vary the parameter k continuously from k = -2 to k = 2, thus showing dynamically the changes in the solution curves (1) in response to changes in k.

Figure 7.4.19 shows snapshots of the interactive display in Fig. 7.4.18 corresponding to the values -1, $\frac{1}{2}$, and 2 for the parameter k. Based on this progression, how would you expect the solution curves in Eq. (1) to look when k = 1? Does Eq. (1) corroborate your guess?



FIGURE 7.4.19. Snapshots of the interactive display in Fig. 7.4.18 with the initial conditions held fixed and the parameter k equal to -1, $\frac{1}{2}$, and 2, respectively.

As another example, a version of the *Mathematica* commands

```
a = {{-5, 17}, {-8, 7}};
x[t_] := {x1[t], x2[t]};
Manipulate[
    soln = DSolve[{x'[t] == a.x[t],
        x[0] == pt[[1]]}, x[t], t];
ParametricPlot[x[t]/.soln, {t, -3.5, 10},
```



FIGURE 7.4.18. Interactive display of the solution curves in Eq. (1). Using the slider, the value of k can be varied continuously from -2 to 2.

was used to generate Fig. 7.4.20, which (like Figs. 7.4.13 and 7.4.14) shows the solution curve of the initial value problem

$$\mathbf{x}' = \begin{bmatrix} -5 & 17\\ -8 & 7 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 4\\ 2 \end{bmatrix}$$
(2)

from Example 13 of the preceding section. However, in Fig. 7.4.20 the initial condition (4, 2) is attached to a "locator point" which can be freely dragged to any desired position in the phase plane, with the corresponding solution curve being instantly redrawn—thus illustrating dynamically the effect of varying the initial conditions.





FIGURE 7.4.20. Interactive display of the initial value problem in Eq. (2). As the "locator point" is dragged to different positions, the solution curve is immediately redrawn, showing the effect of changing the initial conditions.

FIGURE 7.4.21. Interactive display of the initial value problem $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with **A** given by Eq. (3). Both the initial conditions and the value of the parameter *k* can be varied dynamically.

Finally, Fig. 7.4.21 shows a more sophisticated, yet perhaps more revealing, demonstration. As you can verify, the matrix

$$\mathbf{A} = \frac{1}{10} \begin{bmatrix} k+9 & 3-3k\\ 3-3k & 9k+1 \end{bmatrix}$$
(3)

has the *variable* eigenvalues 1 and k but with *fixed* associated eigenvectors $\begin{bmatrix} 3 & 1 \end{bmatrix}^T$ and $\begin{bmatrix} 1 & -3 \end{bmatrix}^T$, respectively. Figure 7.4.21, which was generated by a version of the *Mathematica* commands

shows the phase portrait of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with \mathbf{A} given by Eq. (3). Not only are the initial conditions of the individual trajectories controlled independently by the "locator points," but using the slider we can also vary the value of k continuously from -1 to 1, with the solution curves being instantly redrawn. Thus for a fixed value of k we can experiment with changing initial conditions throughout the phase plane, or, conversely, we can hold the initial conditions fixed and observe the effect of changing the value of k.

As a further example of what such a display can reveal, Fig. 7.4.22 consists of a series of snapshots of Fig. 7.4.21 where the initial conditions are held fixed and k progresses through the specific values -1, -0.25, 0, 0.5, 0.65, and 1. The result is a "video" showing stages in a transition from a saddle point with "hyperbolic" trajectories, to a pair of parallel lines, to an improper nodal source with "parabolic" trajectories, and finally to the exploding star pattern of a proper nodal source with straight-line trajectories. Perhaps these frames provide a new interpretation of the description "dynamical system" for a collection of interdependent differential equations.



FIGURE 7.4.22. Snapshots of the interactive display in Fig. 7.4.21 with the initial conditions held fixed and the parameter k increasing from -1 to 1.

7.5 Second-Order Systems and Mechanical Applications*

In this section we apply the matrix methods of Sections 7.2 and 7.3 to investigate the oscillations of typical mass-and-spring systems having two or more degrees of freedom. Our examples are chosen to illustrate phenomena that are generally characteristic of complex mechanical systems.

* This optional section may be omitted without loss of continuity. It provides a sample of the more technical applications of eigenvalues to physics and engineering problems.

7.5 SECOND-ORDER SYSTEMS AND MECHANICAL APPLICATIO



FIGURE 7.5.1. Three spring-coupled masses.

Figure 7.5.1 shows three masses connected to each other and to two walls by the four indicated springs. We assume that the masses slide without friction and that each spring obeys Hooke's law—its extension or compression x and force F of reaction are related by the formula F = -kx. We further assume that when the system is at equilibrium (so the masses are stationary) none of the springs is stretched or compressed. If the rightward displacements x_1 , x_2 , and x_3 of the three masses (from their respective equilibrium positions) are all positive, then

- The first spring is stretched the distance x_1 ;
- The second spring is stretched the distance $x_2 x_1$;
- The third spring is stretched the distance $x_3 x_2$;
- The fourth spring is compressed the distance *x*₃.

Therefore, application of Newton's law F = ma to the three masses (as in Example 1 of Section 7.1) yields their equations of motion:

$$m_1 x_1'' = -k_1 x_1 + k_2 (x_2 - x_1),$$

$$m_2 x_2'' = -k_2 (x_2 - x_1) + k_3 (x_3 - x_2),$$

$$m_3 x_3'' = -k_3 (x_3 - x_2) - k_4 x_3.$$
(1)

Although we assumed in writing these equations that the displacements of the masses are all positive, they actually follow similarly from Hooke's and Newton's laws, whatever the signs of these displacements.

In terms of the **displacement vector x** = $\begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix}^T$, the **mass matrix**

$$\mathbf{M} = \begin{bmatrix} m_1 & 0 & 0\\ 0 & m_2 & 0\\ 0 & 0 & m_3 \end{bmatrix}$$
(2)

and the stiffness matrix

$$\mathbf{K} = \begin{bmatrix} -(k_1 + k_2) & k_2 & 0\\ k_2 & -(k_2 + k_3) & k_3\\ 0 & k_3 & -(k_3 + k_4) \end{bmatrix},$$
(3)

the system in (1) takes the matrix form

$$\mathbf{M}\mathbf{x}'' = \mathbf{K}\mathbf{x}.\tag{4}$$



FIGURE 7.5.2. A system of *n* spring-coupled masses.

The notation in Eqs. (1) through (4) generalizes in a natural way to the system of n spring-coupled masses shown in Fig. 7.5.2. We need only write

$$\mathbf{M} = \begin{bmatrix} m_1 & 0 & \cdots & 0 \\ 0 & m_2 & \cdot & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & m_n \end{bmatrix}$$
(5)

and

>

$$\mathbf{K} = \begin{bmatrix} -(k_1 + k_2) & k_2 & 0 & \cdots & 0 \\ k_2 & -(k_2 + k_3) & k_3 & \cdots & 0 \\ 0 & k_3 & -(k_3 + k_4) & \cdots & 0 \\ 0 & 0 & k_4 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & -(k_{n-1} + k_n) & k_n \\ 0 & 0 & \cdots & k_n & -(k_n + k_{n+1}) \end{bmatrix}$$
(6)

for the mass and stiffness matrices in Eq. (4).

The diagonal matrix **M** is obviously nonsingular; to get its inverse \mathbf{M}^{-1} we need only replace each diagonal element with its reciprocal. Hence multiplication of each side in Eq. (4) by \mathbf{M}^{-1} yields the homogeneous **second-order system**

$$\mathbf{x}^{\prime\prime} = \mathbf{A}\mathbf{x},\tag{7}$$

where $\mathbf{A} = \mathbf{M}^{-1}\mathbf{K}$. There is a wide variety of *frictionless* mechanical systems for which a displacement or position vector \mathbf{x} , a nonsingular mass matrix \mathbf{M} , and a stiffness matrix \mathbf{K} satisfying Eq. (4) can be defined.

Solution of Second-Order Systems

To seek a solution of Eq. (7), we substitute (as in Section 7.3 for a first-order system) a trial solution of the form

$$\mathbf{x}(t) = \mathbf{v}e^{\alpha t},\tag{8}$$

where **v** is a constant vector. Then $\mathbf{x}'' = \alpha^2 \mathbf{v} e^{\alpha t}$, so substitution of Eq. (8) in (7) gives

$$\alpha^2 \mathbf{v} e^{\alpha t} = \mathbf{A} \mathbf{v} e^{\alpha t},$$

which implies that

$$\mathbf{A}\mathbf{v} = \alpha^2 \mathbf{v}.\tag{9}$$

Therefore $\mathbf{x}(t) = \mathbf{v}e^{\alpha t}$ is a solution of $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ if and only if $\alpha^2 = \lambda$, an eigenvalue of the matrix \mathbf{A} , and \mathbf{v} is an associated eigenvector.

If $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ models a mechanical system, then it is typical that the eigenvalues of **A** are *negative* real numbers. If

$$\alpha^2 = \lambda = -\omega^2 < 0,$$

then $\alpha = \pm \omega i$. In this case the solution given by Eq. (8) is

$$\mathbf{x}(t) = \mathbf{v}e^{i\omega t} = \mathbf{v}(\cos\omega t + i\sin\omega t).$$

The real and imaginary parts

$$\mathbf{x}_1(t) = \mathbf{v}\cos\omega t$$
 and $\mathbf{x}_2(t) = \mathbf{v}\sin\omega t$ (10)

of $\mathbf{x}(t)$ are then linearly independent *real-valued* solutions of the system. This analysis leads to the following theorem.

THEOREM 1 Second-Order Homogeneous Linear Systems

If the $n \times n$ matrix **A** has distinct negative eigenvalues $-\omega_1^2, -\omega_2^2, \ldots, -\omega_n^2$ with associated [real] eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$, then a general solution of

$$\mathbf{x}^{\prime\prime} = \mathbf{A}\mathbf{x}$$

is given by

•
$$\mathbf{x}(t) = \sum_{i=1}^{n} (a_i \cos \omega_i t + b_i \sin \omega_i t) \mathbf{v}_i$$
(11)

with a_i and b_i arbitrary constants. In the special case of a nonrepeated zero eigenvalue λ_0 with associated eigenvector \mathbf{v}_0 ,

$$\mathbf{x}_0(t) = (a_0 + b_0 t) \mathbf{v}_0 \tag{12}$$

is the corresponding part of the general solution.

Remark The nonzero vector \mathbf{v}_0 is an eigenvector corresponding to $\lambda_0 = 0$ provided that $\mathbf{A}\mathbf{v}_0 = \mathbf{0}$. If $\mathbf{x}(t) = (a_0 + b_0 t)\mathbf{v}_0$, then

$$\mathbf{x}'' = 0 \cdot \mathbf{v}_0 = (a_0 + b_0 t) \cdot \mathbf{0} = (a_0 + b_0 t) \cdot (\mathbf{A} \mathbf{v}_0) = \mathbf{A} \mathbf{x},$$

thus verifying the form in Eq. (12).

Mass-spring system Consider the mass-and-spring system with n = 2 shown in Fig. 7.5.3. Because there is no third spring connected to a right-hand wall, we set $k_3 = 0$. If $m_1 = 2$, $m_2 = 1$, $k_1 = 100$, and $k_2 = 50$, then the equation $\mathbf{Mx''} = \mathbf{Kx}$ is

$$\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x}^{\prime\prime} = \begin{bmatrix} -150 & 50 \\ 50 & -50 \end{bmatrix} \mathbf{x},$$
(13)

which reduces to $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ with

$$\mathbf{A} = \begin{bmatrix} -75 & 25\\ 50 & -50 \end{bmatrix}.$$

The characteristic equation of A is

$$(-75 - \lambda)(-50 - \lambda) - 50 \cdot 25 = \lambda^2 + 125\lambda + 2500$$

= $(\lambda + 25)(\lambda + 100) = 0.000$

so **A** has the negative eigenvalues $\lambda_1 = -25$ and $\lambda_2 = -100$. By Theorem 1, the system in (13) therefore has solutions with [circular] frequencies $\omega_1 = 5$ and $\omega_2 = 10$.

CASE 1: $\lambda_1 = -25$. The eigenvector equation $(A - \lambda I)v = 0$ is

$$\begin{bmatrix} -50 & 25\\ 50 & -25 \end{bmatrix} \begin{bmatrix} a\\ b \end{bmatrix} = \begin{bmatrix} 0\\ 0 \end{bmatrix},$$

so an eigenvector associated with $\lambda_1 = -25$ is $\mathbf{v}_1 = \begin{bmatrix} 1 & 2 \end{bmatrix}^T$.



FIGURE 7.5.3. The mass-and-spring system of Example 1.



$$\begin{bmatrix} 25 & 25\\ 50 & 50 \end{bmatrix} \begin{bmatrix} a\\ b \end{bmatrix} = \begin{bmatrix} 0\\ 0 \end{bmatrix}$$

so an eigenvector associated with $\lambda_2 = -100$ is $\mathbf{v}_2 = \begin{bmatrix} 1 & -1 \end{bmatrix}^T$.

By Eq. (11) it follows that a general solution of the system in (13) is given by

$$\mathbf{x}(t) = (a_1 \cos 5t + b_1 \sin 5t)\mathbf{v}_1 + (a_2 \cos 10t + b_2 \sin 10t)\mathbf{v}_2.$$
(14)

The two terms on the right in Eq. (14) represent **free oscillations** of the mass-and-spring system. They describe the physical system's two **natural modes of oscillation** at its two [circular] **natural frequencies** $\omega_1 = 5$ and $\omega_2 = 10$. The natural mode

$$\mathbf{x}_1(t) = (a_1 \cos 5t + b_1 \sin 5t)\mathbf{v}_1 = c_1 \cos(5t - \alpha_1) \begin{bmatrix} 1\\2 \end{bmatrix}$$

(with $c_1 = \sqrt{a_1^2 + b_1^2}$, $\cos \alpha_1 = a_1/c_1$, and $\sin \alpha_1 = b_1/c_1$) has the scalar component equations

$$x_1(t) = c_1 \cos(5t - \alpha_1), x_2(t) = 2c_1 \cos(5t - \alpha_1),$$
(15)

and therefore describes a free oscillation in which the two masses move in synchrony in the same direction and with the same frequency $\omega_1 = 5$, but with the amplitude of motion of m_2 twice that of m_1 (see Fig. 7.5.4). The natural mode

$$\mathbf{x}_2(t) = (a_2 \cos 10t + b_2 \sin 10t)\mathbf{v}_2 = c_2 \cos(10t - \alpha_2) \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

has the scalar component equations

$$\begin{aligned} x_1(t) &= c_2 \cos(10t - \alpha_2), \\ x_2(t) &= -c_2 \cos(10t - \alpha_2), \end{aligned}$$
 (16)

and therefore describes a free oscillation in which the two masses move in synchrony in opposite directions with the same frequency $\omega_2 = 10$ and with equal amplitudes of oscillation (see Fig. 7.5.5).

Railway cars Figure 7.5.6 shows three railway cars connected by buffer springs that react when compressed, but disengage instead of stretching. With n = 3, $k_2 = k_3 = k$, and $k_1 = k_4 = 0$ in Eqs. (2) through (4), we get the system

$$\begin{bmatrix} m_1 & 0 & 0\\ 0 & m_2 & 0\\ 0 & 0 & m_3 \end{bmatrix} \mathbf{x}'' = \begin{bmatrix} -k & k & 0\\ k & -2k & k\\ 0 & k & -k \end{bmatrix} \mathbf{x},$$
 (17)

which is equivalent to

 $\mathbf{x}'' = \begin{bmatrix} -c_1 & c_1 & 0\\ c_2 & -2c_2 & c_2\\ 0 & c_3 & -c_3 \end{bmatrix} \mathbf{x}$ (18)

with

$$c_i = \frac{k}{m_i}$$
 (*i* = 1, 2, 3). (19)

If we assume further that $m_1 = m_3$, so that $c_1 = c_3$, then a brief computation gives

$$-\lambda(\lambda + c_1)(\lambda + c_1 + 2c_2) = 0$$
(20)



FIGURE 7.5.4. Oscillations in the same direction with frequency $\omega_1 = 5$; the amplitude of motion of mass 2 is twice that of mass 1.



FIGURE 7.5.5. Oscillations in opposite directions with frequency $\omega_2 = 10$; the amplitudes of motion of the two masses are the same.

 k_2 k_3 m_3

Example 2

FIGURE 7.5.6. The three railway cars of Example 2.

for the characteristic equation of the coefficient matrix A in Eq. (18). Hence the matrix A has eigenvalues

$$\lambda_1 = 0, \quad \lambda_2 = -c_1, \quad \lambda_3 = -c_1 - 2c_2$$
 (21a)

corresponding to the natural frequencies

$$\omega_1 = 0, \quad \omega_2 = \sqrt{c_1}, \quad \omega_3 = \sqrt{c_1 + 2c_2}$$
 (21b)

of the physical system.

For a numerical example, suppose that the first and third railway cars weigh 12 tons each, that the middle car weighs 8 tons, and that the spring constant is k = 1.5 tons/ft; i.e., k = 3000 lb/ft. Then, using fps units with mass measured in slugs (a weight of 32 pounds has a mass of 1 slug), we have

$$m_1 = m_3 = 750, \quad m_2 = 500,$$

and

$$c_1 = \frac{3000}{750} = 4, \quad c_2 = \frac{3000}{500} = 6.$$

Hence the coefficient matrix A is

$$\mathbf{A} = \begin{bmatrix} -4 & 4 & 0\\ 6 & -12 & 6\\ 0 & 4 & -4 \end{bmatrix},$$
 (22)

and the eigenvalue-frequency pairs given by (21a) and (21b) are $\lambda_1 = 0$, $\omega_1 = 0$; $\lambda_2 = -4$, $\omega_2 = 2$; and $\lambda_3 = -16$, $\omega_3 = 4$.

CASE 1: $\lambda_1 = 0$, $\omega_1 = 0$. The eigenvector equation $(A - \lambda I)v = 0$ is

$$\mathbf{A}\mathbf{v} = \begin{bmatrix} -4 & 4 & 0\\ 6 & -12 & 6\\ 0 & 4 & -4 \end{bmatrix} \begin{bmatrix} a\\ b\\ c \end{bmatrix} = \begin{bmatrix} 0\\ 0\\ 0 \end{bmatrix},$$

so it is clear that $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^T$ is an eigenvector associated with $\lambda_1 = 0$. According to Theorem 1, the corresponding part of a general solution of $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ is

$$\mathbf{x}_1(t) = (a_1 + b_1 t) \mathbf{v}_1.$$

CASE 2: $\lambda_2 = -4$, $\omega_2 = 2$. The eigenvector equation $(A - \lambda I)v = 0$ is

$$(\mathbf{A} + 4\mathbf{I})\mathbf{v} = \begin{bmatrix} 0 & 4 & 0 \\ 6 & -8 & 6 \\ 0 & 4 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

so it is clear that $\mathbf{v}_2 = \begin{bmatrix} 1 & 0 & -1 \end{bmatrix}^T$ is an eigenvector associated with $\lambda_2 = -4$. According to Theorem 1, the corresponding part of a general solution of $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ is

$$\mathbf{x}_2(t) = (a_2 \cos 2t + b_2 \sin 2t)\mathbf{v}_2.$$

CASE 3: $\lambda_3 = -16$, $\omega_3 = 4$. The eigenvector equation $(A - \lambda I)v = 0$ is

$$(\mathbf{A} + 16\mathbf{I})\mathbf{v} = \begin{bmatrix} 12 & 4 & 0\\ 6 & 4 & 6\\ 0 & 4 & 12 \end{bmatrix} \begin{bmatrix} a\\ b\\ c \end{bmatrix} = \begin{bmatrix} 0\\ 0\\ 0 \end{bmatrix},$$

so it is clear that $\mathbf{v}_3 = \begin{bmatrix} 1 & -3 & 1 \end{bmatrix}^T$ is an eigenvector associated with $\lambda_3 = -16$. According to Theorem 1, the corresponding part of a general solution of $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ is

$$\mathbf{x}_3(t) = (a_3\cos 4t + b_3\sin 4t)\mathbf{v}_3.$$

The general solution $\mathbf{x} = \mathbf{x}_1 + \mathbf{x}_2 + \mathbf{x}_3$ of $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ is therefore given by

$$\mathbf{x}(t) = a_1 \begin{bmatrix} 1\\1\\1 \end{bmatrix} + b_1 t \begin{bmatrix} 1\\1\\1 \end{bmatrix} + a_2 \begin{bmatrix} 1\\0\\-1 \end{bmatrix} \cos 2t + b_2 \begin{bmatrix} 1\\0\\-1 \end{bmatrix} \sin 2t + a_3 \begin{bmatrix} 1\\-3\\1 \end{bmatrix} \cos 4t + b_3 \begin{bmatrix} 1\\-3\\1 \end{bmatrix} \sin 4t.$$
(23)

To determine a particular solution, let us suppose that the leftmost car is moving to the right with velocity v_0 and at time t = 0 strikes the other two cars, which are together but at rest. The corresponding initial conditions are

$$x_1(0) = x_2(0) = x_3(0) = 0,$$
(24a)

$$x'_{1}(0) = v_{0}, \quad x'_{2}(0) = x'_{3}(0) = 0.$$
 (24b)

Then substitution of (24a) in (23) gives the scalar equations

$$a_1 + a_2 + a_3 = 0,$$

 $a_1 - 3a_3 = 0,$
 $a_1 - a_2 + a_3 = 0,$

which readily yield $a_1 = a_2 = a_3 = 0$. Hence the position functions of the three cars are

$$x_{1}(t) = b_{1}t + b_{2}\sin 2t + b_{3}\sin 4t,$$

$$x_{2}(t) = b_{1}t - 3b_{3}\sin 4t,$$

$$x_{3}(t) = b_{1}t - b_{2}\sin 2t + b_{3}\sin 4t,$$

(25)

and their velocity functions are

$$x'_{1}(t) = b_{1} + 2b_{2}\cos 2t + 4b_{3}\cos 4t,$$

$$x'_{2}(t) = b_{1} - 12b_{3}\cos 4t,$$

$$x'_{3}(t) = b_{1} - 2b_{2}\cos 2t + 4b_{3}\cos 4t.$$
(26)

Substitution of (24b) in (26) gives the equations

$$b_1 + 2b_2 + 4b_3 = v_0,$$

$$b_1 - 12b_3 = 0,$$

$$b_1 - 2b_2 + 4b_3 = 0$$

that readily yield $b_1 = \frac{3}{8}v_0$, $b_2 = \frac{1}{4}v_0$, and $b_3 = \frac{1}{32}v_0$. Finally, the position functions in (25) are

$$x_{1}(t) = \frac{1}{32}v_{0}(12t + 8\sin 2t + \sin 4t),$$

$$x_{2}(t) = \frac{1}{32}v_{0}(12t - 3\sin 4t),$$

$$x_{3}(t) = \frac{1}{32}v_{0}(12t - 8\sin 2t + \sin 4t).$$

(27)

But these equations hold only so long as the two buffer springs remain compressed; that is, while both

$$x_2 - x_1 < 0$$
 and $x_3 - x_2 < 0$.

To discover what this implies about t, we compute

$$\begin{aligned} x_2(t) - x_1(t) &= \frac{1}{32} v_0 (-8 \sin 2t - 4 \sin 4t) \\ &= -\frac{1}{32} v_0 (8 \sin 2t + 8 \sin 2t \cos 2t) \\ &= -\frac{1}{4} v_0 (\sin 2t) (1 + \cos 2t) \end{aligned}$$

7.5 SECOND-ORDER SYSTEMS AND MECHANICAL APPLICATIONS1

and, similarly,

$$x_3(t) - x_2(t) = -\frac{1}{4}v_0(\sin 2t)(1 - \cos 2t).$$

It follows that $x_2 - x_1 < 0$ and $x_3 - x_2 < 0$ until $t = \pi/2 \approx 1.57$ (seconds), at which time the equations in (26) and (27) give the values

$$x_1\left(\frac{\pi}{2}\right) = x_2\left(\frac{\pi}{2}\right) = x_3\left(\frac{\pi}{2}\right) = \frac{3\pi v_0}{16},$$
$$x_1'\left(\frac{\pi}{2}\right) = x_2'\left(\frac{\pi}{2}\right) = 0, \quad x_3'\left(\frac{\pi}{2}\right) = v_0.$$

We conclude that the three railway cars remain engaged and moving to the right until disengagement occurs at time $t = \pi/2$. Thereafter, cars 1 and 2 remain at rest (!), while car 3 continues to the right with speed v_0 . If, for instance, $v_0 = 48$ feet per second (about 33 miles per hour), then the three cars travel a distance of $9\pi \approx 28.27$ (ft) during their 1.57 seconds of engagement, and

$$x_1(t) = x_2(t) = 9\pi, \quad x_3(t) = 48t - 15\pi$$
 (27')

for $t > \pi/2$. Figure 7.5.7 illustrates the "before" and "after" situations, and Fig. 7.5.8 shows the graphs of the functions $x_1(t)$, $x_2(t)$, and $x_3(t)$ in Eqs. (27) and (27').



FIGURE 7.5.7. (a) Before; (b) after.

FIGURE 7.5.8. Position functions of the three railway cars of Example 2.

Forced Oscillations and Resonance

Suppose now that the *i*th mass of the mass-and-spring system in Fig. 7.5.2 is subject to an external force F_i (i = 1, 2, ..., n) in addition to the forces exerted by the springs attached to it. Then the homogeneous equation $\mathbf{Mx''} = \mathbf{Kx}$ is replaced with the nonhomogeneous equation

$$\mathbf{M}\mathbf{x}'' = \mathbf{K}\mathbf{x} + \mathbf{F},\tag{28}$$

where $\mathbf{F} = \begin{bmatrix} F_1 & F_2 & \dots & F_n \end{bmatrix}^T$ is the **external force vector** for the system. Multiplication by \mathbf{M}^{-1} yields

$$\mathbf{x}^{\prime\prime} = \mathbf{A}\mathbf{x} + \mathbf{f},\tag{29}$$

where \mathbf{f} is the external force vector *per unit mass*. We are especially interested in the case of a *periodic external force*

$$\mathbf{f}(t) = \mathbf{F}_0 \cos \omega t \tag{30}$$

(where \mathbf{F}_0 is a constant vector). We then anticipate a periodic particular solution

$$\mathbf{x}_{p}(t) = \mathbf{c}\cos\omega t \tag{31}$$

with the known external frequency ω and with a coefficient vector **c** yet to be determined. Because $\mathbf{x}''_p = -\omega^2 \mathbf{c} \cos \omega t$, substitution of (30) and (31) in (29), followed by cancellation of the common factor $\cos \omega t$, gives the linear system

$$(\mathbf{A} + \omega^2 \mathbf{I})\mathbf{c} = -\mathbf{F}_0 \tag{32}$$

to be solved for c.

Observe that the matrix $\mathbf{A} + \omega^2 \mathbf{I}$ is nonsingular—in which case Eq. (32) can be solved for **c**—unless $-\omega^2 = \lambda$, an eigenvalue of **A**. Thus a periodic particular solution of the form in Eq. (31) exists provided that the external forcing frequency does *not* equal one of the natural frequencies $\omega_1, \omega_2, \ldots, \omega_n$ of the system. The case in which ω is a natural frequency corresponds to the phenomenon of **resonance** discussed in Section 5.6.

Example 3 Mass-spring resonance Suppose that the second mass in Example 1 is subjected to the external periodic force $50 \cos \omega t$. Then with $m_1 = 2$, $m_2 = 1$, $k_1 = 100$, $k_2 = 50$, and $F_0 = 50$ in Fig. 7.5.9, Eq. (29) takes the form

$$\mathbf{x}'' = \begin{bmatrix} -75 & 25\\ 50 & -50 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 0\\ 50 \end{bmatrix} \cos \omega t, \tag{33}$$



FIGURE 7.5.9. The forced mass-and-spring system of Example 3.



FIGURE 7.5.10. Frequency–amplitude plot for Example 3.

and the substitution $\mathbf{x} = \mathbf{c} \cos \omega t$ leads to the equation

$$\begin{bmatrix} \omega^2 - 75 & 25\\ 50 & \omega^2 - 50 \end{bmatrix} \mathbf{c} = \begin{bmatrix} 0\\ -50 \end{bmatrix}$$
(34)

for the coefficient vector $\mathbf{c} = \begin{bmatrix} c_1 & c_2 \end{bmatrix}^T$. This system is readily solved for

$$c_1 = \frac{1250}{(\omega^2 - 25)(\omega^2 - 100)}, \quad c_2 = -\frac{50(\omega^2 - 75)}{(\omega^2 - 25)(\omega^2 - 100)}.$$
 (35)

For instance, if the external squared frequency is $\omega^2 = 50$, then (35) yields $c_1 = -1$, $c_2 = -1$. The resulting forced periodic oscillation is described by

$$x_1(t) = -\cos\omega t, \quad x_2(t) = -\cos\omega t.$$

Thus the two masses oscillate in synchrony with equal amplitudes and in the same direction. If the external squared frequency is $\omega^2 = 125$, then (35) yields $c_1 = \frac{1}{2}$, $c_2 = -1$. The resulting forced periodic oscillation is described by

$$x_1(t) = \frac{1}{2}\cos\omega t, \quad x_2(t) = -\cos\omega t,$$

and now the two masses oscillate in synchrony in opposite directions, but with the amplitude of motion of m_2 twice that of m_1 .

It is evident from the denominators in (35) that c_1 and c_2 approach $+\infty$ as ω approaches either of the two natural frequencies $\omega_1 = 5$ and $\omega_2 = 10$ (found in Example 1). Figure 7.5.10 shows a plot of the amplitude $\sqrt{c_1^2 + c_2^2}$ of the forced periodic solution $\mathbf{x}(t) = \mathbf{c} \cos \omega t$ as a function of the forced frequency ω . The peaks at $\omega_2 = 5$ and $\omega_2 = 10$ exhibit visually the phenomenon of resonance.

Periodic and Transient Solutions

It follows from Theorem 4 of Section 7.2 that a particular solution of the forced system

$$\mathbf{x}'' = \mathbf{A}\mathbf{x} + \mathbf{F}_0 \cos \omega t \tag{36}$$

will be of the form

≻

$$\mathbf{x}(t) = \mathbf{x}_c(t) + \mathbf{x}_p(t), \tag{37}$$

where $\mathbf{x}_p(t)$ is a particular solution of the nonhomogeneous system and $\mathbf{x}_c(t)$ is a solution of the corresponding homogeneous system. It is typical for the effects of frictional resistance in mechanical systems to damp out the complementary function solution $\mathbf{x}_c(t)$, so that

$$\mathbf{x}_c(t) \to \mathbf{0} \quad \text{as} \quad t \to +\infty.$$
 (38)

Hence $\mathbf{x}_c(t)$ is a **transient solution** that depends only on the initial conditions; it dies out with time, leaving the **steady periodic solution** $\mathbf{x}_p(t)$ resulting from the external driving force:

$$\mathbf{x}(t) \to \mathbf{x}_p(t) \quad \text{as} \quad t \to +\infty.$$
 (39)

As a practical matter, every physical system includes frictional resistance (however small) that damps out transient solutions in this manner.

7.5 Problems

Problems 1 through 7 deal with the mass-and-spring system shown in Fig. 7.5.11 with stiffness matrix

$$\mathbf{K} = \begin{bmatrix} -(k_1 + k_2) & k_2 \\ k_2 & -(k_2 + k_3) \end{bmatrix}$$

and with the given mks values for the masses and spring constants. Find the two natural frequencies of the system and describe its two natural modes of oscillation.

FIGURE 7.5.11. The mass-and-spring system for Problems 1 through 6.

1. $m_1 = m_2 = 1$; $k_1 = 0, k_2 = 2, k_3 = 0$ (no walls) 2. $m_1 = m_2 = 1$; $k_1 = 1, k_2 = 4, k_3 = 1$ 3. $m_1 = 1, m_2 = 2$; $k_1 = 1, k_2 = k_3 = 2$ 4. $m_1 = m_2 = 1$; $k_1 = 1, k_2 = 2, k_3 = 1$ 5. $m_1 = m_2 = 1$; $k_1 = 2, k_2 = 1, k_3 = 2$ 6. $m_1 = 1, m_2 = 2$; $k_1 = 2, k_2 = k_3 = 4$ 7. $m_1 = m_2 = 1$; $k_1 = 4, k_2 = 6, k_3 = 4$

In Problems 8 through 10 the indicated mass-and-spring system is set in motion from rest $(x'_1(0) = x'_2(0) = 0)$ in its equi-

librium position $(x_1(0) = x_2(0) = 0)$ with the given external forces $F_1(t)$ and $F_2(t)$ acting on the masses m_1 and m_2 , respectively. Find the resulting motion of the system and describe it as a superposition of oscillations at three different frequencies.

- 8. The mass-and-spring system of Problem 2, with $F_1(t) = 96 \cos 5t$, $F_2(t) \equiv 0$
- 9. The mass-and-spring system of Problem 3, with $F_1(t) \equiv 0$, $F_2(t) = 120 \cos 3t$
- 10. The mass-and-spring system of Problem 7, with $F_1(t) = 30 \cos t$, $F_2(t) = 60 \cos t$
- 11. Consider a mass-and-spring system containing two masses $m_1 = 1$ and $m_2 = 1$ whose displacement functions x(t) and y(t) satisfy the differential equations

$$x'' = -40x + 8y,$$

$$y'' = 12x - 60y.$$

(a) Describe the two fundamental modes of free oscillation of the system. (b) Assume that the two masses start in motion with the initial conditions

$$x(0) = 19, \quad x'(0) = 12$$

and

$$y(0) = 3, \quad y'(0) = 6$$

and are acted on by the same force, $F_1(t) = F_2(t) = -195 \cos 7t$. Describe the resulting motion as a superposition of oscillations at three different frequencies.

In Problems 12 and 13, find the natural frequencies of the three-mass system of Fig. 7.5.1, using the given masses and spring constants. For each natural frequency ω , give the ratio $a_1:a_2:a_3$ of amplitudes for a corresponding natural mode $x_1 = a_1 \cos \omega t$, $x_2 = a_2 \cos \omega t$, $x_3 = a_3 \cos \omega t$.

12. $m_1 = m_2 = m_3 = 1$; $k_1 = k_2 = k_3 = k_4 = 1$ **13.** $m_1 = m_2 = m_3 = 1$; $k_1 = k_2 = k_3 = k_4 = 2$

(*Hint*: One eigenvalue is $\lambda = -4$.)

14. In the system of Fig. 7.5.12, assume that $m_1 = 1$, $k_1 = 50$, $k_2 = 10$, and $F_0 = 5$ in mks units, and that $\omega = 10$. Then find m_2 so that in the resulting steady periodic oscillations, the mass m_1 will remain at rest(!). Thus the effect of the second mass-and-spring pair will be to neutralize the effect of the force on the first mass. This is an example of a *dynamic damper*. It has an electrical analogy that some cable companies use to prevent your reception of certain cable channels.



FIGURE 7.5.12. The mechanical system of Problem 14.

15. Suppose that $m_1 = 2$, $m_2 = \frac{1}{2}$, $k_1 = 75$, $k_2 = 25$, $F_0 = 100$, and $\omega = 10$ (all in mks units) in the forced mass-andspring system of Fig. 7.5.9. Find the solution of the system $\mathbf{Mx''} = \mathbf{Kx} + \mathbf{F}$ that satisfies the initial conditions $\mathbf{x}(0) = \mathbf{x'}(0) = \mathbf{0}$.

In Problems 16 through 19 we apply the analysis of Example 2 to a system of two railway cars.

16. Figure 7.5.13 shows two railway cars with a buffer spring. We want to investigate the transfer of momentum that occurs after car 1 with initial velocity v_0 impacts car 2 at rest. The analog of Eq. (18) in the text is

$$\mathbf{x}^{\prime\prime} = \begin{bmatrix} -c_1 & c_1 \\ c_2 & -c_2 \end{bmatrix} \mathbf{x}$$

with $c_i = k/m_i$ for i = 1, 2. Show that the eigenvalues of the coefficient matrix **A** are $\lambda_1 = 0$ and $\lambda_2 = -c_1 - c_2$, with associated eigenvectors $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} c_1 & -c_2 \end{bmatrix}^T$.



FIGURE 7.5.13. The two railway cars of Problems 16 through 19.

- 17. If the two cars of Problem 16 both weigh 16 tons (so that $m_1 = m_2 = 1000$ (slugs)) and k = 1 ton/ft (that is, 2000 lb/ft), show that the cars separate after $\pi/2$ seconds, and that $x'_1(t) = 0$ and $x'_2(t) = v_0$ thereafter. Thus the original momentum of car 1 is completely transferred to car 2.
- 18. If cars 1 and 2 weigh 8 and 16 tons, respectively, and k = 3000 lb/ft, show that the two cars separate after $\pi/3$ seconds, and that

$$x'_1(t) = -\frac{1}{3}v_0$$
 and $x'_2(t) = +\frac{2}{3}v_0$

thereafter. Thus the two cars rebound in opposite directions.

19. If cars 1 and 2 weigh 24 and 8 tons, respectively, and k = 1500 lb/ft, show that the cars separate after $\pi/2$ seconds, and that

$$x'_1(t) = +\frac{1}{2}v_0$$
 and $x'_2(t) = +\frac{3}{2}v_0$

thereafter. Thus both cars continue in the original direction of motion, but with different velocities.

Problems 20 through 23 deal with the same system of three railway cars (same masses) and two buffer springs (same spring constants) as shown in Fig. 7.5.6 and discussed in Example 2. The cars engage at time t = 0 with $x_1(0) = x_2(0) = x_3(0) = 0$ and with the given initial velocities (where $v_0 = 48$ ft/s). Show that the railway cars remain engaged until $t = \pi/2$ (s), after which time they proceed in their respective ways with constant velocities. Determine the values of these constant final velocities $x'_1(t), x'_2(t), and x'_3(t)$ of the three cars for $t > \pi/2$. In each problem you should find (as in Example 2) that the first and third railway cars exchange behaviors in some appropriate sense.

20. $x'_1(0) = v_0, x'_2(0) = 0, x'_3(0) = -v_0$

21.
$$x'_1(0) = 2v_0, x'_2(0) = 0, x'_3(0) = -v_0$$

22.
$$x'_1(0) = v_0, x'_2(0) = v_0, x'_3(0) = -2v_0$$

- **23.** $x'_1(0) = 3v_0, x'_2(0) = 2v_0, x'_3(0) = 2v_0$
- 24. In the three-railway-car system of Fig. 7.5.6, suppose that cars 1 and 3 each weigh 32 tons, that car 2 weighs 8 tons, and that each spring constant is 4 tons/ft. If $x'_1(0) = v_0$ and $x'_2(0) = x'_3(0) = 0$, show that the two springs are compressed until $t = \pi/2$ and that

$$x'_1(t) = -\frac{1}{9}v_0$$
 and $x'_2(t) = x'_3(t) = +\frac{8}{9}v_0$

thereafter. Thus car 1 rebounds, but cars 2 and 3 continue with the same velocity.

The Two-Axle Automobile

In Example 4 of Section 5.6 we investigated the vertical oscillations of a one-axle car—actually a unicycle. Now we can analyze a more realistic model: a car with two axles and with separate front and rear suspension systems. Figure 7.5.14 represents the suspension system of such a car. We assume that the car body acts as would a solid bar of mass m and length $L = L_1 + L_2$. It has moment of inertia I about its center of mass C, which is at distance L_1 from the front of the car. The car has front and back suspension springs with Hooke's constants k_1 and k_2 , respectively. When the car is in motion, let x(t) denote the vertical displacement of the center of mass of the car from equilibrium; let $\theta(t)$ denote its angular displacement (in radians) from the horizontal. Then Newton's laws of motion for linear and angular acceleration can be used to derive the equations

$$mx'' = -(k_1 + k_2)x + (k_1L_1 - k_2L_2)\theta,$$

$$I\theta'' = (k_1L_1 - k_2L_2)x - (k_1L_1^2 + k_2L_2^2)\theta.$$
(40)



FIGURE 7.5.14. Model of the two-axle automobile.

25. Suppose that m = 75 slugs (the car weighs 2400 lb), $L_1 = 7$ ft, $L_2 = 3$ ft (it's a rear-engine car), $k_1 = k_2 = 2000$ lb/ft, and I = 1000 ft·lb·s². Then the equations in (40) take the form

 $75x'' + 4000x - 8000\theta = 0,$

$$1000\theta'' - 8000x + 116,000\theta = 0.$$

(a) Find the two natural frequencies ω_1 and ω_2 of the car. (b) Now suppose that the car is driven at a speed of v feet per second along a washboard surface shaped like a sine curve with a wavelength of 40 ft. The result is a periodic force on the car with frequency $\omega = 2\pi v/40 = \pi v/20$. Resonance occurs when $\omega = \omega_1$ or $\omega = \omega_2$. Find the corresponding two critical speeds of the car (in feet per second and in miles per hour).

26. Suppose that $k_1 = k_2 = k$ and $L_1 = L_2 = \frac{1}{2}L$ in Fig. 7.5.14 (the symmetric situation). Then show that every free oscillation is a combination of a vertical oscillation with frequency

$$\omega_1 = \sqrt{2k/m}$$

and an angular oscillation with frequency

$$\omega_2 = \sqrt{kL^2/(2I)}.$$

In Problems 27 through 29, the system of Fig. 7.5.14 is taken as a model for an undamped car with the given parameters in fps units. (a) Find the two natural frequencies of oscillation (in hertz). (b) Assume that this car is driven along a sinusoidal washboard surface with a wavelength of 40 ft. Find the two critical speeds.

27. $m = 100, I = 800, L_1 = L_2 = 5, k_1 = k_2 = 2000$ **28.** $m = 100, I = 1000, L_1 = 6, L_2 = 4, k_1 = k_2 = 2000$ **29.** $m = 100, I = 800, L_1 = L_2 = 5, k_1 = 1000, k_2 = 2000$

7.5 Application Earthquake-Induced Vibrations of Multistory Buildings





FIGURE 7.5.15. The seven-story building.

In this application you are to investigate the response to transverse earthquake ground oscillations of the seven-story building illustrated in Fig. 7.5.15. Suppose that each of the seven (above-ground) floors weighs 16 tons, so the mass of each is m = 1000 (slugs). Also assume a horizontal restoring force of k = 5 (tons per foot) between adjacent floors. That is, the internal forces in response to horizontal displacements of the individual floors are those shown in Fig. 7.5.16. It follows that the free transverse oscillations indicated in Fig. 7.5.15 satisfy the equation $\mathbf{Mx}'' = \mathbf{Kx}$ with n = 7, $m_i = 1000$ (for each i), and $k_i = 10,000$ (lb/ft) for $1 \le i \le 7$. The system then reduces to the form $\mathbf{x}'' = \mathbf{Ax}$ with

$$\mathbf{A} = \begin{bmatrix} -20 & 10 & 0 & 0 & 0 & 0 & 0 \\ 10 & -20 & 10 & 0 & 0 & 0 & 0 \\ 0 & 10 & -20 & 10 & 0 & 0 & 0 \\ 0 & 0 & 10 & -20 & 10 & 0 & 0 \\ 0 & 0 & 0 & 10 & -20 & 10 & 0 \\ 0 & 0 & 0 & 0 & 10 & -20 & 10 \\ 0 & 0 & 0 & 0 & 0 & 10 & -10 \end{bmatrix}.$$
 (1)

Once the matrix **A** has been entered, the TI-Nspire command eigVI(A) instantly computes the seven eigenvalues shown in the λ -column of the table in Fig. 7.5.17. Alternatively, you can use the *Maple* command eigenvals(A), the MATLAB command eig(A), or the *Mathematica* command Eigenvalues[A].

	Eigenvalue	Frequency	Period	
i	λ	$\omega = \sqrt{-\lambda}$	$P = \frac{2\pi}{\omega} (\text{sec})$	
1	-38.2709	6.1863	1.0157	
2	-33.3826	5.7778	1.0875	
3	-26.1803	5.1167	1.2280	
4	-17.9094	4.2320	1.4847	
5	-10.0000	3.1623	1.9869	
6	-3.8197	1.9544	3.2149	
7	-0.4370	0.6611	9.5042	



 $k(x_i - x_{i-1}) \neq$

 $k(x_{i+1} - x_i)$

FIGURE 7.5.16. Forces on the *i*th floor.

FIGURE 7.5.17. Frequencies and periods of natural oscillations of the seven-story building.

Then calculate the entries in the remaining columns of the table showing the natural frequencies and periods of oscillation of the seven-story building. Note that a typical earthquake producing ground oscillations with a period of 2 seconds is uncomfortably close to the fifth natural frequency (with period 1.9869 seconds) of the building.

A horizontal earthquake oscillation $E \cos \omega t$ of the ground, with amplitude Eand acceleration $a = -E\omega^2 \cos \omega t$, produces an opposite inertial force $F = ma = mE\omega^2 \cos \omega t$ on each floor of the building. The resulting nonhomogeneous system is

$$\mathbf{x}'' = \mathbf{A}\mathbf{x} + (E\omega^2 \cos \omega t)\mathbf{b},\tag{2}$$



FIGURE 7.5.18. Resonance vibrations of a seven-story building—maximal amplitude as a function of period.

where $\mathbf{b} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix}^T$ and \mathbf{A} is the matrix of Eq. (1). Figure 7.5.18 shows a plot of *maximal amplitude* (for the forced oscillations of any single floor) versus the *period* of the earthquake vibrations. The spikes correspond to the first six of the seven resonant frequencies. We see, for instance, that whereas an earthquake with period 2 (s) likely would produce destructive resonance vibrations in the building, it probably would be unharmed by an earthquake with period 2.5 (s). Different buildings have different natural frequencies of vibration, and so a given earthquake may demolish one building but leave untouched the one next door. This seeming anomaly was observed in Mexico City after the devastating earthquake of September 19, 1985.

For your personal seven-story building to investigate, let the weight (in tons) of each story equal the largest digit of your student ID number and let k (in tons/ft) equal the smallest nonzero digit. Produce numerical and graphical results like those illustrated in Figs. 7.5.17 and 7.5.18. Is your building susceptible to likely damage from an earthquake with period in the 2- to 3-second range?

You might like to begin by working manually the following warm-up problems.

- 1. Find the periods of the natural vibrations of a building with two above-ground floors, each weighing 16 tons and with each restoring force being k = 5 tons/ft.
- 2. Find the periods of the natural vibrations of a building with three aboveground floors, with each weighing 16 tons and with each restoring force being k = 5 tons/ft.

- **3.** Find the natural frequencies and natural modes of vibration of a building with three above-ground floors as in Problem 2, except that the upper two floors weigh 8 tons each instead of 16. Give the ratios of the amplitudes A, B, and C of oscillations of the three floors in the form A:B:C with A = 1.
- 4. Suppose that the building of Problem 3 is subject to an earthquake in which the ground undergoes horizontal sinusoidal oscillations with a period of 3 s and an amplitude of 3 in. Find the amplitudes of the resulting steady periodic oscillations of the three above-ground floors. Assume the fact that a motion $E \sin \omega t$ of the ground, with acceleration $a = -E\omega^2 \sin \omega t$, produces an opposite inertial force $F = -ma = mE\omega^2 \sin \omega t$ on a floor of mass m.

7.6 Multiple Eigenvalue Solutions

In Section 7.3 we saw that if the $n \times n$ matrix **A** has *n* distinct (real or complex) eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$ with respective associated eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$, then a general solution of the system

$$\frac{d\mathbf{x}}{dt} = \mathbf{A}\mathbf{x} \tag{1}$$

is given by

3

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{\lambda_1 t} + c_2 \mathbf{v}_2 e^{\lambda_2 t} + \dots + c_n \mathbf{v}_n e^{\lambda_n t}$$
(2)

with arbitrary constants c_1, c_2, \ldots, c_n . In this section we discuss the situation when the characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \tag{3}$$

does not have n distinct roots, and thus has at least one repeated root.

An eigenvalue is of **multiplicity** k if it is a k-fold root of Eq. (3). For each eigenvalue λ , the eigenvector equation

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0} \tag{4}$$

has at least one nonzero solution **v**, so there is at least one eigenvector associated with λ . But an eigenvalue of multiplicity k > 1 may have *fewer* than k linearly independent associated eigenvectors. In this case we are unable to find a "complete set" of n linearly independent eigenvectors of **A**, as needed to form the general solution in (2).

Let us call an eigenvalue of multiplicity k complete if it has k linearly independent associated eigenvectors. If every eigenvalue of the matrix A is complete, then—because eigenvectors associated with different eigenvalues are linearly independent—it follows that A does have a complete set of n linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ associated with the eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$ (each repeated with its multiplicity). In this case a general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is still given by the usual combination in (2).

$$\mathbf{x}' = \begin{bmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{bmatrix} \mathbf{x}.$$
 (5)

438 CHAPTER 7 LINEAR SYSTEMS OF DIFFERENTIAL EQUATIONS

Solution The characteristic equation of the coefficient matrix A in Eq. (5) is

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{vmatrix} 9 - \lambda & 4 & 0 \\ -6 & -1 - \lambda & 0 \\ 6 & 4 & 3 - \lambda \end{vmatrix}$$
$$= (3 - \lambda)[(9 - \lambda)(-1 - \lambda) + 24]$$
$$= (3 - \lambda)(15 - 8\lambda + \lambda^2)$$
$$= (5 - \lambda)(3 - \lambda)^2 = 0.$$

Thus **A** has the distinct eigenvalue $\lambda_1 = 5$ and the repeated eigenvalue $\lambda_2 = 3$ of multiplicity k = 2.

CASE 1: $\lambda_1 = 5$. The eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$, where $\mathbf{v} = \begin{bmatrix} a & b & c \end{bmatrix}^T$, is

$$(\mathbf{A} - 5\mathbf{I})\mathbf{v} = \begin{bmatrix} 4 & 4 & 0 \\ -6 & -6 & 0 \\ 6 & 4 & -2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

Each of the first two equations, 4a + 4b = 0 and -6a - 6b = 0, yields b = -a. Then the third equation reduces to 2a - 2c = 0, so that c = a. The choice a = 1 then yields the eigenvector

$$\mathbf{v}_1 = \begin{bmatrix} 1 & -1 & 1 \end{bmatrix}^T$$

associated with the eigenvalue $\lambda_1 = 5$.

CASE 2: $\lambda_2 = 3$. Now the eigenvector equation is

$$(\mathbf{A} - 3\mathbf{I})\mathbf{v} = \begin{bmatrix} 6 & 4 & 0 \\ -6 & -4 & 0 \\ 6 & 4 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

so the nonzero vector $\mathbf{v} = \begin{bmatrix} a & b & c \end{bmatrix}^T$ is an eigenvector if and only if

$$6a + 4b = 0;$$
 (6)

that is, $b = -\frac{3}{2}a$. The fact that Eq. (6) does not involve *c* means that *c* is arbitrary, subject to the condition $\mathbf{v} \neq \mathbf{0}$. If c = 1, then we may choose a = b = 0; this gives the eigenvector

$$\mathbf{v}_2 = \begin{bmatrix} 0 & 0 & 1 \end{bmatrix}^T$$

associated with $\lambda_2 = 3$. If c = 0, then we must choose *a* to be nonzero. For instance, if a = 2 (to avoid fractions), then b = -3, so

$$\mathbf{v}_3 = \begin{bmatrix} 2 & -3 & 0 \end{bmatrix}^T$$

is a second linearly independent eigenvector associated with the multiplicity 2 eigenvalue $\lambda_2 = 3$.

Thus we have found a complete set v_1 , v_2 , v_3 of three eigenvectors associated with the eigenvalues 5, 3, 3. The corresponding general solution of Eq. (5) is

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{5t} + c_2 \mathbf{v}_2 e^{3t} + c_3 \mathbf{v}_3 e^{3t}$$
(7)
= $c_1 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} e^{5t} + c_2 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} e^{3t} + c_3 \begin{bmatrix} 2 \\ -3 \\ 0 \end{bmatrix} e^{3t},$

with scalar component functions given by

$$x_{1}(t) = c_{1}e^{5t} + 2c_{3}e^{3t},$$

$$x_{2}(t) = -c_{1}e^{5t} - 3c_{3}e^{3t},$$

$$x_{3}(t) = c_{1}e^{5t} + c_{2}e^{3t}.$$

Remark Our choice in Example 1 of the two eigenvectors

$$\mathbf{v}_2 = \begin{bmatrix} 0 & 0 & 1 \end{bmatrix}^T$$
 and $\mathbf{v}_3 = \begin{bmatrix} 2 & -3 & 0 \end{bmatrix}^T$

associated with the repeated eigenvalue $\lambda_2 = 3$ bears comment. The fact that $b = -\frac{3}{2}a$ for any eigenvector associated with $\lambda_2 = 3$ means that any such eigenvector can be written as

$$\mathbf{v} = \begin{bmatrix} a \\ -\frac{3}{2}a \\ c \end{bmatrix} = c \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \frac{1}{2}a \begin{bmatrix} 2 \\ -3 \\ 0 \end{bmatrix} = c\mathbf{v}_2 + \frac{1}{2}a\mathbf{v}_3,$$

and thus is a linear combination of v_2 and v_3 . Therefore, given *a* and *c* not both zero, we could choose v rather than v_3 as our third eigenvector, and the new general solution

$$\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{5t} + c_2 \mathbf{v}_2 e^{3t} + c_3 \mathbf{v} e^{3t}$$

would be equivalent to the one in Eq. (7). Thus we need not worry about making the "right" choice of independent eigenvectors associated with a multiple eigenvalue. Any choice will do; we generally make the simplest one we can.

Defective Eigenvalues

The following example shows that—unfortunately—not all multiple eigenvalues are complete.

Example 2 The matrix

$$\mathbf{A} = \begin{bmatrix} 1 & -3\\ 3 & 7 \end{bmatrix} \tag{8}$$

has characteristic equation

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{vmatrix} 1 - \lambda & -3 \\ 3 & 7 - \lambda \end{vmatrix}$$
$$= (1 - \lambda)(7 - \lambda) + 9$$
$$= \lambda^2 - 8\lambda + 16 = (\lambda - 4)^2 = 0.$$

Thus **A** has the single eigenvalue $\lambda_1 = 4$ of multiplicity 2. The eigenvector equation

$$(\mathbf{A} - 4\mathbf{I})\mathbf{v} = \begin{bmatrix} -3 & -3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

then amounts to the equivalent scalar equations

$$-3a - 3b = 0, \quad 3a + 3b = 0.$$

Hence b = -a if $\mathbf{v} = \begin{bmatrix} a & b \end{bmatrix}^T$ is to be an eigenvector of **A**. Therefore, any eigenvector associated with $\lambda_1 = 4$ is a nonzero multiple of $\mathbf{v} = \begin{bmatrix} 1 & -1 \end{bmatrix}^T$. Thus the multiplicity 2 eigenvalue $\lambda_1 = 4$ has only one independent eigenvector, and hence is incomplete.

An eigenvalue λ of multiplicity k > 1 is called **defective** if it is not complete. If λ has only p < k linearly independent eigenvectors, then the number

$$d = k - p \tag{9}$$

of "missing" eigenvectors is called the **defect** of the defective eigenvalue λ . Thus the defective eigenvalue $\lambda_1 = 4$ in Example 2 has multiplicity k = 2 and defect d = 1, because we saw that it has only p = 1 associated eigenvector.

If the eigenvalues of the $n \times n$ matrix **A** are not all complete, then the eigenvalue method as yet described will produce *fewer* than the needed *n* linearly independent solutions of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. We therefore need to discover how to find the "missing solutions" corresponding to a defective eigenvalue λ of multiplicity k > 1.

The Case of Multiplicity k = 2

Let us begin with the case k = 2, and suppose that we have found (as in Example 2) that there is only a single eigenvector \mathbf{v}_1 associated with the defective eigenvalue λ . Then at this point we have found only the single solution

$$\mathbf{x}_1(t) = \mathbf{v}_1 e^{\lambda t} \tag{10}$$

of $\mathbf{x}' = \mathbf{A}\mathbf{x}$. By analogy with the case of a repeated characteristic root for a single linear differential equation (Section 5.3), we might hope to find a second solution of the form

$$\mathbf{x}_2(t) = (\mathbf{v}_2 t)e^{\lambda t} = \mathbf{v}_2 t e^{\lambda t}.$$
(11)

When we substitute $\mathbf{x} = \mathbf{v}_2 t e^{\lambda t}$ in $\mathbf{x}' = \mathbf{A}\mathbf{x}$, we get the equation

$$\mathbf{v}_2 e^{\lambda t} + \lambda \mathbf{v}_2 t e^{\lambda t} = \mathbf{A} \mathbf{v}_2 t e^{\lambda t}.$$

But because the coefficients of both $e^{\lambda t}$ and $te^{\lambda t}$ must balance, it follows that $\mathbf{v}_2 = \mathbf{0}$, and hence that $\mathbf{x}_2(t) \equiv \mathbf{0}$. This means that—contrary to our hope—the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ does not have a *nontrivial* solution of the form assumed in (11).

Instead of simply giving up on the idea behind Eq. (11), let us extend it slightly and replace $\mathbf{v}_2 t$ with $\mathbf{v}_1 t + \mathbf{v}_2$. Thus we explore the possibility of a second solution of the form

$$\mathbf{x}_2(t) = (\mathbf{v}_1 t + \mathbf{v}_2)e^{\lambda t} = \mathbf{v}_1 t e^{\lambda t} + \mathbf{v}_2 e^{\lambda t}, \tag{12}$$

where \mathbf{v}_1 and \mathbf{v}_2 are nonzero constant vectors. When we substitute $\mathbf{x} = \mathbf{v}_1 t e^{\lambda t} + \mathbf{v}_2 e^{\lambda t}$ in $\mathbf{x}' = \mathbf{A}\mathbf{x}$, we get the equation

$$\mathbf{v}_1 e^{\lambda t} + \lambda \mathbf{v}_1 t e^{\lambda t} + \lambda \mathbf{v}_2 e^{\lambda t} = \mathbf{A} \mathbf{v}_1 t e^{\lambda t} + \mathbf{A} \mathbf{v}_2 e^{\lambda t}.$$
 (13)

We equate coefficients of $e^{\lambda t}$ and $te^{\lambda t}$ here, and thereby obtain the two equations

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_1 = \mathbf{0} \tag{14}$$

and

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_2 = \mathbf{v}_1 \tag{15}$$

that the vectors \mathbf{v}_1 and \mathbf{v}_2 must satisfy in order for (12) to give a solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

Note that Eq. (14) merely confirms that \mathbf{v}_1 is an eigenvector of \mathbf{A} associated with the eigenvalue λ . Then Eq. (15) says that the vector \mathbf{v}_2 satisfies the equation

$$(\mathbf{A} - \lambda \mathbf{I})^2 \mathbf{v}_2 = (\mathbf{A} - \lambda \mathbf{I})[(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_2] = (\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_1 = \mathbf{0}$$

It follows that, in order to solve simultaneously the two equations in (14) and (15), it suffices to find a solution \mathbf{v}_2 of the single equation $(\mathbf{A} - \lambda \mathbf{I})^2 \mathbf{v}_2 = \mathbf{0}$ such that the resulting vector $\mathbf{v}_1 = (\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_2$ is *nonzero*. It turns out that this is always possible if the defective eigenvalue λ of \mathbf{A} is of multiplicity 2. Consequently, the procedure described in the following algorithm always succeeds in finding two independent solutions associated with such an eigenvalue.

ALGORITHM Defective Multiplicity 2 Eigenvalues

1. First find a nonzero solution v_2 of the equation

$$(\mathbf{A} - \lambda \mathbf{I})^2 \mathbf{v}_2 = \mathbf{0} \tag{16}$$

such that

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_2 = \mathbf{v}_1 \tag{17}$$

is nonzero, and therefore is an eigenvector \mathbf{v}_1 associated with λ .

2. Then form the two independent solutions

$$\mathbf{x}_1(t) = \mathbf{v}_1 e^{\lambda t} \tag{18}$$

and

$$\mathbf{x}_2(t) = (\mathbf{v}_1 t + \mathbf{v}_2)e^{\lambda t} \tag{19}$$

of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ corresponding to λ .

Example 3 Find a general solution of the system

$$\mathbf{x}' = \begin{bmatrix} 1 & -3\\ 3 & 7 \end{bmatrix} \mathbf{x}.$$
 (20)

Solution In Example 2 we found that the coefficient matrix A in Eq. (20) has the defective eigenvalue $\lambda = 4$ of multiplicity 2. We therefore begin by calculating

$$\left(\mathbf{A} - 4\mathbf{I}\right)^2 = \begin{bmatrix} -3 & -3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} -3 & -3 \\ 3 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Hence Eq. (16) is

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \mathbf{v}_2 = \mathbf{0},$$

and therefore is satisfied by *any* choice of \mathbf{v}_2 . In principle, it could happen that $(\mathbf{A} - 4\mathbf{I})\mathbf{v}_2$ is nonzero (as desired) for some choices of \mathbf{v}_2 though not for others. If we try $\mathbf{v}_2 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ we find that

$$(\mathbf{A} - 4\mathbf{I})\mathbf{v}_2 = \begin{bmatrix} -3 & -3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -3 \\ 3 \end{bmatrix} = \mathbf{v}_1$$

is nonzero, and therefore is an eigenvector associated with $\lambda = 4$. (It is -3 times the eigenvector found in Example 2.) Therefore, the two solutions of Eq. (20) given by Eqs. (18) and (19) are

$$\mathbf{x}_{1}(t) = \mathbf{v}_{1}e^{4t} = \begin{bmatrix} -3\\ 3 \end{bmatrix}e^{4t},$$
$$\mathbf{x}_{2}(t) = (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{4t} = \begin{bmatrix} -3t+1\\ 3t \end{bmatrix}e^{4t}.$$

The resulting general solution

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t)$$

has scalar component functions

$$x_1(t) = (-3c_2t + c_2 - 3c_1)e^{4t},$$

$$x_2(t) = (3c_2t + 3c_1)e^{4t}.$$



FIGURE 7.6.1. Direction field and solution curves for the linear system $x'_1 = x_1 - 3x_2, x'_2 = 3x_1 + 7x_2$ of Example 3.

With $c_2 = 0$ these solution equations reduce to the equations $x_1(t) = -3c_1e^{4t}$, $x_2(t) = 3c_1e^{4t}$, which parametrize the line $x_1 = -x_2$ in the x_1x_2 -plane. The point $(x_1(t), x_2(t))$ then recedes along this line away from the origin as $t \to +\infty$, to the northwest if $c_1 > 0$ and to the southeast if $c_1 < 0$. As indicated in Fig. 7.6.1, each solution curve with $c_2 \neq 0$ is tangent to the line $x_1 = -x_2$ at the origin; the point $(x_1(t), x_2(t))$ approaches the origin as $t \to -\infty$ and approaches $+\infty$ along the solution curve as $t \to +\infty$.

Generalized Eigenvectors

The vector \mathbf{v}_2 in Eq. (16) is an example of a generalized eigenvector. If λ is an eigenvalue of the matrix **A**, then a **rank** *r* **generalized eigenvector** associated with λ is a vector **v** such that

$$(\mathbf{A} - \lambda \mathbf{I})^r \mathbf{v} = \mathbf{0} \quad \text{but} \quad (\mathbf{A} - \lambda \mathbf{I})^{r-1} \mathbf{v} \neq \mathbf{0}.$$
(21)

If r = 1, then (21) simply means that **v** is an eigenvector associated with λ (recalling the convention that the 0th power of a square matrix is the identity matrix). Thus a rank 1 generalized eigenvector is an ordinary eigenvector. The vector **v**₂ in (16) is a rank 2 generalized eigenvector (and not an ordinary eigenvector).

The multiplicity 2 method described earlier boils down to finding a pair $\{\mathbf{v}_1, \mathbf{v}_2\}$ of generalized eigenvectors, one of rank 1 and one of rank 2, such that $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_2 = \mathbf{v}_1$. Higher multiplicity methods involve longer "chains" of generalized eigenvectors. A length *k* chain of generalized eigenvectors based on the eigenvector \mathbf{v}_1 is a set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ of *k* generalized eigenvectors such that

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_{k} = \mathbf{v}_{k-1},$$

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_{k-1} = \mathbf{v}_{k-2},$$

$$\vdots$$

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_{2} = \mathbf{v}_{1}.$$
(22)

Because \mathbf{v}_1 is an ordinary eigenvector, $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_1 = \mathbf{0}$. Therefore, it follows from (22) that

$$(\mathbf{A} - \lambda \mathbf{I})^k \mathbf{v}_k = \mathbf{0}.$$
 (23)

If $\{v_1, v_2, v_3\}$ is a length 3 chain of generalized eigenvectors associated with the multiple eigenvalue λ of the matrix **A**, then it is easy to verify that three linearly independent solutions of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ are given by

>

$$\mathbf{x}_{1}(t) = \mathbf{v}_{1}e^{\lambda t},$$

$$\mathbf{x}_{2}(t) = (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{\lambda t},$$

$$\mathbf{x}_{3}(t) = (\frac{1}{2}\mathbf{v}_{1}t^{2} + \mathbf{v}_{2}t + \mathbf{v}_{3})e^{\lambda t}.$$
(24)

For instance, the equations in (22) give

$$\mathbf{A}\mathbf{v}_3 = \mathbf{v}_2 + \lambda \mathbf{v}_3, \quad \mathbf{A}\mathbf{v}_2 = \mathbf{v}_1 + \lambda \mathbf{v}_2, \quad \mathbf{A}\mathbf{v}_1 = \lambda \mathbf{v}_1,$$

so

$$\begin{aligned} \mathbf{A}\mathbf{x}_{3} &= \left[\frac{1}{2}\mathbf{A}\mathbf{v}_{1}t^{2} + \mathbf{A}\mathbf{v}_{2}t + \mathbf{A}\mathbf{v}_{3}\right]e^{\lambda t} \\ &= \left[\frac{1}{2}\lambda\mathbf{v}_{1}t^{2} + (\mathbf{v}_{1} + \lambda\mathbf{v}_{2})t + (\mathbf{v}_{2} + \lambda\mathbf{v}_{3})\right]e^{\lambda t} \\ &= (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{\lambda t} + \lambda\left(\frac{1}{2}\mathbf{v}_{1}t^{2} + \mathbf{v}_{2}t + \mathbf{v}_{3}\right)e^{\lambda t} \\ &= \mathbf{x}_{3}^{\prime}. \end{aligned}$$

Therefore, $\mathbf{x}_3(t)$ in (24) does, indeed, define a solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

>

7.6 MULTIPLE EIGENVALUE SOLUTION\$43

Consequently, in order to "handle" a multiplicity 3 eigenvalue λ , it suffices to find a length 3 chain $\{v_1, v_2, v_3\}$ of generalized eigenvalues associated with λ . Looking at Eq. (23), we see that we need only find a solution v_3 of

$$(\mathbf{A} - \lambda \mathbf{I})^3 \mathbf{v}_3 = \mathbf{0}$$

such that the vectors

$$\mathbf{v}_2 = (\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_3$$
 and $\mathbf{v}_1 = (\mathbf{A} - \lambda \mathbf{I})\mathbf{v}_2$

are both nonzero (although, as we will see, this is not always possible).

Example 4 Find three linearly independent solutions of the system

$$\mathbf{x}' = \begin{bmatrix} 0 & 1 & 2\\ -5 & -3 & -7\\ 1 & 0 & 0 \end{bmatrix} \mathbf{x}.$$
 (25)

Solution The characteristic equation of the coefficient matrix in Eq. (25) is

$$|\mathbf{A} - \lambda \mathbf{I}| = \begin{bmatrix} -\lambda & 1 & 2\\ -5 & -3 - \lambda & -7\\ 1 & 0 & -\lambda \end{bmatrix}$$
$$= 1 \cdot [-7 - 2 \cdot (-3 - \lambda)] + (-\lambda)[(-\lambda)(-3 - \lambda) + 5]$$
$$= -\lambda^3 - 3\lambda^2 - 3\lambda - 1 = -(\lambda + 1)^3 = 0,$$

and thus A has the eigenvalue $\lambda = -1$ of multiplicity 3. The eigenvector equation $(A - \lambda I)v =$ **0** for an eigenvector $\mathbf{v} = \begin{bmatrix} a & b & c \end{bmatrix}^T$ is

$$(\mathbf{A} + \mathbf{I})\mathbf{v} = \begin{bmatrix} 1 & 1 & 2 \\ -5 & -2 & -7 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

The third row a + c = 0 gives c = -a; then the first row a + b + 2c = 0 gives b = a. Thus, to within a constant multiple, the eigenvalue $\lambda = -1$ has only the single associated eigenvector $\mathbf{v} = \begin{bmatrix} a & a & -a \end{bmatrix}^T$ with $a \neq 0$, and so the defect of $\lambda = -1$ is 2. To apply the method described here for triple eigenvalues, we first calculate

$$(\mathbf{A} + \mathbf{I})^2 = \begin{bmatrix} 1 & 1 & 2 \\ -5 & -2 & -7 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ -5 & -2 & -7 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -2 & -1 & -3 \\ -2 & -1 & -3 \\ 2 & 1 & 3 \end{bmatrix}$$

and

$$(\mathbf{A} + \mathbf{I})^3 = \begin{bmatrix} 1 & 1 & 2 \\ -5 & -2 & -7 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} -2 & -1 & -3 \\ -2 & -1 & -3 \\ 2 & 1 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Thus any nonzero vector \mathbf{v}_3 will be a solution of the equation $(\mathbf{A} + \mathbf{I})^3 \mathbf{v}_3 = \mathbf{0}$. Beginning with $\mathbf{v}_3 = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}^T$, for instance, we calculate

$$\mathbf{v}_{2} = (\mathbf{A} + \mathbf{I})\mathbf{v}_{3} = \begin{bmatrix} 1 & 1 & 2 \\ -5 & -2 & -7 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ -5 \\ 1 \end{bmatrix},$$
$$\mathbf{v}_{1} = (\mathbf{A} + \mathbf{I})\mathbf{v}_{2} = \begin{bmatrix} 1 & 1 & 2 \\ -5 & -2 & -7 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -5 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \\ 2 \end{bmatrix}$$

Note that \mathbf{v}_1 is the previously found eigenvector \mathbf{v} with a = -2; this agreement serves as a check of the accuracy of our matrix computations.

Thus we have found a length 3 chain { v_1 , v_2 , v_3 } of generalized eigenvectors associated with the triple eigenvalue $\lambda = -1$. Substitution in (24) now yields the linearly independent solutions

$$\mathbf{x}_{1}(t) = \mathbf{v}_{1}e^{-t} = \begin{bmatrix} -2\\ -2\\ 2 \end{bmatrix}e^{-t},$$
$$\mathbf{x}_{2}(t) = (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{-t} = \begin{bmatrix} -2t+1\\ -2t-5\\ 2t+1 \end{bmatrix}e^{-t},$$
$$\mathbf{x}_{3}(t) = \left(\frac{1}{2}\mathbf{v}_{1}t^{2} + \mathbf{v}_{2}t + \mathbf{v}_{3}\right)e^{-t} = \begin{bmatrix} -t^{2}+t+1\\ -t^{2}-5t\\ t^{2}+t \end{bmatrix}e^{-t}$$

of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

The General Case

A fundamental theorem of linear algebra states that every $n \times n$ matrix **A** has *n* linearly independent generalized eigenvectors. These *n* generalized eigenvectors may be arranged in chains, with the sum of the lengths of the chains associated with a given eigenvalue λ equal to the multiplicity of λ . But the structure of these chains depends on the defect of λ , and can be quite complicated. For instance, a multiplicity 4 eigenvalue can correspond to

- Four length 1 chains (defect 0);
- Two length 1 chains and a length 2 chain (defect 1);
- Two length 2 chains (defect 2);
- A length 1 chain and a length 3 chain (defect 2); or
- A length 4 chain (defect 3).

Note that, in each of these cases, the length of the longest chain is at most d + 1, where d is the defect of the eigenvalue. Consequently, once we have found all the ordinary eigenvectors associated with a multiple eigenvalue λ , and therefore know the defect d of λ , we can begin with the equation

≻

$$(\mathbf{A} - \lambda \mathbf{I})^{d+1} \mathbf{u} = \mathbf{0}$$
⁽²⁶⁾

to start building the chains of generalized eigenvectors associated with λ .

ALGORITHM Chains of Generalized Eigenvectors

Begin with a nonzero solution \mathbf{u}_1 of Eq. (26) and successively multiply by the matrix $\mathbf{A} - \lambda \mathbf{I}$ until the zero vector is obtained. If

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{u}_1 = \mathbf{u}_2 \neq \mathbf{0},$$

$$\vdots$$

$$(\mathbf{A} - \lambda \mathbf{I})\mathbf{u}_{k-1} = \mathbf{u}_k \neq \mathbf{0}.$$

but $(\mathbf{A} - \lambda \mathbf{I})\mathbf{u}_k = \mathbf{0}$, then the vectors

$$\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} = \{\mathbf{u}_k, \mathbf{u}_{k-1}, \dots, \mathbf{u}_2, \mathbf{u}_1\}$$

(listed in reverse order of their appearance) form a length k chain of generalized eigenvectors based on the (ordinary) eigenvector v_1 .

7.6 MULTIPLE EIGENVALUE SOLUTION \$45

Each length k chain $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ of generalized eigenvectors (with \mathbf{v}_1 an ordinary eigenvector associated with λ) determines a set of k independent solutions of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ corresponding to the eigenvalue λ :

$$\mathbf{x}_{1}(t) = \mathbf{v}_{1}e^{\lambda t},$$

$$\mathbf{x}_{2}(t) = (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{\lambda t},$$

$$\mathbf{x}_{3}(t) = \left(\frac{1}{2}\mathbf{v}_{1}t^{2} + \mathbf{v}_{2}t + \mathbf{v}_{3}\right)e^{\lambda t},$$

$$\vdots$$

$$\mathbf{x}_{k}(t) = \left(\frac{\mathbf{v}_{1}t^{k-1}}{(k-1)!} + \dots + \frac{\mathbf{v}_{k-2}t^{2}}{2!} + \mathbf{v}_{k-1}t + \mathbf{v}_{k}\right)e^{\lambda t}.$$
(27)

Note that (27) reduces to Eqs. (18) through (19) and (24) in the cases k = 2 and k = 3, respectively.

To ensure that we obtain *n* generalized eigenvectors of the $n \times n$ matrix **A** that are actually linearly independent, and therefore produce a complete set of *n* linearly independent solutions of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ when we amalgamate all the "chains of solutions" corresponding to different chains of generalized eigenvectors, we may rely on the following two facts:

- Any chain of generalized eigenvectors constitutes a linearly independent set of vectors.
- If two chains of generalized eigenvectors are based on linearly independent eigenvectors, then the union of these two chains is a linearly independent set of vectors (whether the two base eigenvectors are associated with different eigenvalues or with the same eigenvalue).
- **Example 5** Suppose that the 6×6 matrix **A** has two multiplicity 3 eigenvalues $\lambda_1 = -2$ and $\lambda_2 = 3$ with defects 1 and 2, respectively. Then λ_1 must have an associated eigenvector \mathbf{u}_1 and a length 2 chain $\{\mathbf{v}_1, \mathbf{v}_2\}$ of generalized eigenvectors (with the eigenvectors \mathbf{u}_1 and \mathbf{v}_1 being linearly independent), whereas λ_2 must have a length 3 chain $\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\}$ of generalized eigenvector \mathbf{w}_1 . The six generalized eigenvectors $\mathbf{u}_1, \mathbf{v}_1, \mathbf{v}_2, \mathbf{w}_1, \mathbf{w}_2, \mathbf{and}, \mathbf{w}_3$ are then linearly independent and yield the following six independent solutions of $\mathbf{x}' = \mathbf{A}\mathbf{x}$:

$$\begin{aligned} \mathbf{x}_{1}(t) &= \mathbf{u}_{1}e^{-2t}, \\ \mathbf{x}_{2}(t) &= \mathbf{v}_{1}e^{-2t}, \\ \mathbf{x}_{3}(t) &= (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{-2t}, \\ \mathbf{x}_{4}(t) &= \mathbf{w}_{1}e^{3t}, \\ \mathbf{x}_{5}(t) &= (\mathbf{w}_{1}t + \mathbf{w}_{2})e^{3t}, \\ \mathbf{x}_{6}(t) &= \left(\frac{1}{2}\mathbf{w}_{1}t^{2} + \mathbf{w}_{2}t + \mathbf{w}_{3}\right)e^{3t}. \end{aligned}$$

As Example 5 illustrates, the computation of independent solutions corresponding to different eigenvalues and chains of generalized eigenvalues is a routine matter. The determination of the chain structure associated with a given multiple eigenvalue can be more interesting (as in Example 6).

An Application



FIGURE 7.6.2. The railway cars of Example 6.

Figure 7.6.2 shows two railway cars that are connected with a spring (permanently attached to both cars) and with a damper that exerts opposite forces on the two cars, of magnitude $c(x'_1 - x'_2)$ proportional to their relative velocity. The two cars are also subject to frictional resistance forces $c_1x'_1$ and $c_2x'_2$ proportional to their respective velocities. An application of Newton's law ma = F (as in Example 1 of Section 7.1) yields the equations of motion

$$m_1 x_1'' = k(x_2 - x_1) - c_1 x_1' - c(x_1' - x_2'),$$

$$m_2 x_2'' = k(x_1 - x_2) - c_2 x_2' - c(x_2' - x_1').$$
(28)

In terms of the position vector $\mathbf{x}(t) = \begin{bmatrix} x_1(t) & x_2(t) \end{bmatrix}^T$, these equations can be written in the matrix form

$$\mathbf{M}\mathbf{x}'' = \mathbf{K}\mathbf{x} + \mathbf{R}\mathbf{x}',\tag{29}$$

where \mathbf{M} and \mathbf{K} are mass and stiffness matrices (as in Eqs. (2) and (3) of Section 7.4), and

$$\mathbf{R} = \begin{bmatrix} -(c+c_1) & c \\ c & -(c+c_2) \end{bmatrix}$$

is the *resistance matrix*. Unfortunately, because of the presence of the term involving \mathbf{x}' , the methods of Section 7.4 cannot be used.

Instead, we write (28) as a first-order system in the four unknown functions $x_1(t), x_2(t), x_3(t) = x'_1(t)$, and $x_4(t) = x'_2(t)$. If $m_1 = m_2 = 1$ we get

$$\mathbf{x}' = \mathbf{A}\mathbf{x},\tag{30}$$

where now $\mathbf{x} = \begin{bmatrix} x_1 & x_2 & x_3 & x_4 \end{bmatrix}^T$ and

$$\mathbf{A} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -k & k & -(c+c_1) & c \\ k & -k & c & -(c+c_2) \end{bmatrix}.$$
 (31)

Example 6 Two railway cars With $m_1 = m_2 = c = 1$ and $k = c_1 = c_2 = 2$, the system in Eq. (30) is

$$\mathbf{x}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & 2 & -3 & 1 \\ 2 & -2 & 1 & -3 \end{bmatrix} \mathbf{x}.$$
 (32)

It is not too tedious to calculate manually—although a computer algebra system such as *Maple, Mathematica*, or MATLAB is useful here—the characteristic equation

$$\lambda^4 + 6\lambda^3 + 12\lambda^2 + 8\lambda = \lambda(\lambda+2)^3 = 0$$

of the coefficient matrix **A** in Eq. (32). Thus **A** has the distinct eigenvalue $\lambda_0 = 0$ and the triple eigenvalue $\lambda_1 = -2$.

CASE 1: $\lambda_0 = 0$. The eigenvalue equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ for the eigenvector $\mathbf{v} = \begin{bmatrix} a & b & c & d \end{bmatrix}^T$ is

$$\mathbf{Av} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & 2 & -3 & 1 \\ 2 & -2 & 1 & -3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

The first two rows give c = d = 0; then the last two rows yield a = b. Thus

$$\mathbf{v}_0 = \begin{bmatrix} 1 & 1 & 0 & 0 \end{bmatrix}^T$$

is an eigenvector associated with $\lambda_0 = 0$.

CASE 2:
$$\lambda_1 = -2$$
. The eigenvalue equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{v} = \mathbf{0}$ is

$$(\mathbf{A} + 2\mathbf{I})\mathbf{v} = \begin{bmatrix} 2 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ -2 & 2 & -1 & 1 \\ 2 & -2 & 1 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

The third and fourth scalar equations here are the differences of the first and second equations, and therefore are redundant. Hence v is determined by the first two equations,

$$2a + c = 0$$
 and $2b + d = 0$.

We can choose *a* and *b* independently, then solve for *c* and *d*. Thereby we obtain *two* eigenvectors associated with the triple eigenvalue $\lambda_1 = -2$. The choice a = 1, b = 0 yields c = -2, d = 0 and thereby the eigenvector

$$\mathbf{u}_1 = \begin{bmatrix} 1 & 0 & -2 & 0 \end{bmatrix}^T$$

The choice a = 0, b = 1 yields c = 0, d = -2 and thereby the eigenvector

$$\mathbf{u}_2 = \begin{bmatrix} 0 & 1 & 0 & -2 \end{bmatrix}^T.$$

Because $\lambda_1 = -2$ has defect 1, we need a generalized eigenvector of rank 2, and hence a nonzero solution \mathbf{v}_2 of the equation

$$(\mathbf{A} + 2\mathbf{I})^2 \mathbf{v}_2 = \begin{bmatrix} 2 & 2 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \mathbf{v}_2 = \mathbf{0}.$$

Obviously,

$$\mathbf{v}_2 = \begin{bmatrix} 0 & 0 & 1 & -1 \end{bmatrix}^T$$

is such a vector, and we find that

$$(\mathbf{A} + 2\mathbf{I})\mathbf{v}_2 = \begin{bmatrix} 2 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ -2 & 2 & -1 & 1 \\ 2 & -2 & 1 & -1 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -2 \\ 2 \end{bmatrix} = \mathbf{v}_1$$

is nonzero, and therefore is an eigenvector associated with $\lambda_1 = -2$. Then $\{v_1, v_2\}$ is the length 2 chain we need.

The eigenvector \mathbf{v}_1 just found is neither of the two eigenvectors \mathbf{u}_1 and \mathbf{u}_2 found previously, but we observe that $\mathbf{v}_1 = \mathbf{u}_1 - \mathbf{u}_2$. For a length 1 chain \mathbf{w}_1 to complete the picture, we can choose any linear combination of \mathbf{u}_1 and \mathbf{u}_2 that is independent of \mathbf{v}_1 . For instance, we could choose either $\mathbf{w}_1 = \mathbf{u}_1$ or $\mathbf{w}_1 = \mathbf{u}_2$. However, we will see momentarily that the particular choice

$$\mathbf{w}_1 = \mathbf{u}_1 + \mathbf{u}_2 = \begin{bmatrix} 1 & 1 & -2 & -2 \end{bmatrix}^T$$

yields a solution of the system that is of physical interest.

Finally, the chains $\{v_0\}$, $\{w_1\}$, and $\{v_1, v_2\}$ yield the four independent solutions

$$\mathbf{x}_{1}(t) = \mathbf{v}_{0}e^{0\cdot t} = \begin{bmatrix} 1 & 1 & 0 & 0 \end{bmatrix}^{T},$$

$$\mathbf{x}_{2}(t) = \mathbf{w}_{1}e^{-2t} = \begin{bmatrix} 1 & 1 & -2 & -2 \end{bmatrix}^{T}e^{-2t},$$

$$\mathbf{x}_{3}(t) = \mathbf{v}_{1}e^{-2t} = \begin{bmatrix} 1 & -1 & -2 & 2 \end{bmatrix}^{T}e^{-2t},$$

$$\mathbf{x}_{4}(t) = (\mathbf{v}_{1}t + \mathbf{v}_{2})e^{-2t}$$

$$= \begin{bmatrix} t & -t & -2t + 1 & 2t - 1 \end{bmatrix}^{T}e^{-2t}$$
(33)

of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ in (32).

The four scalar components of the general solution

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + c_3 \mathbf{x}_3(t) + c_4 \mathbf{x}_4(t)$$

are described by the equations

$$x_{1}(t) = c_{1} + e^{-2t}(c_{2} + c_{3} + c_{4}t),$$

$$x_{2}(t) = c_{1} + e^{-2t}(c_{2} - c_{3} - c_{4}t),$$

$$x_{3}(t) = e^{-2t}(-2c_{2} - 2c_{3} + c_{4} - 2c_{4}t),$$

$$x_{4}(t) = e^{-2t}(-2c_{2} + 2c_{3} - c_{4} + 2c_{4}t).$$
(34)

Recall that $x_1(t)$ and $x_2(t)$ are the position functions of the two masses, whereas $x_3(t) = x'_1(t)$ and $x_4(t) = x'_2(t)$ are their respective velocity functions.

For instance, suppose that $x_1(0) = x_2(0) = 0$ and that $x'_1(0) = x'_2(0) = v_0$. Then the equations

$$x_{1}(0) = c_{1} + c_{2} + c_{3} = 0,$$

$$x_{2}(0) = c_{1} + c_{2} - c_{3} = 0,$$

$$x'_{1}(0) = -2c_{2} - 2c_{3} + c_{4} = v_{0},$$

$$x'_{2}(0) = -2c_{2} + 2c_{3} - c_{4} = v_{0}$$

are readily solved for $c_{1} = \frac{1}{2}v_{0}, c_{2} = -\frac{1}{2}v_{0},$ and $c_{3} = c_{4} = 0$, so

$$x_1(t) = x_2(t) = \frac{1}{2}v_0 \left(1 - e^{-2t}\right),$$

$$x_1'(t) = x_2'(t) = v_0 e^{-2t}.$$

In this case the two railway cars continue in the same direction with equal but exponentially damped velocities, approaching the displacements $x_1 = x_2 = \frac{1}{2}v_0$ as $t \to +\infty$.

It is of interest to interpret physically the individual generalized eigenvector solutions given in (33). The degenerate ($\lambda_0 = 0$) solution

$$\mathbf{x}_1(t) = \begin{bmatrix} 1 & 1 & 0 & 0 \end{bmatrix}^T$$

describes the two masses at rest with position functions $x_1(t) \equiv 1$ and $x_2(t) \equiv 1$. The solution

$$\mathbf{x}_2(t) = \begin{bmatrix} 1 & 1 & -2 & -2 \end{bmatrix}^T e^{-2t}$$

corresponding to the carefully chosen eigenvector \mathbf{w}_1 describes damped motions $x_1(t) = e^{-2t}$ and $x_2(t) = e^{-2t}$ of the two masses, with equal velocities in the same direction. Finally, the solutions $\mathbf{x}_3(t)$ and $\mathbf{x}_4(t)$ resulting from the length 2 chain { $\mathbf{v}_1, \mathbf{v}_2$ } both describe damped motion with the two masses moving in opposite directions.

The methods of this section apply to complex multiple eigenvalues just as to real multiple eigenvalues (although the necessary computations tend to be somewhat lengthy). Given a complex conjugate pair $\alpha \pm \beta i$ of eigenvalues of multiplicity k, we work with one of them (say, $\alpha - \beta i$) as if it were real to find k independent complex-valued solutions. The real and imaginary parts of these complex-valued solutions then provide 2k real-valued solutions associated with the two eigenvalues $\lambda = \alpha - \beta i$ and $\overline{\lambda} = \alpha + \beta i$ each of multiplicity k. See Problems 33 and 34.

The Jordan Normal Form

In Section 6.2 we saw that if the $n \times n$ matrix **A** has a complete set $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of *n* linearly independent eigenvectors, then it is similar to a diagonal matrix. In particular,

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D},\tag{35}$$

where $\mathbf{P} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{bmatrix}$ and the diagonal elements of the diagonal matrix **D** are the corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ (not necessarily distinct). This result is a special case of the following general theorem, a proof of which can be found in Appendix B of Gilbert Strang, *Linear Algebra and Its Applications* (4th edition, Brooks Cole, 2006).

THEOREM 1 Jordan Normal Form

If the $n \times n$ matrix **A** has *s* linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_s$, then it is similar to a block-diagonal matrix of the *Jordan normal form*

$$\mathbf{J} = \begin{bmatrix} \mathbf{J}_{1} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{J}_{2} & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & \mathbf{J}_{n} \end{bmatrix},$$
(36)

where each submatrix J_i is a $k \times k$ Jordan block of the form

$$\mathbf{J}_{i} = \begin{bmatrix} \lambda_{i} & 1 & 0 & \cdots & 0 \\ 0 & \lambda_{i} & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_{i} & 1 \\ 0 & 0 & 0 & \cdots & \lambda_{i} \end{bmatrix},$$
(37)

with λ_i being the eigenvalue of **A** corresponding to the eigenvector \mathbf{v}_i ; if k = 1, then $\mathbf{J}_i = [\lambda_i]$.

If the Jordan block \mathbf{J}_i in (37) is of size $k \times k$, then it corresponds to a length k chain of generalized eigenvectors based on the (ordinary) eigenvector \mathbf{v}_i . If all these generalized eigenvectors are arranged as column vectors in proper order corresponding to the appearance of the Jordan blocks in (36), the result is a nonsingular $n \times n$ matrix \mathbf{Q} such that

$$\mathbf{Q}^{-1}\mathbf{A}\mathbf{Q} = \mathbf{J}.$$
 (38)

The block-diagonal matrix **J** such that $\mathbf{A} = \mathbf{Q}\mathbf{J}\mathbf{Q}^{-1}$ is called the **Jordan normal** form of the matrix **A** and is unique [except for the order of appearance of the Jordan blocks in (36)].

Example 7 In Example 3 we saw that the matrix

$$\mathbf{A} = \begin{bmatrix} 1 & -3 \\ 3 & 7 \end{bmatrix}$$

has the single eigenvalue $\lambda = 4$ and the associated length 2 chain of generalized eigenvectors $\{\mathbf{v}_1, \mathbf{v}_2\}$, where $\mathbf{v}_1 = \begin{bmatrix} -3 & 3 \end{bmatrix}^T$ and $\mathbf{v}_2 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$, based on the eigenvector $\mathbf{v}_1 = \mathbf{A}\mathbf{v}_2$.

If we define

$$\mathbf{Q} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 \end{bmatrix} = \begin{bmatrix} -3 & 1 \\ 3 & 0 \end{bmatrix},$$

then we find that the Jordan normal form of A is

$$\mathbf{J} = \mathbf{Q}^{-1}\mathbf{A}\mathbf{Q} = \begin{bmatrix} 4 & 1 \\ 0 & 4 \end{bmatrix}.$$

Here $\mathbf{J} = \mathbf{J}_1$ is a single 2 × 2 Jordan block corresponding to the single eigenvalue $\lambda = 4$ of **A**.

Example 8

In Example 6 we saw that the matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -2 & 2 & -3 & 1 \\ 2 & -2 & 1 & -3 \end{bmatrix}$$

has the distinct eigenvalue $\lambda_1 = 0$ corresponding to the eigenvector

$$\mathbf{v}_1 = \begin{bmatrix} 1 & 1 & 0 & 0 \end{bmatrix}^T$$

and the triple eigenvalue $\lambda_2 = -2$ corresponding both to the eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 & 0 & -2 & 0 \end{bmatrix}^T$ and to the length 2 chain of generalized eigenvectors $\{\mathbf{v}_3, \mathbf{v}_4\}$ with $\mathbf{v}_3 = \begin{bmatrix} 1 & -1 & -2 & 2 \end{bmatrix}^T$ and $\mathbf{v}_4 = \begin{bmatrix} 0 & 0 & 1 & -1 \end{bmatrix}^T$, such that $\mathbf{v}_3 = (\mathbf{A} + 2\mathbf{I})\mathbf{v}_4$. If we define

$$\mathbf{Q} = \begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 & \mathbf{v}_4 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & -1 & 0 \\ 0 & -2 & -2 & 1 \\ 0 & 0 & 2 & -1 \end{bmatrix}$$

then (using a computer algebra system to ease the labor of calculation) we find that the Jordan normal form of A is $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

$$\mathbf{J} = \mathbf{Q}^{-1} \mathbf{A} \mathbf{Q} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & -2 \end{bmatrix}$$

Here we see the 1 × 1 Jordan block $\mathbf{J}_1 = \begin{bmatrix} 0 \end{bmatrix}$ corresponding to the eigenvalue $\lambda_1 = 0$ of \mathbf{A} , as well as the two Jordan blocks $\mathbf{J}_2 = \begin{bmatrix} -2 \end{bmatrix}$ and $\mathbf{J}_3 = \begin{bmatrix} -2 & 1 \\ 0 & -2 \end{bmatrix}$ corresponding to the two linearly independent eigenvectors \mathbf{v}_2 and \mathbf{v}_3 associated with the eigenvalue $\lambda_2 = -2$.

The General Cayley-Hamilton Theorem

In Section 6.3, we showed that every diagonalizable matrix **A** satisfies its characteristic equation $p(\lambda) = |\mathbf{A} - \lambda \mathbf{I}| = 0$, that is, $p(\mathbf{A}) = \mathbf{0}$. We can now use the Jordan normal form to show that this is true whether or not **A** is diagonalizable.

If $\mathbf{J} = \mathbf{Q}^{-1}\mathbf{A}\mathbf{Q}$ is the Jordan normal form of the matrix \mathbf{A} , then $p(\mathbf{A}) = \mathbf{Q}p(\mathbf{J})\mathbf{Q}^{-1}$, just as in the conclusion to the proof of Theorem 1 in Section 6.3. It therefore suffices to show that $p(\mathbf{J}) = \mathbf{0}$. If the Jordan blocks $\mathbf{J}_1, \mathbf{J}_2, \dots, \mathbf{J}_s$ in (36) have sizes k_1, k_2, \dots, k_s —that is, \mathbf{J}_i is a $k_i \times k_i$ matrix—and corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$ (respectively), then

$$p(\lambda) = (\lambda_1 - \lambda)^{k_1} (\lambda_2 - \lambda)^{k_2} \cdots (\lambda_s - \lambda)^{k_s},$$

and so

$$p(\mathbf{J}) = (\lambda_1 \mathbf{I} - \mathbf{J})^{k_1} (\lambda_2 \mathbf{I} - \mathbf{J})^{k_2} \cdots (\lambda_s \mathbf{I} - \mathbf{J})^{k_s}.$$
(39)

Х

Х

Х

Now $p(\mathbf{J})$ has the same block-diagonal structure as \mathbf{J} itself, and we see from (39) that the *i*th block of $p(\mathbf{J})$ involves the factor

$$(\lambda_i \mathbf{I} - \mathbf{J}_i)^{k_i} = \begin{bmatrix} 0 & -1 & 0 & \cdots & 0 \\ 0 & 0 & -1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & -1 \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix}^{k_i},$$

where the $k_i \times k_i$ matrix $\lambda_i \mathbf{I} - \mathbf{J}_i$ has all elements zero except on its first superdiagonal. It is easily seen that $(\lambda_i \mathbf{I} - \mathbf{J}_i)^2$ then has nonzero elements only on its second superdiagonal; $(\lambda_i \mathbf{I} - \mathbf{J}_i)^3$ has nonzero elements only on its third superdiagonal; and so on, in turn, until we see that $(\lambda_i \mathbf{I} - \mathbf{J}_i)^{k_i} = \mathbf{0}$. Hence it follows from (39) that the *i*th block of $p(\mathbf{J})$ is the $k_i \times k_i$ zero matrix. This being true for each i = 1, 2, ..., s, we conclude that $p(\mathbf{J}) = \mathbf{0}$, as desired. This completes the proof of the general case of the Cayley-Hamilton theorem.

13. $\mathbf{x}' = \begin{bmatrix} -1 & 0 & 1 \\ 0 & 1 & -4 \\ 0 & 1 & -3 \end{bmatrix} \mathbf{x}$

7.6 Problems

Find general solutions of the systems in Problems 1 through 22. In Problems 1 through 6, use a computer system or graphing calculator to construct a direction field and typical solution curves for the given system.

1. $\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} \mathbf{x}$	2. $\mathbf{x}' = \begin{bmatrix} 3 & -1 \\ 1 & -1 \end{bmatrix} \mathbf{x}$	14. $\mathbf{x}' = \begin{bmatrix} 0 & 0 & 1 \\ -5 & -1 & -5 \\ 4 & 1 & 2 \end{bmatrix} \mathbf{x}$
$\begin{bmatrix} -1 & -4 \end{bmatrix}$ 3. $\mathbf{x}' = \begin{bmatrix} 1 & -2 \\ 2 & 5 \end{bmatrix} \mathbf{x}$	$4. \ \mathbf{x}' = \begin{bmatrix} 3 & -1 \\ 1 & 5 \end{bmatrix} \mathbf{x}$	$15. \ \mathbf{x}' = \begin{bmatrix} -2 & -9 & 0 \\ 1 & 4 & 0 \\ 1 & 3 & 1 \end{bmatrix} \mathbf{x}$
5. $\mathbf{x}' = \begin{bmatrix} 7 & 1 \\ -4 & 3 \end{bmatrix} \mathbf{x}$	$6. \ \mathbf{x}' = \begin{bmatrix} 1 & -4 \\ 4 & 9 \end{bmatrix} \mathbf{x}$	16. $\mathbf{x}' = \begin{bmatrix} 1 & 0 & 0 \\ -2 & -2 & -3 \\ 2 & 3 & 4 \end{bmatrix} \mathbf{x}$
7. $\mathbf{x}' = \begin{bmatrix} -7 & 9 & 7 \\ 0 & 0 & 2 \end{bmatrix} \mathbf{x}$		17. $\mathbf{x}' = \begin{bmatrix} 1 & 0 & 0 \\ 18 & 7 & 4 \\ -27 & -9 & -5 \end{bmatrix} \mathbf{x}$
$8. \ \mathbf{x}' = \begin{bmatrix} 25 & 12 & 0 \\ -18 & -5 & 0 \\ 6 & 6 & 13 \end{bmatrix} \mathbf{x}$		$18. \ \mathbf{x}' = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 3 & 1 \\ 2 & 0 & 0 \end{bmatrix} \mathbf{x}$
$\mathbf{9. x'} = \begin{bmatrix} -19 & 12 & 84 \\ 0 & 5 & 0 \\ -8 & 4 & 33 \end{bmatrix} \mathbf{x}$		$\mathbf{19. x'} = \begin{bmatrix} 1 & -4 & 0 & -2 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$
10. $\mathbf{x}' = \begin{bmatrix} -13 & 40 & -48 \\ -8 & 23 & -24 \\ 0 & 0 & 3 \end{bmatrix} \mathbf{x}$		$\begin{bmatrix} 6 & -12 & -1 & -6 \\ 0 & -4 & 0 & -1 \\ \hline 2 & 1 & 0 & 1 \end{bmatrix}$
11. $\mathbf{x}' = \begin{bmatrix} -3 & 0 & -4 \\ -1 & -1 & -1 \\ 1 & 0 & 1 \end{bmatrix} \mathbf{x}$		$\mathbf{20. x'} = \begin{bmatrix} 0 & 2 & 1 & 0 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 2 \end{bmatrix}$
12. $\mathbf{x}' = \begin{bmatrix} -1 & 0 & 1 \\ 0 & -1 & 1 \\ 1 & -1 & -1 \end{bmatrix} \mathbf{x}$		21. $\mathbf{x}' = \begin{bmatrix} -1 & -4 & 0 & 0 \\ 1 & 3 & 0 & 0 \\ 1 & 2 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$

452 CHAPTER 7 LINEAR SYSTEMS OF DIFFERENTIAL EQUATIONS

22.
$$\mathbf{x}' = \begin{bmatrix} 1 & 3 & 7 & 0 \\ 0 & -1 & -4 & 0 \\ 0 & 1 & 3 & 0 \\ 0 & -6 & -14 & 1 \end{bmatrix} \mathbf{x}$$

In Problems 23 through 32 the eigenvalues of the coefficient matrix **A** are given. Find a general solution of the indicated system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Especially in Problems 29 through 32, use of a computer algebra system (as in the application material for this section) may be useful.

23.
$$\mathbf{x}' = \begin{bmatrix} 39 & 8 & -16 \\ -36 & -5 & 16 \\ 72 & 16 & -29 \end{bmatrix} \mathbf{x}; \quad \lambda = -1, 3, 3$$

24. $\mathbf{x}' = \begin{bmatrix} 28 & 50 & 100 \\ 15 & 33 & 60 \\ -15 & -30 & -57 \end{bmatrix} \mathbf{x}; \quad \lambda = -2, 3, 3$
25. $\mathbf{x}' = \begin{bmatrix} -2 & 17 & 4 \\ -1 & 6 & 1 \\ 0 & 1 & 2 \end{bmatrix} \mathbf{x}; \quad \lambda = 2, 2, 2$
26. $\mathbf{x}' = \begin{bmatrix} 5 & -1 & 1 \\ 1 & 3 & 0 \\ -3 & 2 & 1 \end{bmatrix} \mathbf{x}; \quad \lambda = 3, 3, 3$
27. $\mathbf{x}' = \begin{bmatrix} -3 & 5 & -5 \\ 3 & -1 & 3 \\ 8 & -8 & 10 \end{bmatrix} \mathbf{x}; \quad \lambda = 2, 2, 2$
28. $\mathbf{x}' = \begin{bmatrix} -15 & -7 & 4 \\ 34 & 16 & -11 \\ 17 & 7 & 5 \end{bmatrix} \mathbf{x}; \quad \lambda = 2, 2, 2$
29. $\mathbf{x}' = \begin{bmatrix} -1 & 1 & 1 & -2 \\ 7 & -4 & -6 & 11 \\ 5 & -1 & 1 & 3 \\ 6 & -2 & -2 & 6 \end{bmatrix} \mathbf{x}; \quad \lambda = -1, -1, 2, 2$
30. $\mathbf{x}' = \begin{bmatrix} 2 & 1 & -2 & 1 \\ 0 & 3 & -5 & 3 \\ 0 & -13 & 22 & -12 \\ 0 & -27 & 45 & -25 \end{bmatrix} \mathbf{x}; \quad \lambda = -1, -1, 2, 2$
31. $\mathbf{x}' = \begin{bmatrix} 35 & -12 & 4 & 30 \\ 22 & -8 & 3 & 19 \\ -10 & 3 & 0 & -9 \\ -27 & 9 & -3 & -23 \end{bmatrix} \mathbf{x}; \quad \lambda = 1, 1, 1, 1$
32. $\mathbf{x}' = \begin{bmatrix} 11 & -1 & 26 & 6 & -3 \\ 0 & 3 & 0 & 0 & 0 \\ -9 & 0 & -24 & -6 & 3 \\ 3 & 0 & 9 & 5 & -1 \\ -48 & -3 & -138 & -30 & 18 \end{bmatrix} \mathbf{x};$

33. The characteristic equation of the coefficient matrix **A** of the system

$$\mathbf{x}' = \begin{bmatrix} 3 & -4 & 1 & 0 \\ 4 & 3 & 0 & 1 \\ 0 & 0 & 3 & -4 \\ 0 & 0 & 4 & 3 \end{bmatrix} \mathbf{x}$$

is

$$\phi(\lambda) = (\lambda^2 - 6\lambda + 25)^2 = 0.$$

Therefore, **A** has the repeated complex conjugate pair $3 \pm 4i$ of eigenvalues. First show that the complex vectors

$$\mathbf{v}_1 = \begin{bmatrix} 1 & i & 0 & 0 \end{bmatrix}^T$$
 and $\mathbf{v}_2 = \begin{bmatrix} 0 & 0 & 1 & i \end{bmatrix}^T$

form a length 2 chain $\{\mathbf{v}_1, \mathbf{v}_2\}$ associated with the eigenvalue $\lambda = 3 - 4i$. Then calculate the real and imaginary parts of the complex-valued solutions

$$\mathbf{v}_1 e^{\lambda t}$$
 and $(\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t}$

to find four independent real-valued solutions of $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

34. The characteristic equation of the coefficient matrix **A** of the system

$$\mathbf{x}' = \begin{bmatrix} 2 & 0 & -8 & -3\\ -18 & -1 & 0 & 0\\ -9 & -3 & -25 & -9\\ 33 & 10 & 90 & 32 \end{bmatrix} \mathbf{x}$$

is

$$\phi(\lambda) = (\lambda^2 - 4\lambda + 13)^2 = 0.$$

Therefore, **A** has the repeated complex conjugate pair $2 \pm 3i$ of eigenvalues. First show that the complex vectors

$$\mathbf{v}_1 = \begin{bmatrix} -i & 3+3i & 0 & -1 \end{bmatrix}^T,$$
$$\mathbf{v}_2 = \begin{bmatrix} 3 & -10+9i & -i & 0 \end{bmatrix}^T$$

form a length 2 chain $\{\mathbf{v}_1, \mathbf{v}_2\}$ associated with the eigenvalue $\lambda = 2 + 3i$. Then calculate (as in Problem 33) four independent real-valued solutions of $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

35. Railway cars Find the position functions $x_1(t)$ and $x_2(t)$ of the railway cars of Fig. 7.5.2 if the physical parameters are given by

$$m_1 = m_2 = c_1 = c_2 = c = k = 1$$

and the initial conditions are

$$x_1(0) = x_2(0) = 0, \quad x'_1(0) = x'_2(0) = v_0.$$

How far do the cars travel before stopping?

- **36.** Railway cars Repeat Problem 35 under the assumption that car 1 is shielded from air resistance by car 2, so now $c_1 = 0$. Show that, before stopping, the cars travel twice as far as those of Problem 35.
- **37–46.** Use the method of Examples 7 and 8 to find the Jordan normal form **J** of each coefficient matrix **A** given in Problems 23 through 32 (respectively).

7.6 Application Defective Eigenvalues and Generalized Eigenvectors



computing resources including *Maple/Mathematica/*MATLAB.

A typical computer algebra system can calculate both the eigenvalues of a given matrix **A** and the linearly independent (ordinary) eigenvectors associated with each eigenvalue. For instance, consider the 4×4 matrix

$$\mathbf{A} = \begin{bmatrix} 35 & -12 & 4 & 30\\ 22 & -8 & 3 & 19\\ -10 & 3 & 0 & -9\\ -27 & 9 & -3 & -23 \end{bmatrix} \tag{1}$$

of Problem 31 in this section. When the matrix \mathbf{A} has been entered, the *Maple* calculation

```
with(linalg): eigenvectors(A);
[1, 4, {[-1, 0, 1, 1], [0, 1, 3, 0]}]
```

or the Mathematica calculation

Eigensystem[A] {{1,1,1,1}, {{-3,-1,0,3}, {0,1,3,0}, {0,0,0,0}, {0,0,0,0}}}

reveals that the matrix A in Eq. (1) has the single eigenvalue $\lambda = 1$ of multiplicity 4 with only two independent associated eigenvectors \mathbf{v}_1 and \mathbf{v}_2 . The MATLAB command

[V, D] = eig(sym(A))

provides the same information. The eigenvalue $\lambda = 1$ therefore has defect d = 2. If **B** = **A** – (1)**I**, you should find that **B**² \neq **0** but **B**³ = **0**. If

$$\mathbf{u}_1 = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}^T, \quad \mathbf{u}_2 = \mathbf{B}\mathbf{u}_1, \quad \mathbf{u}_3 = \mathbf{B}\mathbf{u}_2,$$

then $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ should be a length 3 chain of generalized eigenvectors based on the ordinary eigenvector \mathbf{u}_3 (which should be a linear combination of the original eigenvectors \mathbf{v}_1 and \mathbf{v}_2). Use your computer algebra system to carry out this construction, and finally write four linearly independent solutions of the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

For a more exotic matrix to investigate, consider the MATLAB's gallery (5) example matrix

	-9	11	-21	63	-252	
	70	-69	141	-421	1684	
$\mathbf{A} =$	-575	575	-1149	3451	-13801	(2)
	3891	-3891	7782	-23345	93365	
	1024	-1024	2048	-6144	24572	

Use appropriate commands like those illustrated here to show that **A** has a single eigenvalue $\lambda = 0$ of multiplicity 5 and defect 4. Noting that $\mathbf{A} - (0)\mathbf{I} = \mathbf{A}$, you should find that $\mathbf{A}^4 \neq \mathbf{0}$ but that $\mathbf{A}^5 = \mathbf{0}$. Hence calculate the vectors

$$\mathbf{u}_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \end{bmatrix}^T$$
, $\mathbf{u}_2 = \mathbf{A}\mathbf{u}_1$, $\mathbf{u}_3 = \mathbf{A}\mathbf{u}_2$, $\mathbf{u}_4 = \mathbf{A}\mathbf{u}_3$, $\mathbf{u}_5 = \mathbf{A}\mathbf{u}_4$.

You should find that \mathbf{u}_5 is a nonzero vector such that $A\mathbf{u}_5 = \mathbf{0}$, and is therefore an (ordinary) eigenvector of **A** associated with the eigenvalue $\lambda = 0$. Thus $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4, \mathbf{u}_5\}$ is a length 5 chain of generalized eigenvectors of the matrix **A** in Eq. (2), and you can finally write five linearly independent solutions of the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.
7.7 Numerical Methods for Systems

We now discuss the numerical approximation of solutions of systems of differential equations. Our goal is to apply the methods of Sections 2.4 through 2.6 to the initial value problem

$$\mathbf{x}' = \mathbf{f}(t, \mathbf{x}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \tag{1}$$

for a system of m first-order differential equations. In (1) the independent variable is the scalar t, and

$$\mathbf{x} = (x_1, x_2, \dots, x_m)$$
 and $\mathbf{f} = (f_1, f_2, \dots, f_m)$

are vector-valued functions. If the component functions of **f** and their first-order partial derivatives are all continuous in a neighborhood of the point (t_0, \mathbf{x}_0) , then Theorems 3 and 4 of Appendix A guarantee the existence and uniqueness of a solution $\mathbf{x} = \mathbf{x}(t)$ of (1) on some subinterval [of the *t*-axis] containing t_0 . With this assurance we can proceed to discuss the numerical approximation of this solution.

Beginning with step size *h*, we want to approximate the values of $\mathbf{x}(t)$ at the points t_1, t_2, t_3, \ldots , where $t_{n+1} = t_n + h$ for $n \ge 0$. Suppose that we have already computed the *approximations*

$$\mathbf{X}_1, \quad \mathbf{X}_2, \quad \mathbf{X}_3, \quad \dots, \quad \mathbf{X}_n$$

to the actual values

$$\mathbf{x}(t_1), \quad \mathbf{x}(t_2), \quad \mathbf{x}(t_3), \quad \dots, \quad \mathbf{x}(t_n)$$

of the exact solution of the system in (1). We can then make the step from \mathbf{x}_n to the next approximation $\mathbf{x}_{n+1} \approx \mathbf{x}(t_{n+1})$ by any one of the methods of Sections 2.4 through 2.6. Essentially all that is required is to write the iterative formula of the selected method in the vector notation of the present discussion.

Euler Methods for Systems

For example, the iterative formula of Euler's method for systems is

 $\mathbf{x}_{n+1} = \mathbf{x}_n + h\mathbf{f}(t, \mathbf{x}_n).$ (2)

To examine the case m = 2 of a pair of first-order differential equations, let us write

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix} \quad \text{and} \quad \mathbf{f} = \begin{bmatrix} f \\ g \end{bmatrix}.$$

Then the initial value problem in (1) is

$$\begin{aligned} x' &= f(t, x, y), \quad x(t_0) = x_0, \\ y' &= g(t, x, y), \quad y(t_0) = y_0, \end{aligned}$$
 (3)

and the scalar components of the vector formula in (2) are

$$x_{n+1} = x_n + hf(t_n, x_n, y_n),$$

$$y_{n+1} = y_n + hg(t_n, x_n, y_n).$$
(4)

Note that each iterative formula in (4) has the form of a single Euler iteration, but with y_n inserted like a parameter in the first formula (for x_{n+1}) and with x_n inserted like a parameter in the second formula (for y_{n+1}). The generalization to the system in (3) of each of the other methods in Sections 2.4 through 2.6 follows a similar pattern.

The improved Euler method for systems consists at each step of calculating first the predictor

$$\mathbf{u}_{n+1} = \mathbf{x}_n + h\mathbf{f}(t_n, \mathbf{x}_n) \tag{5}$$

and then the corrector

 \succ

$$\mathbf{x}_{n+1} = \mathbf{x}_n + \frac{h}{2} [\mathbf{f}(t_n, \mathbf{x}_n) + \mathbf{f}(t_{n+1}, \mathbf{u}_{n+1})].$$
(6)

For the case of the two-dimensional initial value problem in (3), the scalar components of the formulas in (5) and (6) are

$$u_{n+1} = x_n + hf(t_n, x_n, y_n), v_{n+1} = y_n + hg(t_n, x_n, y_n)$$
(7)

and

$$x_{n+1} = x_n + \frac{h}{2} [f(t_n, x_n, y_n) + f(t_{n+1}, u_{n+1}, v_{n+1})],$$

$$y_{n+1} = y_n + \frac{h}{2} [g(t_n, x_n, y_n) + g(t_{n+1}, u_{n+1}, v_{n+1})].$$
(8)

cample 1 Consider the initial value problem

$$\begin{aligned} x' &= 3x - 2y, & x(0) = 3; \\ y' &= 5x - 4y, & y(0) = 6. \end{aligned}$$
 (9)

The exact solution of the system in (9) is

$$x(t) = 2e^{-2t} + e^t, \qquad y(t) = 5e^{-2t} + e^t.$$
 (10)

Here we have f(x, y) = 3x - 2y and g(x, y) = 5x - 4y in (3), so the Euler iterative formulas in (4) are

$$x_{n+1} = x_n + h \cdot (3x_n - 2y_n), \quad y_{n+1} = y_n + h \cdot (5x_n - 4y_n).$$

With step size h = 0.1 we calculate

$$x_1 = 3 + (0.1) \cdot [3 \cdot 3 - 2 \cdot 6] = 2.7,$$

$$y_1 = 6 + (0.1) \cdot [5 \cdot 3 - 4 \cdot 6] = 5.1$$

and

$$x_2 = 2.7 + (0.1) \cdot [3 \cdot (2.7) - 2 \cdot (5.1)] = 2.49,$$

$$y_2 = 5.1 + (0.1) \cdot [5 \cdot (2.7) - 4 \cdot (5.1)] = 4.41.$$

The actual values at $t_2 = 0.2$ given by (10) are $x(0.2) \approx 2.562$ and $y(0.2) \approx 4.573$.

To compute the improved Euler approximations to x(0.2) and y(0.2) with a single step of size h = 0.2, we first calculate the predictors

$$u_1 = 3 + (0.2) \cdot [3 \cdot 3 - 2 \cdot 6] = 2.4,$$

$$v_1 = 6 + (0.2) \cdot [5 \cdot 3 - 4 \cdot 6] = 4.2.$$

Then the corrector formulas in (8) yield

$$x_1 = 3 + (0.1) \cdot ([3 \cdot 3 - 2 \cdot 6] + [3 \cdot (2.4) - 2 \cdot (4.2)]) = 2.58,$$

$$y_1 = 6 + (0.1) \cdot ([5 \cdot 3 - 4 \cdot 6] + [5 \cdot (2.4) - 4 \cdot (4.2)]) = 4.62.$$

As we would expect, a single improved Euler step gives better accuracy than two ordinary Euler steps.

The Runge–Kutta Method and Second-Order Equations

The vector version of the iterative formula for the Runge-Kutta method is

>
$$\mathbf{x}_{n+1} = \mathbf{x}_n + \frac{h}{6}(\mathbf{k}_1 + 2\mathbf{k}_2 + 2\mathbf{k}_3 + \mathbf{k}_4),$$
 (11)

where the vectors \mathbf{k}_1 , \mathbf{k}_2 , \mathbf{k}_3 , and \mathbf{k}_4 are defined (by analogy with Eqs. (5a)–(5d) of Section 2.6) as follows:

$$\mathbf{k}_{1} = \mathbf{f}(t_{n}, \mathbf{x}_{n}),$$

$$\mathbf{k}_{2} = \mathbf{f}\left(t_{n} + \frac{1}{2}h, \mathbf{x}_{n} + \frac{1}{2}h\mathbf{k}_{1}\right),$$

$$\mathbf{k}_{3} = \mathbf{f}\left(t_{n} + \frac{1}{2}h, \mathbf{x}_{n} + \frac{1}{2}h\mathbf{k}_{2}\right),$$

$$\mathbf{k}_{4} = \mathbf{f}(t_{n} + h, \mathbf{x}_{n} + h\mathbf{k}_{3}).$$
(12)

To describe in scalar notation the Runge-Kutta method for the twodimensional initial value problem

$$\begin{aligned} x' &= f(t, x, y), & x(t_0) = x_0, \\ y' &= g(t, x, y), & y(t_0) = y_0, \end{aligned}$$
 (3)

let us write

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}, \quad \mathbf{f} = \begin{bmatrix} f \\ g \end{bmatrix}, \text{ and } \mathbf{k}_i = \begin{bmatrix} F_i \\ G_i \end{bmatrix}.$$

Then the Runge–Kutta iterative formulas for the step from (x_n, y_n) to the next approximation $(x_{n+1}, y_{n+1}) \approx (x(t_{n+1}), y(t_{n+1}))$ are

$$x_{n+1} = x_n + \frac{h}{6}(F_1 + 2F_2 + 2F_3 + F_4),$$

$$y_{n+1} = y_n + \frac{h}{6}(G_1 + 2G_2 + 2G_3 + G_4),$$
(13)

where the values F_1 , F_2 , F_3 , and F_4 of the function f are

$$F_{1} = f(t_{n}, x_{n}, y_{n}),$$

$$F_{2} = f\left(t_{n} + \frac{1}{2}h, x_{n} + \frac{1}{2}hF_{1}, y_{n} + \frac{1}{2}hG_{1}\right),$$

$$F_{3} = f\left(t_{n} + \frac{1}{2}h, x_{n} + \frac{1}{2}hF_{2}, y_{n} + \frac{1}{2}hG_{2}\right),$$

$$F_{4} = f(t_{n} + h, x_{n} + hF_{3}, y_{n} + hG_{3});$$
(14)

 G_1, G_2, G_3 , and G_4 are the similarly defined values of the function g.

Perhaps the most common application of the two-dimensional Runge-Kutta method is to the numerical solution of second-order initial value problems of the form

$$\begin{aligned} x'' &= g(t, x, x'), \\ x(t_0) &= x_0, \quad x'(t_0) = y_0. \end{aligned} \tag{15}$$

If we introduce the auxiliary variable y = x', then the problem in (15) translates into the two-dimensional first-order problem

$$\begin{aligned} x' &= y, & x(t_0) = x_0, \\ y' &= g(t, x, y), & y(t_0) = y_0. \end{aligned}$$
 (16)

This is a problem of the form in (3) with f(t, x, y) = y.

7.7 NUMERICAL METHODS FOR SYSTEMS457

If the functions f and g are not too complicated, then it is feasible to carry out manually a reasonable number of steps of the two-dimensional Runge–Kutta method described here. But the first operating electronic computers were constructed (during World War II) specifically to implement methods similar to the Runge–Kutta method for the numerical computation of trajectories of artillery projectiles. The application material for this section lists TI-Nspire CX CAS and Python versions of Program RK2DIM that can be used with two-dimensional systems.

Example 2

The exact solution of the initial value problem

$$x'' = -x; \quad x(0) = 0, \quad x'(0) = 1$$
 (17)

is $x(t) = \sin t$. The substitution y = x' translates (17) into the two-dimensional problem

$$x' = y,$$
 $x(0) = 0;$
 $y' = -x,$ $y(0) = 1,$
(18)

which has the form in (3) with f(t, x, y) = y and g(t, x, y) = -x. The table in Fig. 7.7.1 shows the results produced for $0 \le t \le 5$ (radians) using Program RK2DIM with step size h = 0.05. The values shown for $x = \sin t$ and $y = \cos t$ are all accurate to five decimal places.

Example 3

t	$x = \sin t$	$y = \cos t$
0.5	+0.47943	+0.87758
1.0	+0.84147	+0.54030
1.5	+0.99749	+0.07074
2.0	+0.90930	-0.41615
2.5	+0.59847	-0.80114
3.0	+0.14112	-0.98999
3.5	-0.35078	-0.93646
4.0	-0.75680	-0.65364
4.5	-0.97753	-0.21080
5.0	-0.95892	+0.28366

FIGURE 7.7.1. Runge–Kutta values (with h = 0.05) for the problem in Eq. (18).

Lunar lander In Example 4 of Section 2.3 we considered a lunar lander that initially is falling freely toward the surface of the moon. Its retrorockets, when fired, provide a deceleration of $T = 4 \text{ m/s}^2$. We found that a soft touchdown at the lunar surface is achieved by igniting these retrorockets when the lander is at a height of 41,870 meters (just over 26 miles) above the surface and is then descending at the rate of 450 m/s.

Now we want to compute the *descent time* of the lunar lander. Let the distance x(t) of the lander from the center of the moon be measured in meters and measure time t in seconds. According to the analysis in Section 2.3 (where we used r(t) instead of x(t)), x(t) satisfies the initial value problem

$$\frac{d^2x}{dt^2} = T - \frac{GM}{x^2} = 4 - \frac{4.9044 \times 10^{12}}{x^2},$$
(19)
 $x(0) = R + 41870 = 1,781,870, \quad x'(0) = -450$

where $G \approx 6.6726 \times 10^{-11} \text{ N} \cdot (\text{m/kg})^2$ is the universal gravitational constant and $M = 7.35 \times 10^{22} \text{ kg}$ and $R = 1.74 \times 10^6 \text{ m}$ are the mass and radius of the moon. We seek the value of t when x(t) = R = 1,740,000.

The problem in (19) is equivalent to the first-order system

$$\frac{dx}{dt} = y, \quad x(0) = 1,781,870;$$

$$\frac{dy}{dt} = 4 - \frac{4.9044 \times 10^{12}}{x^2}, \quad y(0) = -450.$$
(20)

The table in Fig. 7.7.2 shows the result of a Runge–Kutta approximation with step size h = 1 (the indicated data agreeing with those obtained with step size h = 2). Evidently, touchdown on the lunar surface (x = 1,740,000) occurs at some time between t = 180 and t = 190 seconds. The table in Fig. 7.7.3 shows a second Runge–Kutta approximation with t(0) = 180, x(0) = 1,740,059, y(0) = -16.83, and h = 0.1. Now it is apparent that the lander's time of descent to the lunar surface is very close to 187 seconds; that is, 3 min 7 s. (The final velocity terms in these two tables are positive because the lander would begin to ascend if its retrorockets were not turned off at touchdown.)

(s)	<i>x</i> (m)	v (m/s)
0	1.781.870	-450.00
20	1,773,360	-401.04
40	1,765,826	-352.37
60	1,759,264	-303.95
80	1,753,667	-255.74
100	1,749,033	-207.73
120	1,745,357	-159.86
140	1,742,637	-112.11
160	1,740,872	-64.45
180	1,740,059	-16.83
200	1,740,199	30.77

FIGURE 7.7.2. The lander's descent to the lunar surface.

FIGURE 7.7.3. Focusing on the lunar lander's soft touchdown.

Higher-Order Systems

As we saw in Section 7.1, any system of higher-order differential equations can be replaced with an equivalent system of first-order differential equations. For example, consider the system

$$\begin{aligned} x'' &= F(t, x, y, x', y'), \\ y'' &= G(t, x, y, x', y') \end{aligned}$$
 (21)

of second-order equations. If we substitute

$$x = x_1$$
, $y = x_2$, $x' = x_3 = x'_1$, $y' = x_4 = x'_2$,

then we get the equivalent system

$$\begin{aligned}
x_1' &= x_3, \\
x_2' &= x_4, \\
x_3' &= F(t, x_1, x_2, x_3, x_4), \\
x_4' &= G(t, x_1, x_2, x_3, x_4)
\end{aligned}$$
(22)

of four first-order equations in the unknown functions $x_1(t) = x(t)$, $x_2(t) = y(t)$, $x_3(t)$, and $x_4(t)$. It would be a routine (if slightly tedious) matter to write a fourdimensional version of program RK2DIM for the purpose of solving such a system. But in a programming language that accommodates vectors, an *n*-dimensional Runge-Kutta program is scarcely more complicated than a one-dimensional program. For instance, the application material for this section lists the *n*-dimensional MATLAB program **rkn** that closely resembles the one-dimensional program **rk** of Fig. 2.6.11.

Example 4

Batted baseball Suppose that a batted ball starts at $x_0 = 0$, $y_0 = 0$ with initial velocity $v_0 = 160$ ft/s and with initial angle of inclination $\theta = 30^\circ$. If air resistance is ignored, we find by the elementary methods of Section 1.2 that the baseball travels a [horizontal] distance of $400\sqrt{3}$ ft (approximately 693 ft) in 5 s before striking the ground. Now suppose that in addition to a downward gravitational acceleration (g = 32 ft/s²), the baseball experiences an acceleration due to air resistance of $(0.0025)v^2$ feet per second per second, directed opposite to its instantaneous direction of motion. Determine how far the baseball will travel horizontally under these conditions.

7.7 NUMERICAL METHODS FOR SYSTEMS459

Solution According to Problem 30 of Section 7.1, the equations of motion of the baseball are

$$\frac{d^2x}{dt^2} = -cv\frac{dx}{dt}, \quad \frac{d^2y}{dt^2} = -cv\frac{dy}{dt} - g$$
(23)

where $v = \sqrt{(x')^2 + (y')^2}$ is the speed of the ball, and where c = 0.0025 and g = 32 in fps units. We convert to a first-order system as in (22) and thereby obtain the system

$$x'_{1} = x_{3},$$

$$x'_{2} = x_{4},$$

$$x'_{3} = -cx_{3}\sqrt{x_{3}^{2} + x_{4}^{2}},$$

$$x'_{4} = -cx_{4}\sqrt{x_{3}^{2} + x_{4}^{2}} - g$$
(24)

of four first-order differential equations with

$$x_1(0) = x_2(0) = 0,$$

$$x_3(0) = 80\sqrt{3}, \quad x_4(0) = 80.$$
(25)

Note that $x_3(t)$ and $x_4(t)$ are simply the x- and y-components of the baseball's velocity vector, so $v = \sqrt{x_3^2 + x_4^2}$. We proceed to apply the Runge–Kutta method to investigate the motion of the batted baseball described by the initial value problem in (24) and (25), first taking c = 0 to ignore air resistance and then using c = 0.0025 to take air resistance into account.

WITHOUT AIR RESISTANCE: Figure 7.7.4 shows the numerical results obtained when a Runge-Kutta program such as **rkn** is applied with step size h = 0.1 and with c = 0 (no air resistance). For convenience in interpreting the results, the printed output at each selected step consists of the horizontal and vertical coordinates x and y of the baseball, its velocity v, and the angle of inclination α of its velocity vector (in degrees measured from the horizontal). These results agree with the exact solution when c = 0. The ball travels a horizontal distance of $400\sqrt{3} \approx 692.82$ ft in exactly 5 s, having reached a maximum height of 100 ft after 2.5 s. Note also that the ball strikes the ground at the same angle and with the same speed as its initial angle and speed.

t	x	У	v	α
0.0	0.00	0.00	160.00	+30
0.5	69.28	36.00	152.63	+25
1.0	138.56	64.00	146.64	+19
1.5	207.85	84.00	142.21	+13
2.0	277.13	96.00	139.48	+7
2.5	346.41	100.00	138.56	+0
3.0	415.69	96.00	139.48	-7
3.5	484.97	84.00	142.21	-13
4.0	554.26	64.00	146.64	-19
4.5	623.54	36.00	152.63	-25
5.0	692.82	0.00	160.00	-30

FIGURE 7.7.4. The batted baseball with no air resistance (c = 0).

t	x	у	v	α
0.0	0.00	0.00	160.00	+30
0.5	63.25	32.74	127.18	+24
1.0	117.11	53.20	104.86	+17
1.5	164.32	63.60	89.72	+8
2.0	206.48	65.30	80.17	-3
2.5	244.61	59.22	75.22	-15
3.0	279.29	46.05	73.99	-27
3.5	310.91	26.41	75.47	-37
4.0	339.67	0.91	78.66	-46

t	x	у	v	α
1.5	164.32	63.60	89.72	+8
1.6	173.11	64.60	87.40	+5
1.7	181.72	65.26	85.29	+3
1.8	190.15	65.60	83.39	+1
1.9	198.40	65.61	81.68	-1 Apex
2.0	206.48	65.30	80.17	-3
÷	÷	•	:	÷
3.8	328.50	11.77	77.24	-42
3.9	334.14	6.45	77.93	-44
4.0	339.67	0.91	78.66	-46 J
4.1	345.10	-4.84	79.43	-47 Impact
4.2	350.41	-10.79	80.22	-49

FIGURE 7.7.5. The batted baseball with air resistance (c = 0.0025).

FIGURE 7.7.6. The batted ball's apex and its impact with the ground.



FIGURE 7.7.7. An "easy out" or a home run?

WITH AIR RESISTANCE: Figure 7.7.5 shows the results obtained with the fairly realistic value of c = 0.0025 for the air resistance for a batted baseball. To within a hundredth of a foot in either direction, the same results are obtained with step sizes h = 0.05 and h = 0.025. We now see that with air resistance the ball travels a distance well under 400 ft in just over 4 s. The more refined data in Fig. 7.7.6 show that the ball travels horizontally only about 340 ft and that its maximum height is only about 66 ft. As illustrated in Fig. 7.7.7, air resistance has converted a massive home run into a routine fly ball (if hit straightaway to center field). Note also that when the ball strikes the ground, it has slightly under *half* its initial speed (only about 79 ft/s) and is falling at a steeper angle (about 46°). Every baseball fan has observed empirically these aspects of the trajectory of a fly ball.

Variable Step Size Methods

The Runge–Kutta method for a large system requires an appreciable amount of computational labor, even when a computer is employed. Therefore, just as the step size h should not be so large that the resulting error in the solution is unacceptable, h ought not to be so small that too many steps are needed, hence requiring an unacceptable amount of computation. Thus the practical numerical solution of differential equations involves a tradeoff between accuracy and efficiency.

To facilitate this tradeoff, modern *variable step size methods* vary the step size h as the solution process proceeds. Large steps are taken in regions where the dependent variables are changing slowly; smaller steps are taken when these variables are changing rapidly, in order to prevent large errors.

An *adaptable* or *variable step size* Runge–Kutta method employs both a preassigned minimum error tolerance *MinTol* and a maximum error tolerance *MaxTol* to attempt to ensure that the error made in the typical step from \mathbf{x}_n to \mathbf{x}_{n+1} is neither too large (and hence inaccurate) nor too small (and hence inefficient). A fairly simple scheme for doing this may be outlined as follows:

• Having reached \mathbf{x}_n with a Runge–Kutta step of length $t_n - t_{n-1} = h$, let $\mathbf{x}^{(1)}$ denote the result of a further Runge–Kutta step of length h and let $\mathbf{x}^{(2)}$ denote the result of *two* successive Runge–Kutta steps each of length h/2.

• On the grounds that $\mathbf{x}^{(2)}$ should be a more accurate approximation to $\mathbf{x}(t_n + h)$ than is $\mathbf{x}^{(1)}$, take

$$Err = |\mathbf{x}^{(1)} - \mathbf{x}^{(2)}|$$

as an estimate of the error in $\mathbf{x}^{(1)}$.

- If $MinTol \leq Err \leq MaxTol$, then let $\mathbf{x}_{n+1} = \mathbf{x}^{(1)}$, $t_{n+1} = t_n + h$, and proceed to the next step.
- If Err < MinTol, then the error is too small! Hence let $\mathbf{x}_{n+1} = \mathbf{x}^{(1)}$, $t_{n+1} = t_n + h$, but *double* the step size to 2h before making the next step.
- If Err > MaxTol, then the error is too large. Hence reject $\mathbf{x}^{(1)}$ and start afresh at \mathbf{x}_n with the *halved* step size h/2.

The detailed implementation of such a scheme can be complicated. For a much more complete but readable discussion of adaptive Runge–Kutta methods, see Section 17.2 of William H. Press et al., *Numerical Recipes: The Art of Scientific Computing* (Cambridge: Cambridge University Press, 2007).

Several widely available scientific computing packages (such as *Maple, Mathematica*, and MATLAB) include sophisticated variable step size programs that will accommodate an essentially arbitrary number of simultaneous differential equations. Such a general-purpose program might be used, for example, to model numerically the major components of the solar system: the sun and the nine (known) major planets. If m_i denotes the mass and $\mathbf{r}_i = (x_i, y_i, z_i)$ denotes the position vector of the *i*th one of these 10 bodies, then—by Newton's laws—the equation of motion of m_i is

$$m_i \mathbf{r}_i'' = \sum_{j \neq i} \frac{Gm_i m_j}{(r_{ij})^3} (\mathbf{r}_j - \mathbf{r}_i),$$
(26)

where $r_{ij} = |\mathbf{r}_j - \mathbf{r}_i|$ denotes the distance between m_i and m_j . For each i = 1, 2, ..., 10, the summation in Eq. (26) is over all values of $j \neq i$ from 1 to 10. The 10 vector equations in (26) constitute a system of 30 second-order scalar equations, and the equivalent first-order system consists of 60 differential equations in the coordinates and velocity components of the 10 major bodies in the solar system. Mathematical models that involve this many (or more) differential equations—and that require so-phisticated software and hardware for their numerical analysis—are quite common in science, engineering, and applied technology.



FIGURE 7.7.8. The Earth–Moon center-of-mass coordinate system.

Earth–Moon Satellite Orbits

For an example of a program whose efficient solution requires adaptive step size methods, we consider an Apollo satellite in orbit about the Earth *E* and Moon *M*. Figure 7.7.8 shows an x_1x_2 -coordinate system whose origin lies at the center of mass of the Earth and the Moon and which rotates at the rate of one revolution per "moon month" of approximately $\tau = 27.32$ days, so the Earth and Moon remain fixed in their positions on the x_1 -axis. If we take as unit distance the distance (about 384,000 kilometers, assumed constant) between the Earth and Moon centers, then their coordinates are $E(-\mu, 0)$ and $M(1 - \mu, 0)$, where $\mu = m_M/(m_E + m_M)$ in terms of the Earth mass m_E and Moon mass m_M . If we take the total mass $m_E + m_M$ as the unit of mass and $\tau/(2\pi) \approx 4.35$ days as the unit of time, then the gravitational constant is G = 1 in Eq. (26), and the equations of motion of the

satellite position $S(x_1, x_2)$ are

$$x_1'' = x_1 + 2x_2' - \frac{(1-\mu)(x_1+\mu)}{(r_E)^3} - \frac{\mu(x_1-1+\mu)}{(r_M)^3},$$

$$x_2'' = x_2 - 2x_1' - \frac{(1-\mu)x_2}{(r_E)^3} - \frac{\mu x_2}{(r_M)^3},$$
(27)

where r_E and r_M denote the satellite's distance to the Earth and Moon (indicated in Fig. 7.7.8). The initial two terms on the right-hand side of each equation result from the rotation of the coordinate system. In the system of units described here, the lunar mass is approximately $m_M = 0.012277471$. The second-order system in (27) can be converted to an equivalent first-order system (of four differential equations) by substituting

$$x'_1 = x_3, \quad x'_2 = x_4, \quad \text{so that} \quad x''_1 = x'_3, \quad x''_2 = x'_4$$

Suppose that the satellite initially is in a clockwise circular orbit of radius about 2400 kilometers about the Moon. At its farthest point from the Earth $(x_1 = 0.994)$ it is "launched" into Earth–Moon orbit with initial velocity v_0 . The corresponding initial conditions are

$$x_1(0) = 0.994, \quad x_2(0) = 0, \quad x_3(0) = 0, \quad x_4(0) = -v_0$$

An adaptive step size method (ode45) in the MATLAB software system was used to solve numerically the system in (27). The orbits in Figs. 7.7.9 and 7.7.10 were obtained with

$$v_0 = 2.031732629557$$
 and $v_0 = 2.001585106379$

respectively. [In the system of units used here, the unit of velocity is approximately 3680 km/h.] In each case a closed but multilooped periodic trajectory about the Earth and the Moon—a so-called bus orbit—is obtained, but a relatively small change in the initial velocity changes the number of loops! For more information, see NASA Contractor Report CR-61139, "Study of the Methods for the Numerical Solution of Ordinary Differential Equations," prepared by O. B. Francis, Jr. et al. for the NASA–George C. Marshall Space Flight Center, June 7, 1966.

So-called Moon–Earth "bus orbits" are periodic—that is, are closed trajectories traversed repeatedly by the satellite—only in a rotating x_1x_2 -coordinate system as discussed above. The satellite of Fig. 7.7.9 traverses its closed orbit and returns to rendezvous with the Moon about 48.4 days after its insertion into orbit. Figures 7.7.11 and 7.7.12 illustrate the motion of the same satellite—but in an ordinary nonrotating x_y -coordinate system centered at the Earth, in which the Moon encircles the Earth counterclockwise in a near-circular orbit, completing one revolution in about 27.3 days. The Moon starts at point *S*, and after 48.4 days it has completed a bit over 1.75 revolutions about the Earth and reaches the point *R* at which its rendezvous with the satellite occurs. Figure 7.7.11 shows the positions of Moon and satellite a day and a half after the satellite's insertion into its orbit, each traveling in a generally counterclockwise direction around the Earth. Figure 7.7.12 shows their positions a day and a half before their rendezvous at point *R*, the satellite meanwhile having encircled the Earth about 2.5 times in an orbit that (in the indicated xy-coordinate system) appears to resemble a slowly varying ellipse.



FIGURE 7.7.9. Apollo Moon–Earth bus orbit with insertion velocity $v_0 = 7476 \text{ km/h.}$



FIGURE 7.7.10. Apollo Moon–Earth bus orbit with insertion velocity $v_0 = 7365$ km/h.



FIGURE 7.7.11. The moon and satellite in a nonrotating coordinate system, 1.5days after orbital insertion of the satellite at starting point *S*.



FIGURE 7.7.12. The moon and satellite in a nonrotating coordinate system, 1.5 days before their rendezvous at point *R*.

7.7 Problems

A hand-held calculator will suffice for Problems 1 through 8. In each problem an initial value problem and its exact solution are given. Approximate the values of x(0.2) and y(0.2) in three ways: (a) by the Euler method with two steps of size h = 0.1; (b) by the improved Euler method with a single step of size h = 0.2; and (c) by the Runge–Kutta method with a single step of size h = 0.2. Compare the approximate values with the actual values x(0.2) and y(0.2).

1. x' = x + 2y, x(0) = 0, y' = 2x + y, y(0) = 2; $x(t) = e^{3t} - e^{-t}, y(t) = e^{3t} + e^{-t}$ **2.** x' = 2x + 3y, x(0) = 1, y' = 2x + y, y(0) = -1; $x(t) = e^{-t}, y(t) = -e^{-t}$ 3. x' = 3x + 4y, x(0) = 1, y' = 3x + 2y, y(0) = 1; $x(t) = \frac{1}{7}(8e^{6t} - e^{-t}), y(t) = \frac{1}{7}(6e^{6t} + e^{-t})$ 4. x' = 9x + 5y, x(0) = 1, y' = -6x - 2y, y(0) = 0; $x(t) = -5e^{3t} + 6e^{4t}, y(t) = 6e^{3t} - 6e^{4t}$ 5. x' = 2x - 5y, x(0) = 2. y' = 4x - 2y, y(0) = 3; $x(t) = 2\cos 4t - \frac{11}{4}\sin 4t, \ y(t) = 3\cos 4t + \frac{1}{2}\sin 4t$ 6. x' = x - 2y, x(0) = 0, y' = 2x + y, y(0) = 4; $x(t) = -4e^t \sin 2t, \ y(t) = 4e^t \cos 2t$ 7. x' = 3x - y, x(0) = 2, v' = x + y, y(0) = 1; $x(t) = (t+2)e^{2t}, y(t) = (t+1)e^{2t}$ 8. x' = 5x - 9y, x(0) = 0, y' = 2x - y, y(0) = -1; $x(t) = 3e^{2t} \sin 3t, \ y(t) = e^{2t} (\sin 3t - \cos 3t)$

A computer will be required for the remaining problems in this section. In Problems 9 through 12, an initial value problem and its exact solution are given. In each of these four problems, use the Runge–Kutta method with step sizes h = 0.1 and

h = 0.05 to approximate to five decimal places the values x(1) and y(1). Compare the approximations with the actual values.

- 9. x' = 2x y, x(0) = 1, y' = x + 2y, y(0) = 0; $x(t) = e^{2t} \cos t$, $y(t) = e^{2t} \sin t$
- **10.** x' = x + 2y, x(0) = 0, $y' = x + e^{-t}$, y(0) = 0; $x(t) = \frac{1}{9}(2e^{2t} - 2e^{-t} + 6te^{-t})$, $y(t) = \frac{1}{9}(e^{2t} - e^{-t} + 6te^{-t})$
- **11.** $x' = -x + y (1 + t^3)e^{-t}$, x(0) = 0, $y' = -x - y - (t - 3t^2)e^{-t}$, y(0) = 1; $x(t) = e^{-t}(\sin t - t)$, $y(t) = e^{-t}(\cos t + t^3)$
- **12.** $x'' + x = \sin t$, x(0) = 0; $x(t) = \frac{1}{2}(\sin t - t \cos t)$
- 13. Crossbow bolt Suppose that a crossbow bolt is shot straight upward with initial velocity 288 ft/s. If its deceleration due to air resistance is (0.04)v, then its height x(t) satisfies the initial value problem

x'' = -32 - (0.04)x'; x(0) = 0, x'(0) = 288.

Find the maximum height that the bolt attains and the time required for it to reach this height.

- 14. Repeat Problem 13, but assume instead that the deceleration of the bolt due to air resistance is $(0.0002)v^2$.
- **15.** Height of projectile Suppose that a projectile is fired straight upward with initial velocity v_0 from the surface of the earth. If air resistance is not a factor, then its height x(t) at time t satisfies the initial value problem

$$\frac{d^2x}{dt^2} = -\frac{gR^2}{(x+R)^2}; \quad x(0) = 0, \quad x'(0) = v_0$$

Use the values $g = 32.15 \text{ ft/s}^2 \approx 0.006089 \text{ mi/s}^2$ for the gravitational acceleration of the earth at its surface and R = 3960 mi as the radius of the earth. If $v_0 = 1 \text{ mi/s}$, find the maximum height attained by the projectile and its time of ascent to this height.

Batted Baseball

Problems 16 through 18 deal with the batted baseball of Example 4, having initial velocity 160 ft/s and air resistance coefficient c = 0.0025.

- 16. Find the *range*—the horizontal distance the ball travels before it hits the ground—and its total time of flight with initial inclination angles 40° , 45° , and 50° .
- 17. Find (to the nearest degree) the initial inclination that maximizes the range. If there were no air resistance it would be exactly 45° , but your answer should be less than 45° .
- **18.** Find (to the nearest half degree) the initial inclination angle greater than 45° for which the range is 300 ft.
- **19.** Home run Find the initial velocity of a baseball hit by Babe Ruth (with c = 0.0025 and initial inclination 40°) if it hit the bleachers at a point 50 ft high and 500 horizontal feet from home plate.

- **20.** Crossbow bolt Consider the crossbow bolt of Problem 14, fired with the same initial velocity of 288 ft/s and with the air resistance deceleration $(0.0002)v^2$ directed opposite its direction of motion. Suppose that this bolt is fired from ground level at an initial angle of 45°. Find how high vertically and how far horizontally it goes, and how long it remains in the air.
- **21.** Artillery projectile Suppose that an artillery projectile is fired from ground level with initial velocity 3000 ft/s and initial inclination angle 40° . Assume that its air resistance deceleration is $(0.0001)v^2$. (a) What is the range of the projectile and what is its total time of flight? What is its speed at impact with the ground? (b) What is the maximum altitude of the projectile, and when is that altitude attained? (c) You will find that the projectile is still losing speed at the apex of its trajectory. What is the *minimum* speed that it attains during its descent?

7.7 Application Comets and Spacecraft

Go to goo.gl/1g94fA to download this application's computing resources including *Maple/Mathematica/*MATLAB. Figure 7.7.13 lists TI-Nspire CX CAS and Python versions of the two-dimensional Runge–Kutta program RK2DIM. You should note that it closely parallels the onedimensional Runge–Kutta program listed in Fig. 2.6.11, with a single line there replaced (where appropriate) with two lines here to calculate a *pair* of *x*- and *y*-values or slopes. Note also that the notation used is essentially that of Eqs. (13) and (14) in this section. The first several lines define the functions and initial data needed for Example 1.

Figure 7.7.14 exhibits an *n*-dimensional MATLAB implementation of the Runge-Kutta method. The MATLAB function \mathbf{f} defines the vector of right-hand sides of the differential equations in the system $\mathbf{x}' = \mathbf{f}(t, \mathbf{x})$ to be solved. The **rkn** function then takes as input the initial *t*-value \mathbf{t} , the column vector \mathbf{x} of initial *x*-values, the final *t*-value $\mathbf{t}1$, and the desired number \mathbf{n} of subintervals. As output it produces the resulting column vector \mathbf{T} of *t*-values and the matrix \mathbf{x} whose rows give the corresponding *x*-values. For instance, with \mathbf{f} as indicated in the figure, the MATLAB command

[T,X] = rkn(0, [0;1], 5, 50)

then generates the data shown in the table of Fig. 7.7.1 (which lists only every fifth value of each variable).

You can use Examples 1 through 3 in this section to test your own implementation of the Runge–Kutta method. Then investigate the comet and spacecraft problems described next. Additional application material at the Expanded Applications site indicated in the margin describes additional numerical ODE investigations ranging from batted baseballs to the Apollo orbits shown in Figs. 7.7.9 and 7.7.10.

Your Spacecraft Landing

Your spacecraft is traveling at constant velocity V, approaching a distant earthlike planet with mass M and radius R. When activated, your deceleration system provides a constant thrust T until impact with the surface of the planet. During the period of deceleration, your distance x(t) from the center of the planet satisfies the

7.7 NUMERICAL METHODS FOR SYSTEMS465

TI-Nspire CX CAS	Python	Comment
Define rk2dim()=Prgm	# Program RK2DIM	Program title
f(t,x,y) := y	def F(T,X,Y): return Y	Define function f
g(t,x,y) := -x	def G(T,X,Y): return -X	Define function g
n:=50	N = 50	No. of steps
t:=0.0	T = 0.0	Initial t
x:=0.0	X = 0.0	Initial x
y:=1.0	Y = 1.0	Initial y
t1:=5.0	T1 = 5	Final t
h:=(t1-t)/n	H = (T1-T) / N	Step size
For i,1,n	for I in range(N):	Begin loop
t0:=t	то = т	Save previous t
x0:=x	$x_0 = x$	Save previous x
y0:=y	Y0 = Y	Save previous y
f1:=f(t,x,y)	F1 = F(T, X, Y)	First f-slope
g1:=g(t,x,y)	G1 = G(T, X, Y)	First g-slope
t:=t0+h/2	T = T0 + H/2	Midpoint t
x:=x0+(h*f1)/2	X = X0 + H*F1/2	Midpt x-predictor
y:=y0+(h*g1)/2	Y = Y0 + H*G1/2	Midpt y-predictor
f2:=f(t,x,y)	F2 = F(T, X, Y)	Second f-slope
g2:=g(t,x,y)	G2 = G(T, X, Y)	Second g-slope
x:=x0+(h*f2)/2	X = X0 + H*F2/2	Midpt x-predictor
y:=y0+(h*g2)/2	Y = Y0 + H*G2/2	Midpt y-predictor
f3:=f(t,x,y)	F3 = F(T,X,Y)	Third f-slope
g3:=g(t,x,y)	G3 = G(T, X, Y)	Third g-slope
t:=t0+h	T = T0 + H	New t
x:=x0+h*f3	X = X0 + H*F3	Endpt x-predictor
y:=y0+h*g3	Y = Y0 + H*G3	Endpt y-predictor
f4:=f(t,x,y)	F4 = F(T, X, Y)	Fourth f-slope
g4:=g(t,x,y)	G4 = G(T, X, Y)	Fourth g-slope
fa:=(f1+2*f2+2*f3+f4)/6	FA = (F1+2*F2+2*F3+F4)/6	Average f-slope
ga:=(g1+2*g2+2*g3+g4)/6	GA = (G1+2*G2+2*G3+G4)/6	Average g-slope
x:=x0+h*fa	X = Y0 + H*FA	x-corrector
y:=y0+h*ga	Y = Y0 + H*GA	y-corrector
Disp t,x,y	print (T,X,Y)	Display results
EndFor	# END	End loop
EndPrgm		

FIGURE 7.7.13. TI-Nspire CX CAS and Python two-dimensional Runge–Kutta programs.

```
function xp = f(t,x)
xp = x;
xp(1) = x(2);
xp(2) = -x(1);
function [T,Y] = rkn(t,x,t1,n)
h = (t1 - t)/n;
                                   % step size
T = t;
                                   % initial t
X = x';
                                   % initial x-vector
for i = 1:n
                                   % begin loop
    k1 = f(t,x);
                                   % first k-vector
    k2 = f(t+h/2,x+h*k1/2);
                                   % second k-vector
    k3 = f(t+h/2,x+h*k2/2);
                                   % third k-vector
    k4 = f(t+h, x+h*k3);
                                   % fourth k-vector
    k = (k1+2*k2+2*k3+k4)/6;
                                   % average k-vector
    t = t + h;
                                   % new t
    x = x + h*k;
                                   % new x
                                   % update t-column
    T = [T;t];
    X = [X; x'];
                                   % update x-matrix
                                   % end loop
    end
```

FIGURE 7.7.14. MATLAB implementation of the Runge–Kutta method.

differential equation

$$\frac{d^2x}{dt^2} = T - \frac{GM}{x^2},\tag{1}$$

where $G \approx 6.6726 \times 10^{-11} \text{ N} \cdot (\text{m/kg})^2$ as in Example 3. Your question is this: At what altitude above the surface should your deceleration system be activated in order to achieve a soft touchdown? For a reasonable problem, you can take

$$M = 5.97 \times 10^{24} \text{ (kg)},$$

$$R = 6.38 \times 10^{6} \text{ (m)},$$

$$V = p \times 10^{4} \text{ (km/h)},$$

$$T = g + q \text{ (m/s2)}$$

where $g = GM/R^2$ is the surface gravitational acceleration of the planet. Choose *p* to be the smallest nonzero digit and *q* the next-to-smallest nonzero digit in your ID number. Find the "ignition altitude" accurate to the nearest meter and the resulting "descent time" accurate to the nearest tenth of a second.

Kepler's Law of Planetary (or Satellite) Motion

Consider a satellite in elliptical orbit around a planet of mass M, and suppose that physical units are so chosen that GM = 1 (where G is the gravitational constant). If the planet is located at the origin in the xy-plane, then the equations of motion of the satellite are

$$\frac{d^2x}{dt^2} = -\frac{x}{(x^2 + y^2)^{3/2}}, \quad \frac{d^2y}{dt^2} = -\frac{y}{(x^2 + y^2)^{3/2}}.$$
 (2)

7.7 NUMERICAL METHODS FOR SYSTEMS467

Let *T* denote the period of revolution of the satellite. Kepler's third law says that the *square* of *T* is proportional to the *cube* of the major semiaxis *a* of its elliptical orbit. In particular, if GM = 1, then

$$T^2 = 4\pi^2 a^3. (3)$$

(For details, see Section 11.6 of Edwards and Penney, *Calculus: Early Transcenden*tals, 7th ed., Hoboken, NJ: Pearson, 2008).) If the satellite's x- and y-components of velocity, $x_3 = x' = x'_1$ and $x_4 = y' = x'_2$, are introduced, then the system in (2) translates into a system of four first-order differential equations having the form of those in Eq. (22) of this section.

(a) Solve this 4×4 system numerically with the initial conditions

$$x(0) = 1$$
, $y(0) = 0$, $x'(0) = 0$, $y'(0) = 1$

that correspond theoretically to a circular orbit of radius a = 1, so Eq. (3) gives $T = 2\pi$. Is this what you get?

(b) Now solve the system numerically with the initial conditions

$$x(0) = 1$$
, $y(0) = 0$, $x'(0) = 0$, $y'(0) = \frac{1}{2}\sqrt{6}$

that correspond theoretically to an elliptical orbit with major semiaxis a = 2, so Eq. (3) gives $T = 4\pi\sqrt{2}$. Is this what you get?

Halley's Comet

Halley's comet last reached perihelion (its point of closest approach to the sun at the origin) on February 9, 1986. Its position and velocity components at that time were

$$\mathbf{p}_0 = (0.325514, -0.459460, 0.166229)$$
 and
 $\mathbf{v}_0 = (-9.096111, -6.916686, -1.305721)$

(respectively), with position in AU (astronomical units, in which the unit of distance is the major semiaxis of the earth's orbit) and time in years. In this system, the threedimensional equations of motion of the comet are

$$\frac{d^2x}{dt^2} = -\frac{\mu x}{r^3}, \quad \frac{d^2y}{dt^2} = -\frac{\mu y}{r^3}, \quad \frac{d^2z}{dt^2} = -\frac{\mu z}{r^3}$$
(4)

where

$$\mu = 4\pi^2$$
 and $r = \sqrt{x^2 + y^2 + z^2}$.

Solve the equations in (4) numerically to verify the appearance of the yz-projection of the orbit of Halley's comet shown in Fig. 7.7.15. Plot the xy- and xz-projections as well.

Figure 7.7.16 shows the graph of the distance r(t) of Halley's comet from the sun. Inspection of this graph indicates that Halley's comet reaches a maximum distance (at aphelion) of about 35 AU in a bit less than 40 years and returns to perihelion after about three-quarters of a century. The closer look in Fig. 7.7.17 indicates that the period of revolution of Halley's comet is about 76 years. Use your numerical solution to refine these observations. What is your best estimate of the calendar date of the comet's next perihelion passage?



FIGURE 7.7.15. *yz*-projection of the orbit of Halley's comet.



FIGURE 7.7.16. 200-year plot of the distance r(t) of Halley's comet from the sun. Is there a cusp near t = 75?



FIGURE 7.7.17. A closer look at Halley's perihelion passage after about 76 years.

Your Own Comet

The night before your birthday in 2007 you set up your telescope on a nearby mountaintop. It was a clear night, and you had a stroke of luck: At 12:30 A.M. you spotted a new comet. After repeating the observation on successive nights, you were able to calculate its solar system coordinates $\mathbf{p}_0 = (x_0, y_0, z_0)$ and its velocity vector $\mathbf{v}_0 = (v_{x0}, v_{y0}, v_{z0})$ on that first night. Using this information, determine the following:

- the comet's perihelion (point nearest the sun) and aphelion (point farthest from the sun),
- the comet's velocity at perihelion and at aphelion,
- the comet's period of revolution around the sun, and
- the comet's next two dates of perihelion passage.

Using units of length in AU and time in earth years, the equations of motion of your comet are given in (4). For your personal comet, begin with random initial position and velocity vectors with the same order of magnitude as those of Halley's comet. Repeat the random selection of initial position and velocity vectors, if necessary, until you get a plausible eccentric orbit that ranges well outside the earth's orbit (as most real comets do).



8.1 Matrix Exponentials and Linear Systems

The solution vectors of an $n \times n$ homogeneous linear system

$$\mathbf{x}' = \mathbf{A}\mathbf{x} \tag{1}$$

can be used to construct a square matrix $\mathbf{X} = \mathbf{\Phi}(t)$ that satisfies the *matrix differential equation*

$$\mathbf{X}' = \mathbf{A}\mathbf{X} \tag{1'}$$

associated with Eq. (1). Suppose that $\mathbf{x}_1(t), \mathbf{x}_2(t), \dots, \mathbf{x}_n(t)$ are *n* linearly independent solutions of Eq. (1). Then the $n \times n$ matrix



$$\mathbf{\Phi}(t) = \begin{bmatrix} | & | & | \\ \mathbf{x}_1(t) & \mathbf{x}_2(t) & \cdots & \mathbf{x}_n(t) \\ | & | & | \end{bmatrix}$$
(2)

having these solution vectors as its column vectors, is called a **fundamental matrix** for the system in (1).

Fundamental Matrix Solutions

Because the column vector $\mathbf{x} = \mathbf{x}_j(t)$ of the fundamental matrix $\Phi(t)$ in (2) satisfies the differential equation $\mathbf{x}' = \mathbf{A}\mathbf{x}$, it follows (from the definition of matrix multiplication) that the matrix $\mathbf{X} = \Phi(t)$ itself satisfies the matrix differential equation $\mathbf{X}' = \mathbf{A}\mathbf{X}$. Because its column vectors are linearly independent, it also follows that the fundamental matrix $\Phi(t)$ is nonsingular, and therefore has an inverse matrix $\Phi(t)^{-1}$. Conversely, any nonsingular matrix solution $\Psi(t)$ of Eq. (1') has linearly independent column vectors that satisfy Eq. (1), so $\Psi(t)$ is a fundamental matrix for the system in (1). In terms of the fundamental matrix $\Phi(t)$ in (2), the general solution

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$$
(3)

of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ can be written in the form

$$\mathbf{x}(t) = \mathbf{\Phi}(t)\mathbf{c} \tag{4}$$

where $\mathbf{c} = [c_1 \ c_2 \ \dots \ c_n]^T$ is an arbitrary *constant* vector. If $\Psi(t)$ is any other fundamental matrix for (1), then each column vector of $\Psi(t)$ is a linear combination of the column vectors of $\Phi(t)$, so it follows from Eq. (4) that

$$\Psi(t) = \Phi(t)\mathbf{C} \tag{4'}$$

for some $n \times n$ matrix **C** of constants.

In order that the solution $\mathbf{x}(t)$ in (3) satisfy a given initial condition

$$\mathbf{x}(0) = \mathbf{x}_0,\tag{5}$$

it suffices that the coefficient vector **c** in (4) be such that $\Phi(0)\mathbf{c} = \mathbf{x}_0$; that is, that

$$\mathbf{c} = \mathbf{\Phi}(0)^{-1} \mathbf{x}_0. \tag{6}$$

When we substitute (6) in Eq. (4), we get the conclusion of the following theorem.

THEOREM 1 Fundamental Matrix Solutions

Let $\Phi(t)$ be a fundamental matrix for the homogeneous linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Then the [unique] solution of the initial value problem

$$\mathbf{x}' = \mathbf{A}\mathbf{x}, \quad \mathbf{x}(0) = \mathbf{x}_0 \tag{7}$$

is given by

>

$$\mathbf{x}(t) = \mathbf{\Phi}(t)\mathbf{\Phi}(0)^{-1}\mathbf{x}_0.$$
 (8)

Section 7.3 tells us how to find a fundamental matrix for the system

$$\mathbf{x}' = \mathbf{A}\mathbf{x} \tag{9}$$

with constant $n \times n$ coefficient matrix **A**, at least in the case where **A** has a complete set of *n* linearly independent eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ associated with the (not necessarily distinct) eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$, respectively. In this event the corresponding solution vectors of Eq. (9) are given by

$$\mathbf{x}_i(t) = \mathbf{v}_i e^{\lambda_i t}$$

for i = 1, 2, ..., n. Therefore, the $n \times n$ matrix

$$\mathbf{\Phi}(t) = \begin{bmatrix} | & | & | \\ \mathbf{v}_1 e^{\lambda_1 t} & \mathbf{v}_2 e^{\lambda_2 t} & \cdots & \mathbf{v}_n e^{\lambda_n t} \\ | & | & | \end{bmatrix}$$
(10)

having the solutions $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ as column vectors is a fundamental matrix for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$.

In order to apply Eq. (8), we must be able to compute the inverse matrix $\Phi(0)^{-1}$. The inverse of the nonsingular 2 × 2 matrix

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

is

$$\mathbf{A}^{-1} = \frac{1}{\Delta} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix},\tag{11}$$

where $\Delta = \det(\mathbf{A}) = ad - bc \neq 0$. The inverse of the nonsingular 3×3 matrix $\mathbf{A} = [a_{ij}]$ is given by

$$\mathbf{A}^{-1} = \frac{1}{\Delta} \begin{bmatrix} +A_{11} & -A_{12} & +A_{13} \\ -A_{21} & +A_{22} & -A_{23} \\ +A_{31} & -A_{32} & +A_{33} \end{bmatrix}^{T},$$
 (12)

where $\Delta = \det(\mathbf{A}) \neq 0$ and A_{ij} denotes the determinant of the 2 × 2 submatrix of \mathbf{A} obtained by deleting the *i*th row and *j*th column of \mathbf{A} . (Do not overlook the symbol *T* for *transpose* in Eq. (12).) The formula in (12) is also valid upon generalization to $n \times n$ matrices, but in practice inverses of larger matrices are usually computed instead by row reduction methods (see any linear algebra text) or by using a calculator or computer algebra system.

Example 1 Find a fundamental matrix for the system $\begin{array}{l}
x' = 4x + 2y, \\
y' = 3x - y,
\end{array}$ (13)

and then use it to find the solution of (13) that satisfies the initial conditions x(0) = 1, y(0) = -1.

Solution The linearly independent solutions

$$\mathbf{x}_1(t) = \begin{bmatrix} e^{-2t} \\ -3e^{-2t} \end{bmatrix}$$
 and $\mathbf{x}_2(t) = \begin{bmatrix} 2e^{5t} \\ e^{5t} \end{bmatrix}$

found in Example 1 of Section 7.3 yield the fundamental matrix

$$\mathbf{\Phi}(t) = \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix}.$$
 (14)

Then

$$\mathbf{\Phi}(0) = \begin{bmatrix} 1 & 2\\ -3 & 1 \end{bmatrix},$$

and the formula in (11) gives the inverse matrix

$$\Phi(0)^{-1} = \frac{1}{7} \begin{bmatrix} 1 & -2\\ 3 & 1 \end{bmatrix}.$$
 (15)

Hence the formula in (8) gives the solution

$$\mathbf{x}(t) = \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \begin{pmatrix} \frac{1}{7} \end{pmatrix} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{pmatrix} \frac{1}{7} \end{pmatrix} \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \begin{bmatrix} 3 \\ 2 \end{bmatrix},$$

and so

$$\mathbf{x}(t) = \frac{1}{7} \begin{bmatrix} 3e^{-2t} + 4e^{5t} \\ -9e^{-2t} + 2e^{5t} \end{bmatrix}.$$

Thus the solution of the original initial value problem is given by

,

$$x(t) = \frac{3}{7}e^{-2t} + \frac{4}{7}e^{5t}, \quad y(t) = -\frac{9}{7}e^{-2t} + \frac{2}{7}e^{5t}.$$

Remark An advantage of the fundamental matrix approach is this: Once we know the fundamental matrix $\Phi(t)$ and the inverse matrix $\Phi(0)^{-1}$, we can calculate rapidly by matrix multiplication the solutions corresponding to different initial conditions. For example, suppose that we seek the solution of the system in (13) satisfying the new initial conditions x(0) = 77, y(0) = 49. Then substitution of (14) and (15) in (8) gives the new particular solution

$$\mathbf{x}(t) = \frac{1}{7} \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 77 \\ 49 \end{bmatrix}$$
$$= \frac{1}{7} \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \begin{bmatrix} -21 \\ 280 \end{bmatrix} = \begin{bmatrix} -3e^{-2t} + 80e^{5t} \\ 9e^{-2t} + 40e^{5t} \end{bmatrix}.$$

Exponential Matrices

We now discuss the possibility of constructing a fundamental matrix for the constantcoefficient linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ directly from the coefficient matrix \mathbf{A} —that is, without first applying the methods of earlier sections to find a linearly independent set of solution vectors.

We have seen that exponential functions play a central role in the solution of linear differential equations and systems, ranging from the scalar equation x' = kx with solution $x(t) = x_0 e^{kt}$ to the vector solution $\mathbf{x}(t) = \mathbf{v}e^{\lambda t}$ of the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ whose coefficient matrix \mathbf{A} has eigenvalue λ with associated eigenvector \mathbf{v} . We now define exponentials of matrices in such a way that

$$\mathbf{X}(t) = e^{\mathbf{A}t}$$

is a matrix solution of the matrix differential equation

$$\mathbf{X}' = \mathbf{A}\mathbf{X}$$

with $n \times n$ coefficient matrix **A**—in analogy with the fact that the ordinary exponential function $x(t) = e^{at}$ is a scalar solution of the first-order differential equation x' = ax.

The exponential e^z of the complex number z may be defined (as in Section 5.3) by means of the exponential series

$$e^{z} = 1 + z + \frac{z^{2}}{2!} + \frac{z^{3}}{3!} + \dots + \frac{z^{n}}{n!} + \dots$$
 (16)

Similarly, if **A** is an $n \times n$ matrix, then the **exponential matrix** $e^{\mathbf{A}}$ is the $n \times n$ matrix defined by the series

$$e^{\mathbf{A}} = \mathbf{I} + \mathbf{A} + \frac{\mathbf{A}^2}{2!} + \dots + \frac{\mathbf{A}^n}{n!} + \dots, \qquad (17)$$

where \mathbf{I} is the identity matrix. The meaning of the infinite series on the right in (17) is given by

$$\sum_{n=0}^{\infty} \frac{\mathbf{A}^n}{n!} = \lim_{k \to \infty} \left(\sum_{n=0}^k \frac{\mathbf{A}^n}{n!} \right),\tag{18}$$

where $\mathbf{A}^0 = \mathbf{I}$, $\mathbf{A}^2 = \mathbf{A}\mathbf{A}$, $\mathbf{A}^3 = \mathbf{A}\mathbf{A}^2$, and so on; inductively, $\mathbf{A}^{n+1} = \mathbf{A}\mathbf{A}^n$ if $n \ge 0$. It can be shown that the limit in (18) exists for every $n \times n$ square matrix \mathbf{A} . That is, the exponential matrix $e^{\mathbf{A}}$ is defined (by Eq. (17)) for every square matrix \mathbf{A} . Example 2 Consider the 2×2 diagonal matrix

$$\mathbf{A} = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}.$$

Then it is apparent that

$$\mathbf{A}^n = \begin{bmatrix} a^n & 0\\ 0 & b^n \end{bmatrix}$$

for each integer $n \ge 1$. It therefore follows that

$$e^{\mathbf{A}} = \mathbf{I} + \mathbf{A} + \frac{\mathbf{A}^{2}}{2!} + \cdots$$
$$= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} + \begin{bmatrix} a^{2}/2! & 0 \\ 0 & b^{2}/2! \end{bmatrix} + \cdots$$
$$= \begin{bmatrix} 1 + a + a^{2}/2! + \cdots & 0 \\ 0 & 1 + b + b^{2}/2! + \cdots \end{bmatrix}.$$

Thus

so the exponential of the *diagonal* 2×2 matrix **A** is obtained simply by exponentiating each diagonal element of **A**.

 $e^{\mathbf{A}} = \begin{bmatrix} e^{a} & 0\\ 0 & e^{b} \end{bmatrix},$

The $n \times n$ analog of the 2×2 result in Example 2 is established in the same way. The exponential of the $n \times n$ diagonal matrix

$$\mathbf{D} = \begin{bmatrix} a_1 & 0 & \cdots & 0 \\ 0 & a_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & a_n \end{bmatrix}$$
(19)

is the $n \times n$ diagonal matrix

$$e^{\mathbf{D}} = \begin{bmatrix} e^{a_1} & 0 & \cdots & 0 \\ 0 & e^{a_2} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & e^{a_n} \end{bmatrix},$$
(20)

obtained by exponentiating each diagonal element of **D**.

The exponential matrix $e^{\mathbf{A}}$ satisfies most of the exponential relations that are familiar in the case of scalar exponents. For instance, if **0** is the $n \times n$ zero matrix, then Eq. (17) yields

$$e^{\mathbf{0}} = \mathbf{I},\tag{21}$$

the $n \times n$ identity matrix. In Problem 31 we ask you to show that a useful law of exponents holds for $n \times n$ matrices that commute:

If
$$\mathbf{AB} = \mathbf{BA}$$
, then $e^{\mathbf{A} + \mathbf{B}} = e^{\mathbf{A}}e^{\mathbf{B}}$. (22)

In Problem 32 we ask you to conclude that

$$\left(e^{\mathbf{A}}\right)^{-1} = e^{-\mathbf{A}}.$$
(23)

In particular, the matrix $e^{\mathbf{A}}$ is nonsingular for every $n \times n$ matrix \mathbf{A} (reminiscent of the fact that $e^z \neq 0$ for all z). It follows from elementary linear algebra that the column vectors of $e^{\mathbf{A}}$ are always linearly independent.

If t is a scalar variable, then substitution of At for A in Eq. (17) gives

$$e^{\mathbf{A}t} = \mathbf{I} + \mathbf{A}t + \mathbf{A}^2 \frac{t^2}{2!} + \dots + \mathbf{A}^n \frac{t^n}{n!} + \dots$$
 (24)

(Of course, At is obtained simply by multiplying each element of A by t.)

Example 3

 $\mathbf{A} = \begin{bmatrix} 0 & 3 & 4 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix},$

$$\mathbf{A}^2 = \begin{bmatrix} 0 & 0 & 18 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{A}^3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

so $\mathbf{A}^n = \mathbf{0}$ for $n \ge 3$. It therefore follows from Eq. (24) that

e

$$e^{\mathbf{A}t} = \mathbf{I} + \mathbf{A}t + \frac{1}{2}\mathbf{A}^{2}t^{2}$$
$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 3 & 4 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix} t + \frac{1}{2} \begin{bmatrix} 0 & 0 & 18 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} t^{2};$$

that is,

$$\mathbf{A}t = \begin{bmatrix} 1 & 3t & 4t + 9t^2 \\ 0 & 1 & 6t \\ 0 & 0 & 1 \end{bmatrix}.$$

Remark If $\mathbf{A}^n = \mathbf{0}$ for some positive integer *n*, then the exponential series in (24) terminates after a finite number of terms, so the exponential matrix $e^{\mathbf{A}}$ (or $e^{\mathbf{A}t}$) is readily calculated as in Example 3. Such a matrix—with a vanishing power—is said to be **nilpotent**.

Example 4

then

If

$$\mathbf{A} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} + \begin{bmatrix} 0 & 3 & 4 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix} = \mathbf{D} + \mathbf{B},$$

 $\mathbf{A} = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 2 & 6 \\ 0 & 0 & 2 \end{bmatrix},$

where $\mathbf{D} = 2\mathbf{I}$ is a diagonal matrix and \mathbf{B} is the nilpotent matrix of Example 3. Therefore, (20) and (22) give

$$e^{\mathbf{A}t} = e^{(\mathbf{D}+\mathbf{B})t} = e^{\mathbf{D}t}e^{\mathbf{B}t} = \begin{bmatrix} e^{2t} & 0 & 0\\ 0 & e^{2t} & 0\\ 0 & 0 & e^{2t} \end{bmatrix} \begin{bmatrix} 1 & 3t & 4t + 9t^2\\ 0 & 1 & 6t\\ 0 & 0 & 1 \end{bmatrix};$$

thus

$$e^{\mathbf{A}t} = \begin{bmatrix} e^{2t} & 3te^{2t} & (4t+9t^2)e^{2t} \\ 0 & e^{2t} & 6te^{2t} \\ 0 & 0 & e^{2t} \end{bmatrix}.$$

then

If

Matrix Exponential Solutions

It happens that term-by-term differentiation of the series in (24) is valid, with the result

$$\frac{d}{dt}\left(e^{\mathbf{A}t}\right) = \mathbf{A} + \mathbf{A}^{2}t + \mathbf{A}^{3}\frac{t^{2}}{2!} + \dots = \mathbf{A}\left(\mathbf{I} + \mathbf{A}t + \mathbf{A}^{2}\frac{t^{2}}{2!} + \dots\right);$$

that is,

>

>

$$\frac{d}{dt}\left(e^{\mathbf{A}t}\right) = \mathbf{A}e^{\mathbf{A}t},\tag{25}$$

in analogy to the formula $D_t(e^{kt}) = ke^{kt}$ from elementary calculus. Thus the matrix-valued function

$$\mathbf{X}(t) = e^{\mathbf{A}t}$$

satisfies the matrix differential equation

$$\mathbf{X}' = \mathbf{A}\mathbf{X}.$$

Because the matrix $e^{\mathbf{A}t}$ is nonsingular, it follows that *the matrix exponential* $e^{\mathbf{A}t}$ *is a fundamental matrix for the linear system* $\mathbf{x}' = \mathbf{A}\mathbf{x}$. In particular, it is *the* fundamental matrix $\mathbf{X}(t)$ such that $\mathbf{X}(0) = \mathbf{I}$. Therefore, Theorem 1 implies the following result.

THEOREM 2 Matrix Exponential Solutions

If **A** is an $n \times n$ matrix, then the solution of the initial value problem

$$\mathbf{x}' = \mathbf{A}\mathbf{x}, \qquad \mathbf{x}(0) = \mathbf{x}_0 \tag{26}$$

is given by

$$\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{x}_0,\tag{27}$$

and this solution is unique.

Thus the solution of homogeneous linear systems reduces to the task of computing exponential matrices. Conversely, if we already know a fundamental matrix $\mathbf{\Phi}(t)$ for the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$, then the facts that $e^{\mathbf{A}t} = \mathbf{\Phi}(t)\mathbf{C}$ (by Eq. (4')) and $e^{\mathbf{A}\cdot\mathbf{0}} = e^{\mathbf{0}} = \mathbf{I}$ (the identity matrix) yield

$$e^{\mathbf{A}t} = \mathbf{\Phi}(t)\mathbf{\Phi}(0)^{-1}.$$
(28)

So we can find the matrix exponential e^{At} by solving the linear system $\mathbf{x}' = A\mathbf{x}$.

Example 5

In Example 1 we found that the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with

$$\mathbf{A} = \begin{bmatrix} 4 & 2\\ 3 & -1 \end{bmatrix}$$

has fundamental matrix

$$\Phi(t) = \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \text{ with } \Phi(0)^{-1} = \frac{1}{7} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix}.$$

Hence Eq. (28) gives

$$e^{\mathbf{A}t} = \frac{1}{7} \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix}$$
$$= \frac{1}{7} \begin{bmatrix} e^{-2t} + 6e^{5t} & -2e^{-2t} + 2e^{5t} \\ -3e^{-2t} + 3e^{5t} & 6e^{-2t} + e^{5t} \end{bmatrix}.$$

Example 6 Use an exponential matrix to solve the initial value problem

$$\mathbf{x}' = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 2 & 6 \\ 0 & 0 & 2 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 19 \\ 29 \\ 39 \end{bmatrix}.$$
 (29)

Solution The coefficient matrix A in (29) evidently has characteristic equation $(2 - \lambda)^3 = 0$ and thus the triple eigenvalue $\lambda = 2, 2, 2$. It is easy to see that the eigenvector equation

$$(\mathbf{A} - 2\mathbf{I})\mathbf{v} = \begin{bmatrix} 0 & 3 & 4 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

has (to within a constant multiple) the single solution $\mathbf{v} = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}^T$. Thus there is only a single eigenvector associated with the eigenvalue $\lambda = 2$, and so we do not yet have the three linearly independent solutions needed for a fundamental matrix. But we note that **A** is the same matrix whose matrix exponential

$$e^{\mathbf{A}t} = \begin{bmatrix} e^{2t} & 3te^{2t} & (4t+9t^2)e^{2t} \\ 0 & e^{2t} & 6te^{2t} \\ 0 & 0 & e^{2t} \end{bmatrix}$$

was calculated in Example 4. Hence, using Theorem 2, the solution of the initial value problem in (29) is given by

$$\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{x}(0) = \begin{bmatrix} e^{2t} & 3te^{2t} & (4t+9t^2)e^{2t} \\ 0 & e^{2t} & 6te^{2t} \\ 0 & 0 & e^{2t} \end{bmatrix} \begin{bmatrix} 19 \\ 29 \\ 39 \end{bmatrix}$$
$$= \begin{bmatrix} (19+243t+351t^2)e^{2t} \\ (29+234t)e^{2t} \\ 39e^{2t} \end{bmatrix}.$$

Remark The same particular solution $\mathbf{x}(t)$ as in Example 6 could be found using the generalized eigenvector method of Section 7.6. One would start by finding the chain of generalized eigenvectors

$$\mathbf{v}_1 = \begin{bmatrix} 18\\0\\0 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 4\\6\\0 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 0\\0\\1 \end{bmatrix}$$

corresponding to the triple eigenvalue $\lambda = 2$ of the matrix **A**. Then one would—using Eqs. (27) in Section 7.6—assemble the linearly independent solutions

$$\mathbf{x}_1(t) = \mathbf{v}_1 e^{2t}, \quad \mathbf{x}_2(t) = (\mathbf{v}_1 t + \mathbf{v}_2) e^{2t}, \quad \mathbf{x}_3(t) = \left(\frac{1}{2} \mathbf{v}_1 t^2 + \mathbf{v}_2 t + \mathbf{v}_3\right) e^{2t}$$

of the differential equation $\mathbf{x}' = \mathbf{A}\mathbf{x}$ in (29). The final step would be to determine values of the coefficients c_1 , c_2 , c_3 so that the particular solution $\mathbf{x}(t) = c_1\mathbf{x}_1(t) + c_2\mathbf{x}_2(t) + c_3\mathbf{x}_3(t)$ satisfies the initial condition in (29). At this point it should be apparent that—especially if the matrix exponential $e^{\mathbf{A}t}$ is readily available (for instance, from a computer algebra system)— the method illustrated in Example 6 can well be more "computationally routine" than the generalized eigenvector method.

General Matrix Exponentials

The relatively simple calculation of e^{At} carried out in Example 4 (and used in Example 6) was based on the observation that if

$$\mathbf{A} = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 2 & 6 \\ 0 & 0 & 2 \end{bmatrix},$$

then $\mathbf{A} - 2\mathbf{I}$ is nilpotent:

$$(\mathbf{A} - 2\mathbf{I})^3 = \begin{bmatrix} 0 & 3 & 4 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix}^3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \mathbf{0}.$$
 (30)

A similar result holds for any 3×3 matrix **A** having a triple eigenvalue *r*, in which case its characteristic equation reduces to $(\lambda - r)^3 = 0$. For such a matrix, an explicit computation similar to that in Eq. (30) will show that

$$(\mathbf{A} - r\mathbf{I})^3 = \mathbf{0}.\tag{31}$$

(This particular result is a special case of the Cayley-Hamilton theorem of Section 6.3, according to which every matrix satisfies its own characteristic equation.) Thus the matrix $\mathbf{A} - r\mathbf{I}$ is nilpotent, and it follows that

$$e^{\mathbf{A}t} = e^{(r\mathbf{I} + \mathbf{A} - r\mathbf{I})t} = e^{r\mathbf{I}t} \cdot e^{(\mathbf{A} - r\mathbf{I})t} = e^{rt}\mathbf{I} \cdot \left[\mathbf{I} + (\mathbf{A} - r\mathbf{I})t + \frac{1}{2}(\mathbf{A} - r\mathbf{I})^2t^2\right], \quad (32)$$

the exponential series here terminating because of Eq. (31). In this way, we can rather easily calculate the matrix exponential e^{At} for any square matrix having only a single eigenvalue.

The calculation in Eq. (32) motivates a method of calculating $e^{\mathbf{A}t}$ for any $n \times n$ matrix **A** whatsoever. As we saw in Section 7.6, **A** has *n* linearly independent generalized eigenvectors $\mathbf{u}_1, \mathbf{u}_2, \ldots, \mathbf{u}_n$. Each generalized eigenvector **u** is associated with an eigenvalue λ of **A** and has a *rank* $r \ge 1$ such that

$$(\mathbf{A} - \lambda \mathbf{I})^r \mathbf{u} = \mathbf{0}$$
 but $(\mathbf{A} - \lambda \mathbf{I})^{r-1} \mathbf{u} \neq \mathbf{0}$. (33)

(If r = 1, then **u** is an ordinary eigenvector such that $Au = \lambda u$.)

Even if we do not yet know e^{At} explicitly, we can consider the function $\mathbf{x}(t) = e^{At}\mathbf{u}$, which is a linear combination of the column vectors of e^{At} and is therefore a solution of the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ with $\mathbf{x}(0) = \mathbf{u}$. Indeed, we can calculate \mathbf{x} explicitly in terms of \mathbf{A} , \mathbf{u} , λ , and r:

$$\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{u} = e^{(\lambda\mathbf{I}+\mathbf{A}-\lambda\mathbf{I})t}\mathbf{u} = e^{\lambda\mathbf{I}t}e^{(\mathbf{A}-\lambda\mathbf{I})t}\mathbf{u}$$
$$= e^{\lambda t}\mathbf{I}\left[\mathbf{I} + (\mathbf{A}-\lambda\mathbf{I})t + \dots + (\mathbf{A}-\lambda\mathbf{I})^{r-1}\frac{t^{r-1}}{(r-1)!} + \dots\right]\mathbf{u},$$

so

$$\mathbf{x}(t) = e^{\lambda t} \left[\mathbf{u} + (\mathbf{A} - \lambda \mathbf{I})\mathbf{u}t + (\mathbf{A} - \lambda \mathbf{I})^2 \mathbf{u} \frac{t^2}{2!} + \cdots + (\mathbf{A} - \lambda \mathbf{I})^{r-1} \mathbf{u} \frac{t^{r-1}}{(r-1)!} \right],$$
(34)

using (33) and the fact that $e^{\lambda \mathbf{I}t} = e^{\lambda t} \mathbf{I}$.

If the linearly independent solutions $\mathbf{x}_1(t)$, $\mathbf{x}_2(t)$, ..., $\mathbf{x}_n(t)$ of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ are calculated using (34) with the linearly independent generalized eigenvectors \mathbf{u}_1 , \mathbf{u}_2 , ..., \mathbf{u}_n , then the $n \times n$ matrix

$$\mathbf{\Phi}(t) = \begin{bmatrix} \mathbf{x}_1(t) & \mathbf{x}_2(t) & \cdots & \mathbf{x}_n(t) \end{bmatrix}$$
(35)

is a fundamental matrix for the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Finally, the specific fundamental matrix $\mathbf{X}(t) = \mathbf{\Phi}(t)\mathbf{\Phi}(0)^{-1}$ satisfies the initial condition $\mathbf{X}(0) = \mathbf{I}$, and thus is the desired matrix exponential $e^{\mathbf{A}t}$. We have therefore outlined a proof of the following theorem.

THEOREM 3 Computation of e^{At}

Let $\mathbf{u}_1, \mathbf{u}_2, \ldots, \mathbf{u}_n$ be *n* linearly independent generalized eigenvectors of the $n \times n$ matrix **A**. For each *i*, $1 \leq i \leq n$, let $\mathbf{x}_i(t)$ be the solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ given by (34), substituting $\mathbf{u} = \mathbf{u}_i$ and the associated eigenvalue λ and rank *r* of the generalized eigenvector \mathbf{u}_i . If the fundamental matrix $\mathbf{\Phi}(t)$ is defined by (35), then

$$e^{\mathbf{A}t} = \mathbf{\Phi}(t)\mathbf{\Phi}(0)^{-1}.$$
(36)

Example 7 Find $e^{\mathbf{A}t}$ if

$$\mathbf{A} = \begin{bmatrix} 3 & 4 & 5\\ 0 & 5 & 4\\ 0 & 0 & 3 \end{bmatrix}.$$
 (37)

Solution Theorem 3 would apply even if the matrix **A** were not upper triangular. But because **A** is upper triangular, this fact enables us to see quickly that its characteristic equation is

$$(5-\lambda)(3-\lambda)^2 = 0.$$

Thus **A** has the distinct eigenvalue $\lambda_1 = 5$ and the repeated eigenvalue $\lambda_2 = 3$.

CASE 1: $\lambda_1 = 5$. The eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{u} = 0$ for $\mathbf{u} = \begin{bmatrix} a & b & c \end{bmatrix}^T$ is

$$(\mathbf{A} - 5\mathbf{I})\mathbf{u} = \begin{bmatrix} -2 & 4 & 5\\ 0 & 0 & 4\\ 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} a\\ b\\ c \end{bmatrix} = \begin{bmatrix} 0\\ 0\\ 0 \end{bmatrix}.$$

The last two scalar equations 4c = 0 and -2c = 0 give c = 0. Then the first equation -2a + 4b = 0 is satisfied by a = 2 and b = 1. Thus the eigenvalue $\lambda_1 = 5$ has the (ordinary) eigenvector $\mathbf{u}_1 = \begin{bmatrix} 2 & 1 & 0 \end{bmatrix}^T$. The corresponding solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is

$$\mathbf{x}_{1}(t) = e^{5t} \mathbf{u}_{1} = e^{5t} \begin{bmatrix} 2 & 1 & 0 \end{bmatrix}^{T}.$$
(38)

CASE 2: $\lambda_2 = 3$. The eigenvector equation $(\mathbf{A} - \lambda \mathbf{I})\mathbf{u} = 0$ for $\mathbf{u} = \begin{bmatrix} a & b & c \end{bmatrix}^T$ is

$$(\mathbf{A} - 3\mathbf{I})\mathbf{u} = \begin{bmatrix} 0 & 4 & 5 \\ 0 & 2 & 4 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

The first two equations 4b + 5c = 0 and 2b + 4c = 0 imply that b = c = 0, but leave *a* arbitrary. Thus the eigenvalue $\lambda_2 = 3$ has the single (ordinary) eigenvector $\mathbf{u}_2 = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}^T$. The corresponding solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is

$$\mathbf{x}_{2}(t) = e^{3t} \mathbf{u}_{2} = e^{3t} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}^{T}.$$
 (39)

8.1 MATRIX EXPONENTIALS AND LINEAR SYSTEMS 479

To look for a generalized eigenvector of rank r = 2 in Eq. (33), we consider the equation

$$(\mathbf{A} - 3\mathbf{I})^2 \mathbf{u} = \begin{bmatrix} 0 & 8 & 16 \\ 0 & 4 & 8 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

The first two equations 8b + 16c = 0 and 4b + 8c = 0 are satisfied by b = 2 and c = -1, but leave *a* arbitrary. With a = 0 we get the generalized eigenvector $\mathbf{u}_3 = \begin{bmatrix} 0 & 2 & -1 \end{bmatrix}^T$ of rank r = 2 associated with the eigenvalue $\lambda = 3$. Because $(\mathbf{A} - 3\mathbf{I})^2 \mathbf{u} = \mathbf{0}$, Eq. (34) yields the third solution

$$\mathbf{x}_{3}(t) = e^{3t} \begin{bmatrix} \mathbf{u}_{3} + (\mathbf{A} - 3\mathbf{I})\mathbf{u}_{3}t \end{bmatrix}$$
$$= e^{3t} \left(\begin{bmatrix} 0\\2\\-1 \end{bmatrix} + \begin{bmatrix} 0 & 4 & 5\\0 & 2 & 4\\0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0\\2\\-1 \end{bmatrix} t \right) = e^{3t} \begin{bmatrix} 3t\\2\\-1 \end{bmatrix}.$$
(40)

With the solutions listed in Eqs. (39) and (40), the fundamental matrix

$$\mathbf{\Phi}(t) = \begin{bmatrix} \mathbf{x}_1(t) & \mathbf{x}_2(t) & \mathbf{x}_3(t) \end{bmatrix}$$

defined by Eq. (35) is

$$\Phi(t) = \begin{bmatrix} 2e^{5t} & e^{3t} & 3te^{3t} \\ e^{5t} & 0 & 2e^{3t} \\ 0 & 0 & -e^{3t} \end{bmatrix} \text{ with } \Phi(0)^{-1} = \begin{bmatrix} 0 & 1 & 2 \\ 1 & -2 & -4 \\ 0 & 0 & -1 \end{bmatrix}.$$

Hence Theorem 3 finally yields

$$e^{\mathbf{A}t} = \mathbf{\Phi}(t)\mathbf{\Phi}(0)^{-1}$$

$$= \begin{bmatrix} 2e^{5t} & e^{3t} & 3te^{3t} \\ e^{5t} & 0 & 2e^{3t} \\ 0 & 0 & -e^{3t} \end{bmatrix} \begin{bmatrix} 0 & 1 & 2 \\ 1 & -2 & -4 \\ 0 & 0 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} e^{3t} & 2e^{5t} - 2e^{3t} & 4e^{5t} - (4+3t)e^{3t} \\ 0 & e^{5t} & 2e^{5t} - 2e^{3t} \\ 0 & 0 & e^{3t} \end{bmatrix}.$$

Remark As in Example 7, Theorem 3 suffices for the computation of e^{At} provided that a basis consisting of generalized eigenvectors of **A** can be found. Alternatively, a computer algebra system can be used as indicated in the project material for this section.

8.1 **Problems**

Find a fundamental matrix of each of the systems in Problems 1 through 8, then apply Eq. (8) to find a solution satisfying the given initial conditions.

1.
$$\mathbf{x}' = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$$

2. $\mathbf{x}' = \begin{bmatrix} 2 & -1 \\ -4 & 2 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$
3. $\mathbf{x}' = \begin{bmatrix} 2 & -5 \\ 4 & -2 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$
4. $\mathbf{x}' = \begin{bmatrix} 3 & -1 \\ 1 & 1 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

5.
$$\mathbf{x}' = \begin{bmatrix} -3 & -2 \\ 9 & 3 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

6. $\mathbf{x}' = \begin{bmatrix} 7 & -5 \\ 4 & 3 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$
7. $\mathbf{x}' = \begin{bmatrix} 5 & 0 & -6 \\ 2 & -1 & -2 \\ 4 & -2 & -4 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$
8. $\mathbf{x}' = \begin{bmatrix} 3 & 2 & 2 \\ -5 & -4 & -2 \\ 5 & 5 & 3 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$

Compute the matrix exponential e^{At} for each system $\mathbf{x}' = A\mathbf{x}$ given in Problems 9 through 20.

9. $x'_1 = 5x_1 - 4x_2, x'_2 = 2x_1 - x_2$ 10. $x'_1 = 6x_1 - 6x_2, x'_2 = 4x_1 - 4x_2$ 11. $x'_1 = 5x_1 - 3x_2, x'_2 = 2x_1$ 12. $x'_1 = 5x_1 - 4x_2, x'_2 = 3x_1 - 2x_2$ 13. $x'_1 = 9x_1 - 8x_2, x'_2 = 6x_1 - 5x_2$ 14. $x'_1 = 10x_1 - 6x_2, x'_2 = 12x_1 - 7x_2$ 15. $x'_1 = 6x_1 - 10x_2, x'_2 = 2x_1 - 3x_2$ 16. $x'_1 = 11x_1 - 15x_2, x'_2 = 6x_1 - 8x_2$ 17. $x'_1 = 3x_1 + x_2, x'_2 = x_1 + 3x_2$ 18. $x'_1 = 4x_1 + 2x_2, x'_2 = 2x_1 + 4x_2$ 19. $x'_1 = 9x_1 + 2x_2, x'_2 = 2x_1 + 6x_2$ 20. $x'_1 = 13x_1 + 4x_2, x'_2 = 4x_1 + 7x_2$

In Problems 21 through 24, show that the matrix A is nilpotent and then use this fact to find (as in Example 3) the matrix exponential e^{At} .

21.
$$\mathbf{A} = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$$

22. $\mathbf{A} = \begin{bmatrix} 6 & 4 \\ -9 & -6 \end{bmatrix}$
23. $\mathbf{A} = \begin{bmatrix} 1 & -1 & -1 \\ 1 & -1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$
24. $\mathbf{A} = \begin{bmatrix} 3 & 0 & -3 \\ 5 & 0 & 7 \\ 3 & 0 & -3 \end{bmatrix}$

Each coefficient matrix **A** in Problems 25 through 30 is the sum of a nilpotent matrix and a multiple of the identity matrix. Use this fact (as in Example 6) to solve the given initial value problem.

25.
$$\mathbf{x}' = \begin{bmatrix} 2 & 5 \\ 0 & 2 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$

26. $\mathbf{x}' = \begin{bmatrix} 7 & 0 \\ 11 & 7 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 5 \\ -10 \end{bmatrix}$
27. $\mathbf{x}' = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$
28. $\mathbf{x}' = \begin{bmatrix} 5 & 0 & 0 \\ 10 & 5 & 0 \\ 20 & 30 & 5 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 40 \\ 50 \\ 60 \end{bmatrix}$
29. $\mathbf{x}' = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 1 & 6 & 3 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$

30.
$$\mathbf{x}' = \begin{bmatrix} 3 & 0 & 0 & 0 \\ 6 & 3 & 0 & 0 \\ 9 & 6 & 3 & 0 \\ 12 & 9 & 6 & 3 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- **31.** Suppose that the $n \times n$ matrices **A** and **B** commute; that is, that AB = BA. Prove that $e^{A+B} = e^A e^B$. (*Suggestion:* Group the terms in the product of the two series on the right-hand side to obtain the series on the left.)
- **32.** Deduce from the result of Problem 31 that, for every square matrix **A**, the matrix $e^{\mathbf{A}}$ is nonsingular with $(e^{\mathbf{A}})^{-1} = e^{-\mathbf{A}}$.
- **33.** Suppose that

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Show that $A^{2n} = I$ and that $A^{2n+1} = A$ if *n* is a positive integer. Conclude that

$$e^{\mathbf{A}t} = \mathbf{I}\cosh t + \mathbf{A}\sinh t,$$

and apply this fact to find a general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$. Verify that it is equivalent to the general solution found by the eigenvalue method.

34. Suppose that

$$\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}.$$

Show that $e^{\mathbf{A}t} = \mathbf{I}\cos 2t + \frac{1}{2}\mathbf{A}\sin 2t$. Apply this fact to find a general solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$, and verify that it is equivalent to the solution found by the eigenvalue method.

Apply Theorem 3 to calculate the matrix exponential $e^{\mathbf{A}t}$ for each of the matrices in Problems 35 through 40.

35.
$$\mathbf{A} = \begin{bmatrix} 3 & 4 \\ 0 & 3 \end{bmatrix}$$

36. $\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix}$
37. $\mathbf{A} = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$
38. $\mathbf{A} = \begin{bmatrix} 5 & 20 & 30 \\ 0 & 10 & 20 \\ 0 & 0 & 5 \end{bmatrix}$
39. $\mathbf{A} = \begin{bmatrix} 1 & 3 & 3 & 3 \\ 0 & 1 & 3 & 3 \\ 0 & 0 & 2 & 3 \\ 0 & 0 & 0 & 2 \end{bmatrix}$
40. $\mathbf{A} = \begin{bmatrix} 2 & 4 & 4 & 4 \\ 0 & 2 & 4 & 4 \\ 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & 3 \end{bmatrix}$

8.1 Application Automated Matrix Exponential Solutions

If **A** is an $n \times n$ matrix, then a computer algebra system can be used first to calculate the fundamental matrix e^{At} for the system

$$\mathbf{x}' = \mathbf{A}\mathbf{x},\tag{1}$$

then to calculate the matrix product $\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{x}_0$ to obtain a solution satisfying the initial condition $\mathbf{x}(0) = \mathbf{x}_0$. For instance, suppose that we want to solve the initial

Go to goo.gl/cSFGFU to download this application's computing resources including *Maple/Mathematica/*MATLAB. value problem

$$x'_{1} = 13x_{1} + 4x_{2},$$

$$x'_{2} = 4x_{1} + 7x_{2};$$

$$x_{1}(0) = 11, \quad x_{2}(0) = 23$$

After the matrices

$$\mathbf{A} = \begin{bmatrix} 13 & 4\\ 4 & 7 \end{bmatrix}, \quad \mathbf{x}_0 = \begin{bmatrix} 11\\ 23 \end{bmatrix}$$
(2)

have been entered, either the Maple command

with(linalg): exponential(A*t)

the Mathematica command

MatrixExp[A t]

or the MATLAB command

syms t, expm(A*t)

yields the matrix exponential

$$\exp \mathbf{At} = \frac{1}{5} \begin{bmatrix} e^{5t} + 4e^{15t} & -2e^{5t} + 2e^{15t} \\ -2e^{5t} + 2e^{15t} & 4e^{5t} + e^{15t} \end{bmatrix}.$$

Then either the *Maple* product multiply(expAt,x0), the *Mathematica* product expAt.x0, or the MATLAB product expAt*x0 gives the solution vector

$$\mathbf{x} = \begin{bmatrix} -7e^{5t} + 18e^{15t} \\ 14e^{5t} + 9e^{15t} \end{bmatrix}.$$

Obviously this, finally, is the way to do it!

Matrix exponentials also allow for convenient interactive exploration of the system (1). For example, a version of the *Mathematica* commands

```
A = {{13, 4}, {4, 7}};
field = VectorPlot[A.{x, y}, {x, -25, 25},
    {y, -25, 25}];
Manipulate[
    curves = ParametricPlot[
        MatrixExp[A t, #]&/@pt, {t, -1, 1},
        PlotRange -> 25];
Show[curves, field],
    {{pt, {{11, 23}, {20, -10}, {-20, -10},
        {-20, 10}}}
```

was used to generate Fig. 8.1.1, which shows four solution curves of the system (1) with the matrix **A** chosen as in Eq. (2). The initial conditions for each solution curve—one of which initially passes through the point (11, 23) of our initial value problem, while the other three pass through the points (20, -10), (-20, -10), and (20, 10)—are specified by a "locator point" which can be freely dragged to any desired position in the phase plane, with the corresponding solution curve being instantly redrawn.



FIGURE 8.1.1. Interactive display of the linear system (1). As the "locator points" are dragged to different positions, the solution curves are immediately redrawn, illustrating the behavior of the system.

Experimenting with such interactive displays can shed considerable light on the behavior of linear systems. For example, notice the straight line solution in Fig. 8.1.1; if you could drag the corresponding locator point around the phase plane, what other straight line solution could you find? How could you have predicted this by examining the matrix **A**?

For a three-dimensional example, solve the initial value problem

$$\begin{aligned} x_1' &= -149x_1 - 50x_2 - 154x_3, \\ x_2' &= 537x_1 + 180x_2 + 546x_3, \\ x_3' &= -27x_1 - 9x_2 - 25x_3; \\ x_1(0) &= 17, \quad x_2(0) = 43, \quad x_3(0) = 79 \end{aligned}$$

And here's a four-dimensional problem:

$$\begin{aligned} x_1' &= 4x_1 + x_2 + x_3 + 7x_4, \\ x_2' &= x_1 + 4x_2 + 10x_3 + x_4, \\ x_3' &= x_1 + 10x_2 + 4x_3 + x_4, \\ x_4' &= 7x_1 + x_2 + x_3 + 4x_4; \end{aligned}$$
$$\begin{aligned} x_1(0) &= 15, \quad x_2(0) = 35, \quad x_3(0) = 55, \quad x_4(0) = 75. \end{aligned}$$

If at this point you're having too much fun with matrix exponentials to stop, make up some problems of your own. For instance, choose any homogeneous linear system appearing in this chapter and experiment with different initial conditions. The exotic 5×5 matrix **A** of the Section 7.6 application may suggest some interesting possibilities.

8.2 Nonhomogeneous Linear Systems

In Section 5.5 we exhibited two techniques for finding a single particular solution of a single nonhomogeneous nth-order linear differential equation—the method of undetermined coefficients and the method of variation of parameters. Each of these may be generalized to nonhomogeneous linear systems. In a linear system modeling a physical situation, nonhomogeneous terms typically correspond to external influences, such as the inflow of liquid to a cascade of brine tanks or an external force acting on a mass-and-spring system.

Given the nonhomogeneous first-order linear system

$$\mathbf{x}' = \mathbf{A}\mathbf{x} + \mathbf{f}(t) \tag{1}$$

(2)

where **A** is an $n \times n$ constant matrix and the "nonhomogeneous term" **f**(*t*) is a given continuous vector-valued function, we know from Theorem 4 of Section 7.2 that a general solution of Eq. (1) has the form

 $\mathbf{x}(t) = \mathbf{x}_c(t) + \mathbf{x}_p(t),$

>

where

- $\mathbf{x}_c(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$ is a general solution of the associated *homogeneous* system $\mathbf{x}' = \mathbf{A}\mathbf{x}$, and
- **x**_p(t) is a single particular solution of the original nonhomogeneous system in (1).

Preceding sections have dealt with $\mathbf{x}_c(t)$, so our task now is to find $\mathbf{x}_p(t)$.

Undetermined Coefficients

First we suppose that the nonhomogeneous term $\mathbf{f}(t)$ in (1) is a linear combination (with constant vector coefficients) of products of polynomials, exponential functions, and sines and cosines. Then the method of undetermined coefficients for systems is essentially the same as for a single linear differential equation. We make an intelligent guess as to the *general form* of a particular solution \mathbf{x}_p , then attempt to determine the coefficients in \mathbf{x}_p by substitution in Eq. (1). Moreover, the choice of this general form is essentially the same as in the case of a single equation (discussed in Section 5.5); we modify it only by using undetermined *vector* coefficients rather than undetermined scalars. We will therefore confine the present discussion to illustrative examples.

Example 1 Find a particular solution of the nonhomogeneous system

$$\mathbf{x}' = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 3 \\ 2t \end{bmatrix}.$$
(3)

Solution The nonhomogeneous term $\mathbf{f} = \begin{bmatrix} 3 & 2t \end{bmatrix}^T$ is linear, so it is reasonable to select a linear trial particular solution of the form

$$\mathbf{x}_{p}(t) = \mathbf{a}t + \mathbf{b} = \begin{bmatrix} a_{1} \\ a_{2} \end{bmatrix} t + \begin{bmatrix} b_{1} \\ b_{2} \end{bmatrix}.$$
 (4)

Upon substitution of $\mathbf{x} = \mathbf{x}_p$ in Eq. (3), we get

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \begin{bmatrix} a_1t + b_1 \\ a_2t + b_2 \end{bmatrix} + \begin{bmatrix} 3 \\ 2t \end{bmatrix}$$
$$= \begin{bmatrix} 3a_1 + 2a_2 \\ 7a_1 + 5a_2 + 2 \end{bmatrix} t + \begin{bmatrix} 3b_1 + 2b_2 + 3 \\ 7b_1 + 5b_2 \end{bmatrix}.$$

We equate the coefficients of t and the constant terms (in both x_1 - and x_2 -components) and thereby obtain the equations

$$3a_1 + 2a_2 = 0,$$

$$7a_1 + 5a_2 + 2 = 0,$$

$$3b_1 + 2b_2 + 3 = a_1,$$

$$7b_1 + 5b_2 = a_2.$$

(5)

We solve the first two equations in (5) for $a_1 = 4$ and $a_2 = -6$. With these values we can then solve the last two equations in (5) for $b_1 = 17$ and $b_2 = -25$. Substitution of these coefficients in Eq. (4) gives the particular solution $\mathbf{x} = \begin{bmatrix} x_1 & x_2 \end{bmatrix}^T$ of (3) described in scalar form by

$$x_1(t) = 4t + 17,$$

 $x_2(t) = -6t - 25.$

Example 2 Cascading brine tanks Figure 8.2.1 shows the system of three brine tanks investigated in Example 2 of Section 7.3. The volumes of the three tanks are $V_1 = 20$, $V_2 = 40$, and $V_3 = 50$ (gal), and the common flow rate is r = 10 (gal/min). Suppose that all three tanks contain fresh water initially, but that the inflow to tank 1 is brine containing 2 pounds of salt per gallon, so that 20 pounds of salt flow into tank 1 per minute. Referring to Eq. (18) in Section 7.3, we see that the vector $\mathbf{x}(t) = \begin{bmatrix} x_1(t) & x_2(t) & x_3(t) \end{bmatrix}^T$ of amounts of salt (in pounds) in the three tanks at time t satisfies the nonhomogeneous initial value problem

$$\frac{d\mathbf{x}}{dt} = \begin{bmatrix} -0.5 & 0 & 0\\ 0.5 & -0.25 & 0\\ 0 & 0.25 & -0.2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 20\\ 0\\ 0 \end{bmatrix}, \quad \mathbf{x}(0) = \begin{bmatrix} 0\\ 0\\ 0 \end{bmatrix}.$$
(6)



FIGURE 8.2.1. The three brine tanks of Example 2.

The nonhomogeneous term $\mathbf{f} = \begin{bmatrix} 20 & 0 & 0 \end{bmatrix}^T$ here corresponds to the 20 lb/min inflow of salt to tank 1, with no (external) inflow of salt into tanks 2 and 3.

Because the nonhomogeneous term is constant, we naturally select a constant trial function $\mathbf{x}_p = \begin{bmatrix} a_1 & a_2 & a_3 \end{bmatrix}^T$, for which $\mathbf{x}'_p \equiv \mathbf{0}$. Then substitution of $\mathbf{x} = \mathbf{x}_p$ in (6) yields the system

$$\begin{bmatrix} 0\\0\\0 \end{bmatrix} = \begin{bmatrix} -0.5 & 0 & 0\\0.5 & -0.25 & 0\\0 & 0.25 & -0.2 \end{bmatrix} \begin{bmatrix} a_1\\a_2\\a_3 \end{bmatrix} + \begin{bmatrix} 20\\0\\0 \end{bmatrix}$$

that we readily solve for $a_1 = 40$, $a_2 = 80$, and $a_3 = 100$ in turn. Thus our particular solution is $\mathbf{x}_p(t) = \begin{bmatrix} 40 & 80 & 100 \end{bmatrix}^T$.

In Example 2 of Section 7.3 we found the general solution

$$\mathbf{x}_{c}(t) = c_{1} \begin{bmatrix} 3\\-6\\5 \end{bmatrix} e^{-t/2} + c_{2} \begin{bmatrix} 0\\1\\-5 \end{bmatrix} e^{-t/4} + c_{3} \begin{bmatrix} 0\\0\\1 \end{bmatrix} e^{-t/5}$$

of the associated homogeneous system, so a general solution $\mathbf{x} = \mathbf{x}_c + \mathbf{x}_p$ of the nonhomogeneous system in (6) is given by

$$\mathbf{x}(t) = c_1 \begin{bmatrix} 3\\-6\\5 \end{bmatrix} e^{-t/2} + c_2 \begin{bmatrix} 0\\1\\-5 \end{bmatrix} e^{-t/4} + c_3 \begin{bmatrix} 0\\0\\1 \end{bmatrix} e^{-t/5} + \begin{bmatrix} 40\\80\\100 \end{bmatrix}.$$
(7)

When we apply the zero initial conditions in (6), we get the scalar equations

$$3c_1 + 40 = 0,$$

$$-6c_1 + c_2 + 80 = 0,$$

$$5c_1 - 5c_2 + c_3 + 100 = 0$$

that are readily solved for $c_1 = -\frac{40}{3}$, $c_2 = -160$, and $c_3 = -\frac{2500}{3}$. Substituting these coefficients in Eq. (7), we find that the amounts of salt in the three tanks at time t are given by

$$x_{1}(t) = 40 - 40e^{-t/2},$$

$$x_{2}(t) = 80 + 80e^{-t/2} - 160e^{-t/4},$$

$$x_{3}(t) = 100 + \frac{100}{3} \left(-2e^{-t/2} + 24e^{-t/4} - 25e^{-t/5} \right).$$
(8)

As illustrated in Fig. 8.2.2, we see the salt in each of the three tanks approaching, as $t \to +\infty$, a uniform density of 2 lb/gal—the same as the salt density in the inflow to tank 1.

In the case of duplicate expressions in the complementary function and the nonhomogeneous terms, there is one difference between the method of undetermined coefficients for systems and for single equations (Rule 2 in Section 5.5). For a system, the usual first choice for a trial solution must be multiplied not only by the smallest integral power of t that will eliminate duplication, but also by all lower (nonnegative integral) powers of t as well, and all the resulting terms must be included in the trial solution.

Example 3

Consider the nonhomogeneous system

$$\mathbf{x}' = \begin{bmatrix} 4 & 2\\ 3 & -1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 15\\ 4 \end{bmatrix} t e^{-2t}.$$
(9)

In Example 1 of Section 7.3 we found the solution

$$\mathbf{x}_{c}(t) = c_{1} \begin{bmatrix} 1\\ -3 \end{bmatrix} e^{-2t} + c_{2} \begin{bmatrix} 2\\ 1 \end{bmatrix} e^{5t}$$
(10)



FIGURE 8.2.2. Solution curves for the amount of salt defined in (8).

of the associated homogeneous system. A preliminary trial solution $\mathbf{x}_p(t) = \mathbf{a}te^{-2t} + \mathbf{b}e^{-2t}$ exhibits duplication with the complementary function in (10). We would therefore select

$$\mathbf{x}_p(t) = \mathbf{a}t^2 e^{-2t} + \mathbf{b}t e^{-2t} + \mathbf{c}e^{-2t}$$

as our trial solution, and we would then have six scalar coefficients to determine. It is simpler to use the method of variation of parameters, our next topic.

Variation of Parameters

Recall from Section 5.5 that the method of variation of parameters may be applied to a linear differential equation with variable coefficients and is not restricted to nonhomogeneous terms involving only polynomials, exponentials, and sinusoidal functions. The method of variation of parameters for systems enjoys the same flexibility and has a concise matrix formulation that is convenient for both practical and theoretical purposes.

We want to find a particular solution \mathbf{x}_p of the nonhomogeneous linear system

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t),\tag{11}$$

given that we have already found a general solution

$$\mathbf{x}_c(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t) + \dots + c_n \mathbf{x}_n(t)$$
(12)

of the associated homogeneous system

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x}.\tag{13}$$

We first use the fundamental matrix $\Phi(t)$ with column vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ to rewrite the complementary function in (12) as

$$\mathbf{x}_c(t) = \mathbf{\Phi}(t)\mathbf{c},\tag{14}$$

where **c** denotes the column vector whose entries are the coefficients c_1, c_2, \ldots, c_n . Our idea is to replace the vector "parameter" **c** with a variable vector **u**(*t*). Thus we seek a particular solution of the form

$$\mathbf{x}_{p}(t) = \mathbf{\Phi}(t)\mathbf{u}(t). \tag{15}$$

We must determine $\mathbf{u}(t)$ so that \mathbf{x}_p does, indeed, satisfy Eq. (11). The derivative of $\mathbf{x}_p(t)$ is (by the product rule)

$$\mathbf{x}'_{p}(t) = \mathbf{\Phi}'(t)\mathbf{u}(t) + \mathbf{\Phi}(t)\mathbf{u}'(t).$$
(16)

Hence substitution of Eqs. (15) and (16) in (11) yields

$$\mathbf{\Phi}'(t)\mathbf{u}(t) + \mathbf{\Phi}(t)\mathbf{u}'(t) = \mathbf{P}(t)\mathbf{\Phi}(t)\mathbf{u}(t) + \mathbf{f}(t).$$
(17)

But

$$\mathbf{\Phi}'(t) = \mathbf{P}(t)\mathbf{\Phi}(t) \tag{18}$$

because each column vector of $\Phi(t)$ satisfies Eq. (13). Therefore, Eq. (17) reduces to

$$\mathbf{\Phi}(t)\mathbf{u}'(t) = \mathbf{f}(t). \tag{19}$$

Thus it suffices to choose $\mathbf{u}(t)$ so that

$$\mathbf{u}'(t) = \mathbf{\Phi}(t)^{-1}\mathbf{f}(t); \tag{20}$$

that is, so that

$$\mathbf{u}(t) = \int \mathbf{\Phi}(t)^{-1} \mathbf{f}(t) \, dt.$$
(21)

Upon substitution of (21) in (15), we finally obtain the desired particular solution, as stated in the following theorem.

THEOREM 1 Variation of Parameters

If $\Phi(t)$ is a fundamental matrix for the homogeneous system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ on some interval where $\mathbf{P}(t)$ and $\mathbf{f}(t)$ are continuous, then a particular solution of the non-homogeneous system

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t)$$

is given by

>

$$\mathbf{x}_{p}(t) = \mathbf{\Phi}(t) \int \mathbf{\Phi}(t)^{-1} \mathbf{f}(t) dt.$$
(22)

This is the **variation of parameters formula** for first-order linear systems. If we add this particular solution and the complementary function in (14), we get the general solution

$$\mathbf{x}(t) = \mathbf{\Phi}(t)\mathbf{c} + \mathbf{\Phi}(t)\int \mathbf{\Phi}(t)^{-1}\mathbf{f}(t) dt$$
(23)

of the nonhomogeneous system in (11).

The choice of the constant of integration in Eq. (22) is immaterial, for we need only a single particular solution. In solving initial value problems it often is convenient to choose the constant of integration so that $\mathbf{x}_p(a) = \mathbf{0}$, and thus integrate from *a* to *t*:

$$\mathbf{x}_p(t) = \mathbf{\Phi}(t) \int_a^t \mathbf{\Phi}(s)^{-1} \mathbf{f}(s) \, ds.$$
(24)

If we add the particular solution of the nonhomogeneous problem

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t), \quad \mathbf{x}(a) = \mathbf{0}$$

in (24) to the solution $\mathbf{x}_c(t) = \mathbf{\Phi}(t)\mathbf{\Phi}(a)^{-1}\mathbf{x}_a$ of the associated homogeneous problem $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}, \mathbf{x}(a) = \mathbf{x}_a$, we get the solution

$$\mathbf{x}(t) = \mathbf{\Phi}(t)\mathbf{\Phi}(a)^{-1}\mathbf{x}_a + \mathbf{\Phi}(t)\int_a^t \mathbf{\Phi}(s)^{-1}\mathbf{f}(s)\,ds \tag{25}$$

of the nonhomogeneous initial value problem

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t), \quad \mathbf{x}(a) = \mathbf{x}_a.$$
(26)

Equations (22) and (25) hold for any fundamental matrix $\Phi(t)$ of the homogeneous system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$. In the constant-coefficient case $\mathbf{P}(t) \equiv \mathbf{A}$ we can use for $\Phi(t)$ the exponential matrix $e^{\mathbf{A}t}$ —that is, the particular fundamental matrix such that $\Phi(0) = \mathbf{I}$. Then, because $(e^{\mathbf{A}t})^{-1} = e^{-\mathbf{A}t}$, substitution of $\Phi(t) = e^{\mathbf{A}t}$ in (22) yields the particular solution

$$\mathbf{x}_{p}(t) = e^{\mathbf{A}t} \int e^{-\mathbf{A}t} \mathbf{f}(t) dt$$
(27)

of the nonhomogeneous system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t)$. Similarly, substitution of $\mathbf{\Phi}(t) = e^{\mathbf{A}t}$ in Eq. (25) with a = 0 yields the solution

$$\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{x}_0 + e^{\mathbf{A}t}\int_0^t e^{-\mathbf{A}t}\mathbf{f}(t)\,dt \tag{28}$$

of the initial value problem

>

≻

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t), \quad \mathbf{x}(0) = \mathbf{x}_0.$$
⁽²⁹⁾

Remark If we retain t as the independent variable but use s for the variable of integration, then the solutions in (27) and (28) can be rewritten in the forms

$$\mathbf{x}_p(t) = \int e^{-\mathbf{A}(s-t)} \mathbf{f}(s) \, ds \quad \text{and} \quad \mathbf{x}(t) = e^{\mathbf{A}t} \mathbf{x}_0 + \int_0^t e^{-\mathbf{A}(s-t)} \mathbf{f}(s) \, ds.$$

Example 4

Solve the initial value problem

$$\mathbf{x}' = \begin{bmatrix} 4 & 2\\ 3 & -1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 15\\ 4 \end{bmatrix} t e^{-2t}, \quad \mathbf{x}(0) = \begin{bmatrix} 7\\ 3 \end{bmatrix}.$$
(30)

Solution The solution of the associated homogeneous system is displayed in Eq. (10). It gives the fundamental matrix

$$\Phi(t) = \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \text{ with } \Phi(0)^{-1} = \frac{1}{7} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix}.$$

It follows by Eq. (28) in Section 8.1 that the matrix exponential for the coefficient matrix **A** in (30) is

$$e^{\mathbf{A}t} = \mathbf{\Phi}(t)\mathbf{\Phi}(0)^{-1} = \begin{bmatrix} e^{-2t} & 2e^{5t} \\ -3e^{-2t} & e^{5t} \end{bmatrix} \cdot \frac{1}{7} \begin{bmatrix} 1 & -2 \\ 3 & 1 \end{bmatrix}$$
$$= \frac{1}{7} \begin{bmatrix} e^{-2t} + 6e^{5t} & -2e^{-2t} + 2e^{5t} \\ -3e^{-2t} + 3e^{5t} & 6e^{-2t} + e^{5t} \end{bmatrix}.$$

Then the variation of parameters formula in Eq. (28) gives

$$e^{-\mathbf{A}t}\mathbf{x}(t) = \mathbf{x}_{0} + \int_{0}^{t} e^{-\mathbf{A}s}\mathbf{f}(s) ds$$

$$= \begin{bmatrix} 7\\3 \end{bmatrix} + \int_{0}^{t} \frac{1}{7} \begin{bmatrix} e^{2s} + 6e^{-5s} & -2e^{2s} + 2e^{-5s} \\ -3e^{2s} + 3e^{-5s} & 6e^{2s} + e^{-5s} \end{bmatrix} \begin{bmatrix} -15se^{-2s} \\ -4se^{-2s} \end{bmatrix} ds$$

$$= \begin{bmatrix} 7\\3 \end{bmatrix} + \int_{0}^{t} \begin{bmatrix} -s - 14se^{-7s} \\ 3s - 7se^{-7s} \end{bmatrix} ds$$

$$= \begin{bmatrix} 7\\3 \end{bmatrix} + \frac{1}{14} \begin{bmatrix} -4 - 7t^{2} + 4e^{-7t} + 28te^{-7t} \\ -2 + 21t^{2} + 2e^{-7t} + 14te^{-7t} \end{bmatrix}.$$

488 CHAPTER 8 MATRIX EXPONENTIAL METHODS

Therefore,

$$e^{-\mathbf{A}t}\mathbf{x}(t) = \frac{1}{14} \begin{bmatrix} 94 - 7t^2 + 4e^{-7t} + 28te^{-7t} \\ 40 + 21t^2 + 2e^{-7t} + 14te^{-7t} \end{bmatrix}$$

Upon multiplication of the right-hand side here by $e^{\mathbf{A}t}$, we find that the solution of the initial value problem in (30) is given by

$$\mathbf{x}(t) = \frac{1}{7} \begin{bmatrix} e^{-2t} + 6e^{5t} & -2e^{-2t} + 2e^{5t} \\ -3e^{-2t} + 3e^{5t} & 6e^{-2t} + e^{5t} \end{bmatrix} \cdot \frac{1}{14} \begin{bmatrix} 94 - 7t^2 + 4e^{-7t} + 28te^{-7t} \\ 40 + 21t^2 + 2e^{-7t} + 14te^{-7t} \end{bmatrix}$$
$$= \frac{1}{14} \begin{bmatrix} (6 + 28t - 7t^2)e^{-2t} + 92e^{5t} \\ (-4 + 14t + 21t^2)e^{-2t} + 46e^{5t} \end{bmatrix}.$$

In conclusion, let us investigate how the variation of parameters formula in (22) "reconciles" with the variation of parameters formula in Theorem 1 of Section 5.5 for the second-order linear differential equation

$$y'' + Py' + Qy = f(t).$$
 (31)

If we write $y = x_1$, $y' = x'_1 = x_2$, $y'' = x''_1 = x'_2$, then the single equation in (31) is equivalent to the linear system $x'_1 = x_2$, $x'_2 = -Qx_1 - Px_2 + f(t)$, that is,

$$\mathbf{x}' = \mathbf{P}(t)\mathbf{x} + \mathbf{f}(t),\tag{32}$$

where

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} y \\ y' \end{bmatrix}, \quad \mathbf{P}(t) = \begin{bmatrix} 0 & 1 \\ -Q & -P \end{bmatrix}, \quad \text{and} \quad \mathbf{f}(t) = \begin{bmatrix} 0 \\ f(t) \end{bmatrix}.$$

Now two linearly independent solutions y_1 and y_2 of the homogeneous system y'' + Py' + Qy = 0 associated with (31) provide two linearly independent solutions

$$\mathbf{x}_1 = \begin{bmatrix} y_1 \\ y'_1 \end{bmatrix}$$
 and $\mathbf{x}_2 = \begin{bmatrix} y_2 \\ y'_2 \end{bmatrix}$

of the homogeneous system $\mathbf{x}' = \mathbf{P}(t)\mathbf{x}$ associated with (32). Observe that the determinant of the fundamental matrix $\mathbf{\Phi} = \begin{bmatrix} \mathbf{x}_1 & \mathbf{x}_2 \end{bmatrix}$ is simply the Wronskian

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$$

of the solutions y_1 and y_2 , so the inverse fundamental matrix is

$$\Phi^{-1} = \frac{1}{W} \begin{vmatrix} y_2' & -y_2 \\ -y_1' & y_1 \end{vmatrix}.$$

Therefore the variation of parameters formula $\mathbf{x}_p = \mathbf{\Phi} \int \mathbf{\Phi}^{-1} \mathbf{f} dt$ in (22) yields

$$\begin{bmatrix} y_p \\ y'_p \end{bmatrix} = \begin{bmatrix} y_1 & y_2 \\ y'_1 & y'_2 \end{bmatrix} \int \frac{1}{W} \begin{bmatrix} y'_2 & -y_2 \\ -y'_1 & y_1 \end{bmatrix} \begin{bmatrix} 0 \\ f \end{bmatrix} dt$$
$$= \begin{bmatrix} y_1 & y_2 \\ y'_1 & y'_2 \end{bmatrix} \int \frac{1}{W} \begin{bmatrix} -y_2 f \\ y_1 f \end{bmatrix} dt.$$

The first component of this column vector is

$$y_p = \begin{bmatrix} y_1 & y_2 \end{bmatrix} \int \frac{1}{W} \begin{bmatrix} -y_2 f \\ y_1 f \end{bmatrix} dt = -y_1 \int \frac{y_2 f}{W} dt + y_2 \int \frac{y_1 f}{W} dt$$

If, finally, we supply the independent variable t throughout, the final result on the right-hand side here is simply the variation of parameters formula in Eq. (33) of Section 5.5 (where, however, the independent variable is denoted by x).

8.2 **Problems**

Apply the method of undetermined coefficients to find a particular solution of each of the systems in Problems 1 through 14. If initial conditions are given, find the particular solution that satisfies these conditions. Primes denote derivatives with respect to t.

1.
$$x' = x + 2y + 3$$
, $y' = 2x + y - 2$
2. $x' = 2x + 3y + 5$, $y' = 2x + y - 2t$
3. $x' = 3x + 4y$, $y' = 3x + 2y + t^{2}$; $x(0) = y(0) = 0$
4. $x' = 4x + y + e^{t}$, $y' = 6x - y - e^{t}$; $x(0) = y(0) = 1$
5. $x' = 6x - 7y + 10$, $y' = x - 2y - 2e^{-t}$
6. $x' = 9x + y + 2e^{t}$, $y' = -8x - 2y + te^{t}$
7. $x' = -3x + 4y + \sin t$, $y' = 6x - 5y$; $x(0) = 1$, $y(0) = 0$
8. $x' = x - 5y + 2\sin t$, $y' = x - y - 3\cos t$
9. $x' = x - 5y + \cos 2t$, $y' = x - y$
10. $x' = x - 2y$, $y' = 2x - y + e^{t}\sin t$
11. $x' = 2x + 4y + 2$, $y' = x + 2y + 3$; $x(0) = 1$, $y(0) = -1$
12. $x' = x + y + 2t$, $y' = x + y - 2t$
13. $x' = 2x + y + 2e^{t}$, $y' = x + 2y - 3e^{t}$
14. $x' = 2x + y + 1$, $y' = 4x + 2y + e^{4t}$

Two Brine Tanks

Problems 15 and 16 are similar to Example 2, but with two brine tanks (having volumes V_1 and V_2 gallons as in Fig. 8.2.1) instead of three tanks. Each tank initially contains fresh water, and the inflow to tank 1 at the rate of r gallons per minute has a salt concentration of c_0 pounds per gallon. (a) Find the amounts $x_1(t)$ and $x_2(t)$ of salt in the two tanks after t minutes. (b) Find the limiting (long-term) amount of salt in each tank. (c) Find how long it takes for each tank to reach a salt concentration of 1 lb/gal.

15. $V_1 = 100, V_2 = 200, r = 10, c_0 = 2$ **16.** $V_1 = 200, V_2 = 100, r = 10, c_0 = 3$

In Problems 17 through 34, use the method of variation of parameters (and perhaps a computer algebra system) to solve the initial value problem

$$\mathbf{x}' = \mathbf{A}\mathbf{x} + \mathbf{f}(t), \quad \mathbf{x}(a) = \mathbf{x}_a.$$

In each problem we provide the matrix exponential e^{At} as provided by a computer algebra system.

17. $\mathbf{A} = \begin{bmatrix} 6 & -7 \\ 1 & -2 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 60 \\ 90 \end{bmatrix}, \mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$ $e^{\mathbf{A}t} = \frac{1}{6} \begin{bmatrix} -e^{-t} + 7e^{5t} & 7e^{-t} - 7e^{5t} \\ -e^{-t} + e^{5t} & 7e^{-t} - e^{5t} \end{bmatrix}$

18. Repeat Problem 17, but with $\mathbf{f}(t)$ replaced with $\begin{bmatrix} 100t \\ -50t \end{bmatrix}$.

19.
$$\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 2 & -2 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 180t \\ 90 \end{bmatrix}, \mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

 $e^{\mathbf{A}t} = \frac{1}{5} \begin{bmatrix} e^{-3t} + 4e^{2t} & -2e^{-3t} + 2e^{2t} \\ -2e^{-3t} + 2e^{2t} & 4e^{-3t} + e^{2t} \end{bmatrix}$

20. Repeat Problem 19, but with $\mathbf{f}(t)$ replaced with $\begin{bmatrix} 75e^{2t} \\ 0 \end{bmatrix}$.

21.
$$\mathbf{A} = \begin{bmatrix} 4 & -1 \\ 5 & -2 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 18e^{2t} \\ 30e^{2t} \end{bmatrix}, \mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$e^{\mathbf{A}t} = \frac{1}{4} \begin{bmatrix} -e^{-t} + 5e^{3t} & e^{-t} - e^{3t} \\ -5e^{-t} + 5e^{3t} & 5e^{-t} - e^{3t} \end{bmatrix}$$

22. Repeat Problem 21, but with $\mathbf{f}(t)$ replaced with $\begin{bmatrix} 28e^{-t} \\ 20e^{3t} \end{bmatrix}$.

23.
$$\mathbf{A} = \begin{bmatrix} 3 & -1 \\ 9 & -3 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 7 \\ 5 \end{bmatrix}, \mathbf{x}(0) = \begin{bmatrix} 3 \\ 5 \end{bmatrix}$$

 $e^{\mathbf{A}t} = \begin{bmatrix} 1+3t & -t \\ 9t & 1-3t \end{bmatrix}$

24. Repeat Problem 23, but with $\mathbf{f}(t) = \begin{bmatrix} 0 \\ t^{-2} \end{bmatrix}$ and $\mathbf{x}(1) = \begin{bmatrix} 3 \end{bmatrix}$

$$\begin{bmatrix} 7 \end{bmatrix}^{\cdot}$$
25. $\mathbf{A} = \begin{bmatrix} 2 & -5 \\ 1 & -2 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 4t \\ 1 \end{bmatrix}, \mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$

$$e^{\mathbf{A}t} = \begin{bmatrix} \cos t + 2\sin t & -5\sin t \\ \sin t & \cos t - 2\sin t \end{bmatrix}$$

26. Repeat Problem 25, but with
$$\mathbf{f}(t) = \begin{bmatrix} 4 \cos t \\ 6 \sin t \end{bmatrix}$$
 and $\mathbf{x}(0) = \begin{bmatrix} 3 \end{bmatrix}$

$$\begin{bmatrix} 5 \end{bmatrix}^{\cdot}$$
27. $\mathbf{A} = \begin{bmatrix} 2 & -4 \\ 1 & -2 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 36t^2 \\ 6t \end{bmatrix}, \mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, e^{\mathbf{A}t} = \begin{bmatrix} 1+2t & -4t \\ t & 1-2t \end{bmatrix}$
28. Repeat Problem 27, but with
$$\mathbf{f}(t) = \begin{bmatrix} 4 \ln t \\ t^{-1} \end{bmatrix}$$
 and $\mathbf{x}(1) = \begin{bmatrix} e^{t} & 3te^{t} & (-13 - 9t)e^{t} + 13e^{2t} \\ 0 & e^{t} & -3e^{t} + 3e^{2t} \\ 0 & 0 & e^{2t} \end{bmatrix}$
29. $\mathbf{A} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, $\mathbf{f}(t) = \begin{bmatrix} \sec t \\ 0 \end{bmatrix}$, $\mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$,
 $e^{\mathbf{A}t} = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix}$
30. $\mathbf{A} = \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix}$, $\mathbf{f}(t) = \begin{bmatrix} t \cos 2t \\ t \sin 2t & \cos 2t \end{bmatrix}$, $\mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$,
 $e^{\mathbf{A}t} = \begin{bmatrix} \cos 2t & -\sin 2t \\ \sin 2t & \cos 2t \end{bmatrix}$
31. $\mathbf{A} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$, $\mathbf{f}(t) = \begin{bmatrix} 0 \\ 0 \\ 6e^{t} \end{bmatrix}$, $\mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$,
 $e^{\mathbf{A}t} = \begin{bmatrix} e^{t} & 2te^{t} & (3t + 2t^{2})e^{t} \\ 0 & e^{t} & 2te^{t} \end{bmatrix}$, $\mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$,
 $e^{\mathbf{A}t} = \begin{bmatrix} e^{t} & 2te^{t} & (3t + 2t^{2})e^{t} \\ 0 & 0 & e^{t} \end{bmatrix}$
32. $\mathbf{A} = \begin{bmatrix} 1 & 3 & 4 \\ 0 & 1 & 3 \\ 0 & 0 & 2 \end{bmatrix}$, $\mathbf{f}(t) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 2e^{2t} \end{bmatrix}$, $\mathbf{x}(0) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$,
 $e^{\mathbf{A}t} = \begin{bmatrix} 1 & 4t & 4(-1 + e^{2t}) & 16t(-1 + e^{2t}) \\ 0 & 1 & 0 & 4(-1 + e^{2t}) \\ 0 & 0 & 0 & e^{2t} \end{bmatrix}$, $\mathbf{x}(0) = \begin{bmatrix} 4 \\ 2 \\ 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$,
 $e^{\mathbf{A}t} = \begin{bmatrix} 1 & 4t & 4(-1 + e^{2t}) & 16t(-1 + e^{2t}) \\ 0 & 1 & 0 & 4(-1 + e^{2t}) \\ 0 & 0 & 0 & e^{2t} \end{bmatrix}$

8.2 Application Automated Variation of Parameters



The application of the variation of parameters formula in Eq. (28) encourages so mechanical an approach as to encourage especially the use of a computer algebra system. The following *Mathematica* commands were used to check the results in Example 4 of this section.

```
A = {{4,2}, {3,-1}};
x0 = {{7}, {3}};
f[t_] := {{-15 t Exp[-2t]}, {-4 t Exp[-2t]};
exp[A_] := MatrixExp[A]
x = exp[A*t].(x0 + Integrate[exp[-A*s].f[s], {s,0,t}])
```

The matrix exponential commands illustrated in the Section 5.6 application provide the basis for analogous *Maple* and MATLAB computations. You can then check routinely the answers for Problems 17 through 34 of this section.

8.3 Spectral Decomposition Methods

Here, we present an alternative approach to the computation of the matrix exponential $e^{\mathbf{A}t}$, one that does not require that eigenvectors (including generalized ones) of the $n \times n$ matrix **A** be found first. Assume that the characteristic polynomial of **A** is written in the form

$$p(\lambda) = (-1)^n |\mathbf{A} - \lambda \mathbf{I}|, \tag{1}$$

with leading term $+\lambda^n$. [Compare Eqs. (4) and (5) in Section 6.1.] If the (not necessarily distinct) eigenvalues of **A** are $\lambda_1, \lambda_2, \ldots, \lambda_n$, then

$$p(\lambda) = (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n).$$
⁽²⁾

The Cayley-Hamilton theorem (Section 6.3) says that any matrix **A** satisfies its own characteristic equation; that is,

$$p(\mathbf{A}) = \prod_{i=1}^{n} (\mathbf{A} - \lambda_i \mathbf{I}) = \mathbf{0}$$
(3)

(where **I** denotes the $n \times n$ identity matrix). This crucial fact is the key to our method in this section.

The way we proceed to calculate the matrix exponential e^{At} depends on whether or not the eigenvalues of **A** are distinct.

The Case of Distinct Eigenvalues

If the eigenvalues of **A** are distinct (whether real or complex), the reciprocal $1/p(\lambda)$ has a partial fraction decomposition of the form

$$\frac{1}{p(\lambda)} = \frac{a_1}{\lambda - \lambda_1} + \frac{a_2}{\lambda - \lambda_2} + \dots + \frac{a_n}{\lambda - \lambda_n},\tag{4}$$

where the numerators $a_1, a_2, ..., a_n$ are constants. Indeed, these constants can be found by multiplying Eq. (4) by $p(\lambda)$ to get

$$1 = \sum_{i=1}^{n} a_i b_i(\lambda) = a_1 b_1(\lambda) + a_2 b_2(\lambda) + \dots + a_n b_n(\lambda),$$
(5)

where the polynomial

$$b_i(\lambda) = \frac{p(\lambda)}{\lambda - \lambda_i} = \prod_{j \neq i} (\lambda - \lambda_j)$$
(6)

is obtained from the characteristic polynomial $p(\lambda)$ upon deletion of the factor $(\lambda - \lambda_i)$ corresponding to the *i*th eigenvalue.

It follows immediately from (6) that $b_i(\lambda_i) \neq 0$, whereas $b_j(\lambda_i) = 0$ if $i \neq j$ [because in this case $b_j(\lambda_i)$ includes the factor $(\lambda_i - \lambda_i)$]. Consequently, substitution of $\lambda = \lambda_i$ into Eq. (5) gives the value

$$a_i = \frac{1}{b_i(\lambda_i)} = \frac{1}{\prod_{j \neq i} (\lambda_i - \lambda_j)}$$
(7)

for i = 1, 2, ..., n. Given these coefficient values, we define the *projection matrices* $\mathbf{P}_1, \mathbf{P}_2, ..., \mathbf{P}_n$ of the fixed $n \times n$ matrix **A** by writing

$$\mathbf{P}_{i} = a_{i}b_{i}(\mathbf{A}) = a_{i}\prod_{j\neq i}(\mathbf{A} - \lambda_{j}\mathbf{I}) = \frac{\prod_{j\neq i}(\mathbf{A} - \lambda_{j}\mathbf{I})}{\prod_{j\neq i}(\lambda_{i} - \lambda_{j})}.$$
(8)

These matrices have very special properties that are summarized by the following proposition.

PROPOSITION 1 The Projection Matrices of A

If the eigenvalues of **A** are distinct, then the projection matrices $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_n$ defined by (8) satisfy the relations

(i)
$$\mathbf{P}_1 + \mathbf{P}_2 + \dots + \mathbf{P}_n = \mathbf{I};$$
 (9)

(ii)
$$\mathbf{P}_i \mathbf{P}_j = \mathbf{0} \text{ if } i \neq j;$$
 (10)

(iii) $\mathbf{P}_i^2 = \mathbf{P}_i$ for each $i = 1, 2, \dots, n$; (11)

(iv)
$$\mathbf{P}_i \mathbf{A} = \lambda_i \mathbf{P}_i$$
 for each $i = 1, 2, \dots, n$. (12)

Remark In terms of the Kronecker delta defined by

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

the conditions in (10) and (11) say that

$$\mathbf{P}_i \mathbf{P}_j = \delta_{ij} \mathbf{P}_i. \tag{13}$$

Also, note that (12) says that \mathbf{P}_i is effectively an "eigenmatrix" associated with the eigenvalue λ_i of the matrix **A**—that is, multiplication of \mathbf{P}_i by the matrix **A** (on the right) gives a scalar multiple of \mathbf{P}_i .

Proof: We get (9) when we substitute **A** for λ in the algebraic identity (5) and recall the definition in (8). If $i \neq j$, then the product

$$\mathbf{P}_i \mathbf{P}_j = a_i b_i(\mathbf{A}) \cdot a_j b_j(\mathbf{A}) = a_i a_j \cdot \prod_{r \neq i} (\mathbf{A} - \lambda_r \mathbf{I}) \cdot \prod_{s \neq j} (\mathbf{A} - \lambda_s \mathbf{I})$$

evidently has the product $p(\mathbf{A}) = \prod_{r=1}^{n} (\mathbf{A} - \lambda_r \mathbf{I})$ as a factor. Hence (10) follows from the Cayley-Hamilton result $p(\mathbf{A}) = \mathbf{0}$ in (3). Next, (11) follows when we multiply (9) by \mathbf{P}_i :

$$\mathbf{P}_i = \mathbf{P}_i \mathbf{I} = \mathbf{P}_i (\mathbf{P}_1 + \mathbf{P}_2 + \dots + \mathbf{P}_n) = \sum_{j=1}^n \mathbf{P}_i \mathbf{P}_j = \mathbf{P}_i^2,$$

because of (10). Finally, to verify (12), we note that

$$\mathbf{P}_{i}\mathbf{A} - \lambda_{i}\mathbf{P}_{i} = \mathbf{P}_{i} \cdot (\mathbf{A} - \lambda_{i}\mathbf{I})$$

$$= a_{i} \prod_{j \neq i} (\mathbf{A} - \lambda_{j}\mathbf{I}) \cdot (\mathbf{A} - \lambda_{i}\mathbf{I})$$

$$= a_{i} \prod_{j=1}^{n} (\mathbf{A} - \lambda_{j}\mathbf{I})$$

$$= a_{i} p(\mathbf{A})$$

$$\mathbf{P}_{i}\mathbf{A} - \lambda_{i}\mathbf{P}_{i} = \mathbf{0},$$

by the use first of (8) and then of (2).

The following theorem expresses the matrix \mathbf{A} in terms of its projection matrices.

THEOREM 1 The Spectral Decomposition of A

If the $n \times n$ matrix **A** has distinct eigenvalues $\lambda_1, \lambda_2, ..., \lambda_n$, and if the projection matrices $\mathbf{P}_1, \mathbf{P}_2, ..., \mathbf{P}_n$ are defined as in (8), then

$$\mathbf{A} = \sum_{i=1}^{n} \lambda_i \mathbf{P}_i = \lambda_1 \mathbf{P}_1 + \lambda_2 \mathbf{P}_2 + \dots + \lambda_n \mathbf{P}_n.$$
(14)

Proof: First, we multiply both sides in (9) on the right by A and get

$$\mathbf{A} = \mathbf{P}_1 \mathbf{A} + \mathbf{P}_2 \mathbf{A} + \dots + \mathbf{P}_n \mathbf{A}$$

= $\mathbf{P}_1 (\mathbf{A} - \lambda_1 \mathbf{I} + \lambda_1 \mathbf{I}) + \mathbf{P}_2 (\mathbf{A} - \lambda_2 \mathbf{I} + \lambda_2 \mathbf{I}) + \dots + \mathbf{P}_n (\mathbf{A} - \lambda_n \mathbf{I} + \lambda_n \mathbf{I}).$

Then, we notice using (12) that

$$\mathbf{P}_i(\mathbf{A} - \lambda_i \mathbf{I} + \lambda_i \mathbf{I}) = \mathbf{P}_i(\mathbf{A} - \lambda_i \mathbf{I}) + \lambda_i \mathbf{P}_i \mathbf{I} = \mathbf{0} + \lambda_i \mathbf{P}_i = \lambda_i \mathbf{P}_i;$$

so (14) follows.

Remark 1 If we square both sides in (14) and then use (13), we get

$$(\lambda_1 \mathbf{P}_1 + \lambda_2 \mathbf{P}_2 + \dots + \lambda_n \mathbf{P}_n) \cdot (\lambda_1 \mathbf{P}_1 + \lambda_2 \mathbf{P}_2 + \dots + \lambda_n \mathbf{P}_n) = \sum_{i,j=1}^n \lambda_i \lambda_j \mathbf{P}_i \mathbf{P}_j,$$

or

$$\mathbf{A}^2 = \sum_{i=1}^n \lambda_i^2 \mathbf{P}_i = \lambda_1^2 \mathbf{P}_2 + \lambda_2^2 \mathbf{P}_2 + \dots + \lambda_n^2 \mathbf{P}_n.$$

Continuing in this fashion, we deduce by repeated multiplication that

$$\mathbf{A}^{k} = \sum_{i=1}^{n} \lambda_{i}^{k} \mathbf{P}_{i} = \lambda_{1}^{k} \mathbf{P}_{1} + \lambda_{2}^{k} \mathbf{P}_{2} + \dots + \lambda_{n}^{k} \mathbf{P}_{n}$$
(15)

for any positive integer k. We will see in the proof of Theorem 2 that this single fact is all we need to express the matrix exponential e^{At} simply and explicitly in terms of the eigenvalues and projection matrices of the matrix A.

Remark 2 The relation in (15) holds also if $k = \frac{1}{2}$. For instance, if n = 2 and $\mathbf{B} = \sqrt{\lambda_1} \mathbf{P}_1 + \sqrt{\lambda_2} \mathbf{P}_2$, then the basic properties of projection matrices (Proposition 1) and (14) yield

$$\mathbf{B}^{2} = \left(\sqrt{\lambda_{1}}\mathbf{P}_{1} + \sqrt{\lambda_{2}}\mathbf{P}_{2}\right)\left(\sqrt{\lambda_{1}}\mathbf{P}_{1} + \sqrt{\lambda_{2}}\mathbf{P}_{2}\right)$$
$$= \lambda_{1}\mathbf{P}_{1}^{2} + \sqrt{\lambda_{1}\lambda_{2}}\mathbf{P}_{1}\mathbf{P}_{2} + \sqrt{\lambda_{2}\lambda_{1}}\mathbf{P}_{2}\mathbf{P}_{1} + \lambda_{2}\mathbf{P}_{2}^{2}$$
$$= \lambda_{1}\mathbf{P}_{1} + \lambda_{2}\mathbf{P}_{2} = \mathbf{A}.$$

It follows that if (for each i) $\sqrt{\lambda_i}$ denotes either square root of the eigenvalue λ_i of **A**, then

$$\sqrt{\mathbf{A}} = \sqrt{\lambda_1} \, \mathbf{P}_1 + \sqrt{\lambda_2} \, \mathbf{P}_2 \tag{16}$$

is a square root of the matrix A.

THEOREM 2 The Matrix Exponential e^{At}

If the $n \times n$ matrix **A** has distinct eigenvalues and has the spectral decomposition in (14), then

$$e^{\mathbf{A}t} = \sum_{i=1}^{n} e^{\lambda_i t} \mathbf{P}_i = e^{\lambda_1 t} \mathbf{P}_1 + e^{\lambda_2 t} \mathbf{P}_2 + \dots + e^{\lambda_n t} \mathbf{P}_n.$$
 (17)

Proof: By using (15) for k = 0, 1, 2, 3, ..., we find that

$$e^{\mathbf{A}t} = \sum_{k=0}^{\infty} \frac{\mathbf{A}^{k} t^{k}}{k!} = \sum_{k=0}^{\infty} \left(\lambda_{1}^{k} \mathbf{P}_{1} + \lambda_{2}^{k} \mathbf{P}_{2} + \dots + \lambda_{n}^{k} \mathbf{P}_{n} \right) \frac{t^{k}}{k!}$$
$$= \left(\sum_{k=0}^{\infty} \frac{\lambda_{1}^{k} t^{k}}{k!} \right) \mathbf{P}_{1} + \left(\sum_{k=0}^{\infty} \frac{\lambda_{2}^{k} t^{k}}{k!} \right) \mathbf{P}_{2} + \dots + \left(\sum_{k=0}^{\infty} \frac{\lambda_{n}^{k} t^{k}}{k!} \right) \mathbf{P}_{n}$$
$$= e^{\lambda_{1}t} \mathbf{P}_{1} + e^{\lambda_{2}t} \mathbf{P}_{2} + \dots + e^{\lambda_{n}t} \mathbf{P}_{n},$$

so

$$e^{\mathbf{A}t} = \sum_{i=1}^{n} e^{\lambda_i t} \mathbf{P}_i,$$
as desired.
$$\mathbf{A}_1 = \sum_{i=1}^{n} e^{\lambda_i t} \mathbf{P}_i,$$
as desired.
$$\mathbf{A}_1 = \frac{1}{\lambda_1 - \lambda_2} \quad \text{and} \quad \lambda_2, \text{ then Eqs. (7), (8), and (17) give first}$$

$$a_1 = \frac{1}{\lambda_1 - \lambda_2} \quad \text{and} \quad a_2 = \frac{1}{\lambda_2 - \lambda_1}, \quad (18)$$
then
$$\mathbf{P}_1 = \frac{\mathbf{A} - \lambda_2 \mathbf{I}}{\lambda_1 - \lambda_2} \quad \text{and} \quad \mathbf{P}_2 = \frac{\mathbf{A} - \lambda_1 \mathbf{I}}{\lambda_2 - \lambda_1}, \quad (19)$$
and finally
$$e^{\mathbf{A}t} = e^{\lambda_1 t} \mathbf{P}_1 + e^{\lambda_2 t} \mathbf{P}_2. \quad (20)$$
Thus the calculation of the matrix exponential $e^{\mathbf{A}t}$ using the projection matrices \mathbf{P}_1 and \mathbf{P}_2
reduces to a routine matter of numerical substitution.
$$\mathbf{P}_1 = \begin{bmatrix} 10 & 1\\ 3 & 8 \end{bmatrix}$$
has the characteristic polynomial
$$p(\lambda) = (10 - \lambda)(8 - \lambda) - 3 = \lambda^2 - 18\lambda + 77 = (\lambda - 7)(\lambda - 11),$$
so its eigenvalues are $\lambda_1 = 7$ and $\lambda_2 = 11$. Hence Eq. (19) gives
$$\mathbf{P}_1 = -\frac{1}{4} \begin{bmatrix} 10 - 11 & 1\\ 3 & 8 - 11 \end{bmatrix} = -\frac{1}{4} \begin{bmatrix} -1 & 1\\ 3 & -3 \end{bmatrix}$$

and

$$\mathbf{P}_2 = \frac{1}{4} \begin{bmatrix} 10-7 & 1\\ 3 & 8-7 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 3 & 1\\ 3 & 1 \end{bmatrix},$$

so Eq. (20) gives

$$e^{\mathbf{A}t} = -\frac{1}{4}e^{7t} \begin{bmatrix} -1 & 1\\ 3 & -3 \end{bmatrix} + \frac{1}{4}e^{11t} \begin{bmatrix} 3 & 1\\ 3 & 1 \end{bmatrix}$$
$$= \frac{1}{4} \begin{bmatrix} e^{7t} + 3e^{11t} & -e^{7t} + e^{11t}\\ -3e^{7t} + 3e^{11t} & 3e^{7t} + e^{11t} \end{bmatrix}.$$

Example 3 The 3 ×

The 3×3 matrix

$$\mathbf{A} = \begin{bmatrix} 3 & 0 & 0 \\ -4 & 6 & 2 \\ 16 & -15 & -5 \end{bmatrix}$$

has the characteristic polynomial

$$p(\lambda) = -|\mathbf{A} - \lambda \mathbf{I}|$$

= -(3 - \lambda)[(6 - \lambda)(-5 - \lambda) + 30]
= -(3 - \lambda)[\lambda^2 - \lambda]
$$p(\lambda) = \lambda(\lambda - 1)(\lambda - 3).$$

Hence the eigenvalues of A are $\lambda_1 = 0$, $\lambda_2 = 1$, and $\lambda_3 = 3$. To find the matrix exponential

$$e^{\mathbf{A}t} = e^{\lambda_1 t} \mathbf{P}_1 + e^{\lambda_2 t} \mathbf{P}_2 + e^{\lambda_3 t} \mathbf{P}_3, \qquad (21)$$

we need only calculate the three projections matrices P_1 , P_2 , and P_3 defined by Eq. (8). But first we need the coefficients

$$a_{1} = \frac{1}{(\lambda_{1} - \lambda_{2})(\lambda_{1} - \lambda_{3})} = \frac{1}{(0 - 1)(0 - 3)} = \frac{1}{3},$$

$$a_{2} = \frac{1}{(\lambda_{2} - \lambda_{1})(\lambda_{2} - \lambda_{3})} = \frac{1}{(1 - 0)(1 - 3)} = -\frac{1}{2}, \text{ and}$$

$$a_{3} = \frac{1}{(\lambda_{3} - \lambda_{1})(\lambda_{3} - \lambda_{2})} = \frac{1}{(3 - 0)(3 - 1)} = \frac{1}{6}$$

that are given by Eq. (7). Then

$$\mathbf{P}_{1} = a_{1}(\mathbf{A} - \lambda_{2}\mathbf{I})(\mathbf{A} - \lambda_{3}\mathbf{I})$$

$$= \frac{1}{3} \begin{bmatrix} 2 & 0 & 0 \\ -4 & 5 & 2 \\ 16 & -15 & -6 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ -4 & 3 & 2 \\ 16 & -15 & -8 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 4 & -5 & 2 \\ -12 & 15 & 6 \end{bmatrix},$$

$$\mathbf{P}_{2} = a_{2}(\mathbf{A} - \lambda_{1}\mathbf{I})(\mathbf{A} - \lambda_{3}\mathbf{I})$$

$$= -\frac{1}{2} \begin{bmatrix} 3 & 0 & 0 \\ -4 & 6 & 2 \\ 16 & -15 & -5 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ -4 & 3 & 2 \\ 16 & -15 & -8 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ -4 & 6 & 2 \\ 10 & -15 & -5 \end{bmatrix},$$

and

$$\mathbf{P}_{3} = a_{3}(\mathbf{A} - \lambda_{1}\mathbf{I})(\mathbf{A} - \lambda_{2}\mathbf{I})$$

= $\frac{1}{6}\begin{bmatrix} 3 & 0 & 0\\ -4 & 6 & 2\\ 16 & -15 & -5 \end{bmatrix}\begin{bmatrix} 2 & 0 & 0\\ -4 & 5 & 2\\ 16 & -15 & -6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0\\ 0 & 0 & 0\\ 2 & 0 & 0 \end{bmatrix}.$

Finally, (17) gives the desired matrix exponential

$$e^{\mathbf{A}t} = e^{0t} \begin{bmatrix} 0 & 0 & 0 \\ 4 & -5 & 2 \\ -12 & 15 & 6 \end{bmatrix} + e^{1t} \begin{bmatrix} 0 & 0 & 0 \\ -4 & 6 & 2 \\ 10 & -15 & -5 \end{bmatrix} + e^{3t} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 2 & 0 & 0 \end{bmatrix}$$
$$= \begin{bmatrix} e^{3t} & 0 & 0 \\ 4 - 4e^t & -5 + 6e^t & -2 + 2e^t \\ -12 + 10e^t + 2e^{3t} & 15 - 15e^t & 6 - 5e^t \end{bmatrix}.$$

As an application, the solution of the initial value problem

$$\mathbf{x}'(t) = \mathbf{A}\mathbf{x}, \quad \mathbf{x}(0) = \mathbf{x}_0 = \begin{bmatrix} 3 & 7 & 11 \end{bmatrix}^T$$

is (by Theorem 2 in Section 8.1) given by

$$\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{x}_{0} = \begin{bmatrix} e^{3t} & 0 & 0 \\ 4 - 4e^{t} & -5 + 6e^{t} & -2 + 2e^{t} \\ -12 + 10e^{t} + 2e^{3t} & 15 - 15e^{t} & 6 - 5e^{t} \end{bmatrix} \begin{bmatrix} 3 \\ 7 \\ 11 \end{bmatrix}$$
$$= \begin{bmatrix} 3e^{3t} \\ -45 + 52e^{t} \\ 135 - 130e^{t} + 6e^{3t} \end{bmatrix}.$$

Second-Order Linear Systems

Spectral decompositions of matrices also yield convenient solutions of second-order matrix linear systems.

THEOREM 3 The Second-Order System x'' = Ax

If $\sqrt{\mathbf{A}}$ is a square root of the $n \times n$ constant coefficient matrix \mathbf{A} , then a general solution of the system $\mathbf{x}''(t) = \mathbf{A}\mathbf{x}$ is given by

$$\mathbf{x}(t) = e^{\sqrt{\mathbf{A}}t}\mathbf{c}_1 + e^{-\sqrt{\mathbf{A}}t}\mathbf{c}_2,$$
(22)

with \mathbf{c}_1 and \mathbf{c}_2 being arbitrary constant column vectors.

Proof: We need only note that, if \mathbf{x} is given by (22), then

$$\mathbf{x}'(t) = \sqrt{\mathbf{A}} e^{\sqrt{\mathbf{A}}t} \mathbf{c}_1 - \sqrt{\mathbf{A}} e^{-\sqrt{\mathbf{A}}t} \mathbf{c}_2,$$

so that

$$\mathbf{x}''(t) = \sqrt{\mathbf{A}} \cdot \sqrt{\mathbf{A}} e^{\sqrt{\mathbf{A}}t} \mathbf{c}_1 - (-\sqrt{\mathbf{A}}) \cdot \sqrt{\mathbf{A}} e^{-\sqrt{\mathbf{A}}t} \mathbf{c}_2$$
$$= \mathbf{A} \left(e^{\sqrt{\mathbf{A}}t} \mathbf{c}_1 + e^{-\sqrt{\mathbf{A}}t} \mathbf{c}_2 \right) = \mathbf{A}\mathbf{x}.$$

Example 4





FIGURE 8.3.1. The mass-and-spring system of Example 4.

Mass-spring system Consider the mass-and-spring system shown in Fig. 8.3.1, with $m_1 = 2$, $m_2 = 1$, $k_1 = 100$, and $k_2 = 50$. As in Example 1 of Section 7.4, the position vector $\mathbf{x}(t) = \begin{bmatrix} x_1(t) & x_2(t) \end{bmatrix}^T$ satisfies the second-order system $\mathbf{x}'' = \mathbf{A}\mathbf{x}$ having coefficient matrix

$$\mathbf{A} = \begin{bmatrix} -75 & 25\\ 50 & -50 \end{bmatrix}$$

We find that **A** has characteristic polynomial $p(\lambda) = \lambda^2 + 125\lambda + 2500 = (\lambda + 25)(\lambda + 100)$ and hence has eigenvalues $\lambda_1 = -25$ and $\lambda_2 = -100$. By using (19), we calculate the projection matrices

$$\mathbf{P}_{1} = \frac{\mathbf{A} - \lambda_{2}\mathbf{I}}{\lambda_{1} - \lambda_{2}} = \frac{1}{75} \begin{bmatrix} 25 & 25\\ 50 & 50 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 & 1\\ 2 & 2 \end{bmatrix} \text{ and}$$
$$\mathbf{P}_{2} = \frac{\mathbf{A} - \lambda_{1}\mathbf{I}}{\lambda_{2} - \lambda_{1}} = -\frac{1}{75} \begin{bmatrix} -50 & 25\\ 50 & -25 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 2 & -1\\ -2 & 1 \end{bmatrix}$$

of A. Starting with the spectral decomposition

$$\sqrt{\mathbf{A}} = \sqrt{\lambda_1} \mathbf{P}_1 + \sqrt{\lambda_2} \mathbf{P}_2 = 5i \mathbf{P}_1 + 10i \mathbf{P}_2$$

of $\sqrt{\mathbf{A}}$, Theorem 2 yields

$$e^{\sqrt{\mathbf{A}}t} = e^{5it}\mathbf{P}_1 + e^{10it}\mathbf{P}_2$$
 and $e^{-\sqrt{\mathbf{A}}t} = e^{-5it}\mathbf{P}_1 + e^{-10it}\mathbf{P}_2$.

8.3 SPECTRAL DECOMPOSITION METHODS 497

Consequently, Eq. (22) in Theorem 3 gives the following general solution:

$$\mathbf{x}(t) = e^{\sqrt{\mathbf{A}}t}\mathbf{c}_{1} + e^{-\sqrt{\mathbf{A}}t}\mathbf{c}_{2} = \left(e^{5it}\mathbf{P}_{1} + e^{10it}\mathbf{P}_{2}\right)\mathbf{c}_{1} + \left(e^{-5it}\mathbf{P}_{1} + e^{-10it}\mathbf{P}_{2}\right)\mathbf{c}_{2}$$

= $\mathbf{P}_{1}\left(\mathbf{c}_{1}e^{5it} + \mathbf{c}_{2}e^{-5it}\right) + \mathbf{P}_{2}\left(\mathbf{c}_{1}e^{10it} + \mathbf{c}_{2}e^{-10it}\right)$
 $\mathbf{x}(t) = \mathbf{P}_{1}(\mathbf{a}\cos 5t + \mathbf{b}\sin 5t) + \mathbf{P}_{2}(\mathbf{a}\cos 10t + \mathbf{b}\sin 10t),$ (23)

where $\mathbf{a} = \mathbf{c}_1 + \mathbf{c}_2$ and $\mathbf{b} = i(\mathbf{c}_1 - \mathbf{c}_2)$. Now

$$\mathbf{P}_{1}\mathbf{a} = \frac{1}{3} \begin{bmatrix} 1 & 1\\ 2 & 2 \end{bmatrix} \begin{bmatrix} a_{1}\\ a_{2} \end{bmatrix} = \frac{1}{3}(a_{1} + a_{2}) \begin{bmatrix} 1\\ 2 \end{bmatrix} \text{ and } \mathbf{P}_{1}\mathbf{b} = \frac{1}{3}(b_{1} + b_{2}) \begin{bmatrix} 1\\ 2 \end{bmatrix},$$
$$\mathbf{P}_{2}\mathbf{a} = \frac{1}{3} \begin{bmatrix} 2 & -1\\ -2 & 1 \end{bmatrix} \begin{bmatrix} a_{1}\\ a_{2} \end{bmatrix} = \frac{1}{3}(2a_{1} - a_{2}) \begin{bmatrix} 1\\ -1 \end{bmatrix}$$

and

$$\mathbf{P}_2\mathbf{b} = \frac{1}{3}(2b_1 - b_2) \begin{bmatrix} 1\\ -1 \end{bmatrix}.$$

Hence (23) finally takes the form

$$\mathbf{x}(t) = (a\cos 5t + b\sin 5t) \begin{bmatrix} 1\\2 \end{bmatrix} + (c\cos 10t + d\sin 10t) \begin{bmatrix} 1\\-1 \end{bmatrix}.$$
 (24)

This last equation expresses the motion $\mathbf{x}(t)$ of the mass-spring system in Fig. 8.3.1 as a linear combination of two natural modes of free oscillation. In the first mode, the two masses move with frequency $\omega_1 = 5$ in the same direction and with the amplitude of m_2 being twice that of m_1 . In the second mode, the two masses move with frequency $\omega_2 = 10$ in opposite directions and with equal amplitudes of motion.

The General Case

Now we want to take into account the possibility of multiple eigenvalues. If the $n \times n$ matrix **A** has distinct eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_q$ having multiplicities m_1, m_2, \ldots, m_q (respectively), then the reciprocal $1/p(\lambda)$ has a partial-fraction decomposition of the form

$$\frac{1}{p(\lambda)} = \frac{a_1(\lambda)}{(\lambda - \lambda_1)^{m_1}} + \frac{a_2(\lambda)}{(\lambda - \lambda_2)^{m_2}} + \dots + \frac{a_q(\lambda)}{(\lambda - \lambda_q)^{m_q}},$$
(25)

where (for each i = 1, 2, ..., q) the numerator $a_i(\lambda)$ is a polynomial in λ of degree at most $m_i - 1$. Once these numerator polynomials have been found, we can define the projection matrices $\mathbf{P}_1, \mathbf{P}_2, ..., \mathbf{P}_q$ of **A** by the formula

$$\mathbf{P}_{i} = a_{i}(\mathbf{A})b_{i}(\mathbf{A}) = a_{i}(\mathbf{A})\prod_{j\neq i}(\mathbf{A}-\lambda_{j}\mathbf{I})^{m_{j}},$$
(26)

analogous to Eq. (8), with the polynomial

$$b_i(\lambda) = \frac{p(\lambda)}{(\lambda - \lambda_i)^{m_i}} = \prod_{j \neq i} (\lambda - \lambda_j)^{m_j}$$
(27)

being obtained from the characteristic polynomial $p(\lambda)$ upon deletion of the factor $(\lambda - \lambda_i)^{m_i}$ corresponding to the *i*th eigenvalue. It follows immediately from (27) that $b_i(\lambda_i) \neq 0$, whereas $b_j(\lambda_i) = 0$ if $i \neq j$ [in which case $b_j(\lambda_i)$ includes the factor $(\lambda_i - \lambda_i)^{m_i}$]. The following properties of the projection matrices $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_q$ are established in essentially the same way as in the proof of Proposition 1.

PROPOSITION 2 The Projection Matrices of A

If the $n \times n$ matrix **A** has *q* distinct eigenvalues, then its projection matrices **P**₁, **P**₂, ..., **P**_q defined by (26) satisfy the conditions

(i)
$$\mathbf{P}_1 + \mathbf{P}_2 + \dots + \mathbf{P}_q = \mathbf{I};$$
 (28)

- (ii) $\mathbf{P}_i^2 = \mathbf{P}_i$ for each $i = 1, 2, \dots, q$; (29)
- (iii) $\mathbf{P}_i \mathbf{P}_j = \mathbf{0} \text{ if } i \neq j.$ (30)

In the distinct-eigenvalue case we saw in part (iv) of Proposition 1 that $\mathbf{P}_i(\mathbf{A} - \lambda_i \mathbf{I}) = \mathbf{0}$ for each *i*. This may no longer be so if $m_i > 1$, so we define

$$\mathbf{N}_i = \mathbf{P}_i (\mathbf{A} - \lambda_i \mathbf{I}) \tag{31}$$

for each i = 1, 2, ..., q. Part (ii) of the following proposition asserts that these matrices are *nilpotent*. Note that $N_i = 0$ if $m_i = 1$ (so that λ_i is a nonrepeated eigenvalue).

PROPOSITION 3 The Nilpotent Matrices of A

If the $n \times n$ matrix **A** has *q* distinct eigenvalues, then the matrices $N_1, N_2, ..., N_q$ defined by (31) satisfy the conditions

- (i) $\mathbf{N}_i \mathbf{N}_j = \mathbf{0} \text{ if } i \neq j;$ (32)
- (ii) $\mathbf{N}_{i}^{m_{i}} = \mathbf{0}$ for each $i = 1, 2, \dots, q$. (33)

Proof: The product

$$\mathbf{N}_{i}\mathbf{N}_{j} = \mathbf{P}_{i}(\mathbf{A} - \lambda_{i}\mathbf{I}) \cdot \mathbf{P}_{j}(\mathbf{A} - \lambda_{j}\mathbf{I})$$

= $a_{i}(\mathbf{A})\prod_{r \neq i} (\mathbf{A} - \lambda_{r}\mathbf{I})^{m_{r}} \cdot (\mathbf{A} - \lambda_{i}\mathbf{I}) \cdot a_{j}(\mathbf{A})\prod_{s \neq j} (\mathbf{A} - \lambda_{s}\mathbf{I})^{m_{s}} \cdot (\mathbf{A} - \lambda_{j}\mathbf{I})$

evidently has the characteristic polynomial

$$p(\mathbf{A}) = \prod_{r=1}^{q} (\mathbf{A} - \lambda_r \mathbf{I})^{m_r} = \mathbf{0}$$

as a factor, so (32) follows from the Cayley-Hamilton theorem. To verify (33), we write

$$\begin{split} \mathbf{N}_{i}^{m_{i}} &= (\mathbf{P}_{i}(\mathbf{A} - \lambda_{i}\mathbf{I}))^{m_{i}} = \mathbf{P}_{i}^{m_{i}} (\mathbf{A} - \lambda_{i}\mathbf{I})^{m_{i}} \\ &= \mathbf{P}_{i} \cdot (\mathbf{A} - \lambda_{i}\mathbf{I})^{m_{i}} \\ &= a_{i}(\mathbf{A}) \prod_{j \neq i} (\mathbf{A} - \lambda_{j}\mathbf{I})^{m_{j}} \cdot (\mathbf{A} - \lambda_{i}\mathbf{I})^{m_{i}} \\ \mathbf{N}_{i}^{m_{i}} &= a_{i}(\mathbf{A}) p(\mathbf{A}) = \mathbf{0}, \end{split}$$

using the Cayley-Hamilton theorem once again.

The following theorem expresses the matrix \mathbf{A} in terms of its projection and nilpotent matrices in the general case (allowing multiple eigenvalues).

THEOREM 4 The Spectral Decomposition of A

If the $n \times n$ matrix **A** has distinct eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_q$, the projection matrices $\mathbf{P}_1, \mathbf{P}_2, \ldots, \mathbf{P}_q$ defined in (26), and the nilpotent matrices $\mathbf{N}_1, \mathbf{N}_2, \ldots, \mathbf{N}_q$ defined in (31), then

$$\mathbf{A} = \sum_{i=1}^{q} (\lambda_i \mathbf{P}_i + \mathbf{N}_i).$$
(34)

(Note that this spectral decomposition reduces to $\mathbf{A} = \sum_{i=1}^{i} \lambda_i \mathbf{P}_i$ in the distincteigenvalue case of Theorem 1, where each $\mathbf{N}_i = \mathbf{0}$.)

Proof: First, we multiply Eq. (28) by A,

$$\mathbf{A} = \mathbf{P}_1 \mathbf{A} + \mathbf{P}_2 \mathbf{A} + \dots + \mathbf{P}_q \mathbf{A}$$

= $\mathbf{P}_1 (\lambda_1 \mathbf{I} + \mathbf{A} - \lambda_1 \mathbf{I}) + \mathbf{P}_2 (\lambda_2 \mathbf{I} + \mathbf{A} - \lambda_2 \mathbf{I}) + \dots + \mathbf{P}_q (\lambda_q \mathbf{I} + \mathbf{A} - \lambda_q \mathbf{I}).$

Then we note from (31) that

$$\mathbf{P}_i(\lambda_i \mathbf{I} + \mathbf{A} - \lambda_i \mathbf{I}) = \lambda_i \mathbf{P}_i \mathbf{I} + \mathbf{P}_i (\mathbf{A} - \lambda_i \mathbf{I}) = \lambda_i \mathbf{P}_i + \mathbf{N}_i,$$

so (34) follows.

1

As in the distinct-eigenvalue case of Theorem 2, the spectral decomposition of Theorem 4 can be used to calculate the matrix exponential e^{At} in the general case.

THEOREM 5 The Matrix Exponential e^{At}

If the $n \times n$ matrix **A** has distinct eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_q$, the projection matrices $\mathbf{P}_1, \mathbf{P}_2, \ldots, \mathbf{P}_q$ defined in (26), and the nilpotent matrices $\mathbf{N}_1, \mathbf{N}_2, \ldots, \mathbf{N}_q$ defined in (31), then

$$e^{\mathbf{A}t} = \sum_{i=1}^{q} \left(e^{\lambda_i t} \mathbf{P}_i \sum_{k=0}^{m_i - 1} (\mathbf{A} - \lambda_i \mathbf{I})^k \frac{t^k}{k!} \right).$$
(35)

Proof: First, we apply the fact that
$$\sum_{i=1}^{q} \mathbf{P}_i = \mathbf{I}$$
 and write

$$e^{\mathbf{A}t} = \mathbf{I} e^{\mathbf{A}t} = \sum_{i=1}^{q} \mathbf{P}_{i} e^{\mathbf{A}t}.$$
(36)

Then, the typical term in (36) is

$$\mathbf{P}_{i} e^{\mathbf{A}t} = \mathbf{P}_{i} e^{(\lambda_{i}\mathbf{I}+\mathbf{A}-\lambda_{i}\mathbf{I})t} = e^{\lambda_{i}t} \mathbf{I} \mathbf{P}_{i} e^{(\mathbf{A}-\lambda_{i}\mathbf{I})t}$$
$$= e^{\lambda_{i}t} \mathbf{P}_{i} \left(\mathbf{I} + \sum_{k=1}^{\infty} (\mathbf{A}-\lambda_{i}\mathbf{I})^{k} \frac{t^{k}}{k!}\right)$$
$$= e^{\lambda_{i}t} \left(\mathbf{P}_{i} + \sum_{k=1}^{\infty} \mathbf{P}_{i} (\mathbf{A}-\lambda_{i}\mathbf{I})^{k} \frac{t^{k}}{k!}\right)$$

$$= e^{\lambda_i t} \left(\mathbf{P}_i + \sum_{k=1}^{\infty} \mathbf{P}_i^k (\mathbf{A} - \lambda_i \mathbf{I})^k \frac{t^k}{k!} \right) \quad (\text{because } \mathbf{P}_i^k = \mathbf{P}_i)$$

$$= e^{\lambda_i t} \left(\mathbf{P}_i + \sum_{k=1}^{\infty} \mathbf{N}_i^k \frac{t^k}{k!} \right) \quad (\text{because } \mathbf{N}_i = \mathbf{P}_i (\mathbf{A} - \lambda_i \mathbf{I}))$$

$$= e^{\lambda_i t} \left(\mathbf{P}_i + \sum_{k=1}^{m_i - 1} \mathbf{N}_i^k \frac{t^k}{k!} \right) \quad (\text{because } \mathbf{N}_i^{m_i} = \mathbf{0})$$

$$= e^{\lambda_i t} \left(\mathbf{P}_i + \sum_{k=1}^{m_i - 1} \mathbf{P}_i (\mathbf{A} - \lambda_i \mathbf{I})^k \frac{t^k}{k!} \right), \quad \text{so}$$

$$\mathbf{P}_i e^{\mathbf{A}t} = e^{\lambda_i t} \mathbf{P}_i \left(\mathbf{I} + \sum_{k=1}^{m_i - 1} (\mathbf{A} - \lambda_i \mathbf{I})^k \frac{t^k}{k!} \right).$$

Finally, we get (35) upon substitution of this last expression for $\mathbf{P}_i e^{\mathbf{A}t}$ in (36).

The complicated formula in (35) looks much simpler in low-dimensional special cases.

Europealo E	
Example 5	If A is a 2 × 2 matrix with a single eigenvalue λ_1 of multiplicity 2 and corresponding projection matrix P ₁ , then (35) reduces to $e^{At} = e^{\lambda_1 t} \mathbf{P}_1[\mathbf{I} + (\mathbf{A} - \lambda_1 \mathbf{I})t]$. In this case, $p(\lambda) = (\lambda - \lambda_1)^2$ so Eqs. (25)–(27) give $a_1(\lambda) = b_1(\lambda) = 1$, and hence P ₁ = I . It therefore
	follows that
	$e^{\mathbf{A}t} = e^{\lambda_1 t} [\mathbf{I} + (\mathbf{A} - \lambda_1 \mathbf{I})t].$
Example 6	If A is a 3×3 matrix with an eigenvalue λ_1 of multiplicity 1, an eigenvalue λ_2 of multiplicity 2, and corresponding projection matrices P ₁ and P ₂ , then (35) reduces to
	$e^{\mathbf{A}t} = e^{\lambda_1 t} \mathbf{P}_1 + e^{\lambda_2 t} \mathbf{P}_2 [\mathbf{I} + (\mathbf{A} - \lambda_2 \mathbf{I})t].$
Example 7	If A is a 5×5 matrix with an eigenvalue λ_1 of multiplicity 2 and an eigenvalue λ_2 of multiplicity 3, then (35) reduces to
	$e^{\mathbf{A}t} = e^{\lambda_1 t} \mathbf{P}_1 \Big[\mathbf{I} + (\mathbf{A} - \lambda_1 \mathbf{I}) t \Big] + e^{\lambda_2 t} \mathbf{P}_2 \Big[\mathbf{I} + (\mathbf{A} - \lambda_2 \mathbf{I}) t + \frac{1}{2} (\mathbf{A} - \lambda_2 \mathbf{I})^2 t^2 \Big].$
	In this case, the principal labor may consist of finding the characteristic polynomial and its partial-fractions decomposition (25), so that Eqs. (26)–(27) can be used to find the projection matrices \mathbf{P}_1 and \mathbf{P}_2 .
Example 8	The matrix
	$\mathbf{A} = \begin{bmatrix} 4 & -2 & 1 \\ 2 & 0 & 1 \\ 2 & -2 & 3 \end{bmatrix} $ (37)
	has the characteristic polynomial
	$p(\lambda) = - \mathbf{A} - \lambda \mathbf{I} = \lambda^3 - 7\lambda^2 + 16\lambda - 12 = (\lambda - 3)(\lambda - 2)^2$
	and thus has eigenvalues $\lambda_1 = 3$ of multiplicity 1 and $\lambda_2 = 2$ of multiplicity 2. The partial-fractions decomposition
	1 1 $1 - \lambda$

$$\frac{1}{(\lambda-3)(\lambda-2)^2} = \frac{1}{\lambda-3} + \frac{1-\lambda}{(\lambda-2)^2}$$

shows that $a_1(\lambda) = 1$ and $a_2 = 1 - \lambda$. Now $b_1(\lambda) = (\lambda - 2)^2$ and $b_2(\lambda) = \lambda - 3$, so (26) gives

$$\mathbf{P}_1 = a_1(\mathbf{A})b_1(\mathbf{A}) = (\mathbf{A} - 2\mathbf{I})^2 = \begin{bmatrix} 2 & -2 & 1 \\ 2 & -2 & 1 \\ 2 & -2 & 1 \end{bmatrix}^2 = \begin{bmatrix} 2 & -2 & 1 \\ 2 & -2 & 1 \\ 2 & -2 & 1 \end{bmatrix}$$

and

$$\mathbf{P}_{2} = a_{2}(\mathbf{A})b_{2}(\mathbf{A}) = (\mathbf{I} - \mathbf{A})(\mathbf{A} - 3\mathbf{I})$$
$$= \begin{bmatrix} -3 & 2 & -1 \\ -2 & 1 & -1 \\ -2 & 2 & -2 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 2 & -3 & 1 \\ 2 & -2 & 0 \end{bmatrix} = \begin{bmatrix} -1 & 2 & -1 \\ -2 & 3 & -1 \\ -2 & 2 & 0 \end{bmatrix}$$

The result of Example 6 therefore gives

$$\begin{split} e^{\mathbf{A}t} &= e^{\lambda_1 t} \mathbf{P}_1 + e^{\lambda_2 t} \mathbf{P}_2 \big[\mathbf{I} + (\mathbf{A} - \lambda_2 \mathbf{I}) t \big] \\ &= e^{3t} \begin{bmatrix} 2 & -2 & 1 \\ 2 & -2 & 1 \\ 2 & -2 & 1 \end{bmatrix} \\ &+ e^{2t} \begin{bmatrix} -1 & 2 & -1 \\ -2 & 3 & -1 \\ -2 & 2 & 0 \end{bmatrix} \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & -2 & 1 \\ 2 & -2 & 1 \\ 2 & -2 & 1 \end{bmatrix} t \right) \\ e^{\mathbf{A}t} &= \begin{bmatrix} 2e^{3t} - e^{2t} & -2e^{3t} + 2e^{2t} & e^{3t} - e^{2t} \\ 2e^{3t} - 2e^{2t} & -2e^{3t} + 3e^{2t} & e^{3t} - e^{2t} \\ 2e^{3t} - 2e^{2t} & -2e^{3t} + 2e^{2t} & e^{3t} \end{bmatrix}. \end{split}$$

For instance, suppose we want to solve the initial value problem

$$\mathbf{x}'(t) = \mathbf{A}\mathbf{x} + \mathbf{f}, \quad \mathbf{x}(0) = \mathbf{x}_0, \tag{38}$$

where **A** is the matrix in (37), $\mathbf{x}_0 = \begin{bmatrix} 29 & 37 & 43 \end{bmatrix}^T$, and $\mathbf{f} = \begin{bmatrix} 60 & 0 & 0 \end{bmatrix}^T$. In this case, the variation of parameters formula in Eq. (28) of Section 8.2 gives

$$e^{-\mathbf{A}t}\mathbf{x}(t) = \mathbf{x}_{0} + \int_{0}^{t} e^{-\mathbf{A}s}\mathbf{f} \, ds$$

$$= \begin{bmatrix} 29\\ 37\\ 43 \end{bmatrix}$$

$$+ \int_{0}^{t} \begin{bmatrix} 2e^{-3s} - e^{-2s} & -2e^{-3s} + 2e^{-2s} & e^{-3s} - e^{-2s} \\ 2e^{-3s} - 2e^{-2s} & -2e^{-3s} + 3e^{-2s} & e^{-3s} - e^{-2s} \\ 2e^{-3s} - 2e^{-2s} & -2e^{-3s} + 2e^{-2s} & e^{-3s} \end{bmatrix} \begin{bmatrix} 60\\ 0\\ 0 \end{bmatrix} ds$$

$$= \begin{bmatrix} 29\\ 37\\ 43 \end{bmatrix} + \int_{0}^{t} \begin{bmatrix} 120e^{-3s} - 60e^{-2s} \\ 120e^{-3s} - 120e^{-2s} \\ 120e^{-3s} - 120e^{-2s} \end{bmatrix} ds$$

$$e^{-\mathbf{A}t}\mathbf{x}(t) = \begin{bmatrix} 29\\ 37\\ 43 \end{bmatrix} + \begin{bmatrix} 10 - 40e^{-3t} + 30e^{-2t} \\ -20 - 40e^{-3t} + 60e^{-2t} \\ -20 - 40e^{-3t} + 60e^{-2t} \end{bmatrix} = \begin{bmatrix} 39 - 40e^{-3t} + 30e^{-2t} \\ 17 - 40e^{-3t} + 60e^{-2t} \\ 23 - 40e^{-3t} + 60e^{-2t} \end{bmatrix}.$$

Finally, multiplication by $e^{\mathbf{A}t}$ yields the solution

$$\mathbf{x}(t) = \begin{bmatrix} 2e^{3t} - e^{2t} & -2e^{3t} + 2e^{2t} & e^{3t} - e^{2t} \\ 2e^{3t} - 2e^{2t} & -2e^{3t} + 3e^{2t} & e^{3t} - e^{2t} \\ 2e^{3t} - 2e^{2t} & -2e^{3t} + 2e^{2t} & e^{3t} \end{bmatrix} \begin{bmatrix} 39 - 40e^{-3t} + 30e^{-2t} \\ 17 - 40e^{-3t} + 60e^{-2t} \\ 23 - 40e^{-3t} + 60e^{-2t} \end{bmatrix}$$
$$= \begin{bmatrix} -10 + 67e^{3t} - 28e^{2t} \\ 20 + 67e^{3t} - 50e^{2t} \\ 20 + 67e^{3t} - 44e^{2t} \end{bmatrix}$$

of the initial value problem in (38).

502 CHAPTER 8 MATRIX EXPONENTIAL METHODS

8.3 Problems

- **1–20.** Use projection matrices to find a fundamental matrix solution $\mathbf{x}(t) = e^{\mathbf{A}t}$ of each of the linear systems $\mathbf{x}' = \mathbf{A}\mathbf{x}$ given in Problems 1 through 20 of Section 7.3.
- **21–30.** Use projection matrices to find a fundamental matrix solution of each of the linear systems given in Problems 1 through 10 of Section 7.5.
- **31–40.** Use projection matrices, as in Example 8 of this section, to find the matrix exponentials and particular solutions desired in Problems 21 through 30 (respectively) of Section 8.2.

In each of Problems 41 through 46, use the spectral decomposition methods of this section to find a fundamental matrix solution $\mathbf{x}(t) = e^{\mathbf{A}t}$ for the linear system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ given in the problem.

41. Problem 23 in Section 7.5

- **42.** Problem 24 in Section 7.5
- **43.** Problem 25 in Section 7.5
- **44.** Problem 26 in Section 7.5
- **45.** Problem 29 in Section 7.5
- **46.** Problem 31 in Section 7.5

Use projection matrices as in Example 4 of this section to solve Problems 47 through 50. You can simplify matters by taking $\mathbf{c}_1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ and $\mathbf{c}_2 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$ to calculate a typical particular solution $\mathbf{x}(t) = e^{\sqrt{\mathbf{A}t}}\mathbf{c}_1 + e^{-\sqrt{\mathbf{A}t}}\mathbf{c}_2$ that exhibits the fundamental frequencies and modes of oscillation.

- **47.** Problem 2 in Section 7.4
- 48. Problem 4 in Section 7.4
- 49. Problem 5 in Section 7.4
- 50. Problem 7 in Section 7.4

Nonlinear Systems and Phenomena

9.1 Stability and the Phase Plane

A wide variety of natural phenomena are modeled by two-dimensional first-order systems of the form

$$\frac{dx}{dt} = F(x, y),$$

$$\frac{dy}{dt} = G(x, y)$$
(1)

in which the independent variable t does not appear explicitly. We usually think of the dependent variables x and y as position variables in the xy-plane and of t as a time variable. We will see that the absence of t on the right-hand sides in (1) makes the system easier to analyze and its solutions easier to visualize. Using the terminology of Section 2.2, such a system of differential equations in which the derivative values are independent (or "autonomous") of time t is often called an **autonomous system**.

We generally assume that the functions *F* and *G* are continuously differentiable in some region *R* of the *xy*-plane. Then according to the existence and uniqueness theorems of Appendix A, given t_0 and any point (x_0, y_0) of *R*, there is a *unique* solution x = x(t), y = y(t) of (1) that is defined on some open interval (a, b) containing t_0 and satisfies the initial conditions

$$x(t_0) = x_0, \quad y(t_0) = y_0.$$
 (2)

The equations x = x(t), y = y(t) then describe a parametrized solution curve in the phase plane. Any such solution curve is called a **trajectory** of the system in (1), and precisely one trajectory passes through each point of the region *R* (Problem 29). A **critical point** of the system in (1) is a point (x_* , y_*) such that

$$F(x_{\star}, y_{\star}) = G(x_{\star}, y_{\star}) = 0.$$
(3)

503

If (x_{\star}, y_{\star}) is a critical point of the system, then the constant-valued functions

$$x(t) \equiv x_{\star}, \quad y(t) \equiv y_{\star} \tag{4}$$

have derivatives $x'(t) \equiv 0$ and $y'(t) \equiv 0$, and therefore automatically satisfy the equations in (1). Such a constant-valued solution is called an **equilibrium solution** of the system. Note that the trajectory of the equilibrium solution in (4) consists of the single point (x_{\star}, y_{\star}) .

In some practical situations these very simple solutions and trajectories are the ones of greatest interest. For example, suppose that the system x' = F(x, y), y' = G(x, y) models two populations x(t) and y(t) of animals that cohabit the same environment, and perhaps compete for the same food or prey on one another; x(t)might denote the number of rabbits and y(t) the number of squirrels present at time t. Then a critical point (x_*, y_*) of the system specifies a *constant* population x_* of rabbits and a *constant* population y_* of squirrels that can coexist with one another in the environment. If (x_0, y_0) is *not* a critical point of the system, then it is *not* possible for constant populations of x_0 rabbits and y_0 squirrels to coexist; one or both must change with time.

Example 1 Find the critical points of the system

$$\frac{dx}{dt} = 14x - 2x^2 - xy,$$

$$\frac{dy}{dt} = 16y - 2y^2 - xy.$$
(5)

Solution When we look at the equations

$$14x - 2x^{2} - xy = x(14 - 2x - y) = 0,$$

$$16y - 2y^{2} - xy = y(16 - 2y - x) = 0$$

that a critical point (x, y) must satisfy, we see that either

$$x = 0$$
 or $14 - 2x - y = 0$, (6a)

and either

$$y = 0$$
 or $16 - 2y - x = 0.$ (6b)

If x = 0 and $y \neq 0$, then the second equation in (6b) gives y = 8. If y = 0 and $x \neq 0$, then the second equation in (6a) gives x = 7. If both x and y are nonzero, then we solve the simultaneous equations

$$2x + y = 14$$
, $x + 2y = 16$

for x = 4, y = 6. Thus the system in (5) has the four critical points (0,0), (0,8), (7,0), and (4,6). If x(t) and y(t) denote the number of rabbits and the number of squirrels, respectively, and if both populations are *constant*, it follows that the equations in (5) allow only three nontrivial possibilities: either no rabbits and 8 squirrels, or 7 rabbits and no squirrels, or 4 rabbits and 6 squirrels. In particular, the critical point (4, 6) describes the *only* possibility for the coexistence of constant nonzero populations of both species.

Phase Portraits

If the initial point (x_0, y_0) is not a critical point, then the corresponding trajectory is a curve in the xy-plane along which the point (x(t), y(t)) moves as t increases. It turns out that any trajectory not consisting of a single point is a nondegenerate curve with no self-intersections (Problem 30). We can exhibit qualitatively the behavior of solutions of the autonomous system in (1) by constructing a picture that shows its critical points together with a collection of typical solution curves or trajectories in the *xy*-plane. Such a picture is called a **phase portrait** (or **phase plane picture**) because it illustrates "phases" or *xy*-states of the system, and indicates how they change with time.

Another way of visualizing the system is to construct a **slope field** in the *xy*-phase plane by drawing typical line segments having slope

$$\frac{dy}{dx} = \frac{y'}{x'} = \frac{G(x, y)}{F(x, y)},$$

or a **direction field** by drawing typical vectors pointing the same direction at each point (x, y) as does the vector (F(x, y), G(x, y)). Such a vector field then indicates which direction along a trajectory to travel in order to "go with the flow" described by the system.

Remark It is worth emphasizing that if our system of differential equations were not autonomous, then its critical points, trajectories, and direction vectors would generally be changing with time. In this event, the concrete visualization afforded by a (fixed) phase portrait or direction field would not be available to us. Indeed, this is a principal reason why an introductory study of nonlinear systems concentrates on autonomous ones.

Figure 9.1.1 shows a direction field and phase portrait for the rabbit-squirrel system of Example 1. The direction field arrows indicate the direction of motion of the point (x(t), y(t)). We see that, given any positive initial numbers $x_0 \neq 4$ and $y_0 \neq 6$ of rabbits and squirrels, this point moves along a trajectory approaching the critical point (4, 6) as t increases.

Example 2 For the system

$$\begin{aligned} x' &= x - y, \\ y' &= 1 - x^2 \end{aligned} \tag{7}$$

we see from the first equation that x = y and from the second that $x = \pm 1$ at each critical point. Thus this system has the two critical points (-1, -1) and (1, 1). The direction field in Fig. 9.1.2 suggests that trajectories somehow "circulate" counterclockwise around the critical point (-1, -1), whereas it appears that some trajectories may approach, while others recede from, the critical point (1, 1). These observations are corroborated by the phase portrait in Fig. 9.1.3 for the system in (7).



FIGURE 9.1.2. Direction field for the system in Eq. (7).

FIGURE 9.1.3. Phase portrait for the system in Eq. (7).

Remark One could carelessly write the critical points in Example 2 as $(\pm 1, \pm 1)$ and then jump to the erroneous conclusion that the system in (7) has four rather than just two critical points. When feasible, a sure-fire way to determine the number of critical points of an



FIGURE 9.1.1. Direction field and phase portrait for the rabbit–squirrel system $x' = 14x - 2x^2 - xy$, $y' = 16y - 2y^2 - xy$ of Example 1.



FIGURE 9.1.4. The two critical points (-1, -1) and (+1, +1) in Example 2 as the intersection of the curves F(x, y) = x - y = 0 and $G(x, y) = 1 - x^2 = 0$.



FIGURE 9.1.5. Close-up view of Fig. 9.1.1 near the critical point (4, 6).

autonomous system is to plot the curves F(x, y) = 0 and G(x, y) = 0 and then note their intersections, each of which represents a critical point of the system. For instance, Fig. 9.1.4 shows the curve (line) F(x, y) = x - y = 0 and the pair of lines x = +1 and x = -1 that constitute the "curve" $G(x, y) = 1 - x^2 = 0$. The (only) *two* points of intersection (-1, -1) and (+1, +1) are then apparent.

Critical Point Behavior

The behavior of the trajectories near an isolated critical point of an autonomous system is of particular interest. Figure 9.1.5 is a close-up view of Fig. 9.1.1 near the critical point (4, 6), and similarly Figs. 9.1.6 and 9.1.7 are close-ups of Fig. 9.1.3 near the critical points (-1, -1) and (1, 1), respectively. We notice immediately that although the systems underlying these phase portraits are *non*linear, each of these three magnifications bears a striking resemblance to one of the cases in our "gallery" Fig. 7.4.16 of phase plane portraits for *linear* constant-coefficient systems. Indeed, the three figures strongly resemble a nodal sink, a spiral source, and a saddle point, respectively.



FIGURE 9.1.6. Close-up view of

Fig. 9.1.3 near the critical point

(-1, -1).



1.5



FIGURE 9.1.7. Close-up view of Fig. 9.1.3 near the critical point (1, 1).

These similarities are not a coincidence. Indeed, as we will explore in detail in the next section, the behavior of a nonlinear system near a critical point is *generally* similar to that of a corresponding linear constant-coefficient system near the origin. For this reason it is useful to extend the language of nodes, sinks, etc., introduced in Section 7.4 for linear constant-coefficient systems to the broader context of critical points of the two-dimensional system (1).

In general, the critical point (x_{\star}, y_{\star}) of the autonomous system in (1) is called a **node** provided that

- *Either* every trajectory approaches (x_{*}, y_{*}) as t → +∞ or every trajectory recedes from (x_{*}, y_{*}) as t → +∞, and
- Every trajectory is tangent at (x_{\star}, y_{\star}) to some straight line through the critical point.

As with linear constant-coefficient systems, a node is said to be **proper** provided that no two different pairs of "opposite" trajectories are tangent to the same straight line through the critical point. On the other hand, the critical point (4, 6) of the system in Eq. (5), shown in Figs. 9.1.1 and 9.1.5, is an **improper** node; as those figures suggest, virtually all of the trajectories approaching this critical point share a common tangent line at that point.

Likewise, a node is further called a **sink** if all trajectories *approach* the critical point, a **source** if all trajectories *recede* (or emanate) from it. Thus the critical point (4, 6) in Figs. 9.1.1 and 9.1.5 is a *nodal sink*, whereas the critical point (-1, -1) in Figs. 9.1.3 and 9.1.6 is a *source* (more specifically, a *spiral source*). The critical point (1, 1) in Figs. 9.1.3 and 9.1.7, on the other hand, is a **saddle point**.

Stability

>

A critical point (x_{\star}, y_{\star}) of the autonomous system in (1) is said to be *stable* provided that if the initial point (x_0, y_0) is sufficiently close to (x_{\star}, y_{\star}) , then (x(t), y(t)) remains close to (x_{\star}, y_{\star}) for all t > 0. In vector notation, with $\mathbf{x}(t) = (x(t), y(t))$, the distance between the initial point $\mathbf{x}_0 = (x_0, y_0)$ and the critical point $\mathbf{x}_{\star} = (x_{\star}, y_{\star})$ is

$$|\mathbf{x}_0 - \mathbf{x}_{\star}| = \sqrt{(x_0 - x_{\star})^2 + (y_0 - y_{\star})^2}$$

Thus the critical point \mathbf{x}_{\star} is **stable** provided that, for each $\epsilon > 0$, there exists $\delta > 0$ such that

$$|\mathbf{x}_0 - \mathbf{x}_{\star}| < \delta$$
 implies that $|\mathbf{x}(t) - \mathbf{x}_{\star}| < \epsilon$ (8)

for all t > 0. Note that the condition in (8) certainly holds if $\mathbf{x}(t) \to \mathbf{x}_{\star}$ as $t \to +\infty$, as in the case of a nodal sink such as the critical point (4, 6) in Figs. 9.1.1 and 9.1.5. Thus this nodal sink can also be described as a *stable node*.

The critical point (x_{\star}, y_{\star}) is called **unstable** if it is not stable. The two critical points at (-1, -1) and (1, 1) shown in Figs. 9.1.3, 9.1.6, and 9.1.7 are both unstable, because, loosely speaking, in neither of these cases can we guarantee that a trajectory will *remain* near the critical point simply by requiring the trajectory to *begin* near the critical point.

If (x_*, y_*) is a critical point, then the equilibrium solution $x(t) \equiv x_*, y(t) \equiv y_*$ is called **stable** or **unstable** depending on the nature of the critical point. In applications the stability of an equilibrium solution is often a crucial matter. For instance, suppose in Example 1 that x(t) and y(t) denote the rabbit and squirrel populations, respectively, *in hundreds*. We will see in Section 9.3 that the critical point (4, 6) in Fig. 9.1.1 is stable. It follows that if we begin with *close to* 400 rabbits and 600 squirrels—rather than exactly these equilibrium values—then for all future time there will remain close to 400 rabbits and close to 600 squirrels. Thus the practical consequence of stability is that slight changes (perhaps due to random births and deaths) in the equilibrium populations will not so upset the equilibrium as to result in large deviations from the equilibrium solutions.

It is possible for trajectories to remain near a stable critical point without approaching it, as Example 3 shows.

Example 3 Undamped mass-spring system Consider a mass *m* that oscillates without damping on a spring with Hooke's constant *k*, so that its position function x(t) satisfies the differential equation $x'' + \omega^2 x = 0$ (where $\omega^2 = k/m$). If we introduce the velocity y = dx/dt of the mass, we get the system

$$\frac{dx}{dt} = y,$$
(9)
$$\frac{dy}{dt} = -\omega^2 x$$

with general solution

$$x(t) = A\cos\omega t + B\sin\omega t, \qquad (10a)$$

$$y(t) = -A\omega\sin\omega t + B\omega\cos\omega t.$$
(10b)



FIGURE 9.1.8. Direction field and elliptical trajectories for the system x' = y, $y' = -\frac{1}{4}x$. The origin is a stable center.



FIGURE 9.1.9. If the initial point (x_0, y_0) lies within distance δ of the origin, then the point (x(t), y(t)) stays within distance ϵ of the origin.

With $C = \sqrt{A^2 + B^2}$, $A = C \cos \alpha$, and $B = C \sin \alpha$, we can rewrite the solution in (10) in the form

$$x(t) = C\cos(\omega t - \alpha), \qquad (11a)$$

$$y(t) = -\omega C \sin(\omega t - \alpha),$$
 (11b)

so it becomes clear that each trajectory other than the critical point (0,0) is an ellipse with equation of the form

$$\frac{x^2}{C^2} + \frac{y^2}{\omega^2 C^2} = 1.$$
 (12)

As illustrated by the phase portrait in Fig. 9.1.8 (where $\omega = \frac{1}{2}$), each point (x_0, y_0) other than the origin in the *xy*-plane lies on exactly one of these ellipses, and each solution (x(t), y(t))traverses the ellipse through its initial point (x_0, y_0) in the clockwise direction with *period* $P = 2\pi/\omega$. (It is clear from (11) that x(t + P) = x(t) and y(t + P) = y(t) for all *t*.) Thus each nontrivial solution of the system in (9) is periodic and its trajectory is a simple closed curve enclosing the critical point at the origin.

Figure 9.1.9 shows a typical elliptical trajectory in Example 3, with its minor semiaxis denoted by δ and its major semiaxis by ϵ . We see that if the initial point (x_0, y_0) lies within distance δ of the origin—so that its elliptical trajectory lies inside the one shown—then the point (x(t), y(t)) always remains within distance ϵ of the origin. Hence the origin (0,0) is a stable critical point of the system x' = y, $y' = -\omega^2 x$, despite the fact that no single trajectory approaches the point (0,0). A stable critical point surrounded by simple closed trajectories representing periodic solutions is called a (stable) center.

Asymptotic Stability

The critical point (x_{\star}, y_{\star}) is called **asymptotically stable** if it is stable and, moreover, every trajectory that begins sufficiently close to (x_{\star}, y_{\star}) also approaches (x_{\star}, y_{\star}) as $t \to +\infty$. That is, there exists $\delta > 0$ such that

$$|\mathbf{x} - \mathbf{x}_{\star}| < \delta$$
 implies that $\lim_{t \to \infty} \mathbf{x}(t) = \mathbf{x}_{\star},$ (13)

where $\mathbf{x}_0 = (x_0, y_0)$, $\mathbf{x}_{\star} = (x_{\star}, y_{\star})$, and $\mathbf{x}(t) = (x(t), y(t))$ is a solution with $\mathbf{x}(0) = \mathbf{x}_0$.

Remark The stable node shown in Figs. 9.1.1 and 9.1.5 is asymptotically stable because every trajectory approaches the critical point (4, 6) as $t \to +\infty$. The center (0, 0) shown in Fig. 9.1.8 is stable but not asymptotically stable, because however small an elliptical trajectory we consider, a point moving around this ellipse does not approach the origin. Thus asymptotic stability is a stronger condition than mere stability.

Now suppose that x(t) and y(t) denote coexisting populations for which (x_{\star}, y_{\star}) is an asymptotically stable critical point. Then if the initial populations x_0 and y_0 are sufficiently close to x_{\star} and y_{\star} , respectively, it follows that both

$$\lim_{t \to \infty} x(t) = x_{\star} \quad \text{and} \quad \lim_{t \to \infty} y(t) = y_{\star}. \tag{14}$$

That is, x(t) and y(t) actually approach the equilibrium populations x_* and y_* as $t \to +\infty$, rather than merely remaining close to those values.

For a mechanical system as in Example 3, a critical point represents an *equilibrium state* of the system—if the velocity y = x' and the acceleration y' = x'' vanish simultaneously, then the mass remains at rest with no net force acting on it. Stability of a critical point concerns the question whether, when the mass is displaced slightly from its equilibrium, it

- 1. Moves back toward the equilibrium point as $t \to +\infty$,
- 2. Merely remains near the equilibrium point without approaching it, or
- **3.** Moves farther away from equilibrium.

In Case 1 the critical [equilibrium] point is asymptotically stable; in Case 2 it is stable but not asymptotically so; in Case 3 it is an unstable critical point. A marble balanced on the top of a soccer ball is an example of an unstable critical point. A mass on a spring with damping illustrates the case of asymptotic stability of a mechanical system. The mass-and-spring without damping in Example 3 is an example of a system that is stable but not asymptotically stable.

Damped mass-spring system Suppose that m = 1 and k = 2 for the mass and spring of Example 3 and that the mass is attached also to a dashpot with damping constant c = 2. Then its displacement function x(t) satisfies the second-order equation

0 0

$$x''(t) + 2x'(t) + 2x(t) = 0.$$
(15)

With y = x' we obtain the equivalent first-order system

$$\frac{dx}{dt} = y,$$
(16)
$$\frac{dy}{dt} = -2x - 2y$$

with critical point (0, 0). The characteristic equation $r^2 + 2r + 2 = 0$ of Eq. (15) has roots -1 + i and -1 - i, so the general solution of the system in (16) is given by

$$x(t) = e^{-t} (A\cos t + B\sin t) = Ce^{-t}\cos(t - \alpha),$$
(17a)

$$y(t) = e^{-t} \left[(B - A) \cos t - (A + B) \sin t \right]$$
(17b)

$$= -C\sqrt{2}e^{-t}\sin\left(t - \alpha + \frac{1}{4}\pi\right),\tag{17b}$$

where $C = \sqrt{A^2 + B^2}$ and $\alpha = \tan^{-1}(B/A)$. We see that x(t) and y(t) oscillate between positive and negative values and that both approach zero as $t \to +\infty$. Thus a typical trajectory spirals inward toward the origin, as illustrated by the spiral in Fig. 9.1.10.

It is clear from (17) that the point (x(t), y(t)) approaches the origin as $t \rightarrow t$ $+\infty$, so it follows that (0,0) is an asymptotically stable critical point for the system x' = y, y' = -2x - 2y of Example 4. Such an asymptotically stable critical point around which the trajectories spiral as they approach it—is called a stable spiral **point** (or a **spiral sink**). In the case of a mass–spring–dashpot system, a spiral sink is the manifestation in the phase plane of the damped oscillations that occur because of resistance.

If the arrows in Fig. 9.1.10 were reversed, we would see a trajectory spiraling *outward* from the origin. An unstable critical point—around which the trajectories spiral as they emanate and recede from it—is called an unstable spiral point (or a **spiral source**). Example 5 shows that it also is possible for a trajectory to spiral into a **closed trajectory**—a simple closed solution curve that represents a periodic solution (like the elliptical trajectories in Fig. 9.1.8).

Example 5 Consider the system

$$\frac{dx}{dt} = -ky + x(1 - x^2 - y^2),$$
(18)
$$\frac{dy}{dt} = kx + y(1 - x^2 - y^2).$$



FIGURE 9.1.10. A stable spiral point and one nearby trajectory.

In Problem 21 we ask you to show that (0, 0) is its only critical point. This system can be solved explicitly by introducing polar coordinates $x = r \cos \theta$, $y = r \sin \theta$, as follows. First note that

$$\frac{d\theta}{dt} = \frac{d}{dt} \left(\arctan \frac{y}{x} \right) = \frac{xy' - x'y}{x^2 + y^2}$$

Then substitute the expressions given in (18) for x' and y' to obtain

$$\frac{d\theta}{dt} = \frac{k(x^2 + y^2)}{x^2 + y^2} = k$$

It follows that

$$\theta(t) = kt + \theta_0, \quad \text{where } \theta_0 = \theta(0).$$
 (19)

Then differentiation of $r^2 = x^2 + y^2$ yields

$$2r\frac{dr}{dt} = 2x\frac{dx}{dt} + 2y\frac{dy}{dt}$$
$$= 2(x^2 + y^2)(1 - x^2 - y^2) = 2r^2(1 - r^2),$$

so r = r(t) satisfies the differential equation

$$\frac{dr}{dt} = r(1 - r^2). \tag{20}$$

In Problem 22 we ask you to derive the solution

$$r(t) = \frac{r_0}{\sqrt{r_0^2 + (1 - r_0^2)e^{-2t}}},$$
(21)

where $r_0 = r(0)$. Thus the typical solution of Eq. (18) may be expressed in the form

$$x(t) = r(t)\cos(kt + \theta_0),$$

$$y(t) = r(t)\sin(kt + \theta_0).$$
(22)

If $r_0 = 1$, then Eq. (21) gives $r(t) \equiv 1$ (the unit circle). Otherwise, if $r_0 > 0$, then Eq. (21) implies that $r(t) \rightarrow 1$ as $t \rightarrow +\infty$. Hence the trajectory defined in (22) spirals in toward the unit circle if $r_0 > 1$ and spirals out toward this closed trajectory if $0 < r_0 < 1$. Figure 9.1.11 shows a trajectory spiraling outward from the origin and four trajectories spiraling inward, all approaching the closed trajectory $r(t) \equiv 1$.

Under rather general hypotheses it can be shown that there are four possibilities for a nondegenerate trajectory of the autonomous system

$$\frac{dx}{dt} = F(x, y), \quad \frac{dy}{dt} = G(x, y)$$

The four possibilities are these:

- **1.** (x(t), y(t)) approaches a critical point as $t \to +\infty$.
- **2.** (x(t), y(t)) is unbounded with increasing t.
- **3.** (x(t), y(t)) is a periodic solution with a closed trajectory.
- **4.** (x(t), y(t)) spirals toward a closed trajectory as $t \to +\infty$.

As a consequence, the qualitative nature of the phase plane picture of the trajectories of an autonomous system is determined largely by the locations of its critical points and by the behavior of its trajectories near its critical points. We will see in Section 9.2 that, subject to mild restrictions on the functions F and G, each isolated critical point of the system x' = F(x, y), y' = G(x, y) resembles qualitatively one of the examples of this section—it is either a node (proper or improper), a saddle point, a center, or a spiral point.



FIGURE 9.1.11. Spiral trajectories of the system in Eq. (18) with k = 5.

9.1 Problems

In Problems 1 through 8, find the critical point or points of the given autonomous system, and thereby match each system with its phase portrait among Figs. 9.1.12 through 9.1.19.

1. $\frac{dx}{dt} = 2x - y$, $\frac{dy}{dt} = x - 3y$ 2. $\frac{dx}{dt} = x - y$, $\frac{dy}{dt} = x + 3y - 4$



FIGURE 9.1.12. Spiral point (-2, 1) and saddle point (2, -1).



FIGURE 9.1.15. Spiral point (0, 0); saddle points (-2, -1) and (2, 1).



FIGURE 9.1.13. Spiral point (1, -1).



FIGURE 9.1.16. Node (1, 1).





FIGURE 9.1.14. Saddle point (0, 0).



FIGURE 9.1.17. Spiral point (-1, -1), saddle point (0, 0), and node (1, -1).



FIGURE 9.1.18. Spiral point $(-2, \frac{2}{3})$ and saddle point $(2, -\frac{2}{5})$.



FIGURE 9.1.19. Stable center (-1, 1).

7.
$$\frac{dx}{dt} = x - 2y$$
, $\frac{dy}{dt} = 4x - x^3$
8. $\frac{dx}{dt} = x - y - x^2 + xy$, $\frac{dy}{dt} = -y - x^2$

In Problems 9 through 12, find each equilibrium solution $x(t) \equiv x_0$ of the given second-order differential equation x'' + f(x, x') = 0. Use a computer system or graphing calculator to construct a phase portrait and direction field for the equivalent first-order system x' = y, y' = -f(x, y). Thereby ascertain whether the critical point $(x_0, 0)$ looks like a center, a saddle point, or a spiral point of this system.

9.
$$x'' + 4x - x^3 = 0$$

10. $x'' + 2x' + x + 4x^3 = 0$
11. $x'' + 3x' + 4\sin x = 0$
12. $x'' + (x^2 - 1)x' + x = 0$

Solve each of the linear systems in Problems 13 through 20 to determine whether the critical point (0,0) is stable, asymptotically stable, or unstable. Use a computer system or graphing calculator to construct a phase portrait and direction field for the given system. Thereby ascertain the stability or instability of each critical point, and identify it visually as a node, a saddle point, a center, or a spiral point.

13.
$$\frac{dx}{dt} = -2x, \quad \frac{dy}{dt} = -2y$$

14.
$$\frac{dx}{dt} = 2x, \quad \frac{dy}{dt} = -2y$$

15.
$$\frac{dx}{dt} = -2x, \quad \frac{dy}{dt} = -y$$

16.
$$\frac{dx}{dt} = x$$
, $\frac{dy}{dt} = 3y$
17. $\frac{dx}{dt} = y$, $\frac{dy}{dt} = -x$

18.
$$\frac{dx}{dt} = -y$$
, $\frac{dy}{dt} = 4x$
19. $\frac{dx}{dt} = 2y$, $\frac{dy}{dt} = -2x$

$$20. \quad \frac{dx}{dt} = y, \quad \frac{dy}{dt} = -5x - 4y$$

- **21.** Verify that (0, 0) is the only critical point of the system in Example 6.
- **22.** Separate variables in Eq. (20) to derive the solution in (21).

In Problems 23 through 26, a system dx/dt = F(x, y), dy/dt = G(x, y) is given. Solve the equation

$$\frac{dy}{dx} = \frac{G(x, y)}{F(x, y)}$$

to find the trajectories of the given system. Use a computer system or graphing calculator to construct a phase portrait and direction field for the system, and thereby identify visually the apparent character and stability of the critical point (0,0) of the given system.

23.
$$\frac{dx}{dt} = y$$
, $\frac{dy}{dt} = -x$
24. $\frac{dx}{dt} = y(1 + x^2 + y^2)$, $\frac{dy}{dt} = x(1 + x^2 + y^2)$

25.
$$\frac{dx}{dt} = 4y(1 + x^2 + y^2), \quad \frac{dy}{dt} = -x(1 + x^2 + y^2)$$

26. $\frac{dx}{dt} = y^3 e^{x+y}, \quad \frac{dy}{dt} = -x^3 e^{x+y}$

27. Let (x(t), y(t)) be a nontrivial solution of the *non*autonomous system

$$\frac{dx}{dt} = y, \quad \frac{dy}{dt} = tx.$$

Suppose that $\phi(t) = x(t + \gamma)$ and $\psi(t) = y(t + \gamma)$, where $\gamma \neq 0$. Show that $(\phi(t), \psi(t))$ is *not* a solution of the system.

Problems 28 through 30 deal with the system

$$\frac{dx}{dt} = F(x, y), \quad \frac{dy}{dt} = G(x, y)$$

in a region where the functions F and G are continuously differentiable, so for each number a and point (x_0, y_0) , there is a unique solution with $x(a) = x_0$ and $y(a) = y_0$.

- **28.** Suppose that (x(t), y(t)) is a solution of the autonomous system and that $\gamma \neq 0$. Define $\phi(t) = x(t + \gamma)$ and $\psi(t) = y(t + \gamma)$. Then show (in contrast with the situation in Problem 27) that $(\phi(t), \psi(t))$ is also a solution of the system. Thus autonomous systems have the simple but important property that a "*t*-translate" of a solution is again a solution.
- **29.** Let $(x_1(t), y_1(t))$ and $(x_2(t), y_2(t))$ be two solutions having trajectories that meet at the point (x_0, y_0) ; thus $x_1(a) = x_2(b) = x_0$ and $y_1(a) = y_2(b) = y_0$ for some values *a* and *b* of *t*. Define

$$x_3(t) = x_2(t + \gamma)$$
 and $y_3(t) = y_2(t + \gamma)$,

where $\gamma = b - a$, so $(x_2(t), y_2(t))$ and $(x_3(t), y_3(t))$ have the same trajectory. Apply the uniqueness theorem to show that $(x_1(t), y_1(t))$ and $(x_3(t), y_3(t))$ are identical solutions. Hence the original two trajectories are identical. Thus no two different trajectories of an autonomous system can intersect.

30. Suppose that the solution $(x_1(t), y_1(t))$ is defined for all *t* and that its trajectory has an apparent self-intersection:

$$x_1(a) = x_1(a + P) = x_0, \quad y_1(a) = y_1(a + P) = y_0$$

for some P > 0. Introduce the solution

$$x_2(t) = x_1(t+P), \quad y_2(t) = y_1(t+P),$$

and then apply the uniqueness theorem to show that

$$x_1(t+P) = x_1(t)$$
 and $y_1(t) = y_1(t+P)$

for *all* t. Thus the solution $(x_1(t), y_1(t))$ is periodic with period P and has a closed trajectory. Consequently a solution of an autonomous system either is periodic with a closed trajectory, or else its trajectory never passes through the same point twice.

9.1 Application Phase Plane Portraits and First-Order Equations

Consider a first-order differential equation of the form

$$\frac{dy}{dx} = \frac{G(x, y)}{F(x, y)},\tag{1}$$

which may be difficult or impossible to solve explicitly. Its solution curves can nevertheless be plotted as trajectories of the corresponding autonomous two-dimensional system

$$\frac{dx}{dt} = F(x, y), \quad \frac{dy}{dt} = G(x, y).$$
(2)

Most ODE plotters can routinely generate phase portraits for autonomous systems. Those appearing in this chapter were plotted using programs that are free for educational use. For instance, the MATLAB program **pplane** illustrated in Fig. 9.1.20 can be found at **math.rice.edu/~dfield**. Another freely available and userfriendly MATLAB-based ODE package with similar graphical capabilities is **Iode** (www.math.uiuc.edu/iode).



FIGURE 9.1.20. MATLAB **pplane** menu entries to plot a direction field and phase portrait for the system x' = y, $y' = -\frac{1}{4}x$ (as shown in Fig. 9.1.8).

For example, to plot solution curves for the differential equation

$$\frac{dy}{dx} = \frac{2xy - y^2}{x^2 - 2xy},$$
(3)

we plot trajectories of the system

$$\frac{dx}{dt} = x^2 - 2xy, \quad \frac{dy}{dt} = 2xy - y^2.$$
 (4)



FIGURE 9.1.21. Phase portrait for the system in Eq. (4).

The result is shown in Fig. 9.1.21.

Plot similarly some solution curves for the following differential equations.

1.
$$\frac{dy}{dx} = \frac{4x - 5y}{2x + 3y}$$

2.
$$\frac{dy}{dx} = \frac{4x - 5y}{2x - 3y}$$

3.
$$\frac{dy}{dx} = \frac{4x - 3y}{2x - 5y}$$

4.
$$\frac{dy}{dx} = \frac{2xy}{x^2 - y^2}$$

5.
$$\frac{dy}{dx} = \frac{x^2 + 2xy}{y^2 + 2xy}$$

Now construct some examples of your own. Homogeneous functions like those in Problems 1 through 5—rational functions with numerator and denominator of the same degree in x and y—work well. The differential equation

$$\frac{dy}{dx} = \frac{25x + y(1 - x^2 - y^2)(4 - x^2 - y^2)}{-25y + x(1 - x^2 - y^2)(4 - x^2 - y^2)}$$
(5)

of this form generalizes Example 5 in this section but would be inconvenient to solve explicitly. Its phase portrait (Fig. 9.1.22) shows two periodic closed trajectories—the circles r = 1 and r = 2. Anyone want to try for three circles?

9.2 Linear and Almost Linear Systems

>

We now discuss the behavior of solutions of the autonomous system

$$\frac{dx}{dt} = f(x, y), \qquad \frac{dy}{dt} = g(x, y) \tag{1}$$

near an isolated critical point (x_0, y_0) where $f(x_0, y_0) = g(x_0, y_0) = 0$. A critical point is called **isolated** if some neighborhood of it contains no other critical point. We assume throughout that the functions f and g are continuously differentiable in a neighborhood of (x_0, y_0) .

We can assume without loss of generality that $x_0 = y_0 = 0$. Otherwise, we make the substitutions $u = x - x_0$, $v = y - y_0$. Then dx/dt = du/dt and dy/dt = dv/dt, so (1) is equivalent to the system

$$\frac{du}{dt} = f(u + x_0, v + y_0) = f_1(u, v),$$

$$\frac{dv}{dt} = g(u + x_0, v + y_0) = g_1(u, v)$$
(2)

that has (0, 0) as an isolated critical point.

The system

$$\frac{dx}{dt} = 3x - x^2 - xy = x(3 - x - y),$$

$$\frac{dy}{dt} = y + y^2 - 3xy = y(1 - 3x + y)$$
(3)



FIGURE 9.1.22. Phase portrait for the system corresponding to Eq. (5).

has (1, 2) as one of its critical points. We substitute u = x - 1, v = y - 2; that is, x = u + 1, y = v + 2. Then

$$3 - x - y = 3 - (u + 1) - (v + 2) = -u - v$$

and

$$-3x + y = 1 - 3(u + 1) + (v + 2) = -3u + v$$

so the system in (3) takes the form

1

$$\frac{du}{dt} = (u+1)(-u-v) = -u - v - u^2 - uv,$$

$$\frac{dv}{dt} = (v+2)(-3u+v) = -6u + 2v + v^2 - 3uv$$
(4)

and has (0, 0) as a critical point. If we can determine the trajectories of the system in (4) near (0, 0), then their translations under the rigid motion that carries (0, 0) to (1, 2) will be the trajectories near (1, 2) of the original system in (3). This equivalence is illustrated by Fig. 9.2.1 (which shows computer-plotted trajectories of the system in (3) near the critical point (1, 2) in the *xy*-plane) and Fig. 9.2.2 (which shows computer-plotted trajectories of the system in (4) near the critical point (0, 0) in the *uv*-plane).



FIGURE 9.2.1. The saddle point (1, 2) for the system $x' = 3x - x^2 - xy$, $y' = y + y^2 - 3xy$ of Example 1.

FIGURE 9.2.2. The saddle point (0, 0) for the equivalent system $u' = -u - v - u^2 - uv$, $v' = -6u + 2v + v^2 - 3uv$.

Figures 9.2.1 and 9.2.2 illustrate the fact that the solution curves of the *xy*-system in (1) are simply the images under the translation $(u, v) \rightarrow (u + x_0, v + y_0)$ of the solution curves of the *uv*-system in (2). Near the two corresponding critical points— (x_0, y_0) in the *xy*-plane and (0, 0) in the *uv*-plane—the two phase portraits therefore look precisely the same.

Linearization Near a Critical Point

Taylor's formula for functions of two variables implies that—if the function f(x, y) is continuously differentiable near the fixed point (x_0, y_0) —then

$$f(x_0 + u, y_0 + v) = f(x_0, y_0) + f_x(x_0, y_0)u + f_y(x_0, y_0)v + r(u, v)$$

where the "remainder term" r(u, v) satisfies the condition

$$\lim_{(u,v)\to(0,0)}\frac{r(u,v)}{\sqrt{u^2+v^2}}=0.$$

(Note that this condition would not be satisfied if r(u, v) were a sum containing either constants or terms linear in u or v. In this sense, r(u, v) consists of the "nonlinear part" of the function $f(x_0 + u, y_0 + v)$ of u and v.)

If we apply Taylor's formula to both f and g in (2) and assume that (x_0, y_0) is an isolated critical point so $f(x_0, y_0) = g(x_0, y_0) = 0$, the result is

$$\frac{du}{dt} = f_x(x_0, y_0)u + f_y(x_0, y_0)v + r(u, v),$$

$$\frac{dv}{dt} = g_x(x_0, y_0)u + g_y(x_0, y_0)v + s(u, v)$$
(5)

where r(u, v) and the analogous remainder term s(u, v) for g satisfy the condition

$$\lim_{(u,v)\to(0,0)} \frac{r(u,v)}{\sqrt{u^2+v^2}} = \lim_{(u,v)\to(0,0)} \frac{s(u,v)}{\sqrt{u^2+v^2}} = 0.$$
 (6)

Then, when the values u and v are small, the remainder terms r(u, v) and s(u, v) are *very* small (being small even in comparison with u and v).

If we drop the presumably small nonlinear terms r(u, v) and s(u, v) in (5), the result is the *linear* system

$$\frac{du}{dt} = f_x(x_0, y_0)u + f_y(x_0, y_0)v,$$

$$\frac{dv}{dt} = g_x(x_0, y_0)u + g_y(x_0, y_0)v$$
(7)

whose constant coefficients (of the variables u and v) are the values $f_x(x_0, y_0)$, $f_y(x_0, y_0)$ and $g_x(x_0, y_0)$, $g_y(x_0, y_0)$ of the functions f and g at the critical point (x_0, y_0) . Because (5) is equivalent to the original (and generally) nonlinear system $u' = f(x_0 + u, y_0 + v)$, $v' = g(x_0 + u, y_0 + v)$ in (2), the conditions in (6) suggest that the **linearized system** in (7) closely approximates the given nonlinear system when (u, v) is close to (0, 0).

Assuming that (0,0) is also an isolated critical point of the linear system, and that the remainder terms in (5) satisfy the condition in (6), the original system x' = f(x, y), y' = g(x, y) is said to be **almost linear** at the isolated critical point (x_0, y_0) . In this case, its **linearization** at (x_0, y_0) is the linear system in (7). In short, this linearization is the linear system $\mathbf{u}' = \mathbf{J}\mathbf{u}$ (where $\mathbf{u} = \begin{bmatrix} u & v \end{bmatrix}^T$) whose coefficient matrix is the so-called **Jacobian matrix**

$$\mathbf{J}(x_0, y_0) = \begin{bmatrix} f_x(x_0, y_0) & f_y(x_0, y_0) \\ g_x(x_0, y_0) & g_y(x_0, y_0) \end{bmatrix}$$
(8)

of the functions f and g, evaluated at the point (x_0, y_0) .

Example 1 Continued In (3) we have
$$f(x, y) = 3x - x^2 - xy$$
 and $g(x, y) = y + y^2 - 3xy$. Then

$$\mathbf{J}(x, y) = \begin{bmatrix} 3 - 2x - y & -x \\ -3y & 1 + 2y - 3x \end{bmatrix}, \text{ so } \mathbf{J}(1, 2) = \begin{bmatrix} -1 & -1 \\ -6 & 2 \end{bmatrix}.$$

Hence the linearization of the system $x' = 3x - x^2 - xy$, $y' = y + y^2 - 3xy$ at its critical point (1, 2) is the linear system

$$u' = -u - v,$$

$$v' = -6u + 2v$$

that we get when we drop the nonlinear (quadratic) terms in (4).

It turns out that in most (though not all) cases, the phase portrait of an almost linear system near an isolated critical point (x_0, y_0) strongly resembles qualitatively—the phase portrait of its linearization near the origin. Consequently, the first step toward understanding general autonomous systems is to characterize the isolated critical points of linear systems.

Isolated Critical Points of Linear Systems

In Section 7.4 we used the eigenvalue-eigenvector method to study the 2×2 linear system

with constant-coefficient matrix **A**. The origin (0, 0) is a critical point of the system regardless of the matrix **A**, but if we further require the origin to be an *isolated* critical point, then (by a standard theorem of linear algebra) the determinant ad - bc of **A** must be nonzero. From this we can conclude that *the eigenvalues* λ_1 and λ_2 of **A** must be nonzero. Indeed, λ_1 and λ_2 are the solutions of the characteristic equation

$$det(\mathbf{A} - \lambda \mathbf{I}) = \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix}$$
$$= (a - \lambda)(d - \lambda) - bc$$
$$= \lambda^2 - (a + d)\lambda + (ad - bc)$$
$$= 0,$$
(10)

and the fact that $ad - bc \neq 0$ implies that $\lambda = 0$ cannot satisfy Eq. (10); hence λ_1 and λ_2 are nonzero. The converse also holds: If the characteristic equation (10) has no zero solution—that is, if all eigenvalues of the matrix **A** are nonzero—then the determinant ad - bc is nonzero. Altogether, we conclude that the origin (0, 0) is an isolated critical point of the system in Eq. (9) if and only if the eigenvalues of **A** are all nonzero. Our study of this critical point can be divided, therefore, into the five cases listed in the table in Fig. 9.2.3. This table also gives the type of each critical point as found in Section 7.4 and shown in our gallery Fig. 7.4.16 of typical phase plane portraits:

Eigenvalues of A	Type of Critical Point
Real, unequal, same sign	Improper node
Real, unequal, opposite sign	Saddle point
Real and equal	Proper or improper node
Complex conjugate	Spiral point
Pure imaginary	Center

FIGURE 9.2.3. Classification of the isolated critical point (0, 0) of the two-dimensional system $\mathbf{x}' = A\mathbf{x}$.

Closer inspection of that gallery, however, reveals a striking connection between the *stability* properties of the critical point and the eigenvalues λ_1 and λ_2 of **A**. For example, if λ_1 and λ_2 are real, unequal, and negative, then the origin represents an improper nodal sink; because all trajectories approach the origin as $t \to +\infty$, the critical point is asymptotically stable. Likewise, if λ_1 and λ_2 are real, equal, and negative, then the origin is a proper nodal sink, and is again asymptotically stable. Further, if λ_1 and λ_2 are complex conjugate with negative real part, then the origin is a spiral sink, and is once more asymptotically stable. All three of these cases can be captured as follows: *If the real parts of* λ_1 *and* λ_2 *are negative, then the origin is an asymptotically stable critical point.* (Of course, if λ_1 and λ_2 are real, then they are themselves their real parts.)

Similar generalizations can be made for other combinations of signs of the real parts of λ_1 and λ_2 . Indeed, as the table in Fig. 9.2.4 shows, the stability properties

Real Parts of λ_1 and λ_2	Type of Critical Point	Stability
Both negative	 Proper or improper nodal sink, <i>or</i> Spiral sink 	Asymptotically stable
Both zero (i.e., λ_1 and λ_2 are given by $\pm iq$ with $q \neq 0$)	• Center	Stable but not asymptotically stable
At least one positive	 Proper or improper nodal source, <i>or</i> Spiral source, <i>or</i> Saddle point 	Unstable

FIGURE 9.2.4. Stability properties of the isolated critical point (0, 0) of the system in Eq. (9) with nonzero eigenvalues λ_1 and λ_2 .

of the isolated critical point (0, 0) of the system in Eq. (9) are always determined by the signs of the real parts of λ_1 and λ_2 . (We invite you to use the gallery in Fig. 7.4.16 to verify the conclusions in the table.)

These findings are summarized in Theorem 1:

THEOREM 1 Stability of Linear Systems

Let λ_1 and λ_2 be the eigenvalues of the coefficient matrix **A** of the twodimensional linear system

$$\frac{dx}{dt} = ax + by,$$

$$\frac{dy}{dt} = cx + dy$$
(11)

with $ad - bc \neq 0$. Then the critical point (0, 0) is

- 1. Asymptotically stable if the real parts of λ_1 and λ_2 are both negative;
- 2. Stable but not asymptotically stable if the real parts of λ_1 and λ_2 are both zero (so that $\lambda_1, \lambda_2 = \pm qi$);
- **3.** Unstable if either λ_1 or λ_2 has a positive real part.

 $\lambda_1 = qi$ $\circ \mu_1 = r + si$ $\lambda_2 = -qi$

FIGURE 9.2.5. The effects of perturbation of pure imaginary roots.

It is worthwhile to consider the effect of small perturbations in the coefficients a, b, c, and d of the linear system in (11), which result in small perturbations of the eigenvalues λ_1 and λ_2 . If these perturbations are sufficiently small, then positive real parts (of λ_1 and λ_2) remain positive and negative real parts remain negative. Hence an asymptotically stable critical point remains asymptotically stable and an unstable critical point remains unstable. Part 2 of Theorem 1 is therefore the only case in which arbitrarily small perturbations can affect the stability of the critical point (0,0). In this case pure imaginary roots $\lambda_1, \lambda_2 = \pm qi$ of the characteristic equation can be changed to nearby complex roots $\mu_1, \mu_2 = r \pm si$, with r either positive or negative (see Fig. 9.2.5). Consequently, a small perturbation of the coefficients of the linear system in (11) can change a stable center to a spiral point that is either unstable or asymptotically stable.

There is one other exceptional case in which the type, though not the stability, of the critical point (0,0) can be altered by a small perturbation of its coefficients.



FIGURE 9.2.6. The effects of perturbation of real equal roots.

This is the case with $\lambda_1 = \lambda_2$, equal roots that (under a small perturbation of the coefficients) can split into two roots μ_1 and μ_2 , which are either complex conjugates or unequal real roots (see Fig. 9.2.6). In either case, the sign of the real parts of the roots is preserved, so the stability of the critical point is unaltered. Its nature may change, however; the table in Fig. 9.2.3 shows that a node with $\lambda_1 = \lambda_2$ can either remain a node (if μ_1 and μ_2 are real) or change to a spiral point (if μ_1 and μ_2 are complex conjugates).

Suppose that the linear system in (11) is used to model a physical situation. It is unlikely that the coefficients in (11) can be measured with total accuracy, so let the unknown precise linear model be

$$\frac{dx}{dt} = a^* x + b^* y,$$

$$\frac{dy}{dt} = c^* x + d^* y.$$
(11*)

If the coefficients in (11) are sufficiently close to those in (11^{*}), it then follows from the discussion in the preceding paragraph that the origin (0, 0) is an asymptotically stable critical point for (11) if it is an asymptotically stable critical point for (11^{*}), and is an unstable critical point for (11) if it is an unstable critical point for (11^{*}). Thus in this case the approximate model in (11) and the precise model in (11^{*}) predict the same qualitative behavior (with respect to asymptotic stability versus instability).

Almost Linear Systems

Recall that we first encountered an almost linear system at the beginning of this section, when we used Taylor's formula to write the nonlinear system (2) in the almost linear form (5) which led to the linearization (7) of the original nonlinear system. In case the nonlinear system x' = f(x, y), y' = g(x, y) has (0, 0) as an isolated critical point, the corresponding almost linear system is

$$\frac{dx}{dt} = ax + by + r(x, y),$$

$$\frac{dy}{dt} = cx + dy + s(x, y),$$
(12)

where $a = f_x(0,0)$, $b = f_y(0,0)$ and $c = g_x(0,0)$, $d = g_y(0,0)$; we assume also that $ad - bc \neq 0$. Theorem 2, which we state without proof, essentially implies that—with regard to the type and stability of the critical point (0,0)—the effect of the small nonlinear terms r(x, y) and s(x, y) is equivalent to the effect of a small perturbation in the coefficients of the associated *linear* system in (11).

THEOREM 2 Stability of Almost Linear Systems

Let λ_1 and λ_2 be the eigenvalues of the coefficient matrix of the linear system in (11) associated with the almost linear system in (12). Then

- 1. If $\lambda_1 = \lambda_2$ are equal real eigenvalues, then the critical point (0, 0) of (12) is either a node or a spiral point, and is asymptotically stable if $\lambda_1 = \lambda_2 < 0$, unstable if $\lambda_1 = \lambda_2 > 0$.
- **2.** If λ_1 and λ_2 are pure imaginary, then (0,0) is either a center or a spiral point, and may be either asymptotically stable, stable, or unstable.

THEOREM 2 Stability of Almost Linear Systems (Continued)

3. Otherwise—that is, unless λ_1 and λ_2 are either real equal or pure imaginary—the critical point (0,0) of the almost linear system in (12) is of the same type and stability as the critical point (0,0) of the associated linear system in (11).

Thus, if $\lambda_1 \neq \lambda_2$ and $\operatorname{Re}(\lambda_1) \neq 0$, then the type and stability of the critical point of the almost linear system in (12) can be determined by analysis of its associated linear system in (11), and only in the case of pure imaginary eigenvalues is the stability of (0, 0) not determined by the linear system. Except in the sensitive cases $\lambda_1 = \lambda_2$ and $\operatorname{Re}(\lambda_i) = 0$, the trajectories near (0, 0) will resemble qualitatively those of the associated linear system—they enter or leave the critical point in the same way, but may be "deformed" in a nonlinear manner. The table in Fig. 9.2.7 summarizes the situation.

Eigenvalues λ_1 , λ_2 for the Linearized System	Type of Critical Point of the Almost Linear System
$\lambda_1 < \lambda_2 < 0$	Stable improper node
$\lambda_1 = \lambda_2 < 0$	Stable node or spiral point
$\lambda_1 < 0 < \lambda_2$	Unstable saddle point
$\lambda_1 = \lambda_2 > 0$	Unstable node or spiral point
$\lambda_1 > \lambda_2 > 0$	Unstable improper node
$\lambda_1, \lambda_2 = a \pm bi (a < 0)$	Stable spiral point
$\lambda_1, \lambda_2 = a \pm bi (a > 0)$	Unstable spiral point
$\lambda_1, \lambda_2 = \pm bi$	Stable or unstable, center or spiral point

FIGURE 9.2.7. Classification of critical points of an almost linear system.

An important consequence of the classification of cases in Theorem 2 is that a *critical point of an almost linear system is asymptotically stable if it is an asymptotically stable critical point of the linearization of the system.* Moreover, a critical point of the almost linear system is unstable if it is an unstable critical point of the linearized system. If an almost linear system is used to model a physical situation, then—apart from the sensitive cases mentioned earlier—it follows that the qualitative behavior of the system near a critical point can be determined by examining its linearization.

Example 2 Determine the type and stability of the critical point (0, 0) of the almost linear system

$$\frac{dx}{dt} = 4x + 2y + 2x^2 - 3y^2,$$
(13)

$$\frac{dy}{dt} = 4x - 3y + 7xy.$$

Solution

The characteristic equation for the associated linear system (obtained simply by deleting the quadratic terms in (13)) is

$$(4 - \lambda)(-3 - \lambda) - 8 = (\lambda - 5)(\lambda + 4) = 0,$$

so the eigenvalues $\lambda_1 = 5$ and $\lambda_2 = -4$ are real, unequal, and have opposite signs. By our discussion of this case we know that (0, 0) is an unstable saddle point of the linear system, and hence by Part 3 of Theorem 2, it is also an unstable saddle point of the almost linear



FIGURE 9.2.8. Trajectories of the linearized system of Example 2.



FIGURE 9.2.9. Trajectories of the original almost linear system of Example 2.



FIGURE 9.2.10. Phase portrait for the almost linear system in Eq. (13).

system in (13). The trajectories of the linear system near (0, 0) are shown in Fig. 9.2.8, and those of the nonlinear system in (13) are shown in Fig. 9.2.9. Figure 9.2.10 shows a phase portrait of the nonlinear system in (13) from a "wider view." In addition to the saddle point at (0, 0), there are spiral points near the points (0.279, 1.065) and (0.933, -1.057), and a node near (-2.354, -0.483).

We have seen that the system x' = f(x, y), y' = g(x, y) with isolated critical point (x_0, y_0) transforms via the substitution $x = u + x_0$, $y = v + y_0$ to an equivalent *uv*-system with corresponding critical point (0, 0) and linearization $\mathbf{u}' = \mathbf{J}\mathbf{u}$, whose coefficient matrix \mathbf{J} is the Jacobian matrix in (8) of the functions f and g at (x_0, y_0) . Consequently we need not carry out the substitution explicitly; instead, we can proceed directly to calculate the eigenvalues of \mathbf{J} preparatory to application of Theorem 2.

Example 3 Determine the type and stability of the critical point (4, 3) of the almost linear system

$$\frac{dx}{dt} = 33 - 10x - 3y + x^2,$$
(14)
$$\frac{dy}{dt} = -18 + 6x + 2y - xy.$$

Solution

With $f(x, y) = 33 - 10x - 3y + x^2$, g(x, y) = -18 + 6x + 2y - xy and $x_0 = 4$, $y_0 = 3$ we have

$$\mathbf{J}(x, y) = \begin{bmatrix} -10 + 2x & -3\\ 6 - y & 2 - x \end{bmatrix}, \text{ so } \mathbf{J}(4, 3) = \begin{bmatrix} -2 & -3\\ 3 & -2 \end{bmatrix}$$

The associated linear system

$$\frac{du}{dt} = -2u - 3v,$$

$$\frac{dv}{dt} = 3u - 2v$$
(15)

has characteristic equation $(\lambda + 2)^2 + 9 = 0$, with complex conjugate roots $\lambda = -2 \pm 3i$. Hence (0,0) is an asymptotically stable spiral point of the linear system in (15), so Theorem 2 implies that (4,3) is an asymptotically stable spiral point of the original almost linear system in (14). Figure 9.2.11 shows some typical trajectories of the linear system in (15), and Fig. 9.2.12 shows how this spiral point fits into the phase portrait for the original almost linear system in (14).



FIGURE 9.2.11. Spiral trajectories of the linear system in Eq. (15).



FIGURE 9.2.12. Phase portrait for the almost linear system in Eq. (14).

9.2 **Problems**

In Problems 1 through 10, apply Theorem 1 to determine the type of the critical point (0,0) and whether it is asymptotically stable, stable, or unstable. Verify your conclusion by using a computer system or graphing calculator to construct a phase portrait for the given linear system.

1.
$$\frac{dx}{dt} = -2x + y$$
, $\frac{dy}{dt} = x - 2y$
2. $\frac{dx}{dt} = 4x - y$, $\frac{dy}{dt} = 2x + y$
3. $\frac{dx}{dt} = x + 2y$, $\frac{dy}{dt} = 2x + y$
4. $\frac{dx}{dt} = 3x + y$, $\frac{dy}{dt} = 5x - y$
5. $\frac{dx}{dt} = x - 2y$, $\frac{dy}{dt} = 2x - 3y$
6. $\frac{dx}{dt} = 5x - 3y$, $\frac{dy}{dt} = 3x - y$
7. $\frac{dx}{dt} = 3x - 2y$, $\frac{dy}{dt} = 4x - y$
8. $\frac{dx}{dt} = x - 3y$, $\frac{dy}{dt} = 6x - 5y$
9. $\frac{dx}{dt} = 2x - 2y$, $\frac{dy}{dt} = 4x - 2y$
10. $\frac{dx}{dt} = x - 2y$, $\frac{dy}{dt} = 5x - y$

Each of the systems in Problems 11 through 18 has a single critical point (x_0, y_0) . Apply Theorem 2 to classify this critical point as to type and stability. Verify your conclusion by using a computer system or graphing calculator to construct a phase portrait for the given system.

11.
$$\frac{dx}{dt} = x - 2y, \quad \frac{dy}{dt} = 3x - 4y - 2$$

12. $\frac{dx}{dt} = x - 2y - 8, \quad \frac{dy}{dt} = x + 4y + 10$
13. $\frac{dx}{dt} = 2x - y - 2, \quad \frac{dy}{dt} = 3x - 2y - 2$

14.	$\frac{dx}{dt} = x + y - 7, \frac{dy}{dt} = 3x - y - 5$
15.	$\frac{dx}{dt} = x - y, \frac{dy}{dt} = 5x - 3y - 2$
16.	$\frac{dx}{dt} = x - 2y + 1, \frac{dy}{dt} = x + 3y - 9$
17.	$\frac{dx}{dt} = x - 5y - 5, \frac{dy}{dt} = x - y - 3$
18.	$\frac{dx}{dt} = 4x - 5y + 3, \frac{dy}{dt} = 5x - 4y + 6$

In Problems 19 through 28, investigate the type of the critical point (0,0) of the given almost linear system. Verify your conclusion by using a computer system or graphing calculator to construct a phase portrait. Also, describe the approximate locations and apparent types of any other critical points that are visible in your figure. Feel free to investigate these additional critical points; you can use the computational methods discussed in the application material for this section.

19.
$$\frac{dx}{dt} = x - 3y + 2xy, \quad \frac{dy}{dt} = 4x - 6y - xy$$

20. $\frac{dx}{dt} = 6x - 5y + x^2, \quad \frac{dy}{dt} = 2x - y + y^2$
21. $\frac{dx}{dt} = x + 2y + x^2 + y^2, \quad \frac{dy}{dt} = 2x - 2y - 3xy$
22. $\frac{dx}{dt} = x + 4y - xy^2, \quad \frac{dy}{dt} = 2x - y + x^2y$
23. $\frac{dx}{dt} = 2x - 5y + x^3, \quad \frac{dy}{dt} = 4x - 6y + y^4$
24. $\frac{dx}{dt} = 5x - 3y + y(x^2 + y^2), \quad \frac{dy}{dt} = 5x + y(x^2 + y^2)$
25. $\frac{dx}{dt} = x - 2y + 3xy, \quad \frac{dy}{dt} = 2x - 3y - x^2 - y^2$
26. $\frac{dx}{dt} = 3x - 2y - x^2 - y^2, \quad \frac{dy}{dt} = 2x - y - 3xy$
27. $\frac{dx}{dt} = x - y + x^4 - y^2, \quad \frac{dy}{dt} = 2x - y + y^4 - x^2$
28. $\frac{dx}{dt} = 3x - y + x^3 + y^3, \quad \frac{dy}{dt} = 13x - 3y + 3xy$

In Problems 29 through 32, find all critical points of the given system, and investigate the type and stability of each. Verify your conclusions by means of a phase portrait constructed using a computer system or graphing calculator.

 $29. \quad \frac{dx}{dt} = x - y, \quad \frac{dy}{dt} = x^2 - y$

30.
$$\frac{dx}{dt} = y - 1$$
, $\frac{dy}{dt} = x^2 - y$

31.
$$\frac{dx}{dt} = y^2 - 1$$
, $\frac{dy}{dt} = x^3 - y$

$$32. \quad \frac{dx}{dt} = xy - 2, \quad \frac{dy}{dt} = x - 2y$$

Bifurcations

The term bifurcation generally refers to something "splitting apart." With regard to differential equations or systems involving a parameter, it refers to abrupt changes in the character of the solutions as the parameter is changed continuously. Problems 33 through 36 illustrate sensitive cases in which small perturbations in the coefficients of a linear or almost linear system can change the type or stability (or both) of a critical point.



FIGURE 9.2.13(a). Stable spiral with $\epsilon = -0.2$.



FIGURE 9.2.13(c). Stable center with $\epsilon = 0$.



FIGURE 9.2.13(d). Unstable spiral with $\epsilon = 0.05$.

33. Consider the linear system

$$\frac{dx}{dt} = \epsilon x - y, \quad \frac{dy}{dt} = x + \epsilon y.$$

Show that the critical point (0, 0) is (a) a stable spiral point if $\epsilon < 0$; (b) a center if $\epsilon = 0$; (c) an unstable spiral point if $\epsilon > 0$. Thus small perturbations of the system x' = -y, y' = x can change both the type and stability of the critical point. Figures 9.2.13(a)–(e) illustrate the loss of stability that occurs at $\epsilon = 0$ as the parameter increases from $\epsilon < 0$ to $\epsilon > 0$.

34. Consider the linear system

$$\frac{dx}{dt} = -x + \epsilon y, \quad \frac{dy}{dt} = x - y.$$

Show that the critical point (0, 0) is (a) a stable spiral point if $\epsilon < 0$; (b) a stable node if $0 \le \epsilon < 1$. Thus small perturbations of the system x' = -x, y' = x - y can change the type of the critical point (0, 0) without changing its stability.

35. This problem deals with the almost linear system

$$\frac{dx}{dt} = y + hx(x^2 + y^2), \quad \frac{dy}{dt} = -x + hy(x^2 + y^2),$$



FIGURE 9.2.13(b). Stable spiral with $\epsilon = -0.05$.



FIGURE 9.2.13(e). Unstable spiral with $\epsilon = 0.2$.

in illustration of the sensitive case of Theorem 2, in which the theorem provides no information about the stability of the critical point (0, 0). (a) Show that (0, 0) is a center of the linear system obtained by setting h = 0. (b) Suppose that $h \neq 0$. Let $r^2 = x^2 + y^2$, then apply the fact that

$$x\frac{dx}{dt} + y\frac{dy}{dt} = r\frac{dr}{dt}$$

to show that $dr/dt = hr^3$. (c) Suppose that h = -1. Integrate the differential equation in (b); then show that $r \to 0$ as $t \to +\infty$. Thus (0,0) is an asymptotically stable critical point of the almost linear system in this case. (d) Suppose that h = +1. Show that $r \to +\infty$ as t increases, so (0,0) is an unstable critical point in this case.

36. This problem presents the famous *Hopf bifurcation* for the almost linear system

$$\frac{dx}{dt} = \epsilon x + y - x(x^2 + y^2),$$
$$\frac{dy}{dt} = -x + \epsilon y - y(x^2 + y^2),$$

which has imaginary characteristic roots $\lambda = \pm i$ if $\epsilon = 0$. (a) Change to polar coordinates as in Example 5 of Section 9.1 to obtain the system $r' = r(\epsilon - r^2)$, $\theta' = -1$. (b) Separate variables and integrate directly to show that if $\epsilon \leq 0$, then $r(t) \rightarrow 0$ as $t \rightarrow +\infty$, so in this case the origin is a stable spiral point. (c) Show similarly that if $\epsilon > 0$, then $r(t) \rightarrow \sqrt{\epsilon}$ as $t \rightarrow +\infty$, so in this case the origin is an unstable spiral point. The circle $r(t) \equiv \sqrt{\epsilon}$ itself is a closed periodic solution or *limit cycle*. Thus a limit cycle of increasing size is spawned as the parameter ϵ increases through the critical value 0.

37. In the case of a two-dimensional system that is *not* almost linear, the trajectories near an isolated critical point can exhibit a considerably more complicated structure than those near the nodes, centers, saddle points, and spiral points discussed in this section. For example, consider the system

$$\frac{dx}{dt} = x(x^3 - 2y^3),$$

$$\frac{dy}{dt} = y(2x^3 - y^3)$$
(16)

having (0,0) as an isolated critical point. This system is not almost linear because (0,0) is not an *isolated* critical point of the trivial associated linear system x' = 0, y' = 0. Solve the homogeneous first-order equation

$$\frac{dy}{dx} = \frac{y(2x^3 - y^3)}{x(x^3 - 2y^3)}$$

to show that the trajectories of the system in (16) are *folia* of Descartes of the form

$$x^3 + y^3 = 3cxy,$$

where c is an arbitrary constant (Fig. 9.2.14).



FIGURE 9.2.14. Trajectories of the system in Eq. (16).

38. First note that the characteristic equation of the 2 × 2 matrix A can be written in the form λ² – Tλ + D = 0, where D is the determinant of A and the trace T of the matrix A is the sum of its two diagonal elements. Then apply Theorem 1 to show that the type of the critical point (0,0) of the system x' = Ax is determined—as indicated in Fig. 9.2.15 —by the location of the point (T, D) in the *trace-determinant plane* with horizontal T-axis and vertical D-axis.



FIGURE 9.2.15. The critical point (0, 0) of the system x' = Ax is a

- spiral sink or source if the point (T, D) lies above the parabola $T^2 = 4D$ but off the *D*-axis;
- stable center if (*T*, *D*) lies on the positive *D*-axis;
- nodal sink or source if (*T*, *D*) lies between the parabola and the *T*-axis;
- saddle point if (T, D) lies beneath the *T*-axis.

9.2 Application Phase Plane Portraits of Almost Linear Systems



Maple/Mathematica/MATLAB.

Interesting and complicated phase portraits often result from simple nonlinear perturbations of linear systems. For instance, Fig. 9.2.16 shows a phase portrait for the almost linear system

$$\frac{dx}{dt} = -y\cos(x+y-1),$$

$$\frac{dy}{dt} = x\cos(x-y+1).$$
(1)

Among the seven critical points marked with dots, we see

- Apparent spiral points in the first and third quadrants of the *xy*-plane;
- Apparent saddle points in the second and fourth quadrants, plus another one on the positive *x*-axis;
- A critical point of undetermined character on the negative y-axis; and
- An apparently "very weak" spiral point at the origin—meaning one that is approached very slowly as *t* increases or decreases (according as it is a sink or a source).

Some ODE software systems can automatically locate and classify critical points. For instance, Fig. 9.2.17 shows a screen produced by the MATLAB program **pplane** that is cited in the Section 9.1 application. It shows that the fourth-quadrant critical point in Fig. 9.2.16 has approximate coordinates (1.5708, -2.1416), and that the coefficient matrix of the associated linear system has the positive eigenvalue $\lambda_1 \approx 2.8949$ and the negative eigenvalue $\lambda_2 \approx -2.3241$. It therefore follows from Theorem 2 that this critical point is, indeed, a saddle point of the almost linear system in (1).

🛿 PPLANE5 Equilibrium point data	_ 🗆 ×
File Window	
There is a saddle point at (1.5708, -2.1416).	
The Jacobian is: 2.1416 2.1416 1.5708 -1.5708	
The eigenvalues and eigenvectors are: 2.8949 (0.94334, 0.33182) -2.3241 (-0.43241, 0.90168)	

FIGURE 9.2.17. The fourth-quadrant saddle point revealed.

With a general computer algebra system such as *Maple* or *Mathematica*, you may have to do a bit of work yourself—or tell the computer precisely what to do—to find and classify a critical point. For instance, the *Maple* command

or the Mathematica command



FIGURE 9.2.16. Phase plane portrait for the system in Eq. (1).
526 CHAPTER 9 NONLINEAR SYSTEMS AND PHENOMENA



FIGURE 9.2.18. TI-NspireTM CX CAS calculation of the fourth-quadrant critical point of the almost linear system (1).



will find the critical-point coordinates a = 1.5708, b = -2.1416 indicated earlier. Similarly, Fig. 9.2.18 shows a TI handheld calculation of this critical point. Then the substitution x = u + a, y = v + b yields the translated system

$$\frac{du}{dt} = (2.1416 - v)\cos(1.5708 - u - v) = f(u, v),$$

$$\frac{dv}{dt} = (1.5708 + u)\cos(4.7124 + u - v) = g(u, v).$$
(2)

If we substitute u = v = 0 in the Jacobian matrix $\partial(f, g)/\partial(u, v)$, we get the coefficient matrix

$$\mathbf{A} = \begin{bmatrix} 2.1416 & 2.1416 \\ 1.5708 & -1.5708 \end{bmatrix}$$

of the linear system corresponding to (2). Then the Maple command

evalf(Eigenvals(A))

or the Mathematica command

Eigenvalues[A]

or the Wolfram Alpha query

```
((2.1416, 2.1416), (1.5708, -1.5708))
```

yields the eigenvalues $\lambda_1 \approx 2.8949$ and $\lambda_2 \approx -2.3241$, thereby verifying that the critical point (1.5708, -2.1416) of (1) is, indeed, a saddle point.

Use a computer algebra system to find and classify the other critical points of (1) indicated in Fig. 9.2.16. Then investigate similarly an almost linear system of your own construction. One convenient way to construct such a system is to begin with a linear system and insert sine or cosine factors resembling the ones in (1).

9.3 Ecological Models: Predators and Competitors

Some of the most interesting and important applications of stability theory involve the interactions between two or more biological populations occupying the same environment. We consider first a **predator-prey** situation involving two species. One species—the **predators**—feeds on the other species—the **prey**—which in turn feeds on some third food item readily available in the environment. A standard example is a population of foxes and rabbits in a woodland; the foxes (predators) eat rabbits (the prey), while the rabbits eat certain vegetation in the woodland. Other examples are sharks (predators) and food fish (prey), bass (predators) and sunfish (prey), ladybugs (predators) and aphids (prey), and beetles (predators) and scale insects (prey).

The classical mathematical model of a predator-prey situation was developed in the 1920s by the Italian mathematician Vito Volterra (1860–1940) in order to analyze the cyclic variations observed in the shark and food-fish populations in the Adriatic Sea. To construct such a model, we denote the number of prey at time t by x(t), the number of predators by y(t), and make the following simplifying assumptions.

1. In the absence of predators, the prey population would grow at a natural rate, with dx/dt = ax, a > 0.

- 2. In the absence of prey, the predator population would decline at a natural rate, with dy/dt = -by, b > 0.
- **3.** When both predators and prey are present, there occurs, in combination with these natural rates of growth and decline, a decline in the prey population and a growth in the predator population, each at a rate proportional to the frequency of encounters between individuals of the two species. We assume further that the frequency of such encounters is proportional to the product xy, reasoning that doubling either population alone should double the frequency of encounters, while doubling both populations ought to quadruple the frequency of encounters. Consequently, the consumption of prey by predators results in
 - an interaction rate of decline -pxy in the prey population x, and
 - an interaction rate of growth qxy in the predator population y.

When we combine the natural and interaction rates ax and -pxy for the prey population x, as well as the natural and interaction rates -by and qxy for the predator population y, we get the **predator-prey system**

$$\frac{dx}{dt} = ax - pxy = x(a - py),$$

$$\frac{dy}{dt} = -by + qxy = y(-b + qx),$$
(1)

with the constants *a*, *b*, *p*, and *q* all positive. [*Note:* You may see the predator and prey equations written in either order in (1). It is important to recognize that the predator equation has negative linear term and positive interaction term, whereas the prey equation has positive linear term and negative interaction term.]

Example 1 The critical points A critical point of the general predator–prey system in (1) is a solution (x, y) of the equations

$$x(a - py) = 0, \quad y(-b + qx) = 0.$$
 (2)

The first of these two equations implies that either x = 0 or $y = a/p \neq 0$, and the second implies that either y = 0 or $x = b/q \neq 0$. It follows readily that this predator-prey system has the two (isolated) critical points (0, 0) and (b/q, a/p).

THE CRITICAL POINT (0, 0): The Jacobian matrix of the system in (1) is

$$\mathbf{J}(x,y) = \begin{bmatrix} a - py & -px \\ qy & -b + qx \end{bmatrix}, \text{ so } \mathbf{J}(0,0) = \begin{bmatrix} a & 0 \\ 0 & -b \end{bmatrix}.$$
 (3)

The matrix $\mathbf{J}(0, 0)$ has characteristic equation $(a - \lambda)(-b - \lambda) = 0$ and the eigenvalues $\lambda_1 = a > 0$, $\lambda_2 = -b < 0$ with different signs. Hence it follows from Theorems 1 and 2 in Section 9.2 that the critical point (0, 0) is an unstable saddle point, both of the predator–prey system and of its linearization at (0, 0). The corresponding equilibrium solution $x(t) \equiv 0$, $y(t) \equiv 0$ merely describes simultaneous extinction of the prey (x) and predator (y) populations.

THE CRITICAL POINT (b/q, a/p): The Jacobian matrix

$$\mathbf{J}(b/q, a/p) = \begin{bmatrix} 0 & -\frac{pb}{q} \\ \frac{aq}{p} & 0 \end{bmatrix}$$
(4)

has characteristic equation $\lambda^2 + ab = 0$ and the pure imaginary eigenvalues $\lambda_1, \lambda_2 = \pm i \sqrt{ab}$. It follows from Theorem 1 in Section 9.2 that the linearization of the predator-prey system at (b/q, a/p) has a stable center at the origin. Thus we have the indeterminate case of Theorem 2 in Section 9.2, in which case the critical point can (aside from a stable center) also be either a stable spiral sink or an unstable spiral source of the predator-prey system itself. Hence further investigation is required to determine the actual character of the critical point (b/q, a/p). The corresponding equilibrium solution $x(t) \equiv b/q$, $y(t) \equiv a/p$ describes the only *nonzero* constant prey (x) and predator (y) populations that coexist permanently.

THE PHASE PLANE PORTRAIT: In Problem 1 we ask you to analyze numerically a typical predator–prey system and verify that the linearizations at its two critical points agree qualitatively with the phase plane portrait shown in Fig. 9.3.1—where the nontrivial critical point appears visually to be a stable center. Of course, only the first quadrant of this portrait corresponds to physically meaningful solutions describing nonnegative populations of prey and predators.

In Problem 2 we ask you to derive an exact implicit solution of the predator-prey system of Fig. 9.3.1—a solution that can be used to show that its phase plane trajectories in the first quadrant are, indeed, simple closed curves that encircle the critical point (75, 50) as indicated in the figure. It then follows from Problem 30 in Section 9.1 that the explicit solution functions x(t) and y(t) are both periodic functions of t—thus explaining the periodic fluctuations that are observed empirically in predator-prey populations.



FIGURE 9.3.1. Phase plane portrait for the predator–prey system x' = 200x - 4xy, y' = -150y + 2xy with critical points (0, 0) and (75, 50).



Example 2 Oscillating populations Figure 9.3.2 shows a computer-generated direction field and phase portrait for the predator–prey system

$$\frac{dx}{dt} = (0.2)x - (0.005)xy = (0.005)x(40 - y),$$

$$\frac{dy}{dt} = -(0.5)y + (0.01)xy = (0.01)y(-50 + x),$$
(5)

where x(t) denotes the number of rabbits and y(t) the number of foxes after t months. Evidently the critical point (50, 40) is a stable center representing equilibrium populations of 50 rabbits and 40 foxes. Any other initial point lies on a closed trajectory enclosing this equilibrium point. The direction field indicates that the point (x(t), y(t)) traverses its trajectory in a counterclockwise direction, with the rabbit and fox populations oscillating periodically between their separate maximum and minimum values. A drawback is that the phase plane plot provides no indication as to the speed with which each trajectory is traversed.



FIGURE 9.3.3. Periodic oscillations of the predator and prey populations in Example 2.

This lost "sense of time" is recaptured by graphing the two individual population functions as functions of time t. In Fig. 9.3.3 we have graphed approximate solution functions x(t) and y(t) calculated using the Runge–Kutta method of Section 7.7 with initial values x(0) = 70 and y(0) = 40. We see that the rabbit population oscillates between the extreme values $x_{\text{max}} \approx 72$ and $x_{\text{min}} \approx 33$, while the fox population oscillates (out of phase) between the extreme values $y_{\text{max}} \approx 70$ and $y_{\text{min}} \approx 20$. A careful measurement indicates that the period P of oscillation of each population is slightly over 20 months. One could "zoom in" on the maximum/minimum points on each graph in order to refine these estimates of the period and the maximum and minimum rabbit and fox populations.

Any positive initial conditions $x_0 = x(0)$ and $y_0 = y(0)$ yield a similar picture, with the rabbit and fox populations both surviving in coexistence with each other.

Competing Species

Now we consider two species (of animals, plants, or bacteria, for instance) with populations x(t) and y(t) at time t that compete with each other for the food available in their common environment. This is in marked contrast to the case in which one species preys on the other. To construct a mathematical model that is as realistic as possible, let us assume that in the absence of either species, the other would have a bounded (logistic) population like those considered in Section 2.1. In the absence of any interaction or competition between the two species, their populations x(t) and y(t) would then satisfy the differential equations

$$\frac{dx}{dt} = a_1 x - b_1 x^2,$$

$$\frac{dy}{dt} = a_2 y - b_2 y^2,$$
(6)

each of the form of Eq. (2) of Section 2.1. But in addition, we assume that competition has the effect of a rate of decline in each population that is proportional to their product xy. We insert such terms with *negative* proportionality constants $-c_1$ and $-c_2$ in the equations in (6) to obtain the **competition system**

$$\frac{dx}{dt} = a_1 x - b_1 x^2 - c_1 x y = x(a_1 - b_1 x - c_1 y),$$

$$\frac{dy}{dt} = a_2 y - b_2 y^2 - c_2 x y = y(a_2 - b_2 y - c_2 x),$$
(7)

where the coefficients a_1 , a_2 , b_1 , b_2 , c_1 , and c_2 are all positive.

The almost linear system in (7) has four critical points. Upon setting the righthand sides of the two equations equal to zero, we see that if x = 0, then either y = 0or $y = a_2/b_2$, whereas if y = 0, then either x = 0 or $x = a_1/b_1$. This gives the three critical points (0,0), (0, a_2/b_2), and (a_1/b_1 , 0). The fourth critical point is obtained from the simultaneous solution of the equations

$$b_1 x + c_1 y = a_1, \quad c_2 x + b_2 y = a_2.$$
 (8)

We assume that, as in most interesting applications, these equations have a single solution and that the corresponding critical point lies in the first quadrant of the *xy*-plane. This point (x_E, y_E) is then the fourth critical point of the system in (7), and it represents the possibility of coexistence of the two species, with constant nonzero equilibrium populations $x(t) \equiv x_E$ and $y(t) \equiv y_E$.

We are interested in the stability of the critical point (x_E, y_E) . This turns out to depend on whether

$$c_1 c_2 < b_1 b_2 \quad \text{or} \quad c_1 c_2 > b_1 b_2.$$
 (9)

Each inequality in (9) has a natural interpretation. Examining the equations in (6), we see that the coefficients b_1 and b_2 represent the inhibiting effect of each population on its own growth (possibly due to limitations of food or space). On the other hand, c_1 and c_2 represent the effect of competition between the two populations. Thus b_1b_2 is a measure of *inhibition*, while c_1c_2 is a measure of *competition*. A general analysis of the system in (7) shows the following:

- **1.** If $c_1c_2 < b_1b_2$, so that competition is small in comparison with inhibition, then (x_E, y_E) is an asymptotically stable critical point that is approached by each solution as $t \to +\infty$. Thus the two species can and do coexist in this case.
- **2.** If $c_1c_2 > b_1b_2$, so that competition is large in comparison with inhibition, then (x_E, y_E) is an unstable critical point, and either x(t) or y(t) approaches zero as $t \to +\infty$. Thus the two species cannot coexist in this case; one survives and the other becomes extinct.

Rather than carrying out this general analysis, we present two examples that illustrate these two possibilities.

Example 3 Survival of a single species Suppose that the populations x(t) and y(t) satisfy the equations

$$\frac{dx}{dt} = 14x - \frac{1}{2}x^2 - xy,$$
(10)
$$\frac{dy}{dt} = 16y - \frac{1}{2}y^2 - xy,$$

in which $a_1 = 14$, $a_2 = 16$, $b_1 = b_2 = \frac{1}{2}$, and $c_1 = c_2 = 1$. Then $c_1c_2 = 1 > \frac{1}{4} = b_1b_2$, so we should expect survival of a single species as predicted in Case 2 previously. We find readily that the four critical points are (0, 0), (0, 32), (28, 0), and (12, 8). We shall investigate them individually.

THE CRITICAL POINT (0, 0): The Jacobian matrix of the system in (10) is

$$\mathbf{J}(x, y) = \begin{bmatrix} 14 - x - y & -x \\ -y & 16 - y - x \end{bmatrix}, \text{ so } \mathbf{J}(0, 0) = \begin{bmatrix} 14 & 0 \\ 0 & 16 \end{bmatrix}.$$
(11)

The matrix $\mathbf{J}(0,0)$ has characteristic equation $(14 - \lambda)(16 - \lambda) = 0$ and has the eigenvalues

$$\lambda_1 = 14$$
 with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$

and

$$\lambda_2 = 16$$
 with eigenvector $\mathbf{v}_2 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$.

Both eigenvalues are positive, so it follows that (0,0) is a nodal source for the system's linearization x' = 14x, y' = 16y at (0,0), and hence—by Theorem 2 in Section 9.2—is also an unstable nodal source for the original system in (10). Figure 9.3.4 shows a phase portrait for the linearized system near (0,0).



FIGURE 9.3.4. Phase plane portrait for the linear system x' = 14x, y' = 16y corresponding to the critical point (0, 0).

THE CRITICAL POINT (0, 32): Substitution of x = 0, y = 32 in the Jacobian matrix J(x, y) shown in (11) yields the Jacobian matrix

$$\mathbf{J}(0,32) = \begin{bmatrix} -18 & 0\\ -32 & -16 \end{bmatrix}$$
(12)

of the nonlinear system (10) at the point (0, 32). Comparing Eqs. (7) and (8) in Section 9.2, we see that this Jacobian matrix corresponds to the linearization

$$\frac{du}{dt} = -18u,$$

$$\frac{dv}{dt} = -32u - 16v$$
(13)

of (10) at (0, 32). The matrix $\mathbf{J}(0, 32)$ has characteristic equation $(-18 - \lambda)(-16 - \lambda) = 0$ and has the eigenvalues $\lambda_1 = -18$ with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 & 16 \end{bmatrix}^T$ and $\lambda_2 = -16$ with eigenvector $\mathbf{v}_2 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$. Because both eigenvalues are negative, it follows that (0, 0) is a nodal sink for the linearized system, and hence—by Theorem 2 in Section 9.2—that (0, 32) is also a stable nodal sink for the original system in (10). Figure 9.3.5 shows a phase portrait for the linearized system near (0, 0).

THE CRITICAL POINT (28, 0): The Jacobian matrix

$$\mathbf{J}(28,0) = \begin{bmatrix} -14 & -28\\ 0 & -12 \end{bmatrix} \tag{14}$$

corresponds to the linearization

$$\frac{du}{dt} = -14u - 28v,$$

$$\frac{dv}{dt} = -12v$$
(15)

of (10) at (28,0). The matrix $\mathbf{J}(28,0)$ has characteristic equation $(-14 - \lambda)(-12 - \lambda) = 0$ and has the eigenvalues $\lambda_1 = -14$ with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ and $\lambda_2 = -12$ with eigenvector $\mathbf{v}_2 = \begin{bmatrix} -14 & 1 \end{bmatrix}^T$. Because both eigenvalues are negative, it follows that (0,0) is a nodal sink for the linearized system, and hence—by Theorem 2 in Section 9.2—that (28,0) is also a stable nodal sink for the original nonlinear system in (10). Figure 9.3.6 shows a phase portrait for the linearized system near (0,0).

THE CRITICAL POINT (12, 8): The Jacobian matrix

$$\mathbf{J}(12,8) = \begin{bmatrix} -6 & -12\\ -8 & -4 \end{bmatrix} \tag{16}$$

corresponds to the linearization

$$\frac{du}{dt} = -6u - 12v,$$

$$\frac{dv}{dt} = -8u - 4v$$
(17)

of (10) at (12, 8). The matrix J(12, 8) has characteristic equation

$$(-6 - \lambda)(-4 - \lambda) - (-8)(-12) = \lambda^2 + 10\lambda - 72 = 0$$

and has the eigenvalues

$$\lambda_1 = -5 - \sqrt{97} < 0$$
 with eigenvector $\mathbf{v}_1 = \begin{bmatrix} \frac{1}{8} (1 + \sqrt{97}) & 1 \end{bmatrix}^T$



FIGURE 9.3.6. Phase plane portrait for the linear system in Eq. (15) corresponding to the critical point (28, 0).



FIGURE 9.3.5. Phase plane portrait

for the linear system in Eq. (13)

(0, 32).

corresponding to the critical point

and



FIGURE 9.3.7. Phase plane portrait for the linear system in Eq. (17) corresponding to the critical point (12, 8).



Now that our local analysis of each of the four critical points is complete, it remains to assemble the information found into a coherent global picture. If we accept the facts that

- Near each critical point, the trajectories for the original system in (10) resemble qualitatively the linearized trajectories shown in Figs. 9.3.4–9.3.7, and
- As t → +∞ each trajectory either approaches a critical point or diverges toward infinity,

then it would appear that the phase plane portrait for the original system must resemble the rough sketch shown in Fig. 9.3.8. This sketch shows a few typical freehand trajectories connecting a nodal source at (0,0), nodal sinks at (0,32) and (28,0), and a saddle point at (12,8), with indicated directions of flow along these trajectories consistent with the known character of these critical points. Figure 9.3.9 shows a more precise computer-generated phase portrait and direction field for the nonlinear system in (10).





FIGURE 9.3.8. Rough sketch consistent with the analysis in Example 3.

FIGURE 9.3.9. Phase plane portrait for the system in Example 3.

The two trajectories that approach the saddle point (12, 8), together with that saddle point, form a **separatrix** that separates Regions I and II in Fig. 9.3.8. It plays a crucial role in determining the long-term behavior of the two populations. If the initial point (x_0, y_0) lies precisely on the separatrix, then (x(t), y(t)) approaches (12, 8) as $t \to +\infty$. Of course, random events make it extremely unlikely that (x(t), y(t)) will remain on the separatrix. If not, peaceful coexistence of the two species is impossible. If (x_0, y_0) lies in Region I above the separatrix, then (x(t), y(t)) approaches (0, 32) as $t \to +\infty$, so the population x(t) decreases to zero. Alternatively, if (x_0, y_0) lies in Region II below the separatrix, then (x(t), y(t)) approaches (28, 0) as $t \to +\infty$, so the population y(t) dies out. In short, whichever population has the initial competitive advantage survives, while the other faces extinction.

Example 4

Peaceful coexistence of two species Suppose that the populations x(t) and y(t) satisfy the competition system

$$\frac{dx}{dt} = 14x - 2x^2 - xy,$$

$$\frac{dy}{dt} = 16y - 2y^2 - xy,$$
(18)

for which $a_1 = 14$, $a_2 = 16$, $b_1 = b_2 = 2$, and $c_1 = c_2 = 1$. Then $c_1c_2 = 1 < 4 = b_1b_2$, so now the effect of inhibition is greater than that of competition. We find readily that the four critical points are (0, 0), (0, 8), (7, 0), and (4, 6). We proceed as in Example 3.

THE CRITICAL POINT (0, 0): When we drop the quadratic terms in (18), we get the same linearization x' = 14x, y' = 16y at (0,0) as in Example 3. Thus its coefficient matrix has the two positive eigenvalues $\lambda_1 = 14$ and $\lambda_2 = 16$, and its phase portrait is the same as that shown in Fig. 9.3.4. Therefore, (0,0) is an unstable nodal source for the original system in (18).

THE CRITICAL POINT (0, 8): The Jacobian matrix of the system in (18) is

$$\mathbf{J}(x, y) = \begin{bmatrix} 14 - 4x - y & -x \\ -y & 16 - 4y - x \end{bmatrix}, \text{ so } \mathbf{J}(0, 8) = \begin{bmatrix} 6 & 0 \\ -8 & -16 \end{bmatrix}.$$
(19)

The matrix J(0, 8) corresponds to the linearization

$$\frac{du}{dt} = 6u,$$

$$\frac{dv}{dt} = -8u - 16v$$
(20)

of (18) at (0,8). It has characteristic equation $(6 - \lambda)(-16 - \lambda) = 0$ and has the positive eigenvalue $\lambda_1 = 6$ with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 11 & -4 \end{bmatrix}^T$ and the negative eigenvalue $\lambda_2 = -16$ with eigenvector $\mathbf{v}_2 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$. It follows that (0,0) is a saddle point for the linearized system, and hence that (0, 8) is an unstable saddle point for the original system in (18). Figure 9.3.10 shows a phase portrait for the linearized system near (0, 0).

THE CRITICAL POINT (7, 0): The Jacobian matrix

$$\mathbf{J}(7,0) = \begin{bmatrix} -14 & -7\\ 0 & 9 \end{bmatrix}$$
(21)

corresponds to the linearization

$$\frac{du}{dt} = -14u - 7v,$$

$$\frac{dv}{dt} = 9v$$
(22)

of (18) at (7,0). The matrix $\mathbf{J}(7,0)$ has characteristic equation $(-14 - \lambda)(9 - \lambda) = 0$ and has the negative eigenvalue $\lambda_1 = -14$ with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$ and the positive eigenvalue $\lambda_2 = 9$ with eigenvector $\mathbf{v}_2 = \begin{bmatrix} -7 & 23 \end{bmatrix}^T$. It follows that (0, 0) is a saddle point for the linearized system, and hence that (7,0) is an unstable saddle point for the original system in (18). Figure 9.3.11 shows a phase portrait for the linearized system near (0, 0).

d

THE CRITICAL POINT (4, 6): The Jacobian matrix

$$\mathbf{J}(4,6) = \begin{bmatrix} -8 & -4\\ -6 & -12 \end{bmatrix}$$
(23)

corresponds to the linearization

$$\frac{du}{dt} = -8u - 4v,$$

$$\frac{dv}{dt} = -6u - 12v$$
(24)

of (18) at (4, 6). The matrix $\mathbf{J}(4, 6)$ has characteristic equation

$$(-8 - \lambda)(-12 - \lambda) - (-6)(-4) = \lambda^2 + 20\lambda + 72 = 0$$



FIGURE 9.3.11. Phase plane portrait for the linear system in Eq. (22) corresponding to the critical point (7, 0).



FIGURE 9.3.10. Phase plane portrait for the linear system in Eq. (20) corresponding to the critical point (0, 8).

and has the two negative eigenvalues

$$\lambda_1 = 2(-5 - \sqrt{7})$$
 with eigenvector $\mathbf{v}_1 = \begin{bmatrix} \frac{1}{3}(-1 + \sqrt{7}) & 1 \end{bmatrix}^T$

and

$$\lambda_2 = 2(-5 + \sqrt{7})$$
 with eigenvector $\mathbf{v}_2 = \begin{bmatrix} \frac{1}{3}(-1 - \sqrt{7}) & 1 \end{bmatrix}^T$

It follows that (0,0) is a nodal sink for the linearized system, and hence that (4,6) is a stable nodal sink for the original system in (18). Figure 9.3.12 shows a phase portrait for the linearized system near (0,0).

Figure 9.3.13 assembles all this local information into a global phase plane portrait for the original system in (18). The notable feature of this system is that—for *any* positive initial population values x_0 and y_0 —the point (x(t), y(t)) approaches the single critical point (4, 6) as $t \to +\infty$. It follows that the two species both survive in stable (peaceful) existence.

Interactions of Logistic Populations

If the coefficients a_1, a_2, b_1, b_2 are positive but $c_1 = c_2 = 0$, then the equations

FIGURE 9.3.12. Phase plane portrait for the linear system in Eq. (24) corresponding to the critical point (4, 6).



FIGURE 9.3.13. Direction field and phase portrait for the competition system $x' = 14x - 2x^2 - xy$, $y' = 16y - 2y^2 - xy$ of Example 4.

$$\frac{dx}{dt} = a_1 x - b_1 x^2 - c_1 xy,$$

$$\frac{dy}{dt} = a_2 y - b_2 y^2 - c_2 xy$$
(25)

describe two separate logistic populations x(t) and y(t) that have no effect on each other. Examples 3 and 4 illustrate cases in which the *xy*-coefficients c_1 and c_2 are both *positive*. The interaction between the two populations is then described as **competition**, because the effect of the *xy*-terms in (25) is to *decrease* the rates of growth of both populations—that is, each population is "hurt" by their mutual interaction.

Suppose, however, that the interaction coefficients c_1 and c_2 in (25) are both *negative*. Then the effect of the *xy*-terms is to increase the rates of growth of both populations—that is, each population is "helped" by their mutual interaction. This type of interaction is aptly described as **cooperation** between the two logistic populations.

Finally, the interaction between the two populations is one of **predation** if the interaction coefficients have *different signs*. For instance, if $c_1 > 0$ but $c_2 < 0$, then the *x*-population is hurt but the *y*-population is helped by their interaction. We may therefore describe x(t) as a prey population and y(t) as a predator population.

If either b_1 or b_2 is zero in (25), then the corresponding population would (in the absence of the other) exhibit exponential growth rather than logistic growth. For instance, suppose that $a_1 > 0$, $a_2 < 0$, $b_1 = b_2 = 0$, and $c_1 > 0$, $c_2 < 0$. Then x(t) is a naturally growing prey population while y(t) is a naturally declining predator population. This is the original predator–prey model with which we began this section.

Problems 26 through 34 illustrate a variety of the possibilities indicated here. The problems and examples in this section illustrate the power of elementary critical-point analysis. But remember that ecological systems in nature are seldom so simple as in these examples. Frequently they involve more than two species, and the growth rates of these populations and the interactions among them often are more complicated than those discussed in this section. Consequently, the mathematical modeling of ecological systems remains an active area of current research.

a



9.3 Problems

Predator–Prey System

Problems 1 and 2 deal with the predator-prey system

$$\frac{dx}{dt} = 200x - 4xy,$$

$$\frac{dy}{dt} = -150y + 2xy$$
(1)

that corresponds to Fig. 9.3.1.

- **1.** Starting with the Jacobian matrix of the system in (1), derive its linearizations at the two critical points (0,0) and (75, 50). Use a graphing calculator or computer system to construct phase plane portraits for these two linearizations that are consistent with the "big picture" shown in Fig. 9.3.1.
- 2. Separate the variables in the quotient

$$\frac{dy}{dx} = \frac{-150y + 2xy}{200x - 4xy}$$

of the two equations in (1), and thereby derive the exact implicit solution

$$200 \ln y + 150 \ln x - 2x - 4y = C$$

of the system. Use the contour plot facility of a graphing calculator or computer system to plot the contour curves of this equation through the points (75, 100), (75, 150), (75, 200), (75, 250), and (75, 300) in the *xy*-plane. Are your results consistent with Fig. 9.3.1?

3. Insect population Let x(t) be a harmful insect population (aphids?) that under natural conditions is held somewhat in check by a benign predator insect population y(t) (ladybugs?). Assume that x(t) and y(t) satisfy the predator-prey equations in (1), so that the stable equilibrium populations are $x_E = b/q$ and $y_E = a/p$. Now suppose that an insecticide is employed that kills (per unit time) the same fraction f < a of each species of insect. Show that the harmful population x_E is increased, while the benign population y_E is decreased, so the use of the insecticide is counterproductive. This is an instance in which mathematical analysis reveals undesirable consequences of a well-intentioned interference with nature.

Competition System

Problems 4 through 7 deal with the competition system

$$\frac{dx}{dt} = 60x - 4x^2 - 3xy,$$

$$\frac{dy}{dt} = 42y - 2y^2 - 3xy,$$
(2)

in which $c_1c_2 = 9 > 8 = b_1b_2$, so the effect of competition should exceed that of inhibition. Problems 4 through 7 imply that the four critical points (0,0), (0,21), (15,0), and (6,12) of the system in (2) resemble those shown in Fig. 9.3.9—a nodal source at the origin, a nodal sink on each coordinate axis, and a saddle point interior to the first quadrant. In each of these problems use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Finally, construct a first-quadrant phase plane portrait for the nonlinear system in (2). Do your local and global portraits look consistent?

- **4.** Show that the coefficient matrix of the linearization x' = 60x, y' = 42y of (2) at (0,0) has positive eigenvalues $\lambda_1 = 60$ and $\lambda_2 = 42$. Hence (0,0) is a nodal source for (2).
- 5. Show that the linearization of (2) at (0, 21) is u' = -3u, v' = -63u 42v. Then show that the coefficient matrix of this linear system has negative eigenvalues $\lambda_1 = -3$ and $\lambda_2 = -42$. Hence (0, 21) is a nodal sink for the system in (2).
- 6. Show that the linearization of (2) at (15,0) is u' = -60u 45v, v' = -3v. Then show that the coefficient matrix of this linear system has negative eigenvalues $\lambda_1 = -60$ and $\lambda_2 = -3$. Hence (15,0) is a nodal sink for the system in (2).
- 7. Show that the linearization of (2) at (6, 12) is u' = -24u 18v, v' = -36u 24v. Then show that the coefficient matrix of this linear system has eigenvalues $\lambda_1 = -24 18\sqrt{2} < 0$ and $\lambda_2 = -24 + 18\sqrt{2} > 0$. Hence (6, 12) is a saddle point for the system in (2).

Competition System

Problems 8 through 10 deal with the competition system

$$\frac{dx}{dt} = 60x - 3x^2 - 4xy,$$

$$\frac{dy}{dt} = 42y - 3y^2 - 2xy,$$
(3)

in which $c_1c_2 = 8 < 9 = b_1b_2$, so the effect of inhibition should exceed that of competition. The linearization of the system in (3) at (0,0) is the same as that of (2). This observation and Problems 8 through 10 imply that the four critical points (0,0), (0,14), (20,0), and (12,6) of (3) resemble those shown in Fig. 9.3.13—a nodal source at the origin, a saddle point on each coordinate axis, and a nodal sink interior to the first quadrant. In each of these problems use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Finally, construct a first-quadrant phase plane portrait for the nonlinear system in (3). Do your local and global portraits look consistent?

- 8. Show that the linearization of (3) at (0, 14) is u' = 4u, v' = -28u 42v. Then show that the coefficient matrix of this linear system has the positive eigenvalue $\lambda_1 = 4$ and the negative eigenvalue $\lambda_2 = -42$. Hence (0, 14) is a saddle point for the system in (3).
- **9.** Show that the linearization of (3) at (20, 0) is u' = -60u 80v, v' = 2v. Then show that the coefficient matrix of this linear system has the negative eigenvalue $\lambda_1 = -60$ and

the positive eigenvalue $\lambda_2 = 2$. Hence (20, 0) is a saddle point for the system in (3).

10. Show that the linearization of (3) at (12, 6) is u' = -36u - 48v, v' = -12u - 18v. Then show that the coefficient matrix of this linear system has eigenvalues $\lambda_1 = -27 + 3\sqrt{73}$ and $\lambda_2 = -27 - 3\sqrt{73}$, both of which are negative. Hence (12, 6) is a nodal sink for the system in (3).

Logistic Prey Population

Problems 11 through 13 deal with the predator-prey system

$$\frac{dx}{dt} = 5x - x^2 - xy,$$

$$\frac{dy}{dt} = -2y + xy,$$
(4)

in which the prey population x(t) is logistic but the predator population y(t) would (in the absence of any prey) decline naturally. Problems 11 through 13 imply that the three critical points (0,0), (5,0), and (2,3) of the system in (4) are as shown in Fig. 9.3.14—with saddle points at the origin and on the positive x-axis, and with a spiral sink interior to the first quadrant. In each of these problems use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Do your local portraits look consistent with Fig. 9.3.14?



FIGURE 9.3.14. Direction field and phase portrait for the predator–prey system of Problems 11 through 13.

- 11. Show that the coefficient matrix of the linearization x' = 5x, y' = -2y of (4) at (0,0) has the positive eigenvalue $\lambda_1 = 5$ and the negative eigenvalue $\lambda_2 = -2$. Hence (0,0) is a saddle point of the system in (4).
- 12. Show that the linearization of (4) at (5,0) is u' = -5u 5v, v' = 3v. Then show that the coefficient matrix of this linear system has the negative eigenvalue $\lambda_1 = -5$ and the positive eigenvalue $\lambda_2 = 3$. Hence (5,0) is a saddle point for the system in (4).
- 13. Show that the linearization of (4) at (2, 3) is u' = -2u 2v, v' = 3u. Then show that the coefficient matrix of this linear system has the complex conjugate eigenvalues λ₁, λ₂ = -1 ± i √5 with negative real part. Hence (2, 3) is a spiral sink for the system in (4).

Doomsday vs. Extinction

Problems 14 through 17 deal with the predator-prey system

$$\frac{dx}{dt} = x^2 - 2x - xy,$$

$$\frac{dy}{dt} = y^2 - 4y + xy.$$
(5)

Here each population—the prey population x(t) and the predator population y(t)—is an unsophisticated population (like the alligators of Section 2.1) for which the only alternatives (in the absence of the other population) are doomsday and extinction. Problems 14 through 17 imply that the four critical points (0,0), (0,4), (2,0), and (3,1) of the system in (5) are as shown in Fig. 9.3.15—a nodal sink at the origin, a saddle point on each coordinate axis, and a spiral source interior to the first quadrant. This is a two-dimensional version of "doomsday versus extinction." If the initial point (x_0, y_0) lies in Region I, then both populations increase without bound (until doomsday), whereas if it lies in Region II, then both populations decrease to zero (and thus both become extinct). In each of these problems use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Do your local portraits look consistent with Fig. 9.3.15?



FIGURE 9.3.15. Direction field and phase portrait for the predator–prey system of Problems 14 through 17.

- 14. Show that the coefficient matrix of the linearization x' = -2x, y' = -4y of the system in (5) at (0, 0) has the negative eigenvalues $\lambda_1 = -2$ and $\lambda_2 = -4$. Hence (0, 0) is a nodal sink for (5).
- 15. Show that the linearization of (5) at (0, 4) is u' = -6u, v' = 4u + 4v. Then show that the coefficient matrix of this linear system has the negative eigenvalue $\lambda_1 = -6$ and the positive eigenvalue $\lambda_2 = 4$. Hence (0, 4) is a saddle point for the system in (5).
- 16. Show that the linearization of (5) at (2, 0) is u' = 2u 2v, v' = -2v. Then show that the coefficient matrix of this linear system has the positive eigenvalue λ₁ = 2 and the negative eigenvalue λ₂ = -2. Hence (2, 0) is a saddle point for the system in (5).
- 17. Show that the linearization of (5) at (3, 1) is u' = 3u 3v, v' = u + v. Then show that the coefficient matrix of this linear system has complex conjugate eigenvalues λ_1 ,

 $\lambda_2 = 2 \pm i\sqrt{2}$ with positive real part. Hence (3, 1) is a spiral source for (5).

Problems 18 through 25 deal with the predator-prey system

$$\frac{dx}{dt} = 2x - xy + \epsilon x(5 - x),$$

$$\frac{dy}{dt} = -5y + xy,$$
(6)

for which a bifurcation occurs at the value $\epsilon = 0$ of the parameter ϵ . Problems 18 and 19 deal with the case $\epsilon = 0$, in which case the system in (6) takes the form

$$\frac{dx}{dt} = 2x - xy, \quad \frac{dy}{dt} = -5x + xy, \tag{7}$$

and these problems suggest that the two critical points (0,0)and (5,2) of the system in (7) are as shown in Fig. 9.3.16 a saddle point at the origin and a center at (5,2). In each problem use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Do your local portraits look consistent with Fig. 9.3.16?



FIGURE 9.3.16. The case $\epsilon = 0$ (Problems 18 and 19).

- **18.** Show that the coefficient matrix of the linearization x' = 2x, y' = -5y of (7) at (0,0) has the positive eigenvalue $\lambda_1 = 2$ and the negative eigenvalue $\lambda_2 = -5$. Hence (0,0) is a saddle point for the system in (7).
- 19. Show that the linearization of the system in (7) at (5, 2) is u' = -5v, v' = 2u. Then show that the coefficient matrix of this linear system has conjugate imaginary eigenvalues λ₁, λ₂ = ±i√10. Hence (0, 0) is a stable center for the linear system. Although this is the indeterminate case of Theorem 2 in Section 9.2, Fig. 9.3.16 suggests that (5, 2) also is a stable center for (7).

Problems 20 through 22 deal with the case $\epsilon = -1$, for which the system in (6) becomes

$$\frac{dx}{dt} = -3x + x^2 - xy, \quad \frac{dy}{dt} = -5y + xy,$$
 (8)

and imply that the three critical points (0, 0), (3, 0), and (5, 2)of (8) are as shown in Fig. 9.3.17—with a nodal sink at the origin, a saddle point on the positive x-axis, and a spiral source at (5, 2). In each problem use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Do your local portraits look consistent with Fig. 9.3.17?



FIGURE 9.3.17. The case $\epsilon = -1$ (Problems 20 through 22).

- **20.** Show that the coefficient matrix of the linearization x' = -3x, y' = -5y of the system in (8) at (0, 0) has the negative eigenvalues $\lambda_1 = -3$ and $\lambda_2 = -5$. Hence (0, 0) is a nodal sink for (8).
- 21. Show that the linearization of the system in (8) at (3,0) is u' = 3u 3v, v' = -2v. Then show that the coefficient matrix of this linear system has the positive eigenvalue λ₁ = 3 and the negative eigenvalue λ₂ = -2. Hence (3,0) is a saddle point for (8).
- **22.** Show that the linearization of (8) at (5, 2) is u' = 5u 5v, v' = 2u. Then show that the coefficient matrix of this linear system has complex conjugate eigenvalues λ_1 , $\lambda_2 = \frac{1}{2}(5 \pm i\sqrt{15})$ with positive real part. Hence (5, 2) is a spiral source for the system in (8).

Problems 23 through 25 deal with the case $\epsilon = 1$, so that the system in (6) takes the form

$$\frac{dx}{dt} = 7x - x^2 - xy, \quad \frac{dy}{dt} = -5y + xy,$$
 (9)

and these problems imply that the three critical points (0,0), (7,0), and (5,2) of the system in (9) are as shown in Fig. 9.3.18 —with saddle points at the origin and on the positive x-axis and with a spiral sink at (5,2). In each problem use a graphing calculator or computer system to construct a phase plane portrait for the linearization at the indicated critical point. Do your local portraits look consistent with Fig. 9.3.18?



FIGURE 9.3.18. The case $\epsilon = +1$ (Problems 23 through 25).

538 CHAPTER 9 NONLINEAR SYSTEMS AND PHENOMENA

- **23.** Show that the coefficient matrix of the linearization x' = 7x, y' = -5y of (9) at (0,0) has the positive eigenvalue $\lambda_1 = 7$ and the negative eigenvalue $\lambda_2 = -5$. Hence (0,0) is a saddle point for the system in (9).
- 24. Show that the linearization of (9) at (7,0) is u' = -7u 7v, v' = 2v. Then show that the coefficient matrix of this linear system has the negative eigenvalue $\lambda_1 = -7$ and the positive eigenvalue $\lambda_2 = 2$. Hence (7,0) is a saddle point for the system in (9).
- **25.** Show that the linearization of (9) at (5, 2) is u' = -5u 5v, v' = 2u. Then show that the coefficient matrix of this linear system has the complex conjugate eigenvalues λ_1 , $\lambda_2 = \frac{1}{2} \left(-5 \pm i \sqrt{15}\right)$ with negative real part. Hence (5, 2) is a spiral sink for the system in (9).

For each two-population system in Problems 26 through 34, first describe the type of x- and y-populations involved (exponential or logistic) and the nature of their interaction competition, cooperation, or predation. Then find and characterize the system's critical points (as to type and stability). Determine what nonzero x- and y-populations can coexist. Finally, construct a phase plane portrait that enables you to describe the long-term behavior of the two populations in terms of their initial populations x(0) and y(0).

26. $\frac{dx}{dt} = 2x - xy$, $\frac{dy}{dt} = 3y - xy$ 27. $\frac{dx}{dt} = 2xy - 4x$, $\frac{dy}{dt} = xy - 3y$ 28. $\frac{dx}{dt} = 2xy - 16x$, $\frac{dy}{dt} = 4y - xy$ 29. $\frac{dx}{dt} = 3x - x^2 - \frac{1}{2}xy$, $\frac{dy}{dt} = 4y - 2xy$ 30. $\frac{dx}{dt} = 3x - x^2 + \frac{1}{2}xy$, $\frac{dy}{dt} = \frac{1}{5}xy - y$ 31. $\frac{dx}{dt} = 3x - x^2 - \frac{1}{4}xy$, $\frac{dy}{dt} = xy - 2y$ 32. $\frac{dx}{dt} = 30x - 3x^2 + xy$, $\frac{dy}{dt} = 60y - 3y^2 + 4xy$ 33. $\frac{dx}{dt} = 30x - 2x^2 - xy$, $\frac{dy}{dt} = 80y - 4y^2 + 2xy$

9.3 Application Your Own Wildlife Conservation Preserve

You own a large wildlife conservation preserve that you originally stocked with F_0 foxes and R_0 rabbits on January 1, 2007. The following differential equations model the numbers R(t) of rabbits and F(t) foxes t months later:

$$\frac{dR}{dt} = (0.01)pR - (0.0001)aRF,$$
$$\frac{dF}{dt} = -(0.01)qF + (0.0001)bRF,$$

where *p* and *q* are the two largest digits (with p < q) and *a* and *b* are the two smallest nonzero digits (with a < b) in your student ID number.

The numbers of foxes and rabbits will oscillate periodically, out of phase with each other (like the functions x(t) and y(t) in Fig. 9.3.3). Choose your initial numbers F_0 of foxes and R_0 of rabbits—perhaps several hundred of each—so that the resulting solution curve in the *RF*-plane is a fairly eccentric closed curve. (The eccentricity may be increased if you begin with a relatively large number of rabbits and a small number of foxes, as any wildlife preserve owner would naturally do—because foxes prev on rabbits.)

Your task is to determine

- **1.** The period of oscillation of the rabbit and fox populations;
- **2.** The maximum and minimum numbers of rabbits, and the calendar dates on which they first occur;
- **3.** The maximum and minimum numbers of foxes, and the calendar dates on which they first occur.

With computer software that can plot both RF-trajectories and tR- and tF-solution curves like those in Figs. 9.3.2 and 9.3.3, you can "zoom in" graphically on the points whose coordinates provide the requested information.

Go to goo.gl/OWzvNu to download this application's computing resources including *Maple/Mathematica/*MATLAB.

9.4 Nonlinear Mechanical Systems



FIGURE 9.4.1. The mass on a spring.

Now we apply the qualitative methods of Sections 9.1 and 9.2 to the analysis of simple mechanical systems like the mass-on-a-spring system shown in Fig. 9.4.1. Let *m* denote the mass in a suitable system of units and let x(t) denote the displacement of the mass at time *t* from its equilibrium position (in which the spring is unstretched). Previously we have always assumed that the force F(x) exerted by the spring on the mass is a *linear* function of *x*: F(x) = -kx (Hooke's law). In reality, however, every spring in nature actually is nonlinear (even if only slightly so). Moreover, springs in some automobile suspension systems deliberately are designed to be nonlinear. Here, then, we are interested specifically in the effects of nonlinearity.

So now we allow the force function F(x) to be nonlinear. Because F(0) = 0 at the equilibrium position x = 0, we may assume that F has a power series expansion of the form

$$F(x) = -kx + \alpha x^2 + \beta x^3 + \cdots .$$
⁽¹⁾

We take k > 0 so that the reaction of the spring is directed opposite to the displacement when x is sufficiently small. If we assume also that the reaction of the spring is symmetric with respect to positive and negative displacements by the same distance, then F(-x) = -F(x), so F is an odd function. In this case it follows that the coefficient of x^n in Eq. (1) is zero if n is even, so the first nonlinear term is the one involving x^3 .

For a simple mathematical model of a nonlinear spring we therefore take

$$F(x) = -kx + \beta x^3,\tag{2}$$

ignoring all terms in Eq. (1) of degree greater than 3. The equation of motion of the mass m is then

$$mx'' = -kx + \beta x^3. \tag{3}$$

The Position–Velocity Phase Plane

If we introduce the *velocity*

>

$$\mathbf{v}(t) = \mathbf{x}'(t) \tag{4}$$

of the mass with *position* x(t), then we get from Eq. (3) the equivalent first-order system

$$\frac{dx}{dt} = y,$$

$$m\frac{dy}{dt} = -kx + \beta x^{3}.$$
(5)

A phase plane trajectory of this system is a position-velocity plot that illustrates the motion of the mass on the spring. We can solve explicitly for the trajectories of this system by writing

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-kx + \beta x^3}{my},$$

whence

$$my\,dy + (kx - \beta x^3)\,dx = 0.$$

Integration then yields

$$\frac{1}{2}my^2 + \frac{1}{2}kx^2 - \frac{1}{4}\beta x^4 = E \tag{6}$$

for the equation of a typical trajectory. We write *E* for the arbitrary constant of integration because $KE = \frac{1}{2}my^2$ is the kinetic energy of the mass with velocity *y*, and it is natural to define

$$PE = \frac{1}{2}kx^2 - \frac{1}{4}\beta x^4 \tag{7}$$

as the potential energy of the spring. Then Eq. (6) takes the form KE + PE = E, so the constant *E* turns out to be the total energy of the mass-spring system. Equation (6) then expresses conservation of energy for the *undamped* motion of a mass on a spring.

The behavior of the mass depends on the sign of the nonlinear term in Eq. (2). The spring is called

- hard if $\beta < 0$,
- *soft* if $\beta > 0$.

We consider the two cases separately.

HARD SPRING OSCILLATIONS: If $\beta < 0$, then the second equation in (5) takes the form $my' = -x (|\beta|x^2 + k)$, so it follows that the only critical point of the system is the origin (0, 0). Each trajectory

$$\frac{1}{2}my^2 + \frac{1}{2}kx^2 + \frac{1}{4}|\beta|x^4 = E > 0 \tag{8}$$

is an oval closed curve like those shown in Fig. 9.4.2, and thus (0,0) is a stable center. As the point (x(t), y(t)) traverses a trajectory in the clockwise direction, the position x(t) and velocity y(t) of the mass oscillate alternately, as illustrated in Fig. 9.4.3. The mass is moving to the right (with x increasing) when y > 0, to the left when y < 0. Thus the behavior of a mass on a nonlinear hard spring resembles qualitatively that of a mass on a linear spring with $\beta = 0$ (as in Example 3 of Section 9.1). But one difference between the linear and nonlinear spring is independent of the initial conditions, the period of a mass on a linear spring depends on its initial position x(0) and initial velocity y(0) (Problems 21 through 26).

Remark The hard spring equation $mx'' = -kx - |\beta|x^3$ has equivalent first-order system

$$x' = y, \quad y' = -\frac{k}{m}x - \frac{|\beta|}{m}x^3$$

with Jacobian matrix

$$\mathbf{J}(x, y) = \begin{bmatrix} 0 & 1\\ -\frac{k}{m} - \frac{3|\beta|}{m}x^2 & 0 \end{bmatrix}, \text{ so } \mathbf{J}(0, 0) = \begin{bmatrix} 0 & 1\\ -\omega^2 & 0 \end{bmatrix}$$

(writing $k/m = \omega^2$ as usual). The latter matrix has characteristic equation $\lambda^2 + \omega^2 = 0$ and pure imaginary eigenvalues $\lambda_1, \lambda_2 = \pm \omega i$. Thus the linearized system $x' = y, y' = -\omega^2 x$ has a stable center at the critical point (0, 0)—as we observed in Example 4 of Section 9.1. However, the nonlinear cubic term in the differential equation has (in effect) replaced the elliptical trajectories (as in Fig. 9.1.8) of the linear system with the "flatter" quartic ovals we see in Fig. 9.4.2.



FIGURE 9.4.2. Position–velocity phase plane portrait for the hard mass-and-spring system with m = k = 2 and $\beta = -4 < 0$.



FIGURE 9.4.3. Position and velocity solution curves for the hard mass-and-spring system with m = k = 2 and $\beta = -4 < 0$.

SOFT SPRING OSCILLATIONS: If $\beta > 0$, then the second equation in (5) takes the form $my' = x (\beta x^2 - k)$, so it follows that the system has the two critical points $(\pm \sqrt{k/\beta}, 0)$ in addition to the critical point (0, 0). These three critical points yield the only solutions for which the mass can remain at rest. The following example illustrates the greater range of possible behaviors of a mass on a soft spring.

Example 1 Undamped soft spring If m = 1, k = 4, and $\beta = 1$, then the equation of motion of the mass is

$$\frac{t^2x}{tt^2} + 4x - x^3 = 0, (9)$$

and Eq. (6) gives the trajectories in the form

$$\frac{1}{2}y^2 + 2x^2 - \frac{1}{4}x^4 = E.$$
(10)

After solving for

$$y = \pm \sqrt{2E - 4x^2 + \frac{1}{2}x^4},\tag{10'}$$

we could select a fixed value of the constant energy E and plot manually a trajectory like one of those shown in the computer-generated position–velocity phase plane portrait in Fig. 9.4.4.

The different types of phase plane trajectories correspond to different values of the energy *E*. If we substitute $x = \pm \sqrt{k/\beta}$ and y = 0 into (6), we get the energy value $E = k^2/(4\beta) = 4$ (because k = 4 and $\beta = 1$) that corresponds to the trajectories that intersect the *x*-axis at the nontrivial critical points (-2, 0) and (2, 0). These emphasized trajectories are called **separatrices** because they separate phase plane regions of different behavior.

The nature of the motion of the mass is determined by which type of trajectory its initial conditions determine. The simple closed trajectories encircling (0,0) in the region bounded by the separatrices correspond to energies in the range 0 < E < 4. These closed trajectories represent *periodic* oscillations of the mass back and forth around the equilibrium point x = 0.

The unbounded trajectories lying in the regions above and below the separatrices correspond to values of *E* greater than 4. These represent motions in which the mass approaches x = 0 with sufficient energy that it continues on through the equilibrium point, never to return again (as indicated in Fig. 9.4.5).



10 5 Velocity y -5 -5 -1 -0.5 0 0 -5 1 -0.5 0 0.5 1

FIGURE 9.4.4. Position–velocity phase plane portrait for the soft mass-and-spring system with m = 1, k = 4, and $\beta = 1 > 0$. The separatrices are emphasized.



The unbounded trajectories opening to the right and left correspond to values of E less than 4. These represent motions in which the mass initially is headed toward the equilibrium point x = 0, but with insufficient energy to reach it. At some point the mass reverses direction and heads back whence it came.

In Fig. 9.4.4 it appears that the critical point (0, 0) is a stable center, whereas the critical points $(\pm 2, 0)$ look like saddle points of the equivalent first-order system

$$x' = y, \quad y' = -4x + x^3 \tag{11}$$

with Jacobian matrix

$$\mathbf{J}(x,y) = \begin{bmatrix} 0 & 1\\ -4+3x^2 & 0 \end{bmatrix}.$$

To check these observations against the usual critical-point analysis, we note first that the Jacobian matrix

$$\mathbf{J}(0,0) = \begin{bmatrix} 0 & 1\\ -4 & 0 \end{bmatrix}$$

at the critical point (0, 0) has characteristic equation $\lambda^2 + 4 = 0$ and pure imaginary eigenvalues $\lambda_1, \lambda_2 = \pm 2i$ consistent with a stable center. Moreover, the Jacobian matrix

$$\mathbf{J}(\pm 2,0) = \begin{bmatrix} 0 & 1\\ 8 & 0 \end{bmatrix}$$

corresponding to the other two critical points has characteristic equation $\lambda^2 - 8 = 0$ and real eigenvalues λ_1 , $\lambda_2 = \pm \sqrt{8}$ of opposite sign, consistent with the saddle-point behavior that we observe near (-2, 0) and (+2, 0).

Remark Figures 9.4.2 and 9.4.4 illustrate a significant qualitative difference between hard springs with $\beta < 0$ and soft springs with $\beta > 0$ in the nonlinear equation $mx'' = kx + \beta x^3$. Whereas the phase plane trajectories for a hard spring are all bounded, a soft spring has unbounded phase plane trajectories (as well as bounded ones). However, we should realize that the unbounded soft-spring trajectories cease to represent physically realistic motions faithfully when they exceed the spring's capability of expansion without breaking.

Damped Nonlinear Vibrations

Suppose now that the mass on a spring is connected also to a dashpot that provides a force of resistance proportional to the velocity y = dx/dt of the mass. If the spring is still assumed nonlinear as in Eq. (2), then the equation of motion of the mass is

$$mx'' = -cx' - kx + \beta x^3, \tag{12}$$

where c > 0 is the resistance constant. If $\beta > 0$, then the equivalent first-order system

$$\frac{dx}{dt} = y, \quad \frac{dy}{dt} = \frac{-kx - cy + \beta x^3}{m} = -\frac{c}{m}y - \frac{k}{m}x\left(1 - \frac{\beta}{k}x^2\right) \tag{13}$$

has critical points (0, 0) and $\left(\pm \sqrt{k/\beta}, 0\right)$ and Jacobian matrix

$$\mathbf{J}(x, y) = \begin{bmatrix} 0 & 1\\ -\frac{k}{m} + \frac{3\beta}{m}x^2 & -\frac{c}{m} \end{bmatrix}.$$

Now the critical point at the origin is the most interesting one. The Jacobian matrix

$$\mathbf{J}(0,0) = \begin{bmatrix} 0 & 1\\ -\frac{k}{m} & -\frac{c}{m} \end{bmatrix}$$

has characteristic equation

$$(-\lambda)\left(-\frac{c}{m}-\lambda\right) + \frac{k}{m} = \frac{1}{m}(m\lambda^2 + c\lambda + k) = 0$$

and eigenvalues

$$\lambda_1, \, \lambda_2 = \frac{-c \pm \sqrt{c^2 - 4km}}{2m}.$$

It follows from Theorem 2 in Section 9.2 in that the critical point (0, 0) of the system in (13) is

- a nodal sink if the resistance is so great that $c^2 > 4km$ (in which case the eigenvalues are negative and unequal), but is
- a spiral sink if $c^2 < 4km$ (in which case the eigenvalues are complex conjugates with negative real part).

The following example illustrates the latter case. (In the borderline case with equal negative eigenvalues, the origin may be either a nodal or a spiral sink.)

Example 2 Damped soft spring Suppose that
$$m = 1, c = 2, k = 5$$
, and $\beta = \frac{5}{4}$. Then the nonlinear system in (13) is

$$\frac{dx}{dt} = y, \quad \frac{dy}{dt} = -5x - 2y + \frac{5}{4}x^3 = -2y - 5x(1 - \frac{1}{4}x^2). \tag{14}$$

It has critical points (0, 0), $(\pm 2, 0)$ and Jacobian matrix

$$\mathbf{J}(x,y) = \begin{bmatrix} 0 & 1\\ -5 + \frac{15}{4}x^2 & -2 \end{bmatrix}$$

At (0, 0): The Jacobian matrix

$$\mathbf{J}(0,0) = \begin{bmatrix} 0 & 1\\ -5 & -2 \end{bmatrix}$$

has characteristic equation $\lambda^2 + 2\lambda + 5 = 0$ and has complex conjugate eigenvalues λ_1 , $\lambda_2 = -1 \pm 2i$ with negative real part. Hence (0, 0) is a spiral sink of the nonlinear system in (14), and the linearized position function of the mass is of the form

$$x(t) = e^{-t} (A\cos 2t + B\sin 2t)$$

that corresponds to an exponentially damped oscillation about the equilibrium position x = 0. At ($\pm 2, 0$): The Jacobian matrix

$$\mathbf{J}(\pm 2,0) = \begin{bmatrix} 0 & 1\\ 10 & -2 \end{bmatrix}$$

has characteristic equation $\lambda^2 + 2\lambda - 10 = 0$ and real eigenvalues $\lambda_1 = -1 - \sqrt{11} < 0$ and $\lambda_2 = -1 + \sqrt{11} > 0$ with different signs. It follows that (-2, 0) and (+2, 0) are both saddle points of the system in (14).

The position-velocity phase plane portrait in Fig. 9.4.6 shows trajectories of (14) and the spiral sink at (0,0), as well as the unstable saddle points at (-2,0) and (2,0). The emphasized separatrices divide the phase plane into regions of different behavior. The behavior of the mass depends on the region in which its initial point (x_0 , y_0) is located. If this initial point lies in

- Region I between the separatrices, then the trajectory spirals into the origin as t → +∞, and hence the periodic oscillations of the undamped case (Fig. 9.4.4) are now replaced with damped oscillations around the stable equilibrium position x = 0;
- Region II, then the mass passes through x = 0 moving from left to right (x increasing);
- Region III, then the mass passes through x = 0 moving from right to left (x decreasing);
- Region IV, then the mass approaches (but does not reach) the unstable equilibrium position x = -2 from the left, but stops and then returns to the left;
- Region V, then the mass approaches (but does not reach) the unstable equilibrium position *x* = 2 from the right, but stops and then returns to the right.



FIGURE 9.4.6. Position–velocity phase plane portrait for the soft mass-and-spring system with $m = 1, k = 5, \beta = \frac{5}{4}$, and resistance constant c = 2. The (black) separatrices are emphasized.

If the initial point (x_0, y_0) lies precisely on one of the separatrices, then the corresponding trajectory either approaches the stable spiral point or recedes to infinity from a saddle point as $t \to +\infty$.

The Nonlinear Pendulum

In Section 5.4 we derived the equation

$$\frac{d^2\theta}{dt^2} + \frac{g}{L}\sin\theta = 0 \tag{15}$$

for the undamped oscillations of the simple pendulum shown in Fig. 9.4.7. There we used the approximation $\sin \theta \approx \theta$ for θ near zero to replace Eq. (15) with the linear model

$$\frac{d^2\theta}{dt^2} + \omega^2\theta = 0, \tag{16}$$

where $\omega^2 = g/L$. The general solution

$$\theta(t) = A\cos\omega t + B\sin\omega t \tag{17}$$

of Eq. (16) describes oscillations around the equilibrium position $\theta = 0$ with circular frequency ω and amplitude $C = (A^2 + B^2)^{1/2}$.

The linear model does not adequately describe the possible motions of the pendulum for large values of θ . For instance, the equilibrium solution $\theta(t) \equiv \pi$ of Eq. (15), with the pendulum standing straight up, does not satisfy the linear equation in (16). Nor does Eq. (17) include the situation in which the pendulum "goes over the top" repeatedly, so that $\theta(t)$ is a steadily increasing rather than an oscillatory function of t. To investigate these phenomena we must analyze the nonlinear equation $\theta'' + \omega^2 \theta = 0$. We also want to include the possibility of resistance proportional to velocity, so we consider the general *nonlinear pendulum equation*



FIGURE 9.4.7. The simple pendulum.



9.4 NONLINEAR MECHANICAL SYSTEMS545

The case c > 0 corresponds to damped motion in which there actually *is* resistance proportional to (angular) velocity. But we examine first the undamped case in which c = 0. With $x(t) = \theta(t)$ and $y(t) = \theta'(t)$ the equivalent first-order system is

$$\frac{dx}{dt} = y, \quad \frac{dy}{dt} = -\omega^2 \sin x. \tag{19}$$

We see that this system is almost linear by writing it in the form

$$\frac{dx}{dt} = y,$$
(20)
$$\frac{dy}{dt} = -\omega^2 x + g(x),$$

where

$$g(x) = -\omega^2(\sin x - x) = \omega^2\left(\frac{x^3}{3!} - \frac{x^5}{5!} + \cdots\right)$$

has only higher-degree terms.

The critical points of the system in (19) are the points $(n\pi, 0)$ with *n* an integer, and its Jacobian matrix is given by

$$\mathbf{J}(x,y) = \begin{bmatrix} 0 & 1\\ -\omega^2 \cos x & 0 \end{bmatrix}.$$
 (21)

The nature of the critical point $(n\pi, 0)$ depends on whether *n* is even or odd.

EVEN CASE: If n = 2m is an even integer, then $\cos n\pi = +1$, so (21) yields the matrix

$$\mathbf{J}(2m\pi,0) = \begin{bmatrix} 0 & 1\\ -\omega^2 & 0 \end{bmatrix}$$

with characteristic equation $\lambda^2 + \omega^2 = 0$ and pure imaginary eigenvalues $\lambda_1, \lambda_2 = \pm \omega i$. The linearization of (19) at $(n\pi, 0)$ is therefore the system

$$\frac{du}{dt} = v, \quad \frac{dv}{dt} = -\omega^2 u \tag{22}$$

for which (0, 0) is the familiar stable center enclosed by elliptical trajectories (as in Example 3 of Section 9.1). Although this is the delicate case for which Theorem 2 of Section 9.2 does not settle the matter, we will see presently that $(2m\pi, 0)$ is also a stable center for the original nonlinear pendulum system in (19).

ODD CASE: If n = 2m + 1 is an odd integer, then $\cos n\pi = -1$, so (21) yields the matrix

$$\mathbf{J}((2m+1)\pi,0) = \begin{bmatrix} 0 & 1\\ \omega^2 & 0 \end{bmatrix}$$

with characteristic equation $\lambda^2 - \omega^2 = 0$ and real eigenvalues $\lambda_1, \lambda_2 = \pm \omega$ with different signs. The linearization of (19) at $((2m + 1)\pi, 0)$ is therefore the system

$$\frac{du}{dt} = v, \quad \frac{dv}{dt} = \omega^2 u \tag{23}$$

for which (0, 0) is a saddle point. It follows from Theorem 2 of Section 9.2 that the critical point $((2m + 1)\pi, 0)$ is a similar saddle point for the original nonlinear pendulum system in (19).

THE TRAJECTORIES: We can see how these "even centers" and "odd saddle points" fit together by solving the system in (19) explicitly for the phase plane trajectories. If we write

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = -\frac{\omega^2 \sin x}{y}$$

and separate the variables,

$$y\,dy + \omega^2 \sin x\,dx = 0,$$

then integration from x = 0 to x = x yields

$$\frac{1}{2}y^2 + \omega^2(1 - \cos x) = E.$$
(24)

We write *E* for the arbitrary constant of integration because, if physical units are so chosen that m = L = 1, then the first term on the left is the kinetic energy and the second term the potential energy of the mass on the end of the pendulum. Then *E* is the total mechanical energy; Eq. (24) thus expresses conservation of mechanical energy for the undamped pendulum.

If we solve Eq. (24) for y and use a half-angle identity, we get the equation

$$y = \pm \sqrt{2E - 4\omega^2 \sin^2 \frac{1}{2}x}$$
 (25)

that defines the phase plane trajectories. Note that the radicand in (25) remains positive if $E > 2\omega^2$. Figure 9.4.8 shows (along with a direction field) the results of plotting these trajectories for various values of the energy *E*.



FIGURE 9.4.8. Position–velocity phase plane portrait for the undamped pendulum system x' = y, $y' = -\sin x$. The (black) separatrices are emphasized.

The emphasized separatrices in Fig. 9.4.8 correspond to the critical value $E = 2\omega^2$ of the energy; they enter and leave the unstable critical points $(n\pi, 0)$ with n an odd integer. Following the arrows along a separatrix, the pendulum theoretically approaches a balanced vertical position $\theta = x = (2m + 1)\pi$ with just enough energy to reach it but not enough to "go over the top." The instability of this equilibrium position indicates that this behavior may never be observed in practice!

9.4 NONLINEAR MECHANICAL SYSTEMS547

The simple closed trajectories encircling the stable critical points—all of which correspond to the downward position $\theta = 2m\pi$ of the pendulum—represent periodic oscillations of the pendulum back and forth around the stable equilibrium position $\theta = 0$. These correspond to energies $E < 2\omega^2$ that are insufficient for the pendulum to ascend to the vertical upward position—so its back-and-forth motion is that which we normally associate with a "swinging pendulum."

The unbounded trajectories with $E > 2\omega^2$ represent whirling motions of the pendulum in which it goes over the top repeatedly—in a clockwise direction if y(t) remains positive, in a counterclockwise direction if y(t) is negative.

Period of Undamped Oscillation

If the pendulum is released from rest with initial conditions

$$x(0) = \theta(0) = \alpha, \quad y(0) = \theta'(0) = 0,$$
 (26)

then Eq. (24) with t = 0 reduces to

$$\omega^2 (1 - \cos \alpha) = E. \tag{27}$$

Hence $E < 2\omega^2$ if $0 < \alpha < \pi$, so a periodic oscillation of the pendulum ensues. To determine the *period* of this oscillation, we subtract Eq. (27) from Eq. (24) and write the result (with $x = \theta$ and $y = d\theta/dt$) in the form

$$\frac{1}{2} \left(\frac{d\theta}{dt}\right)^2 = \omega^2 (\cos\theta - \cos\alpha). \tag{28}$$

The period *T* of time required for one complete oscillation is four times the amount of time required for θ to decrease from $\theta = \alpha$ to $\theta = 0$, one-fourth of an oscillation. Hence we solve Eq. (28) for $dt/d\theta$ and integrate to get

$$T = \frac{4}{\omega\sqrt{2}} \int_0^\alpha \frac{d\theta}{\sqrt{\cos\theta - \cos\alpha}}.$$
 (29)

To attempt to evaluate this integral we first use the identity $\cos \theta = 1 - 2 \sin^2(\theta/2)$ and get

$$T = \frac{2}{\omega} \int_0^\alpha \frac{d\theta}{\sqrt{k^2 - \sin^2(\theta/2)}}$$

where

$$k = \sin \frac{\alpha}{2}$$
.

Next, the substitution $u = (1/k) \sin(\theta/2)$ yields

$$T = \frac{4}{\omega} \int_0^1 \frac{du}{\sqrt{(1 - u^2)(1 - k^2 u^2)}}$$

Finally, the substitution $u = \sin \phi$ gives

$$T = \frac{4}{\omega} \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}}.$$
 (30)

The integral in (30) is the *elliptic integral of the first kind* that is often denoted by $F(k, \pi/2)$. Whereas elliptic integrals normally cannot be evaluated in closed form, this integral can be approximated numerically as follows. First we use the binomial series

$$\frac{1}{\sqrt{1-x}} = 1 + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdots (2n-1)}{2 \cdot 4 \cdots (2n)} x^n$$
(31)

with $x = k^2 \sin^2 \phi < 1$ to expand the integrand in (30). Then we integrate termwise using the tabulated integral formula

$$\int_0^{\pi/2} \sin^{2n} \phi \, d\phi = \frac{\pi}{2} \cdot \frac{1 \cdot 3 \cdots (2n-1)}{2 \cdot 4 \cdots (2n)}.$$
(32)

The final result is the formula

$$T = \frac{2\pi}{\omega} \left[1 + \sum_{n=1}^{\infty} \left(\frac{1 \cdot 3 \cdots (2n-1)}{2 \cdot 4 \cdots (2n)} \right)^2 k^{2n} \right]$$

= $T_0 \left[1 + \left(\frac{1}{2} \right)^2 k^2 + \left(\frac{1 \cdot 3}{2 \cdot 4} \right)^2 k^4 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \right)^2 k^6 + \cdots \right]$ (33)

for the period *T* of the nonlinear pendulum released from rest with initial angle $\theta(0) = \alpha$, in terms of the linearized period $T_0 = 2\pi/\omega$ and $k = \sin(\alpha/2)$.

The infinite series within the second pair of brackets in Eq. (33) gives the factor T/T_0 by which the nonlinear period T is longer than the linearized period. The table in Fig. 9.4.9, obtained by summing this series numerically, shows how T/T_0 increases as α is increased. Thus T is 0.19% greater than T_0 if $\alpha = 10^\circ$, whereas T is 18.03% greater than T_0 if $\alpha = 90^\circ$. But even a 0.19% discrepancy is significant—the calculation

$$(0.0019) \times 3600 \frac{\text{seconds}}{\text{hour}} \times 24 \frac{\text{hours}}{\text{day}} \times 7 \frac{\text{days}}{\text{week}} \approx 1149 \text{ (seconds/week)}$$

shows that the linearized model is quite inadequate for a pendulum clock; a discrepancy of 19 min 9 s after only one week is unacceptable.

Damped Pendulum Oscillations

Finally, we discuss briefly the *damped* nonlinear pendulum. The almost linear first-order system equivalent to Eq. (19) is

$$\frac{dx}{dt} = y,$$

$$\frac{dy}{dt} = -\omega^2 \sin x - cy,$$
(34)

and again the critical points are of the form $(n\pi, 0)$ where *n* is an integer. In Problems 9 through 11 we ask you to verify that

- If n is odd, then (nπ, 0) is an unstable saddle point of (34), just as in the undamped case; but
- If *n* is even and $c^2 > 4\omega^2$, then $(n\pi, 0)$ is a nodal sink; whereas
- If *n* is even and $c^2 < 4\omega^2$, then $(n\pi, 0)$ is a spiral sink.

α	T/T_0
10°	1.0019
20°	1.0077
30°	1.0174
40°	1.0313
50°	1.0498
60°	1.0732
70°	1.1021
80°	1.1375
90°	1.1803

FIGURE 9.4.9. Dependence of the period *T* of a nonlinear pendulum on its initial angle α .

9.4 NONLINEAR MECHANICAL SYSTEMS549

Figure 9.4.10 illustrates the phase plane trajectories for the more interesting underdamped case, $c^2 < 4\omega^2$. Other than the physically unattainable separatrix trajectories that enter unstable saddle points, every trajectory eventually is "trapped" by one of the stable spiral points $(n\pi, 0)$ with *n* an even integer. What this means is that even if the pendulum starts with enough energy to go over the top, after a certain (finite) number of revolutions it has lost enough energy that thereafter it undergoes damped oscillations around its stable (lower) equilibrium position.



FIGURE 9.4.10. Position–velocity phase plane portrait for the damped pendulum system x' = y, $y' = -\sin x - \frac{1}{4}y$. The (black) separatrices are emphasized.

9.4 **Problems**

In Problems 1 through 4, show that the given system is almost linear with (0, 0) as a critical point, and classify this critical point as to type and stability. Use a computer system or graphing calculator to construct a phase plane portrait that illustrates your conclusion.

1. $\frac{dx}{dt} = 1 - e^x + 2y, \frac{dy}{dt} = -x - 4\sin y$ 2. $\frac{dx}{dt} = 2\sin x + \sin y, \frac{dy}{dt} = \sin x + 2\sin y$ (Fig. 9.4.11)



FIGURE 9.4.11. Trajectories of the system in Problem 2.

4.
$$\frac{dx}{dt} = \sin x \cos y - 2y, \frac{dy}{dt} = 4x - 3\cos x \sin y$$

Find and classify each of the critical points of the almost linear systems in Problems 5 through 8. Use a computer system or graphing calculator to construct a phase plane portrait that illustrates your findings.

5.
$$\frac{dx}{dt} = -x + \sin y, \frac{dy}{dt} = 2x$$

6.
$$\frac{dx}{dt} = y, \frac{dy}{dt} = \sin \pi x - y$$

7.
$$\frac{dx}{dt} = 1 - e^{x-y}, \frac{dy}{dt} = 2\sin x$$

8.
$$\frac{dx}{dt} = -2\sin x + y, \frac{dy}{dt} = -2\sin x + y$$

8.
$$\frac{dx}{dt} = 3\sin x + y, \frac{dy}{dt} = \sin x + 2y$$

Critical Points for Damped Pendulum

Problems 9 through 11 deal with the damped pendulum system $x' = y, y' = -\omega^2 \sin x - cy.$

- **9.** Show that if *n* is an odd integer, then the critical point $(n\pi, 0)$ is a saddle point for the damped pendulum system.
- 10. Show that if *n* is an even integer and $c^2 > 4\omega^2$, then the critical point $(n\pi, 0)$ is a nodal sink for the damped pendulum system.

3.
$$\frac{dx}{dt} = e^x + 2y - 1, \frac{dy}{dt} = 8x + e^y - 1$$

11. Show that if n is an even integer and $c^2 < 4\omega^2$, then the critical point $(n\pi, 0)$ is a spiral sink for the damped pendulum system.

Critical Points for Mass-Spring System

In each of Problems 12 through 16, a second-order equation of the form x'' + f(x, x') = 0, corresponding to a certain mass-and-spring system, is given. Find and classify the critical points of the equivalent first-order system.

- 12. $x'' + 20x 5x^3 = 0$: Verify that the critical points resemble those shown in Fig. 9.4.4.
- 13. $x'' + 2x' + 20x 5x^{3} = 0$: Verify that the critical points resemble those shown in Fig. 9.4.6.
- 14. $x'' 8x + 2x^3 = 0$: Here the linear part of the force is repulsive rather than attractive (as for an ordinary spring). Verify that the critical points resemble those shown in Fig. 9.4.12. Thus there are two stable equilibrium points and three types of periodic oscillations.
- 15. $x'' + 4x x^2 = 0$: Here the force function is nonsymmetric. Verify that the critical points resemble those shown in Fig. 9.4.13.
- **16.** $x'' + 4x 5x^3 + x^5 = 0$: The idea here is that terms through the fifth degree in an odd force function have been retained. Verify that the critical points resemble those shown in Fig. 9.4.14.

Critical Points for Physical Systems

In Problems 17 through 20, analyze the critical points of the indicated system, use a computer system to construct an illustrative position-velocity phase plane portrait, and describe the oscillations that occur.

- 17. Example 2 in this section illustrates the case of damped vibrations of a soft mass-spring system. Investigate an example of damped vibrations of a hard mass-spring system by using the same parameters as in Example 2, except now with $\beta = -\frac{5}{4} < 0$.
- 18. Example 2 illustrates the case of damped vibrations of a soft mass-spring system with the resistance proportional to the velocity. Investigate an example of resistance proportional to the square of the velocity by using the same parameters as in Example 2, but with resistance term -cx'|x'| instead of -cx' in Eq. (12).
- 19. Now repeat Example 2 with both the alterations corresponding to Problems 17 and 18. That is, take $\beta = -\frac{5}{4} < 0$ and replace the resistance term in Eq. (12) with -cx'|x'|.
- **20.** The equations x' = y, $y' = -\sin x \frac{1}{4}y|y|$ model a damped pendulum system as in Eqs. (34) and Fig. 9.4.10. But now the resistance is proportional to the square of the angular velocity of the pendulum. Compare the oscillations with those that occur when the resistance is proportional to the angular velocity itself.

Period of Oscillation

Problems 21 through 26 outline an investigation of the period T of oscillation of a mass on a nonlinear spring with equation of motion

$$\frac{d^2x}{dt^2} + \phi(x) = 0.$$
 (35)

If $\phi(x) = kx$ with k > 0, then the spring actually is linear with period $T_0 = 2\pi/\sqrt{k}$.



for Problem 14.

FIGURE 9.4.12. The phase portrait



FIGURE 9.4.13. The phase portrait for Problem 15.



FIGURE 9.4.14. The phase portrait for Problem 16.

9.4 NONLINEAR MECHANICAL SYSTEMS551

21. Integrate once (as in Eq. (6)) to derive the energy equation

$$\frac{1}{2}y^2 + V(x) = E,$$
(36)

where y = dx/dt and

$$V(x) = \int_0^x \phi(u) \, du. \tag{37}$$

22. If the mass is released from rest with initial conditions $x(0) = x_0$, y(0) = 0 and periodic oscillations ensue, conclude from Eq. (36) that $E = V(x_0)$ and that the time *T* required for one complete oscillation is

$$T = \frac{4}{\sqrt{2}} \int_0^{x_0} \frac{du}{\sqrt{V(x_0) - V(u)}}.$$
 (38)

23. If $\phi(x) = kx - \beta x^3$ as in the text, deduce from Eqs. (37) and (38) that

$$T = 4\sqrt{2} \int_0^{x_0} \frac{dx}{\sqrt{(x_0^2 - u^2)(2k - \beta x_0^2 - \beta u^2)}}.$$
 (39)

24. Substitute $u = x_0 \cos \phi$ in (39) to show that

$$T = \frac{2T_0}{\pi\sqrt{1-\epsilon}} \int_0^{\pi/2} \frac{d\phi}{\sqrt{1-\mu\sin^2\phi}},\qquad(40)$$

where $T_0 = 2\pi/\sqrt{k}$ is the linear period,

$$\epsilon = \frac{\beta}{k} x_0^2$$
, and $\mu = -\frac{1}{2} \cdot \frac{\epsilon}{1 - \epsilon}$. (41)

25. Finally, use the binomial series in (31) and the integral formula in (32) to evaluate the elliptic integral in (40) and thereby show that the period *T* of oscillation is given by

$$T = \frac{T_0}{\sqrt{1-\epsilon}} \left(1 + \frac{1}{4}\mu + \frac{9}{64}\mu^2 + \frac{25}{256}\mu^3 + \cdots \right).$$
(42)

26. If $\epsilon = \beta x_0^2/k$ is sufficiently small that ϵ^2 is negligible, deduce from Eqs. (41) and (42) that

$$T \approx T_0 \left(1 + \frac{3}{8} \epsilon \right) = T_0 \left(1 + \frac{3\beta}{8k} x_0^2 \right).$$
(43)

It follows that

- If β > 0, so the spring is *soft*, then T > T₀, and increasing x₀ increases T, so the larger ovals in Fig. 9.4.4 correspond to smaller frequencies.
- If $\beta < 0$, so the spring is *hard*, then $T < T_0$, and increasing x_0 decreases T, so the larger ovals in Fig. 9.4.2 correspond to larger frequencies.

9.4 Application The Rayleigh, van der Pol, and FitzHugh-Nagumo Equations

Go to goo.gl/hlwIzu to download this application's computing resources including *Maple/Mathematica*/MATLAB. Here we present a trio of nonlinear differential equations or systems of equations, drawn from the areas of acoustics, electrical engineering, and neuroscience. Each of these models has been fundamental within its field; taken together, they give some indication of the importance of nonlinear equations across a wide variety of applications.

Rayleigh's Equation

The British mathematical physicist Lord Rayleigh (John William Strutt, 1842–1919) introduced an equation of the form

$$mx'' + kx = ax' - b(x')^3$$
(1)

to model the oscillations of a clarinet reed. With y = x' we get the autonomous system

$$x' = y,$$

$$y' = \frac{-kx + ay - by^3}{m},$$
(2)

whose phase plane portrait is shown in Fig. 9.4.15 (for the case m = k = a = b = 1). The outward and inward spiral trajectories converge to a "limit cycle" solution that corresponds to periodic oscillations of the reed. The period *T* (and hence the frequency) of these oscillations can be measured on a *tx*-solution curve plotted as in Fig. 9.4.16. This period of oscillation depends only on the parameters *m*, *k*, *a*, and *b* in Eq. (1) and is independent of the initial conditions (why?).



2.0 1.5 1.0 0.5 × 0.0 -0.5-1.0-1.5-2.00 5 10 15 20 25 30 35 40

FIGURE 9.4.15. Phase plane portrait for the Rayleigh system in (2) with m = k = a = b = 1.

FIGURE 9.4.16. The *tx*-solution curve with initial conditions x(0) = 0.01, x'(0) = 0.

Choose your own parameters m, k, a, and b (perhaps the least four nonzero digits in your student ID number), and use an available ODE plotting utility to plot trajectories and solution curves as in Figs. 9.4.15 and 9.4.16. Change *one* of your parameters to see how the amplitude and frequency of the resulting periodic oscillations are altered.

Van der Pol's Equation

Figure 9.4.17 shows a simple *RLC* circuit in which the usual (passive) resistance *R* has been replaced with an active element (such as a vacuum tube or semiconductor) across which the voltage drop *V* is given by a known function f(I) of the current *I*. Of course, V = f(I) = IR for a resistor. If we substitute f(I) for *IR* in the well-known *RLC*-circuit equation LI' + RI + Q/C = 0, then differentiation gives the second-order equation

$$LI'' + f'(I)I' + \frac{I}{C} = 0.$$
 (3)

In a 1924 study of oscillator circuits in early commercial radios, Balthasar van der Pol (1889–1959) assumed the voltage drop to be given by a nonlinear function of the form $f(I) = bI^3 - aI$, which with Eq. (3) becomes

$$LI'' + (3bI^2 - a)I' + \frac{I}{C} = 0.$$
 (4)

This equation is closely related to Rayleigh's equation and has phase portraits resembling Fig. 9.4.15. Indeed, differentiation of the second equation in (2) and the resubstitution x' = y yield the equation

$$my'' + (3by^2 - a)y' + ky = 0,$$
(5)

which has the same form as Eq. (4).

If we denote by τ the time variable in Eq. (4) and make the substitutions I = px, $t = \tau/\sqrt{LC}$, the result is

$$\frac{d^2x}{dt^2} + \left(3bp^2x^2 - a\right)\sqrt{\frac{C}{L}}\frac{dx}{dt} + x = 0.$$



FIGURE 9.4.17. A simple circuit with an active element.

9.4 NONLINEAR MECHANICAL SYSTEMS553

With $p = \sqrt{a/(3b)}$ and $\mu = a\sqrt{C/L}$, this gives the standard form

$$x'' + \mu(x^2 - 1)x' + x = 0$$
(6)

of van der Pol's equation.

For every nonnegative value of the parameter μ , the solution of van der Pol's equation with x(0) = 2, x'(0) = 0 is periodic, and the corresponding phase plane trajectory is a limit cycle to which the other trajectories converge (as in Fig. 9.4.15). It will be instructive for you to solve van der Pol's equation numerically and to plot this periodic trajectory for a selection of values from $\mu = 0$ to $\mu = 1000$ or more. With $\mu = 0$ it is a circle of radius 2 (why?). Figure 9.4.18 shows the periodic trajectory with $\mu = 1$, and Fig. 9.4.19 shows the corresponding x(t) and y(t) solution curves. When μ is large, van der Pol's equation is quite "stiff" and the periodic trajectory is more eccentric as in Fig. 9.4.20, which was plotted using MAT-LAB's stiff ODE solver ode15s. The corresponding x(t) and y(t) solution curves in Figs. 9.4.21 and 9.4.22 reveal surprising behavior of these component functions. Each alternates long intervals of very slow change with periods of abrupt change during very short time intervals that correspond to the "quasi-discontinuities" that are visible in Figs. 9.4.21 and 9.4.22. For instance, Fig. 9.4.23 shows that, between t = 1614.28 and t = 1614.29, the value of y(t) zooms from near zero to over 1300 and back again. Perhaps you can measure the distance between x- or y-intercepts



FIGURE 9.4.18. The phase plane trajectory of a periodic solution of van der Pol's equation with $\mu = 1$, as well as some trajectories spiraling in and out.



FIGURE 9.4.19. x(t) and y(t) solution curves defining the periodic solution of van der Pol's equation with $\mu = 1$.



FIGURE 9.4.20. The phase plane trajectory of the periodic solution of van der Pol's equation with $\mu = 1000$.



FIGURE 9.4.21. Graph of x(t) with $\mu = 1000$.



FIGURE 9.4.22. Graph of y(t) with $\mu = 1000$.



FIGURE 9.4.23. The upper spike in the graph of y(t).

to show that the period of circuit around the cycle in Fig. 9.4.20 is approximately T = 1614. Indeed, this calculation and the construction of figures like those shown here may serve as a good test of the robustness of your computer system's ODE solver.

You might also plot other trajectories for $\mu = 10$, 100 or 1000 that (like the trajectories in Fig. 9.4.18) are "attracted" from within and without by the limit cycle. The origin looks like a spiral point in Fig 9.4.18. Indeed, show that (0, 0) is a spiral source for van der Pol's equation if $0 < \mu < 2$ but is a nodal source if $\mu \ge 2$.

The FitzHugh-Nagumo Equations

Since the early experiments of Luigi Galvani (1737–1798) in which electrical stimulus caused the leg muscles of dead frogs to twitch, the electrical properties of *neurons*, the cells that form the building blocks of the nervous system, have been intensively studied. One of the most important of those properties is the *action potential*, an electrical signal that travels from the body of a neuron down along its axon (Fig. 9.4.24). Action potentials are the units of information of the nervous system; when an action potential reaches the end of the axon, chemicals known as neurotransmitters are released from the axon terminals. These neurotransmitters then find receptors in the dendrites of other nerve cells, causing action potentials in those "target" neurons, and thus propagating the "message." Because of the great speed with which action potentials traverse the neuron, they provide a mechanism by which signals can be rapidly transmitted through the nervous system.

Action potentials are particularly known for their *all-or-none* character. If the stimulus received by a target neuron is below a certain threshold, then no action potential is generated. If this threshold is exceeded, however, then the neuron will "fire" an action potential, or perhaps (if the stimulus is sufficiently strong) several action potentials in succession. In this way, the nervous system's method of electrical signaling resembles the binary code used by computers.



FIGURE 9.4.24. The structure of a typical neuron.

In the early 1950's A. F. Huxley (1917–2012) and A. L. Hodgkin (1914–1998) published a landmark series of papers in which they modeled the action potential in the giant axon of the squid as an electrical circuit. The focus of this model was the neuron's *membrane potential*, that is, the voltage difference between the inside and outside of the nerve cell. In its resting state, a typical neuron has a negative membrane potential, that is, the inside of the cell is at a lower voltage than the surrounding medium. (We now know that this voltage difference is largely due to "ion pumps" in the cell membrane, which maintain a lower concentration of positively charged sodium ions inside the cell than in the surrounding medium. These pumps require energy, and indeed a significant portion of the body's metabolic energy is devoted to this task.) Potassium ions also play an important role in neuron electrical activity.

During an action potential, the membrane potential exhibits a characteristic pattern of sudden and rapid changes through positive and negative values as the electrical signal traverses the neuron. A central goal of Hodgkin and Huxley's work was to explain these changes in terms of the sodium and potassium *conductances* of the neuron membrane. Conductance—the reciprocal of resistance—is a measure of the permeability of the membrane to charged ions. An increase in the sodium conductance, for example, allows sodium ions to flow more freely across the membrane, from areas of high concentration to low. Hodgkin and Huxley proposed that during an action potential, an electrical stimulus (arising from another neuron "upstream," for example) causes changes in the neuron membrane's sodium and potassium conductances. This results in a series of flows of charged ions, and thus electrical currents, across the cell membrane.

The researchers applied some of the basic principles of circuit theory to model these currents. The result was a system of four nonlinear differential equations, whose variables are the neuron membrane potential together with three other quantities related to the membrane's sodium and potassium conductances. Not only did the predictions of this model show remarkable accord with experimental results, they also helped point the way to subsequent discoveries in neurophysiology. The Hodgkin-Huxley model was a triumph both of experimental technique and theoretical analysis, and remains today the starting point for mathematical modeling of action potentials. Together with John Eccles, Hodgkin and Huxley were awarded the 1963 Nobel Prize in Physiology or Medicine for their work.

Analysis of the Hodgkin-Huxley model can be challenging, however, because its phase space is four-dimensional, making features such as solution curves difficult to visualize. For this reason, in 1961 Richard FitzHugh (1922–2007) proposed a two-dimensional simplification of the Hodgkin-Huxley model, which was subsequently analyzed in electrical circuit terms by J. Nagumo and others. Whereas the *FitzHugh-Nagumo equations* are not intended to capture the physiological properties of the neuron as directly as the original Hodgkin-Huxley equations do, this simplified model is important because it displays much of the *qualitative* behavior characteristic of neuron electrical activity, while offering the advantage of being considerably easier to study. FitzHugh's model actually is a generalization of van der Pol's equation (6). You can show that introduction of the variable $y = \frac{1}{\mu}x' + \frac{1}{3}x^3 - x$ in van der Pol's equation (which results in better phase plane analysis than does simply taking y = x') leads to the system

$$\begin{aligned} x' &= \mu \left(y + x - \frac{1}{3} x^3 \right), \\ y' &= -\frac{1}{\mu} x, \end{aligned}$$

which FitzHugh generalized by adding terms:

$$x' = \mu \left(y + x - \frac{1}{3}x^3 + I \right),$$

$$y' = -\frac{1}{\mu}(x - a + by).$$
(7)

Here *a* and *b* are constants and *I* is a function of the time *t*. Loosely speaking, x(t) behaves in a manner similar to the neuron membrane potential, I(t) is the electrical stimulus applied to the neuron, and y(t) is a composite of the other three variables in the Hodgkin-Huxley model. To simulate neuron electrical activity, we will use

FitzHugh's values a = 0.7, b = 0.8, and $\mu = 3$, while assigning various constant values to the stimulus I(t).

First, with $I(t) \equiv 0$ (corresponding to the resting state of the neuron), you can verify that the system (7) has exactly one equilibrium point at roughly x = 1.1994and y = -0.6243. What is striking is the way in which the system responds as *I* becomes nonzero (corresponding to electrical stimulation of the neuron). Figure 9.4.25 shows the solution curves of the system corresponding to the three constant values $I(t) \equiv -0.15$, $I(t) \equiv -0.17$, and $I(t) \equiv -0.5$. The graph on the left gives the phase plane for x and y, whereas the graph on the right shows x(t) as a function of t. All curves begin at time t = 0 at the original equilibrium point (1.1994, -0.6243) indicated in the graph on the left.



FIGURE 9.4.25. Solutions of the FitzHugh-Nagumo equations (7) with three constant values of the electrical stimulus I(t), using a = 0.7, b = 0.8, and $\mu = 3$.

When the stimulus I(t) is held at the constant value -0.15, x (which is analogous to the neuron membrane potential) varies slightly from its rest value, but soon finds a new "perturbed" equilibrium at which it then remains (thin blue curves); this mirrors the behavior of the membrane potential when the neuron receives a stimulus insufficiently large to generate an action potential. The response of the system differs dramatically, however, when the constant stimulus is decreased slightly to -0.17; once again x finds a new equilibrium, but only after exhibiting wide swings downward and back upward (dashed curves); this is suggestive of the firing of a single action potential. Finally, when $I(t) \equiv -0.5$, x oscillates repeatedly, in a manner reminiscent of both the Rayleigh and van der Pol systems (thick blue curves); this corresponds to repetitive firing of the neuron.

After using your own computer system's ODE solver to verify these behaviors, you can investigate what happens with other constant nonzero values of the stimulus I(t). Do all such values lead either to oscillatory phase plane solutions or to ones that converge to the same perturbed equilibrium point? Or do certain different values lead to different perturbed equilibrium points? Do all oscillatory solutions correspond to phase plane curves that appear to converge to a single limit cycle?

The FitzHugh-Nagumo system has proven to be a very useful simplification of the Hodgkin-Huxley system, exhibiting a number of characteristics of neuron electrical activity. For a more detailed discussion see the classic work of L. Edelstein-Keshet, *Mathematical Models in Biology* (Society for Industrial and Applied Mathematics, 2005).

10 Laplace Transform Methods

10.1 Laplace Transforms and Inverse Transforms





FIGURE 10.1.1. Transformation of a function: \mathcal{L} in analogy with D.

In Chapter 5 we saw that linear differential equations with constant coefficients have numerous applications and can be solved systematically. There are common situations, however, in which the alternative methods of this chapter are preferable. For example, recall the differential equations

$$mx'' + cx' + kx = F(t)$$
 and $LI'' + RI' + \frac{1}{C}I = E'(t)$

corresponding to a mass-spring-dashpot system and a series *RLC* circuit, respectively. It often happens in practice that the forcing term, F(t) or E'(t), has discontinuities—for example, when the voltage supplied to an electrical circuit is turned off and on periodically. In this case the methods of Chapter 5 can be quite awkward, and the Laplace transform method is more convenient.

The differentiation operator D can be viewed as a transformation which, when applied to the function f(t), yields the new function $D\{f(t)\} = f'(t)$. The Laplace transformation \mathcal{L} involves the operation of integration and yields the new function $\mathcal{L}\{f(t)\} = F(s)$ of a new independent variable s. The situation is diagrammed in Fig. 10.1.1. After learning in this section how to compute the Laplace transform F(s) of a function f(t), we will see in Section 10.2 that the Laplace transform converts a *differential* equation in the unknown function f(t) into an *algebraic* equation in F(s). Because algebraic equations are generally easier to solve than differential equations, this is one method that simplifies the problem of finding the solution f(t). >

DEFINITION The Laplace Transform

Given a function f(t) defined for all $t \ge 0$, the Laplace transform of f is the function F defined as follows:

$$F(s) = \mathcal{L}{f(t)} = \int_0^\infty e^{-st} f(t) dt \tag{1}$$

for all values of *s* for which the improper integral converges.

Recall that an **improper integral** over an infinite interval is defined as a limit of integrals over bounded intervals; that is,

$$\int_{a}^{\infty} g(t) dt = \lim_{b \to \infty} \int_{a}^{b} g(t) dt.$$
 (2)

If the limit in (2) exists, then we say that the improper integral **converges**; otherwise, it **diverges** or fails to exist. Note that the integrand of the improper integral in (1) contains the parameter *s* in addition to the variable of integration *t*. Therefore, when the integral in (1) converges, it converges not merely to a number, but to a *function F* of *s*. As in the following examples, it is typical for the improper integral in the definition of $\mathcal{L}{f(t)}$ to converge for some values of *s* and diverge for others.

Example 1 With $f(t) \equiv 1$ for $t \ge 0$, the definition of the Laplace transform in (1) gives

$$\mathscr{L}\{1\} = \int_0^\infty e^{-st} dt = \left[-\frac{1}{s}e^{-st}\right]_0^\infty = \lim_{b \to \infty} \left[-\frac{1}{s}e^{-bs} + \frac{1}{s}\right].$$

and therefore

$$\mathcal{L}\{1\} = \frac{1}{s} \quad \text{for} \quad s > 0. \tag{3}$$

As in (3), it's good practice to specify the domain of the Laplace transform—in problems as well as in examples. Also, in this computation we have used the common abbreviation

$$\left[g(t)\right]_{a}^{\infty} = \lim_{b \to \infty} \left[g(t)\right]_{a}^{b}.$$
(4)

Remark The limit we computed in Example 1 would not exist if s < 0, for then $(1/s)e^{-bs}$ would become unbounded as $b \to +\infty$. Hence $\mathcal{L}\{1\}$ is defined only for s > 0. This is typical of Laplace transforms; the domain of a transform is normally of the form s > a for some number a.

Example 2

e 2 With
$$f(t) = e^{at}$$
 for $t \ge 0$, we obtain

$$\mathcal{L}\{e^{at}\} = \int_0^\infty e^{-st} e^{at} dt = \int_0^\infty e^{-(s-a)t} dt = \left[-\frac{e^{-(s-a)t}}{s-a}\right]_{t=0}^\infty.$$

If s - a > 0, then $e^{-(s-a)t} \to 0$ as $t \to +\infty$, so it follows that

$$\pounds \{e^{at}\} = \frac{1}{s-a} \quad \text{for} \quad s > a.$$
(5)

Note here that the improper integral giving $\mathcal{L}\{e^{at}\}$ diverges if $s \leq a$. It is worth noting also that the formula in (5) holds if *a* is a complex number. For then, with $a = \alpha + i\beta$,

$$e^{-(s-a)t} = e^{i\beta t}e^{-(s-\alpha)t} \to 0$$

as $t \to +\infty$, provided that $s > \alpha = \operatorname{Re}[a]$; recall that $e^{i\beta t} = \cos\beta t + i\sin\beta t$.

10.1 LAPLACE TRANSFORMS AND INVERSE TRANSFORMS 559

The Laplace transform $\mathcal{L}{t^a}$ of a power function is most conveniently expressed in terms of the **gamma function** $\Gamma(x)$, which is defined for x > 0 by the formula

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt.$$
(6)

For an elementary discussion of $\Gamma(x)$, see the subsection on the gamma function in Section 11.4, where it is shown that

$$\Gamma(1) = 1 \tag{7}$$

and that

>

$$\Gamma(x+1) = x\Gamma(x) \tag{8}$$

for x > 0. It then follows that if *n* is a positive integer, then

$$\Gamma(n+1) = n\Gamma(n)$$

= $n \cdot (n-1)\Gamma(n-1)$
= $n \cdot (n-1) \cdot (n-2)\Gamma(n-2)$
:
= $n(n-1)(n-2) \cdots 2 \cdot \Gamma(2)$
= $n(n-1)(n-2) \cdots 2 \cdot 1 \cdot \Gamma(1);$

thus

$$\Gamma(n+1) = n! \tag{9}$$

if *n* is a positive integer. Therefore, the function $\Gamma(x + 1)$, which is defined and continuous for all x > -1, agrees with the factorial function for x = n, a positive integer.

xample 3 Suppose that
$$f(t) = t^a$$
 where *a* is real and $a > -1$. Then
 $\mathcal{L}\{t^a\} = \int_0^\infty e^{-st} t^a dt.$

If we substitute u = st, t = u/s, and dt = du/s in this integral, we get

$$\mathcal{L}\{t^a\} = \frac{1}{s^{a+1}} \int_0^\infty e^{-u} u^a \, du = \frac{\Gamma(a+1)}{s^{a+1}} \tag{10}$$

for all s > 0 (so that u = st > 0). Because $\Gamma(n + 1) = n!$ if *n* is a nonnegative integer, we see that

$$\mathscr{L}\lbrace t^n\rbrace = \frac{n!}{s^{n+1}} \quad \text{for} \quad s > 0.$$
⁽¹¹⁾

For instance,

$$\mathcal{L}{t} = \frac{1}{s^2}, \quad \mathcal{L}{t^2} = \frac{2}{s^3}, \text{ and } \mathcal{L}{t^3} = \frac{6}{s^4}.$$

As in Problems 1 and 2, these formulas can be derived immediately from the definition, without the use of the gamma function.

Linearity of Transforms

It is not necessary for us to proceed much further in the computation of Laplace transforms directly from the definition. Once we know the Laplace transforms of several functions, we can combine them to obtain transforms of other functions. The reason is that the Laplace transformation is a *linear* operation.

>

THEOREM 1 Linearity of the Laplace Transform

If *a* and *b* are constants, then

$$\mathcal{L}\{af(t) + bg(t)\} = a\mathcal{L}\{f(t)\} + b\mathcal{L}\{g(t)\}$$
(12)

for all s such that the Laplace transforms of the functions f and g both exist.

The proof of Theorem 1 follows immediately from the linearity of the operations of taking limits and of integration:

$$\begin{aligned} \mathcal{L}\{af(t) + bg(t)\} &= \int_0^\infty e^{-st} \left[af(t) + bg(t)\right] dt \\ &= \lim_{c \to \infty} \int_0^c e^{-st} \left[af(t) + bg(t)\right] dt \\ &= a \left(\lim_{c \to \infty} \int_0^c e^{-st} f(t) dt\right) + b \left(\lim_{c \to \infty} \int_0^c e^{-st} g(t) dt\right) \\ &= a \mathcal{L}\{f(t)\} + b \mathcal{L}\{g(t)\}. \end{aligned}$$

The computation of $\mathcal{L}\{t^{n/2}\}$ is based on the known special value

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi} \tag{13}$$

of the gamma function. For instance, it follows that

$$\Gamma\left(\frac{5}{2}\right) = \frac{3}{2}\Gamma\left(\frac{3}{2}\right) = \frac{3}{2} \cdot \frac{1}{2}\Gamma\left(\frac{1}{2}\right) = \frac{3}{4}\sqrt{\pi},$$

using the formula $\Gamma(x + 1) = x\Gamma(x)$ in (9), first with $x = \frac{3}{2}$ and then with $x = \frac{1}{2}$. Now the formulas in (10) through (12) yield

$$\pounds\{3t^2 + 4t^{3/2}\} = 3 \cdot \frac{2!}{s^3} + \frac{4\Gamma\left(\frac{5}{2}\right)}{s^{5/2}} = \frac{6}{s^3} + 3\sqrt{\frac{\pi}{s^5}}.$$

Example 5

Example 4

Recall that $\cosh kt = (e^{kt} + e^{-kt})/2$. If k > 0, then Theorem 1 and Example 2 together give

$$\mathcal{L}\{\cosh kt\} = \frac{1}{2}\mathcal{L}\{e^{kt}\} + \frac{1}{2}\mathcal{L}\{e^{-kt}\} = \frac{1}{2}\left(\frac{1}{s-k} + \frac{1}{s+k}\right)$$

that is,

$$\mathcal{L}\{\cosh kt\} = \frac{s}{s^2 - k^2} \quad \text{for } s > k > 0.$$
(14)

Similarly,

$$\mathcal{L}\{\sinh kt\} = \frac{k}{s^2 - k^2} \quad \text{for } s > k > 0.$$
(15)

Because $\cos kt = (e^{ikt} + e^{-ikt})/2$, the formula in (5) (with a = ik) yields

$$\mathscr{L}\{\cos kt\} = \frac{1}{2} \left(\frac{1}{s - ik} + \frac{1}{s + ik} \right) = \frac{1}{2} \cdot \frac{2s}{s^2 - (ik)^2},$$

and thus

$$\mathcal{L}\{\cos kt\} = \frac{s}{s^2 + k^2} \quad \text{for } s > 0.$$
(16)

(The domain follows from s > Re[ik] = 0.) Similarly,

$$\mathcal{L}\{\sin kt\} = \frac{k}{s^2 + k^2} \quad \text{for } s > 0.$$
⁽¹⁷⁾

Example 6 Applying linearity, the formula in (16), and a familiar trigonometric identity, we get

$$\mathcal{L}\{3e^{2t} + 2\sin^2 3t\} = \mathcal{L}\{3e^{2t} + 1 - \cos 6t\}$$
$$= \frac{3}{s-2} + \frac{1}{s} - \frac{s}{s^2 + 36}$$
$$= \frac{3s^3 + 144s - 72}{s(s-2)(s^2 + 36)} \quad \text{for } s > 0.$$

Inverse Transforms

According to Theorem 3 of this section, no two different functions that are both continuous for all $t \ge 0$ can have the same Laplace transform. Thus if F(s) is the transform of some continuous function f(t), then f(t) is uniquely determined. This observation allows us to make the following definition: If $F(s) = \mathcal{L}{f(t)}$, then we call f(t) the **inverse Laplace transform** of F(s) and write

$$f(t) = \mathcal{L}^{-1}\{F(s)\}.$$
 (18)

Example 7 Using the Laplace transforms derived in Examples 2, 3, and 5 we see that

$$\mathcal{L}^{-1}\left\{\frac{1}{s^3}\right\} = \frac{1}{2}t^2, \quad \mathcal{L}^{-1}\left\{\frac{1}{s+2}\right\} = e^{-2t}, \quad \mathcal{L}^{-1}\left\{\frac{2}{s^2+9}\right\} = \frac{2}{3}\sin 3t$$

and so on.

≻

NOTATION: FUNCTIONS AND THEIR TRANSFORMS. Throughout this chapter we denote functions of t by lowercase letters. The transform of a function will always be denoted by that same letter capitalized. Thus F(s) is the Laplace transform of f(t) and x(t) is the inverse Laplace transform of X(s).

A table of Laplace transforms serves a purpose similar to that of a table of integrals. The table in Fig. 10.1.2 lists the transforms derived in this section; many additional transforms can be derived from these few, using various general properties of the Laplace transformation (which we will discuss in subsequent sections).

Piecewise Continuous Functions

As we remarked at the beginning of this section, we need to be able to handle certain types of discontinuous functions. The function f(t) is said to be **piecewise continuous** on the bounded interval $a \leq t \leq b$ provided that [a, b] can be subdivided into finitely many abutting subintervals in such a way that

- **1.** f is continuous in the interior of each of these subintervals; and
- **2.** f(t) has a finite limit as t approaches each endpoint of each subinterval from its interior.
| f(t) | F(s) | |
|-------------------|-------------------------------|-----------------|
| 1 | $\frac{1}{s}$ | (s > 0) |
| t | $\frac{1}{s^2}$ | (<i>s</i> > 0) |
| $t^n \ (n \ge 0)$ | $\frac{n!}{s^{n+1}}$ | (<i>s</i> > 0) |
| $t^a (a > -1)$ | $\frac{\Gamma(a+1)}{s^{a+1}}$ | (<i>s</i> > 0) |
| e ^{at} | $\frac{1}{s-a}$ | (s > a) |
| $\cos kt$ | $\frac{s}{s^2 + k^2}$ | (s > 0) |
| sin k t | $\frac{k}{s^2 + k^2}$ | (s > 0) |
| $\cosh k t$ | $\frac{s}{s^2 - k^2}$ | (s > k) |
| sinh k t | $\frac{k}{s^2 - k^2}$ | (s > k) |
| u(t-a) | $\frac{e^{-as}}{s}$ | (<i>s</i> > 0) |

FIGURE 10.1.2. A short table of Laplace transforms.

We say that f is piecewise continuous for $t \ge 0$ if it is piecewise continuous on every bounded subinterval of $[0, +\infty)$. Thus a piecewise continuous function has only simple discontinuities (if any) and only at isolated points. At such points the value of the function experiences a finite jump, as indicated in Fig. 10.1.3. The jump in f(t) at the point c is defined to be f(c+) - f(c-), where

$$f(c+) = \lim_{\epsilon \to 0^+} f(c+\epsilon)$$
 and $f(c-) = \lim_{\epsilon \to 0^+} f(c-\epsilon)$.

Perhaps the simplest piecewise continuous (but discontinuous) function is the unit step function, whose graph appears in Fig. 10.1.4. It is defined as follows:

$$u(t) = \begin{cases} 0 & \text{for } t < 0, \\ 1 & \text{for } t \ge 0. \end{cases}$$
(19)

Because u(t) = 1 for $t \ge 0$ and because the Laplace transform involves only the values of a function for $t \ge 0$, we see immediately that

$$\mathcal{L}\{u(t)\} = \frac{1}{s} \quad (s > 0).$$
(20)

The graph of the unit step function $u_a(t) = u(t-a)$ appears in Fig. 10.1.5. Its jump occurs at t = a rather than at t = 0; equivalently,

$$u_{a}(t) = u(t-a) = \begin{cases} 0 & \text{for } t < a, \\ 1 & \text{for } t \ge a. \end{cases}$$
(21)

$$\begin{array}{c} \mathsf{le 8} \\ \mathsf{Find } \mathcal{L}\{u_a(t)\} \\ \end{array}$$

Solution

if a > 0. We begin with the definition of the Laplace transform. We obtain

$$\mathcal{L}\{u_a(t)\} = \int_0^\infty e^{-st} u_a(t) \, dt = \int_a^\infty e^{-st} \, dt = \lim_{b \to \infty} \left[-\frac{e^{-st}}{s} \right]_{t=a}^b;$$

consequently,

$$\mathcal{L}\{u_a(t)\} = \frac{e^{-as}}{s} \quad (s > 0, \ a > 0).$$
⁽²²⁾



FIGURE 10.1.3. The graph of a piecewise continuous function; the solid dots indicate values of the function at discontinuities.



FIGURE 10.1.4. The graph of the unit step function.



FIGURE 10.1.5. The unit step function $u_a(t)$ has a jump at t = a.

General Properties of Transforms

It is a familiar fact from calculus that the integral

$$\int_{a}^{b} g(t) \, dt$$

exists if g is piecewise continuous on the bounded interval [a, b]. Hence if f is piecewise continuous for $t \ge 0$, it follows that the integral

$$\int_0^b e^{-st} f(t) \, dt$$

exists for all $b < +\infty$. But in order for F(s)—the limit of this last integral as $b \to +\infty$ —to exist, we need some condition to limit the rate of growth of f(t) as $t \to +\infty$. The function f is said to be **of exponential order** as $t \to +\infty$ if there exist nonnegative constants M, c, and T such that

$$|f(t)| \le M e^{ct} \quad \text{for } t \ge T.$$
(23)

Thus a function is of exponential order provided that it grows no more rapidly (as $t \to +\infty$) than a constant multiple of some exponential function with a linear exponent. The particular values of M, c, and T are not so important. What is important is that *some* such values exist so that the condition in (23) is satisfied.

The condition in (23) merely says that $f(t)/e^{ct}$ lies between -M and M and is therefore bounded in value for t sufficiently large. In particular, this is true (with c = 0) if f(t) itself is bounded. Thus every *bounded* function—such as $\cos kt$ or $\sin kt$ —is of exponential order.

If p(t) is a polynomial, then the familiar fact that $p(t)e^{-t} \to 0$ as $t \to +\infty$ implies that (23) holds (for T sufficiently large) with M = c = 1. Thus every *polynomial* function is of exponential order.

For an example of an elementary function that is continuous and therefore bounded on every (finite) interval, but nevertheless is not of exponential order, consider the function $f(t) = e^{t^2} = \exp(t^2)$. Whatever the value of c, we see that

$$\lim_{t \to \infty} \frac{f(t)}{e^{ct}} = \lim_{t \to \infty} \frac{e^{t^2}}{e^{ct}} = \lim_{t \to \infty} e^{t^2 - ct} = +\infty$$

because $t^2 - ct \to +\infty$ as $t \to +\infty$. Hence the condition in (23) cannot hold for any (finite) value *M*, so we conclude that the function $f(t) = e^{t^2}$ is *not* of exponential order.

Similarly, because $e^{-st}e^{t^2} \to +\infty$ as $t \to +\infty$, we see that the improper integral $\int_0^\infty e^{-st}e^{t^2} dt$ that would define $\mathcal{L}\{e^{t^2}\}$ does not exist (for any *s*), and therefore that the function e^{t^2} does *not* have a Laplace transform. The following theorem guarantees that piecewise functions of exponential order *do* have Laplace transforms.

THEOREM 2 Existence of Laplace Transforms

If the function f is piecewise continuous for $t \ge 0$ and is of exponential order as $t \to +\infty$, then its Laplace transform $F(s) = \mathcal{L}{f(t)}$ exists. More precisely, if f is piecewise continuous and satisfies the condition in (23), then F(s) exists for all s > c.

Proof: First we note that we can take T = 0 in (23). For by piecewise continuity, |f(t)| is bounded on [0, T]. Increasing M in (23) if necessary, we can therefore assume that $|f(t)| \leq M$ if $0 \leq t \leq T$. Because $e^{ct} \geq 1$ for $t \geq 0$, it then follows that $|f(t)| \leq Me^{ct}$ for all $t \geq 0$.

By a standard theorem on convergence of improper integrals—the fact that absolute convergence implies convergence—it suffices for us to prove that the integral

$$\int_0^\infty |e^{-st} f(t)| \, dt$$

exists for s > c. To do this, it suffices in turn to show that the value of the integral

$$\int_0^b |e^{-st} f(t)| \, dt$$

remains bounded as $b \to +\infty$. But the fact that $|f(t)| \leq Me^{ct}$ for all $t \geq 0$ implies that

$$\int_0^b |e^{-st} f(t)| dt \leq \int_0^b |e^{-st} M e^{ct}| dt = M \int_0^b e^{-(s-c)t} dt$$
$$\leq M \int_0^\infty e^{-(s-c)t} dt = \frac{M}{s-c}$$

if s > c. This proves Theorem 2.

We have shown, moreover, that

$$|F(s)| \leq \int_0^\infty |e^{-st} f(t)| dt \leq \frac{M}{s-c}$$
(24)

if s > c. When we take limits as $s \to +\infty$, we get the following result.

COROLLARY F(s) for s Large

If f(t) satisfies the hypotheses of Theorem 2, then

$$\lim_{s \to \infty} F(s) = 0. \tag{25}$$

The condition in (25) severely limits the functions that can be Laplace transforms. For instance, the function G(s) = s/(s+1) cannot be the Laplace transform of any "reasonable" function because its limit as $s \to +\infty$ is 1, not 0. More generally, a rational function—a quotient of two polynomials—can be (and is, as we shall see) a Laplace transform only if the degree of its numerator is less than that of its denominator.

On the other hand, the hypotheses of Theorem 2 are sufficient, but not necessary, conditions for existence of the Laplace transform of f(t). For example, the function $f(t) = 1/\sqrt{t}$ fails to be piecewise continuous (at t = 0), but nevertheless (Example 3 with $a = -\frac{1}{2} > -1$) its Laplace transform

$$\mathcal{L}\left\{t^{-1/2}\right\} = \frac{\Gamma\left(\frac{1}{2}\right)}{s^{1/2}} = \sqrt{\frac{\pi}{s}}$$

both exists and violates the condition in (24), which would imply that sF(s) remains bounded as $s \to +\infty$.

The remainder of this chapter is devoted largely to techniques for solving a differential equation by first finding the Laplace transform of its solution. It is then vital for us to know that this uniquely determines the solution of the differential equation; that is, that the function of s we have found has only one inverse Laplace transform that could be the desired solution. The following theorem is proved in Chapter 6 of Churchill's Operational Mathematics, 3rd ed. (New York: McGraw-Hill, 1972).

THEOREM 3 Uniqueness of Inverse Laplace Transforms

Suppose that the functions f(t) and g(t) satisfy the hypotheses of Theorem 2, so that their Laplace transforms F(s) and G(s) both exist. If F(s) = G(s) for all s > c (for some c), then f(t) = g(t) wherever on $[0, +\infty)$ both f and g are continuous.

Thus two piecewise continuous functions of exponential order with the same Laplace transform can differ only at their isolated points of discontinuity. This is of no importance in most practical applications, so we may regard inverse Laplace transforms as being essentially unique. In particular, two solutions of a differential equation must both be continuous, and hence must be the same solution if they have the same Laplace transform.

Historical Remark Laplace transforms have an interesting history. The integral in the definition of the Laplace transform probably appeared first in the work of Euler. It is customary in mathematics to name a technique or theorem for the next person after Euler to discover it (else there would be several hundred different examples of "Euler's theorem"). In this case, the next person was the French mathematician Pierre Simon de Laplace (1749–1827), who employed such integrals in his work on probability theory. The so-called operational methods for solving differential equations, which are based on Laplace transforms, were not exploited by Laplace. Indeed, they were discovered and popularized by practicing engineers—notably the English electrical engineer Oliver Heaviside (1850–1925). These techniques were successfully and widely applied before they had been rigorously justified, and around the beginning of the twentieth century their validity was the subject of considerable controversy. One reason is that Heaviside blithely assumed the existence of functions whose Laplace transforms contradict the condition that $F(s) \to 0$ as $s \to 0$, thereby raising questions as to the meaning and nature of functions in mathematics. (This is reminiscent of the way Leibniz two centuries earlier had obtained correct results in calculus using "infinitely small" real numbers, thereby raising questions as to the nature and role of numbers in mathematics.)

10.1Problems

Apply the definition in (1) to find directly the Laplace transforms of the functions described (by formula or graph) in Problems 1 through 10.

- **2.** $f(t) = t^2$ **1.** f(t) = t**4.** $f(t) = \cos t$ 3. $f(t) = e^{3t+1}$
- **5.** $f(t) = \sinh t$

FIGURE 10.1.6.











FIGURE 10.1.9.

Use the transforms in Fig. 10.1.2 to find the Laplace transforms of the functions in Problems 11 through 22. A preliminary integration by parts may be necessary.

11. $f(t) = \sqrt{t} + 3t$ **12.** $f(t) = 3t^{5/2} - 4t^3$ **13.** $f(t) = t - 2e^{3t}$ **14.** $f(t) = t^{3/2} - e^{-10t}$ **15.** $f(t) = 1 + \cosh 5t$ **16.** $f(t) = \sin 2t + \cos 2t$ **17.** $f(t) = \cos^2 2t$ **18.** $f(t) = \sin 3t \cos 3t$ **19.** $f(t) = (1 + t)^3$ **20.** $f(t) = te^t$ **21.** $f(t) = t \cos 2t$ **22.** $f(t) = \sinh^2 3t$

Use the transforms in Fig. 10.1.2 to find the inverse Laplace transforms of the functions in Problems 23 through 32.

- 23. $F(s) = \frac{3}{s^4}$ 24. $F(s) = s^{-3/2}$ 25. $F(s) = \frac{1}{s} - \frac{2}{s^{5/2}}$ 26. $F(s) = \frac{1}{s+5}$ 27. $F(s) = \frac{3}{s-4}$ 28. $F(s) = \frac{3s+1}{s^2+4}$ 29. $F(s) = \frac{5-3s}{s^2+9}$ 30. $F(s) = \frac{9+s}{4-s^2}$
- **31.** $F(s) = \frac{10s 3}{25 s^2}$ **32.** $F(s) = 2s^{-1}e^{-3s}$
- **33.** Derive the transform of $f(t) = \sin kt$ by the method used in the text to derive the formula in (16).
- **34.** Derive the transform of $f(t) = \sinh k t$ by the method used in the text to derive the formula in (14).
- **35.** Use the tabulated integral

$$\int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx) + C$$

to obtain $\mathcal{L}\{\cos kt\}$ directly from the definition of the Laplace transform.

- **36.** Show that the function $f(t) = \sin(e^{t^2})$ is of exponential order as $t \to +\infty$ but that its derivative is not.
- **37.** Given a > 0, let f(t) = 1 if $0 \le t < a$, f(t) = 0 if $t \ge a$. First, sketch the graph of the function f, making clear its value at t = a. Then express f in terms of unit step functions to show that $\mathcal{L}{f(t)} = s^{-1}(1 - e^{-as})$.
- **38.** Given that 0 < a < b, let f(t) = 1 if $a \le t < b$, f(t) = 0 if either t < a or $t \ge b$. First, sketch the graph of the function f, making clear its values at t = a and t = b. Then express f in terms of unit step functions to show that $\mathcal{L}{f(t)} = s^{-1}(e^{-as} e^{-bs})$.

39. The *unit staircase function* is defined as follows:

$$f(t) = n$$
 if $n-1 \le t < n$, $n = 1, 2, 3, ...$

(a) Sketch the graph of f to see why its name is appropriate. (b) Show that

$$f(t) = \sum_{n=0}^{\infty} u(t-n)$$

for all $t \ge 0$. (c) Assume that the Laplace transform of the infinite series in part (b) can be taken termwise (it can). Apply the geometric series to obtain the result

$$\mathcal{L}\{f(t)\} = \frac{1}{s(1 - e^{-s})}$$

40. (a) The graph of the function *f* is shown in Fig. 10.1.10. Show that *f* can be written in the form

$$f(t) = \sum_{n=0}^{\infty} (-1)^n u(t-n).$$

(b) Use the method of Problem 39 to show that



FIGURE 10.1.10. The graph of the function of Problem 40.

41. The graph of the *square-wave function* g(t) is shown in Fig. 10.1.11. Express g in terms of the function f of Problem 40 and hence deduce that



FIGURE 10.1.11. The graph of the function of Problem 41.

42. Given constants *a* and *b*, define h(t) for $t \ge 0$ by

$$h(t) = \begin{cases} a & \text{if } n-1 \leq t < n \text{ and } n \text{ is odd;} \\ b & \text{if } n-1 \leq t < n \text{ and } n \text{ is even.} \end{cases}$$

Sketch the graph of h and apply one of the preceding problems to show that

$$\mathcal{L}{h(t)} = \frac{a+be^{-s}}{s(1+e^{-s})}.$$

10.1 Application Computer Algebra Transforms and Inverse Transforms



computing resources including *Maple/Mathematica/*MATLAB.

If $f(t) = t \cos 3t$, then the definition of the Laplace transform gives the improper integral

$$F(s) = \mathcal{L}{f(t)} = \int_0^\infty t e^{-st} \cos 3t \, dt,$$

whose evaluation would appear to require a tedious integration by parts. Consequently a computer algebra Laplace transforms package is useful for the quick calculation of transforms. *Maple* contains the integral transforms package **inttrans**, and the commands

```
with(inttrans):
f := t*cos(3*t):
F := laplace(f, t, s);
```

yield immediately the Laplace transform $F(s) = (s^2 - 9)/(s^2 + 9)^2$, as do the *Mathematica* commands

```
f = t*Cos[3*t];
F = LaplaceTransform[f, t, s]
```

and the Wolfram|Alpha query

laplace transform t*cos(3t)

We can recover the original function $f(t) = t \cos 3t$ with the *Maple* command

invlaplace(F, s, t);

or the Mathematica command

```
InverseLaplaceTransform[F, s, t]
```

or the Wolfram Alpha query

```
inverse laplace transform (s^2 - 9)/(s^2 + 9)^2
```

Remark Note carefully the order of s and t in the preceding *Maple* and *Mathematica* commands—first t, then s when transforming; first s, then t when inverse transforming.

You can use these computer algebra commands to check the answers to Problems 11 through 32 in this section, as well as a few interesting problems of your own selection.

10.2 Transformation of Initial Value Problems

We now discuss the application of Laplace transforms to solve a linear differential equation with constant coefficients, such as

$$ax''(t) + bx'(t) + cx(t) = f(t),$$
(1)

with given initial conditions $x(0) = x_0$ and $x'(0) = x'_0$. By the linearity of the Laplace transformation, we can transform Eq. (1) by separately taking the Laplace transform of each term in the equation. The transformed equation is

$$a\mathcal{L}\{x''(t)\} + b\mathcal{L}\{x'(t)\} + c\mathcal{L}\{x(t)\} = \mathcal{L}\{f(t)\};$$
(2)

it involves the transforms of the derivatives x' and x'' of the unknown function x(t). The key to the method is Theorem 1, which tells us how to express the transform of the *derivative* of a function in terms of the transform of the function itself.



FIGURE 10.2.1. The discontinuities of f' correspond to "corners" on the graph of f.

THEOREM 1 Transforms of Derivatives

Suppose that the function f(t) is continuous and piecewise smooth for $t \ge 0$ and is of exponential order as $t \to +\infty$, so that there exist nonnegative constants M, c, and T such that

$$|f(t)| \le M e^{ct} \quad \text{for } t \ge T.$$
(3)

Then $\mathcal{L}{f'(t)}$ exists for s > c, and

$$\mathcal{L}\{f'(t)\} = s\mathcal{L}\{f(t)\} - f(0) = sF(s) - f(0).$$
(4)

The function f is called **piecewise smooth** on the bounded interval [a, b] if it is piecewise continuous on [a, b] and differentiable except at finitely many points, with f'(t) being piecewise continuous on [a, b]. We may assign arbitrary values to f(t) at the isolated points at which f is not differentiable. We say that f is piecewise smooth for $t \ge 0$ if it is piecewise smooth on every bounded subinterval of $[0, +\infty)$. Figure 10.2.1 indicates how "corners" on the graph of f correspond to discontinuities in its derivative f'.

The main idea of the proof of Theorem 1 is exhibited best by the case in which f'(t) is continuous (not merely piecewise continuous) for $t \ge 0$. Then, beginning with the definition of $\mathcal{L}{f'(t)}$ and integrating by parts, we get

$$\mathcal{L}\{f'(t)\} = \int_0^\infty e^{-st} f'(t) \, dt = \left[e^{-st} f(t)\right]_{t=0}^\infty + s \int_0^\infty e^{-st} f(t) \, dt.$$

Because of (3), the integrated term $e^{-st} f(t)$ approaches zero (when s > c) as $t \to +\infty$, and its value at the lower limit t = 0 contributes -f(0) to the evaluation of the preceding expression. The integral that remains is simply $\mathcal{L}{f(t)}$; by Theorem 2 of Section 10.1, the integral converges when s > c. Then $\mathcal{L}{f'(t)}$ exists when s > c, and its value is that given in Eq. (4). We will defer the case in which f'(t) has isolated discontinuities to the end of this section.

Solution of Initial Value Problems

In order to transform Eq. (1), we need the transform of the second derivative as well. If we assume that g(t) = f'(t) satisfies the hypotheses of Theorem 1, then that theorem implies that

$$\mathcal{L}{f''(t)} = \mathcal{L}{g'(t)} = s\mathcal{L}{g(t)} - g(0)$$

= $s\mathcal{L}{f'(t)} - f'(0)$
= $s[s\mathcal{L}{f(t)} - f(0)] - f'(0),$

and thus

$$\mathcal{L}\{f''(t)\} = s^2 F(s) - sf(0) - f'(0).$$
(5)

A repetition of this calculation gives

$$\mathcal{L}\{f'''(t)\} = s\mathcal{L}\{f''(t)\} - f''(0) = s^3 F(s) - s^2 f(0) - sf'(0) - f''(0).$$
(6)

After finitely many such steps we obtain the following extension of Theorem 1.

COROLLARY Transforms of Higher Derivatives

Suppose that the functions $f, f', f'', \ldots, f^{(n-1)}$ are continuous and piecewise smooth for $t \ge 0$, and that each of these functions satisfies the conditions in (3) with the same values of M and c. Then $\mathcal{L}\{f^{(n)}(t)\}$ exists when s > c, and

$$\mathcal{L}\{f^{(n)}(t)\} = s^n \mathcal{L}\{f(t)\} - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

= $s^n F(s) - s^{n-1} f(0) - \dots - s f^{(n-2)}(0) - f^{(n-1)}(0).$ (7)

Example 1

Solve the initial value problem

x'' - x' - 6x = 0; x(0) = 2, x'(0) = -1.

Solution

With the given initial values, Eqs. (4) and (5) yield

$$\mathcal{L}\{x'(t)\} = s\mathcal{L}\{x(t)\} - x(0) = sX(s) - 2$$

and

$$\mathcal{L}\{x''(t)\} = s^2 \mathcal{L}\{x(t)\} - sx(0) - x'(0) = s^2 X(s) - 2s + 1,$$

where (according to our convention about notation) X(s) denotes the Laplace transform of the (unknown) function x(t). Hence the transformed equation is

$$\left[s^{2}X(s) - 2s + 1\right] - \left[sX(s) - 2\right] - 6\left[X(s)\right] = 0$$

which we quickly simplify to

$$(s^2 - s - 6)X(s) - 2s + 3 = 0.$$

Thus

$$X(s) = \frac{2s-3}{s^2-s-6} = \frac{2s-3}{(s-3)(s+2)}.$$

By the method of partial fractions (of integral calculus), there exist constants A and B such that

$$\frac{2s-3}{(s-3)(s+2)} = \frac{A}{s-3} + \frac{B}{s+2},$$

and multiplication of both sides of this equation by (s-3)(s+2) yields the identity

$$2s - 3 = A(s + 2) + B(s - 3).$$

If we substitute s = 3, we find that $A = \frac{3}{5}$; substitution of s = -2 shows that $B = \frac{7}{5}$. Hence

$$X(s) = \mathcal{L}\{x(t)\} = \frac{\frac{3}{5}}{s-3} + \frac{\frac{7}{5}}{s+2}.$$

Because $\mathcal{L}^{-1}\{1/(s-a)\} = e^{at}$, it follows that

$$x(t) = \frac{3}{5}e^{3t} + \frac{7}{5}e^{-2t}$$

is the solution of the original initial value problem. Note that we did not first find the general solution of the differential equation. The Laplace transform method directly yields the desired particular solution, automatically taking into account—via Theorem 1 and its corollary—the given initial conditions.

Remark In Example 1 we found the values of the partial-fraction coefficients *A* and *B* by the "trick" of separately substituting the roots s = 3 and s = -2 of the original denominator $s^2 - s - 6 = (s - 3)(s + 2)$ into the equation

$$2s - 3 = A(s + 2) + B(s - 3)$$

that resulted from clearing fractions. In lieu of any such shortcut, the "sure-fire" method is to collect coefficients of powers of *s* on the right-hand side,

$$2s - 3 = (A + B)s + (2A - 3).$$

Then upon equating coefficients of terms of like degree, we get the linear equations

$$\begin{array}{rcl} A + & B = & 2, \\ 2A - 3B = -3, \end{array}$$

which are readily solved for the same values $A = \frac{3}{5}$ and $B = \frac{7}{5}$.

Example 2 Forced mass-spring system Solve the initial value problem

$$x'' + 4x = \sin 3t; \quad x(0) = x'(0) = 0$$

Such a problem arises in the motion of a mass-and-spring system with external force, as shown in Fig. 10.2.2.

n Because both initial values are zero, Eq. (5) yields $\mathcal{L}\{x''(t)\} = s^2 X(s)$. We read the transform of sin 3t from the table in Fig. 10.1.2 (Section 10.1) and thereby get the transformed equation

$$s^2 X(s) + 4X(s) = \frac{3}{s^2 + 9}.$$

Therefore,

$$X(s) = \frac{3}{(s^2 + 4)(s^2 + 9)}$$

The method of partial fractions calls for

$$\frac{3}{(s^2+4)(s^2+9)} = \frac{As+B}{s^2+4} + \frac{Cs+D}{s^2+9}.$$

The sure-fire approach would be to clear fractions by multiplying both sides by the common denominator, and then collect coefficients of powers of s on the right-hand side. Equating coefficients of like powers on the two sides of the resulting equation would then yield four linear equations that we could solve for A, B, C, and D.

However, here we can anticipate that A = C = 0, because neither the numerator nor the denominator on the left involves any odd powers of *s*, whereas nonzero values for *A* or *C* would lead to odd-degree terms on the right. So we replace *A* and *C* with zero before clearing fractions. The result is the identity

$$3 = B(s^{2} + 9) + D(s^{2} + 4) = (B + D)s^{2} + (9B + 4D).$$

When we equate coefficients of like powers of s we get the linear equations

$$B + D = 0,$$

$$9B + 4D = 3.$$

which are readily solved for $B = \frac{3}{5}$ and $D = -\frac{3}{5}$. Hence

$$X(s) = \mathcal{L}\{x(t)\} = \frac{3}{10} \cdot \frac{2}{s^2 + 4} - \frac{1}{5} \cdot \frac{3}{s^2 + 9}.$$

Because $\mathcal{L}{\sin 2t} = 2/(s^2 + 4)$ and $\mathcal{L}{\sin 3t} = 3/(s^2 + 9)$, it follows that

$$x(t) = \frac{3}{10}\sin 2t - \frac{1}{5}\sin 3t$$

Figure 10.2.3 shows the graph of this period 2π position function of the mass. Note that the Laplace transform method again gives the solution directly, without the necessity of first finding the complementary function and a particular solution of the original nonhomogeneous differential equation. Thus nonhomogeneous equations are solved in exactly the same manner as are homogeneous equations.

Examples 1 and 2 illustrate the solution procedure that is outlined in Fig. 10.2.4.



FIGURE 10.2.2. A mass-andspring system satisfying the initial value problem in Example 2. The mass is initially at rest in its equilibrium position.



FIGURE 10.2.3. The position function x(t) in Example 2.

10.2 TRANSFORMATION OF INITIAL VALUE PROBLEMS571



FIGURE 10.2.4. Using the Laplace transform to solve an initial value problem.

Linear Systems

Laplace transforms are used frequently in engineering problems to solve linear systems in which the coefficients are all constants. When initial conditions are specified, the Laplace transform reduces such a linear system of differential equations to a linear system of algebraic equations in which the unknowns are the transforms of the solution functions. As Example 3 illustrates, the technique for a system is essentially the same as for a single linear differential equation with constant coefficients.

Example 3 Dual mass-spring system Solve the system

$$2x'' = -6x + 2y,$$

$$y'' = 2x - 2y + 40\sin 3t,$$
(8)

subject to the initial conditions

$$x(0) = x'(0) = y(0) = y'(0) = 0.$$
 (9)

Thus the force $f(t) = 40 \sin 3t$ is applied to the second mass of Fig. 10.2.5, beginning at time t = 0 when the system is at rest in its equilibrium position.



FIGURE 10.2.5. A mass–and–spring system satisfying the initial value problem in Example 3. Both masses are initially at rest in their equilibrium positions.

Solution

We write
$$X(s) = \mathcal{L}\{x(t)\}$$
 and $Y(s) = \mathcal{L}\{y(t)\}$. Then the initial conditions in (9) imply that
 $\mathcal{L}\{x''(t)\} = s^2 X(s)$ and $\mathcal{L}\{y''(t)\} = s^2 Y(s)$.

Because $\mathcal{L}{\sin 3t} = 3/(s^2 + 9)$, the transforms of the equations in (8) are the equations

$$2s^{2}X(s) = -6X(s) + 2Y(s),$$

$$s^{2}Y(s) = 2X(s) - 2Y(s) + \frac{120}{s^{2} + 9}$$

Thus the transformed system is

$$(s^{2} + 3)X(s) - Y(s) = 0,$$

$$-2X(s) + (s^{2} + 2)Y(s) = \frac{120}{s^{2} + 9}.$$
 (10)

The determinant of this pair of linear equations in X(s) and Y(s) is

$$\begin{vmatrix} s^2 + 3 & -1 \\ -2 & s^2 + 2 \end{vmatrix} = (s^2 + 3)(s^2 + 2) - 2 = (s^2 + 1)(s^2 + 4),$$

and we readily solve—using Cramer's rule, for instance—the system in (10) for

$$X(s) = \frac{120}{(s^2+1)(s^2+4)(s^2+9)} = \frac{5}{s^2+1} - \frac{8}{s^2+4} + \frac{3}{s^2+9}$$
(11a)

and

$$Y(s) = \frac{120(s^2+3)}{(s^2+1)(s^2+4)(s^2+9)} = \frac{10}{s^2+1} + \frac{8}{s^2+4} - \frac{18}{s^2+9}.$$
 (11b)

The partial fraction decompositions in Eqs. (11a) and (11b) are readily found using the method of Example 2. For instance, noting that the denominator factors are linear in s^2 , we can write

$$\frac{120}{(s^2+1)(s^2+4)(s^2+9)} = \frac{A}{s^2+1} + \frac{B}{s^2+4} + \frac{C}{s^2+9}$$

and it follows that

$$120 = A(s^{2} + 4)(s^{2} + 9) + B(s^{2} + 1)(s^{2} + 9) + C(s^{2} + 1)(s^{2} + 4).$$
(12)

Substitution of $s^2 = -1$ (that is, s = i, a zero of the factor $s^2 + 1$) in Eq. (12) gives $120 = A \cdot 3 \cdot 8$, so A = 5. Similarly, substitution of $s^2 = -4$ in Eq. (12) yields B = -8, and substitution of $s^2 = -9$ yields C = 3. Thus we obtain the partial fraction decomposition shown in Eq. (11a).

At any rate, the inverse Laplace transforms of the expressions in Eqs. (11a) and (11b) give the solution

$$x(t) = 5\sin t - 4\sin 2t + \sin 3t,$$

$$y(t) = 10\sin t + 4\sin 2t - 6\sin 3t.$$

Figure 10.2.6 shows the graphs of these two period 2π position functions of the two masses.

The Transform Perspective

Let us regard the general constant-coefficient second-order equation as the equation of motion

$$mx'' + cx' + kx = f(t)$$

of the familiar mass-spring-dashpot system (Fig. 10.2.7). Then the transformed equation is

$$m\left[s^{2}X(s) - sx(0) - x'(0)\right] + c\left[sX(s) - x(0)\right] + kX(s) = F(s).$$
(13)

Note that Eq. (13) is an *algebraic* equation—indeed, a linear equation—in the "un-known" X(s). This is the source of the power of the Laplace transform method:

Linear *differential* equations are transformed into readily solved *algebraic* equations.

If we solve Eq. (13) for X(s), we get

$$X(s) = \frac{F(s)}{Z(s)} + \frac{I(s)}{Z(s)},$$
(14)



FIGURE 10.2.6. The position functions x(t) and y(t) in Example 3.



FIGURE 10.2.7. A mass-springdashpot system with external force f(t).

where

$$Z(s) = ms^{2} + cs + k$$
 and $I(s) = mx(0)s + mx'(0) + cx(0)$

Note that Z(s) depends only on the physical system itself. Thus Eq. (14) presents $X(s) = \mathcal{L}{x(t)}$ as the sum of a term depending only on the external force and one depending only on the initial conditions. In the case of an underdamped system, these two terms are the transforms

$$\mathcal{L}{x_{sp}(t)} = \frac{F(s)}{Z(s)}$$
 and $\mathcal{L}{x_{tr}(t)} = \frac{I(s)}{Z(s)}$

of the steady periodic solution and the transient solution, respectively. The only potential difficulty in finding these solutions is in finding the inverse Laplace transform of the right-hand side in Eq. (14). Much of the remainder of this chapter is devoted to finding Laplace transforms and inverse transforms. In particular, we seek those methods that are sufficiently powerful to enable us to solve problems that—unlike those in Examples 1 and 2—cannot be solved readily by the methods of Chapter 5.

Additional Transform Techniques

Solution If
$$f(t) = te^{at}$$
, then $f(0) = 0$ and $f'(t) = e^{at} + ate^{at}$. Hence Theorem 1 gives
 $\mathcal{L}\{e^{at} + ate^{at}\} = \mathcal{L}\{f'(t)\} = s\mathcal{L}\{f(t)\} = s\mathcal{L}\{te^{at}\}.$
It follows from the linearity of the transform that
 $\mathcal{L}\{e^{at}\} + a\mathcal{L}\{te^{at}\} = s\mathcal{L}\{te^{at}\}.$
Hence
 $\mathcal{L}\{te^{at}\} = \frac{\mathcal{L}\{e^{at}\}}{s-a} = \frac{1}{(s-a)^2}$ (15)
because $\mathcal{L}\{e^{at}\} = 1/(s-a).$

The derivative involves the new function $t \cos kt$, so we note that f'(0) = 0 and differentiate again. The result is

$$f''(t) = 2k\cos kt - k^2t\sin kt.$$

But $\mathcal{L}{f''(t)} = s^2 \mathcal{L}{f(t)}$ by the formula in (5) for the transform of the second derivative, and $\mathcal{L}{\cos kt} = s/(s^2 + k^2)$, so we have

$$\frac{2ks}{s^2+k^2} - k^2 \mathcal{L}\{t\sin kt\} = s^2 \mathcal{L}\{t\sin kt\}.$$

Finally, we solve this equation for

$$\mathcal{L}\{t\sin kt\} = \frac{2ks}{(s^2 + k^2)^2}.$$
(16)

This procedure is considerably more pleasant than the alternative of evaluating the integral

$$\mathcal{L}\{t\sin kt\} = \int_0^\infty t e^{-st} \sin kt \, dt.$$

Examples 4 and 5 exploit the fact that if f(0) = 0, then differentiation of f corresponds to multiplication of its transform by s. It is reasonable to expect the inverse operation of integration (antidifferentiation) to correspond to division of the transform by s.

THEOREM 2 Transforms of Integrals

If f(t) is a piecewise continuous function for $t \ge 0$ and satisfies the condition of exponential order $|f(t)| \le Me^{ct}$ for $t \ge T$, then

$$\mathscr{L}\left\{\int_0^t f(\tau) \, d\tau\right\} = \frac{1}{s} \mathscr{L}\{f(t)\} = \frac{F(s)}{s} \tag{17}$$

for s > c. Equivalently,

$$\mathcal{L}^{-1}\left\{\frac{F(s)}{s}\right\} = \int_0^t f(\tau) \, d\tau.$$
(18)

Proof: Because f is piecewise continuous, the fundamental theorem of calculus implies that

$$g(t) = \int_0^t f(\tau) \, d\tau$$

is continuous and that g'(t) = f(t) where f is continuous; thus g is continuous and piecewise smooth for $t \ge 0$. Furthermore,

$$|g(t)| \leq \int_0^t |f(\tau)| \, d\tau \leq M \int_0^t e^{c\tau} \, d\tau = \frac{M}{c} (e^{ct} - 1) < \frac{M}{c} e^{ct},$$

so g(t) is of exponential order as $t \to +\infty$. Hence we can apply Theorem 1 to g; this gives

$$\mathcal{L}{f(t)} = \mathcal{L}{g'(t)} = s\mathcal{L}{g(t)} - g(0).$$

Now g(0) = 0, so division by *s* yields

$$\mathscr{L}\left\{\int_0^t f(\tau) \ d\tau\right\} = \mathscr{L}\left\{g(t)\right\} = \frac{\mathscr{L}\left\{f(t)\right\}}{s},$$

which completes the proof.

Example 6 Fi

Find the inverse Laplace transform of

$$G(s) = \frac{1}{s^2(s-a)}.$$

Solution

In effect, Eq. (18) means that we can delete a factor of s from the denominator, find the inverse transform of the resulting simpler expression, and finally integrate from 0 to t (to "correct" for the missing factor s). Thus

$$\mathcal{L}^{-1}\left\{\frac{1}{s(s-a)}\right\} = \int_0^t \mathcal{L}^{-1}\left\{\frac{1}{s-a}\right\} d\tau = \int_0^t e^{a\tau} d\tau = \frac{1}{a}(e^{at} - 1).$$

We now repeat the technique to obtain

$$\mathcal{L}^{-1}\left\{\frac{1}{s^2(s-a)}\right\} = \int_0^t \mathcal{L}^{-1}\left\{\frac{1}{s(s-a)}\right\} d\tau = \int_0^t \frac{1}{a}(e^{a\tau}-1) d\tau$$
$$= \left[\frac{1}{a}\left(\frac{1}{a}e^{a\tau}-\tau\right)\right]_0^t = \frac{1}{a^2}(e^{at}-at-1).$$

10.2 TRANSFORMATION OF INITIAL VALUE PROBLEMS575

This technique is often a more convenient way than the method of partial fractions for finding an inverse transform of a fraction of the form $P(s)/[s^n Q(s)]$.

Proof of Theorem 1: We conclude this section with the proof of Theorem 1 in the general case in which f' is merely piecewise continuous. We need to prove that the limit

$$\lim_{b \to \infty} \int_0^b e^{-st} f'(t) \, dt$$

exists and also need to find its value. With *b* fixed, let $t_1, t_2, \ldots, t_{k-1}$ be the points interior to the interval [0, b] at which f' is discontinuous. Let $t_0 = 0$ and $t_k = b$. Then we can integrate by parts on each interval (t_{n-1}, t_n) where f' is continuous. This yields

$$\int_{0}^{b} e^{-st} f'(t) dt = \sum_{n=1}^{k} \int_{t_{n-1}}^{t_n} e^{-st} f'(t) dt$$
$$= \sum_{n=1}^{k} \left[e^{-st} f(t) \right]_{t_{n-1}}^{t_n} + \sum_{n=1}^{k} s \int_{t_{n-1}}^{t_n} e^{-st} f(t) dt.$$
(19)

Now the first summation

$$\sum_{n=1}^{k} \left[e^{-st} f(t) \right]_{t_{n-1}}^{t_n} = \left[-f(t_0) + e^{-st_1} f(t_1) \right] + \left[-e^{-st_1} f(t_1) + e^{-st_2} f(t_2) \right] + \dots + \left[-e^{st_{k-2}} f(t_{k-2}) + e^{-st_{k-1}} f(t_{k-1}) \right]$$
(20)
+ $\left[-e^{st_{k-1}} f(t_{k-1}) + e^{-st_k} f(t_k) \right]$

in (19) telescopes down to $-f(t_0) + e^{-st_k} f(t_k) = -f(0) + e^{-sb} f(b)$, and the second summation adds up to *s* times the integral from $t_0 = 0$ to $t_k = b$. Therefore (19) reduces to

$$\int_0^b e^{-st} f'(t) dt = -f(0) + e^{-sb} f(b) + s \int_0^b e^{-st} f(t) dt.$$

But from Eq. (3) we get

$$\left| e^{-sb} f(b) \right| \leq e^{-sb} \cdot Me^{cb} = Me^{-b(s-c)} \to 0$$

if s > c. Therefore, finally taking limits (with *s* fixed) as $b \to +\infty$ in the preceding equation, we get the desired result

$$\mathcal{L}{f'(t)} = s\mathcal{L}{f(t)} - f(0).$$

Extension of Theorem 1

Now suppose that the function f is only piecewise continuous (instead of continuous), and let t_1, t_2, t_3, \ldots be the points (for t > 0) where either f or f' is discontinuous. The fact that f is piecewise continuous includes the assumption that—within each interval $[t_{n-1}, t_n]$ between successive points of discontinuity—f agrees with a function that is continuous on the whole closed interval and has "endpoint values"

$$f(t_{n-1}^+) = \lim_{t \to t_{n-1}^+} f(t)$$
 and $f(t_n^-) = \lim_{t \to t_n^-} f(t)$

that may not agree with the actual values $f(t_{n-1})$ and $f(t_n)$. The value of an integral on an interval is not affected by changing the values of the integrand at the endpoints. However, if the fundamental theorem of calculus is applied to find the value of the integral, then the antiderivative function must be continuous on the closed interval. We therefore use the "continuous from within the interval" endpoint values above in evaluating (by parts) the integrals on the right in (19). The result is

$$\sum_{n=1}^{k} \left[e^{-st} f(t) \right]_{t_{n-1}}^{t_n} = \left[-f(t_0^+) + e^{-st_1} f(t_1^-) \right] + \left[-e^{-st_1} f(t_1^+) + e^{-st_2} f(t_2^-) \right] + \dots + \left[-e^{st_{k-2}} f(t_{k-2}^+) + e^{-st_{k-1}} f(t_{k-1}^-) \right] + \left[-e^{st_{k-1}} f(t_{k-1}^+) + e^{-st_k} f(t_k^-) \right] = -f(0^+) - \sum_{n=1}^{k-1} j_f(t_n) + e^{-sb} f(b^-),$$
(20')

where

$$j_f(t_n) = f(t_n^+) - f(t_n^-)$$
(21)

denotes the (finite) jump in f(t) at $t = t_n$. Assuming that $\mathcal{L}{f'(t)}$ exists, we therefore get the generalization

$$\mathcal{L}\{f'(t)\} = sF(s) - f(0^+) - \sum_{n=1}^{\infty} e^{-st_n} j_f(t_n)$$
(22)

of $\mathcal{L}{f'(t)} = sF(s) - f(0)$ when we now take the limit in (19) as $b \to +\infty$.

Let f(t) = 1 + [t] be the unit staircase function; its graph is shown in Fig. 10.2.8. Then f(0) = 1, $f'(t) \equiv 0$, and $j_f(n) = 1$ for each integer n = 1, 2, 3, ... Hence Eq. (22) yields

$$0 = sF(s) - 1 - \sum_{n=1}^{\infty} e^{-ns}.$$

so the Laplace transform of f(t) is

$$F(s) = \frac{1}{s} \sum_{n=0}^{\infty} e^{-ns} = \frac{1}{s(1-e^{-s})}.$$

In the last step we used the formula for the sum of a geometric series,

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x},$$

with $x = e^{-s} < 1$.

10.2 **Problems**

Use Laplace transforms to solve the initial value problems in Problems 1 through 16.

- **1.** x'' + 4x = 0; x(0) = 5, x'(0) = 0**2.** x'' + 9x = 0; x(0) = 3, x'(0) = 4
- **3.** x'' x' 2x = 0; x(0) = 0, x'(0) = 2

4. x'' + 8x' + 15x = 0; x(0) = 2, x'(0) = -3 **5.** $x'' + x = \sin 2t; x(0) = 0 = x'(0)$ **6.** $x'' + 4x = \cos t; x(0) = 0 = x'(0)$ **7.** $x'' + x = \cos 3t; x(0) = 1, x'(0) = 0$ **8.** x'' + 9x = 1; x(0) = 0 = x'(0)





10.2 TRANSFORMATION OF INITIAL VALUE PROBLEMS 577

- 9. x'' + 4x' + 3x = 1; x(0) = 0 = x'(0)**10.** x'' + 3x' + 2x = t; x(0) = 0, x'(0) = 2**11.** x' = 2x + y, y' = 6x + 3y; x(0) = 1, y(0) = -2**12.** x' = x + 2y, $y' = x + e^{-t}$; x(0) = y(0) = 0**13.** x' + 2y' + x = 0, x' - y' + y = 0; x(0) = 0, y(0) = 1**14.** x'' + 2x + 4y = 0, y'' + x + 2y = 0; x(0) = y(0) = 0,
- x'(0) = y'(0) = -1**15.** x'' + x' + y' + 2x - y = 0, y'' + x' + y' + 4x - 2y = 0; x(0) = y(0) = 1, x'(0) = y'(0) = 0
- **16.** x' = x + z, y' = x + y, z' = -2x z; x(0) = 1, y(0) = 0, z(0) = 0

Apply Theorem 2 to find the inverse Laplace transforms of the functions in Problems 17 through 24.

- **17.** $F(s) = \frac{1}{s(s-3)}$ **18.** $F(s) = \frac{3}{s(s+5)}$ **19.** $F(s) = \frac{1}{s(s^2+4)}$ **20.** $F(s) = \frac{2s+1}{s(s^2+9)}$ **21.** $F(s) = \frac{1}{s^2(s^2+1)}$ **22.** $F(s) = \frac{1}{s(s^2-9)}$ **23.** $F(s) = \frac{1}{s^2(s^2-1)}$ **24.** $F(s) = \frac{1}{s(s+1)(s+2)}$
- **25.** Apply Theorem 1 to derive $\mathcal{L}{\sin kt}$ from the formula for $\mathcal{L}\{\cos kt\}.$
- **26.** Apply Theorem 1 to derive $\mathcal{L} \{\cosh kt\}$ from the formula for $\mathcal{L}{\sinh kt}$.
- 27. (a) Apply Theorem 1 to show that

$$\mathcal{L}\lbrace t^n e^{at}\rbrace = \frac{n}{s-a}\mathcal{L}\lbrace t^{n-1} e^{at}\rbrace.$$

(**b**) Deduce that $\mathcal{L}{t^n e^{at}} = n!/(s-a)^{n+1}$ for n = 1, 2,3,

Apply Theorem 1 as in Example 5 to derive the Laplace transforms in Problems 28 through 30.

28.
$$\mathcal{L}\{t\cos kt\} = \frac{s^2 - k^2}{(s^2 + k^2)^2}$$

29. $\mathcal{L}\{t\sinh kt\} = \frac{2ks}{(s^2 - k^2)^2}$

30.
$$\mathcal{L}$$
{ $t \cosh kt$ } = $\frac{s^2 + k^2}{(s^2 - k^2)^2}$

31. Apply the results in Example 5 and Problem 28 to show that

$$\mathcal{L}^{-1}\left\{\frac{1}{(s^2+k^2)^2}\right\} = \frac{1}{2k^3}(\sin kt - kt\cos kt).$$

Apply the extension of Theorem 1 in Eq. (22) to derive the Laplace transforms given in Problems 32 through 37.

- **32.** $\mathcal{L}{u(t-a)} = s^{-1}e^{-as}$ for a > 0.
- **33.** If f(t) = 1 on the interval [a, b] (where 0 < a < b) and f(t) = 0 otherwise, then

$$\mathcal{L}{f(t)} = \frac{e^{-as} - e^{-bs}}{s}$$

34. If $f(t) = (-1)^{[[t]]}$ is the square-wave function whose graph is shown in Fig. 10.2.9, then

$$\mathcal{L}{f(t)} = \frac{1}{s} \tanh \frac{s}{2}.$$

(Suggestion: Use the geometric series.)



FIGURE 10.2.9. The graph of the square-wave function of Problem 34.

35. If f(t) is the unit on-off function whose graph is shown in Fig. 10.2.10, then

$$\mathcal{L}{f(t)} = \frac{1}{s(1+e^{-s})}.$$



FIGURE 10.2.10. The graph of the on-off function of Problem 35.

36. If g(t) is the triangular wave function whose graph is shown in Fig. 10.2.11, then





FIGURE 10.2.11. The graph of the triangular wave function of Problem 36.

37. If f(t) is the sawtooth function whose graph is shown in Fig. 10.2.12, then

$$\mathcal{L}{f(t)} = \frac{1}{s^2} - \frac{e^{-s}}{s(1 - e^{-s})}$$

(Suggestion: Note that $f'(t) \equiv 1$ where it is defined.)



FIGURE 10.2.12. The graph of the sawtooth function of Problem 37.

10.2 Application Transforms of Initial Value Problems



The typical computer algebra system knows Theorem 1 and its corollary, hence can transform not only functions (as in the Section 10.1 application), but also entire initial value problems. We illustrate the technique here with *Mathematica* and in the Section 10.3 application with *Maple*. Consider the initial value problem

 $x'' + 4x = \sin 3t$, x(0) = x'(0) = 0

of Example 2. First we define the differential equation with its initial conditions, then load the Laplace transform package.

de = x''[t] + 4*x[t] == Sin[3*t]inits = { $x[0] \rightarrow 0, x'[0] \rightarrow 0$ }

The Laplace transform of the differential equation is given by

```
DE = LaplaceTransform[ de, t, s ]
```

The result of this command—which we do not show explicitly here—is a linear (algebraic) equation in the as yet unknown LaplaceTransform[x[t],t,s]. We proceed to solve for this transform X(s) of the unknown function x(t) and substitute the initial conditions.

```
X = Solve[DE, LaplaceTransform[x[t],t,s]]
X = X // Last // Last // Last
X = X /. inits
\frac{3}{(s^2+4)(s^2+9)}
```

Finally we need only compute an inverse transform to find x(t).

```
x = InverseLaplaceTransform[X,s,t]

\frac{1}{5}(3\cos(t)\sin(t) - \sin(3t))
x /. {Cos[t] sin[t] -> 1/2 sin[2t]}// Expand

\frac{3}{10}\sin(2t) - \frac{1}{5}\sin(3t)
```

Of course we could probably get this result immediately with **DSolve**, but the intermediate output generated by the steps shown here can be quite instructive. You can try it for yourself with the initial value problems in Problems 1 through 16.

10.3 Translation and Partial Fractions

As illustrated by Examples 1 and 2 of Section 10.2, the solution of a linear differential equation with constant coefficients can often be reduced to the matter of finding the inverse Laplace transform of a rational function of the form

$$R(s) = \frac{P(s)}{Q(s)} \tag{1}$$

where the degree of P(s) is less than that of Q(s). The technique for finding $\mathcal{L}^{-1}\{R(s)\}$ is based on the same method of partial fractions that we use in elementary calculus to integrate rational functions. The following two rules describe

the **partial fraction decomposition** of R(s), in terms of the factorization of the denominator Q(s) into linear factors and irreducible quadratic factors corresponding to the real and complex zeros, respectively, of Q(s).

RULE 1 Linear Factor Partial Fractions

The portion of the partial fraction decomposition of R(s) corresponding to the linear factor s - a of multiplicity n is a sum of n partial fractions, having the form

$$\frac{A_1}{s-a} + \frac{A_2}{(s-a)^2} + \dots + \frac{A_n}{(s-a)^n},$$
(2)

where A_1, A_2, \ldots , and A_n are constants.

RULE 2 Quadratic Factor Partial Fractions

The portion of the partial fraction decomposition corresponding to the irreducible quadratic factor $(s - a)^2 + b^2$ of multiplicity *n* is a sum of *n* partial fractions, having the form

$$\frac{A_1s + B_1}{(s-a)^2 + b^2} + \frac{A_2s + B_2}{[(s-a)^2 + b^2]^2} + \dots + \frac{A_ns + B_n}{[(s-a)^2 + b^2]^n},$$
(3)

where $A_1, A_2, \ldots, A_n, B_1, B_2, \ldots$, and B_n are constants.

Finding $\mathcal{L}^{-1}{R(s)}$ involves two steps. First we must find the partial fraction decomposition of R(s), and then we must find the inverse Laplace transform of each of the individual partial fractions of the types that appear in (2) and (3). The latter step is based on the following elementary property of Laplace transforms.

THEOREM 1 Translation on the s-Axis

If $F(s) = \mathcal{L}{f(t)}$ exists for s > c, then $\mathcal{L}{e^{at} f(t)}$ exists for s > a + c, and

$$\mathcal{L}\{e^{at} f(t)\} = F(s-a). \tag{4}$$

Equivalently,

$$\mathcal{L}^{-1}\{F(s-a)\} = e^{at} f(t).$$
(5)

Thus the translation $s \to s - a$ in the transform corresponds to multiplication of the original function of t by e^{at} .

Proof: If we simply replace s with s - a in the definition of $F(s) = \mathcal{L}{f(t)}$, we obtain

$$F(s-a) = \int_0^\infty e^{-(s-a)t} f(t) \, dt = \int_0^\infty e^{-st} \left[e^{at} f(t) \right] dt = \mathcal{L}\{e^{at} f(t)\}.$$

This is Eq. (4), and it is clear that Eq. (5) is the same.

If we apply the translation theorem to the formulas for the Laplace transforms of t^n , $\cos kt$, and $\sin kt$ that we already know—multiplying each of these functions

by e^{at} and replacing s with s - a in the transforms—we get the following additions to the table in Fig. 10.1.2.

$$f(t)$$
 $F(s)$

e

$$\frac{n!}{(s-a)^{n+1}} \qquad (s>a) \tag{6}$$

$$e^{at}\cos kt \qquad \frac{s-a}{(s-a)^2+k^2} \quad (s>a) \tag{7}$$

$$at \sin kt \qquad \frac{k}{(s-a)^2 + k^2} \quad (s > a) \tag{8}$$

For ready reference, all the Laplace transforms derived in this chapter are listed in the table of transforms that appears in the endpapers.

Damped mass-spring system Consider a mass-and-spring system with $m = \frac{1}{2}$, k = 17, and c = 3 in mks units (Fig. 10.3.1). As usual, let x(t) denote the displacement of the mass *m* from its equilibrium position. If the mass is set in motion with x(0) = 3 and x'(0) = 1, find x(t) for the resulting damped free oscillations.

The differential equation is $\frac{1}{2}x'' + 3x' + 17x = 0$, so we need to solve the initial value problem x'' + 6x' + 34x = 0; x(0) = 3, x'(0) = 1. We take the Laplace transform of each term of the differential equation. Because (obviously)

Solution

Example 1



FIGURE 10.3.2. The position function x(t) in Example 1.

k = 17

m =

 $\dot{x}(t)$

FIGURE 10.3.1. The mass–spring– dashpot system of Example 1.

which we solve for

 $\mathcal{L}{0} \equiv 0$, we get the equation

$$X(s) = \frac{3s+19}{s^2+6s+34} = 3 \cdot \frac{s+3}{(s+3)^2+25} + 2 \cdot \frac{5}{(s+3)^2+25}$$

 $\left[s^{2}X(s) - 3s - 1\right] + 6\left[sX(s) - 3\right] + 34X(s) = 0,$

Applying the formulas in (7) and (8) with a = -3 and k = 5, we now see that

 $x(t) = e^{-3t} \left(3\cos 5t + 2\sin 5t \right).$

Figure 10.3.2 shows the graph of this rapidly decaying damped oscillation.

Example 2 illustrates a useful technique for finding the partial fraction coefficients in the case of nonrepeated linear factors.

Example 2 Find the inverse Laplace transform of

$$R(s) = \frac{s^2 + 1}{s^3 - 2s^2 - 8s}.$$

Solution

Note that the denominator of
$$R(s)$$
 factors as $Q(s) = s(s+2)(s-4)$. Hence

$$\frac{s^2 + 1}{s^3 - 2s^2 - 8s} = \frac{A}{s} + \frac{B}{s+2} + \frac{C}{s-4}$$

Multiplication of each term of this equation by Q(s) yields

$$s^{2} + 1 = A(s + 2)(s - 4) + Bs(s - 4) + Cs(s + 2).$$

When we successively substitute the three zeros s = 0, s = -2, and s = 4 of the denominator Q(s) in this equation, we get the results

$$-8A = 1$$
, $12B = 5$, and $24C = 17$.

Thus $A = -\frac{1}{8}$, $B = \frac{5}{12}$, and $C = \frac{17}{24}$, so

$$\frac{s^2+1}{s^3-2s^2-8s} = -\frac{\frac{1}{8}}{s} + \frac{\frac{5}{12}}{s+2} + \frac{\frac{17}{24}}{s-4},$$

and therefore

$$\mathcal{L}^{-1}\left\{\frac{s^2+1}{s^3-2s^2-8s}\right\} = -\frac{1}{8} + \frac{5}{12}e^{-2t} + \frac{17}{24}e^{4t}.$$

Example 3 illustrates a differentiation technique for finding the partial fraction coefficients in the case of repeated linear factors.

Example 3 Solve the initial value problem

$$y'' + 4y' + 4y = t^2;$$
 $y(0) = y'(0) = 0.$

Solution The transformed equation is

$$s^{2}Y(s) + 4sY(s) + 4Y(s) = \frac{2}{s^{3}}.$$

Thus

$$Y(s) = \frac{2}{s^3(s+2)^2} = \frac{A}{s^3} + \frac{B}{s^2} + \frac{C}{s} + \frac{D}{(s+2)^2} + \frac{E}{s+2}.$$
(9)

To find A, B, and C, we multiply both sides by s^3 to obtain

$$\frac{2}{(s+2)^2} = A + Bs + Cs^2 + s^3F(s), \tag{10}$$

where $F(s) = D(s+2)^{-2} + E(s+2)^{-1}$ is the sum of the two partial fractions corresponding to $(s+2)^2$. Substitution of s = 0 in Eq. (10) yields $A = \frac{1}{2}$. To find *B* and *C*, we differentiate Eq. (10) twice to obtain

$$\frac{-4}{(s+2)^3} = B + 2Cs + 3s^2F(s) + s^3F'(s)$$
(11)

and

$$\frac{12}{(s+2)^4} = 2C + 6sF(s) + 6s^2F'(s) + s^3F''(s).$$
(12)

Now substitution of s = 0 in Eq. (11) yields $B = -\frac{1}{2}$, and substitution of s = 0 in Eq. (12) yields $C = \frac{3}{8}$.

To find D and E, we multiply each side in Eq. (9) by $(s + 2)^2$ to get

$$\frac{2}{s^3} = D + E(s+2) + (s+2)^2 G(s), \tag{13}$$

where $G(s) = As^{-3} + Bs^{-2} + Cs^{-1}$, and then differentiate to obtain

$$-\frac{6}{s^4} = E + 2(s+2)G(s) + (s+2)^2G'(s).$$
(14)

Substitution of s = -2 in Eqs. (13) and (14) now yields $D = -\frac{1}{4}$ and $E = -\frac{3}{8}$. Thus

$$Y(s) = \frac{\frac{1}{2}}{s^3} - \frac{\frac{1}{2}}{s^2} + \frac{\frac{3}{8}}{s} - \frac{\frac{1}{4}}{(s+2)^2} - \frac{\frac{3}{8}}{s+2},$$

so the solution of the given initial value problem is

$$y(t) = \frac{1}{4}t^2 - \frac{1}{2}t + \frac{3}{8} - \frac{1}{4}te^{-2t} - \frac{3}{8}e^{-2t}.$$

Examples 4, 5, and 6 illustrate techniques for dealing with quadratic factors in partial fraction decompositions.

Mass-spring-dashpot system Consider the mass-spring-dashpot system as in Example 1, but with initial conditions x(0) = x'(0) = 0 and with the imposed external force F(t) =15 sin 2t. Find the resulting transient motion and steady periodic motion of the mass. The initial value problem we need to solve is

Solution

$$x'' + 6x' + 34x = 30\sin 2t; \quad x(0) = x'(0) = 0.$$

The transformed equation is

$$s^{2}X(s) + 6sX(s) + 34X(s) = \frac{60}{s^{2} + 4}.$$

Hence

$$X(s) = \frac{60}{(s^2+4)[(s+3)^2+25]} = \frac{As+B}{s^2+4} + \frac{Cs+D}{(s+3)^2+25}.$$

When we multiply both sides by the common denominator, we get

$$60 = (As + B)[(s + 3)^{2} + 25] + (Cs + D)(s^{2} + 4).$$
(15)

To find A and B, we substitute the zero s = 2i of the quadratic factor $s^2 + 4$ in Eq. (15); the result is $60 = (2iA + B)[(2i + 3)^2 + 25],$

which we simplify to

$$60 = (-24A + 30B) + (60A + 12B)i.$$

We now equate real parts and imaginary parts on each side of this equation to obtain the two linear equations

-24A + 30B = 60 and 60A + 12B = 0,

which are readily solved for $A = -\frac{10}{29}$ and $B = \frac{50}{29}$.

To find C and D, we substitute the zero s = -3 + 5i of the quadratic factor $(s + 3)^2 + 25$ in Eq. (15) and get

$$60 = [C(-3+5i) + D][(-3+5i)^2 + 4],$$

which we simplify to

$$60 = (186C - 12D) + (30C - 30D)i$$

Again we equate real parts and imaginary parts; this yields the two linear equations

$$186C - 12D = 60$$
 and $30C - 30D = 0$,

and we readily find their solution to be $C = D = \frac{10}{29}$. With these values of the coefficients *A*, *B*, *C*, and *D*, our partial fractions decomposition of X(s) is

$$X(s) = \frac{1}{29} \left(\frac{-10s + 50}{s^2 + 4} + \frac{10s + 10}{(s + 3)^2 + 25} \right)$$
$$= \frac{1}{29} \left(\frac{-10s + 25 \cdot 2}{s^2 + 4} + \frac{10(s + 3) - 4 \cdot 5}{(s + 3)^2 + 25} \right)$$



$$x(t) = \frac{5}{29}(-2\cos 2t + 5\sin 2t) + \frac{2}{29}e^{-3t}(5\cos 5t - 2\sin 5t).$$

The terms of circular frequency 2 constitute the steady periodic forced oscillation of the mass, whereas the exponentially damped terms of circular frequency 5 constitute its transient motion, which disappears very rapidly (see Fig. 10.3.3). Note that the transient motion is nonzero even though both initial conditions are zero.



FIGURE 10.3.3. The periodic forced oscillation $x_{sp}(t)$, damped transient motion $x_{tr}(t)$, and solution $x(t) = x_{sp}(t) + x_{tr}(t)$ in Example 4.

Resonance and Repeated Quadratic Factors

The following two inverse Laplace transforms are useful in inverting partial fractions that correspond to the case of repeated quadratic factors:

$$\mathcal{L}^{-1}\left\{\frac{s}{(s^2+k^2)^2}\right\} = \frac{1}{2k}t\sin kt,$$
(16)

$$\mathcal{L}^{-1}\left\{\frac{1}{(s^2+k^2)^2}\right\} = \frac{1}{2k^3}(\sin kt - kt\cos kt).$$
(17)

These follow from Example 5 and Problem 31 of Section 10.2, respectively. Because of the presence in Eqs. (16) and (17) of the terms $t \sin kt$ and $t \cos kt$, a repeated quadratic factor ordinarily signals the phenomenon of resonance in an undamped mechanical or electrical system.

Example 5 Use Laplace transforms to solve the initial value problem

$$x'' + \omega_0^2 x = F_0 \sin \omega t; \quad x(0) = 0 = x'(0)$$

that determines the undamped forced oscillations of a mass on a spring. When we transform the differential equation, we get the equation

$$s^{2}X(s) + \omega_{0}^{2}X(s) = \frac{F_{0}\omega}{s^{2} + \omega^{2}}, \text{ so } X(s) = \frac{F_{0}\omega}{(s^{2} + \omega^{2})(s^{2} + \omega_{0}^{2})}$$

If $\omega \neq \omega_0$, we find without difficulty that

$$X(s) = \frac{F_0\omega}{\omega^2 - \omega_0^2} \left(\frac{1}{s^2 + \omega_0^2} - \frac{1}{s^2 + \omega^2} \right),$$

so it follows that

Solution

$$x(t) = \frac{F_0\omega}{\omega^2 - \omega_0^2} \left(\frac{1}{\omega_0} \sin \omega_0 t - \frac{1}{\omega} \sin \omega t \right).$$

But if $\omega = \omega_0$, we have

$$X(s) = \frac{F_0 \omega_0}{(s^2 + \omega_0^2)^2},$$

so Eq. (17) yields the resonance solution

$$x(t) = \frac{F_0}{2\omega_0^2} (\sin \omega_0 t - \omega_0 t \cos \omega_0 t).$$
(18)

Remark The solution curve defined in Eq. (18) bounces back and forth (see Fig. 10.3.4) between the "envelope curves" $x = \pm C(t)$ that are obtained by writing (18) in the form

$$x(t) = A(t)\cos\omega_0 t + B(t)\sin\omega_0 t$$

and then defining the usual "amplitude" $C = \sqrt{A^2 + B^2}$. In this case we find that

$$C(t) = \frac{F_0}{2\omega_0^2} \sqrt{\omega_0^2 t^2 + 1}.$$

This technique for constructing envelope curves of resonance solutions is illustrated further in the application material for this section.

Solve the initial value problem

$$y^{(4)} + 2y'' + y = 4te^t;$$
 $y(0) = y'(0) = y''(0) = y^{(3)}(0) = 0$



FIGURE 10.3.4. The resonance solution in (18) with $\omega_0 = \frac{1}{2}$ and $F_0 = 1$, together with its envelope curves $x = \pm C(t)$.

584 CHAPTER 10 LAPLACE TRANSFORM METHODS

Solution First we observe that

$$\mathcal{L}\{y''(t)\} = s^2 Y(s), \quad \mathcal{L}\{y^{(4)}(t)\} = s^4 Y(s), \text{ and } \mathcal{L}\{te^t\} = \frac{1}{(s-1)^2}.$$

Hence the transformed equation is

$$(s^4 + 2s^2 + 1)Y(s) = \frac{4}{(s-1)^2}.$$

Thus our problem is to find the inverse transform of

$$Y(s) = \frac{4}{(s-1)^2(s^2+1)^2}$$
$$= \frac{A}{(s-1)^2} + \frac{B}{s-1} + \frac{Cs+D}{(s^2+1)^2} + \frac{Es+F}{s^2+1}.$$
(19)

If we multiply by the common denominator $(s-1)^2(s^2+1)^2$, we get the equation

$$A(s^{2}+1)^{2} + B(s-1)(s^{2}+1)^{2} + Cs(s-1)^{2} + D(s-1)^{2} + Es(s-1)^{2}(s^{2}+1) + F(s-1)^{2}(s^{2}+1) = 4.$$
 (20)

Upon substituting s = 1 we find that A = 1.

Equation (20) is an identity that holds for all values of s. To find the values of the remaining coefficients, we substitute in succession the values s = 0, s = -1, s = 2, s = -2, and s = 3 in Eq. (20). This yields the system

of five linear equations in *B*, *C*, *D*, *E*, and *F*. With the aid of a calculator programmed to solve linear systems, we find that B = -2, C = 2, D = 0, E = 2, and F = 1.

We now substitute in Eq. (19) the coefficients we have found, and thus obtain

$$Y(s) = \frac{1}{(s-1)^2} - \frac{2}{s-1} + \frac{2s}{(s^2+1)^2} + \frac{2s+1}{s^2+1}$$

Recalling Eq. (16), the translation property, and the familiar transforms of $\cos t$ and $\sin t$, we see finally that the solution of the given initial value problem is

$$y(t) = (t-2)e^{t} + (t+1)\sin t + 2\cos t.$$

10.3 **Problems**

Apply the translation theorem to find the Laplace transforms of the functions in Problems 1 through 4.

1.
$$f(t) = t^4 e^{\pi t}$$
 2. $f(t) = t^{3/2} e^{-4t}$

3.
$$f(t) = e^{-2t} \sin 3\pi t$$
 4. $f(t) = e^{-t/2} \cos 2\left(t - \frac{1}{8}\pi\right)$

Apply the translation theorem to find the inverse Laplace transforms of the functions in Problems 5 through 10.

5.
$$F(s) = \frac{3}{2s-4}$$
 6. $F(s) = \frac{s-1}{(s+1)^3}$

7.
$$F(s) = \frac{1}{s^2 + 4s + 4}$$

8. $F(s) = \frac{s + 2}{s^2 + 4s + 5}$
9. $F(s) = \frac{3s + 5}{s^2 - 6s + 25}$
10. $F(s) = \frac{2s - 3}{9s^2 - 12s + 20}$

Use partial fractions to find the inverse Laplace transforms of the functions in Problems 11 through 22.

11.
$$F(s) = \frac{1}{s^2 - 4}$$

12. $F(s) = \frac{5s - 6}{s^2 - 3s}$
13. $F(s) = \frac{5 - 2s}{s^2 + 7s + 10}$
14. $F(s) = \frac{5s - 4}{s^3 - s^2 - 2s}$

15.
$$F(s) = \frac{1}{s^3 - 5s^2}$$

16. $F(s) = \frac{1}{(s^2 + s - 6)^2}$
17. $F(s) = \frac{1}{s^4 - 16}$
18. $F(s) = \frac{s^3}{(s - 4)^4}$
19. $F(s) = \frac{s^2 - 2s}{s^4 + 5s^2 + 4}$
20. $F(s) = \frac{1}{s^4 - 8s^2 + 16}$
21. $F(s) = \frac{s^2 + 3}{(s^2 + 2s + 2)^2}$
22. $F(s) = \frac{2s^3 - s^2}{(4s^2 - 4s + 5)^2}$

Use the factorization

$$s^4 + 4a^4 = (s^2 - 2as + 2a^2)(s^2 + 2as + 2a^2)$$

to derive the inverse Laplace transforms listed in Problems 23 through 26.

23.
$$\mathcal{L}^{-1}\left\{\frac{s^3}{s^4 + 4a^4}\right\} = \cosh at \cos at$$

24. $\mathcal{L}^{-1}\left\{\frac{s}{s^4 + 4a^4}\right\} = \frac{1}{2a^2}\sinh at \sin at$
25. $\mathcal{L}^{-1}\left\{\frac{s^2}{s^4 + 4a^4}\right\} = \frac{1}{2a}(\cosh at \sin at + \sinh at \cos at)$
26. $\mathcal{L}^{-1}\left\{\frac{1}{s^4 + 4a^4}\right\} = \frac{1}{4a^3}(\cosh at \sin at - \sinh at \cos at)$

Use Laplace transforms to solve the initial value problems in Problems 27 through 38.

27. x'' + 6x' + 25x = 0; x(0) = 2, x'(0) = 3 **28.** x'' - 6x' + 8x = 2; x(0) = x'(0) = 0 **29.** x'' - 4x = 3t; x(0) = x'(0) = 0 **30.** $x'' + 4x' + 8x = e^{-t}; x(0) = x'(0) = 0$ **31.** $x^{(3)} + x'' - 6x' = 0; x(0) = 0, x'(0) = x''(0) = 1$ **32.** $x^{(4)} - x = 0; x(0) = 1, x'(0) = x''(0) = x^{(3)}(0) = 0$ **33.** $x^{(4)} + x = 0; x(0) = x'(0) = x''(0) = 0, x^{(3)}(0) = 1$

10.3 TRANSLATION AND PARTIAL FRACTIONS 585

- **34.** $x^{(4)} + 13x'' + 36x = 0; x(0) = x''(0) = 0, x'(0) = 2, x^{(3)}(0) = -13$
- **35.** $x^{(4)} + 8x'' + 16x = 0; x(0) = x'(0) = x''(0) = 0,$ $x^{(3)}(0) = 1$
- **36.** $x^{(4)} + 2x'' + x = e^{2t}$; $x(0) = x'(0) = x''(0) = x^{(3)}(0) = 0$

37. $x'' + 4x' + 13x = te^{-t}$; x(0) = 0, x'(0) = 2

38. $x'' + 6x' + 18x = \cos 2t$; x(0) = 1, x'(0) = -1

Resonance

Problems 39 and 40 illustrate two types of resonance in a mass–spring–dashpot system with given external force F(t) and with the initial conditions x(0) = x'(0) = 0.

- **39.** Suppose that m = 1, k = 9, c = 0, and $F(t) = 6 \cos 3t$. Use the inverse transform given in Eq. (16) to derive the solution $x(t) = t \sin 3t$. Construct a figure that illustrates the resonance that occurs.
- **40.** Suppose that m = 1, k = 9.04, c = 0.4, and $F(t) = 6e^{-t/5}\cos 3t$. Derive the solution

$$x(t) = te^{-t/5}\sin 3t.$$

Show that the maximum value of the amplitude function $A(t) = te^{-t/5}$ is A(5) = 5/e. Thus (as indicated in Fig. 10.3.5) the oscillations of the mass increase in amplitude during the first 5 s before being damped out as $t \to +\infty$.



FIGURE 10.3.5. The graph of the damped oscillation in Problem 40.

10.3 Application **Damping and Resonance Investigations**

Go to goo.gl/4ntUK9 to download this application's computing resources including *Maple/Mathematica*/MATLAB.

Here we outline a *Maple* investigation of the behavior of the mass-spring-dashpot system

$$mx'' + cx' + kx = F(t), \quad x(0) = x'(0) = 0$$
(1)

with parameter values

in response to a variety of possible external forces:

1.
$$F(t) \equiv 226$$

This should give damped oscillations "leveling off" to a constant solution (why?).

2.
$$F(t) = 901 \cos 3t$$

With this periodic external force you should see a steady periodic oscillation with an exponentially damped transient motion (as illustrated in Fig. 5.6.13).

3.
$$F(t) = 900e^{-t/5}\cos 3t$$

Now the periodic external force is exponentially damped, and the transform X(s) includes a repeated quadratic factor that signals the presence of a resonance phenomenon. The response x(t) is a constant multiple of that shown in Fig. 10.3.5.

4.
$$F(t) = 900te^{-t/5}\cos 3t$$

We have inserted the factor t to make it a bit more interesting. The solution in this case is illustrated below.

5.
$$F(t) = 162t^3 e^{-t/5} \cos 3t$$

In this case you'll find that the transform X(s) involves the *fifth* power of a quadratic factor, and its inverse transform by manual methods would be impossibly tedious.

To illustrate the *Maple* approach, we first set up the differential equation corresponding to Case 4.

```
F := 900*t*exp(-t/5)*cos(3*t);
de := m*diff(x(t),t$2) + c*diff(x(t),t) + k*x(t) = F;
```

Then we apply the Laplace transform and substitute the initial conditions.

```
with(inttrans):
DE := laplace(de, t, s):
X(s) := solve(DE, laplace(x(t), t, s)):
X(s) := simplify(subs(x(0)=0, D(x)(0)=0, X(s)));
```

At this point the command factor(denom(X(s))) shows that

$$X(s) = \frac{22500(25s^2 + 10s - 224)}{(25s^2 + 10s + 226)^3}$$

The cubed quadratic factor would be difficult to handle manually, but the command

x(t) := invlaplace(X(s), s, t);

soon yields

$$x(t) = e^{-t/5} \left(t \cos 3t + \left(3t^2 - \frac{1}{3} \right) \sin 3t \right).$$

The amplitude function for these damped oscillations is defined by

```
C(t) := \exp(-t/5) * \operatorname{sqrt}(t^2 + (3*t^2 - 1/3)^2);
```

and finally the command

 $plot({x(t), C(t), -C(t)}, t=0..40);$

produces the plot shown in Fig. 10.3.6. The resonance resulting from the repeated quadratic factor consists of a temporary buildup before the oscillations are damped out.

For a similar solution in one of the other cases listed previously, you need only enter the appropriate force \mathbf{F} in the initial command above and then re-execute the subsequent commands. To see the advantage of using Laplace transforms, set up the differential equation **de** for Case 5 and examine the result of the command

```
dsolve({de, x(0)=0, D(x)(0)=0}, x(t));
```



FIGURE 10.3.6. The resonance solution and its envelope curves in Case 4.

10.4 DERIVATIVES, INTEGRALS, AND PRODUCTS OF TRANSFORM\$587

Of course you can substitute your own favorite mass-spring-dashpot parameters for those used here. But it will simplify the calculations if you choose m, c, and k so that

$$mr^{2} + cr + k = (pr + a)^{2} + b^{2}$$
⁽²⁾

where p, a, and b are integers. One way is to select the latter integers first, then use Eq. (2) to determine m, c, and k.

10.4 Derivatives, Integrals, and Products of Transforms

The Laplace transform of the (initially unknown) solution of a differential equation is sometimes recognizable as the product of the transforms of two *known* functions. For example, when we transform the initial value problem

$$x'' + x = \cos t; \quad x(0) = x'(0) = 0,$$

we get

$$X(s) = \frac{s}{(s^2 + 1)^2} = \frac{s}{s^2 + 1} \cdot \frac{1}{s^2 + 1} = \mathcal{L}\{\cos t\} \cdot \mathcal{L}\{\sin t\}.$$

This strongly suggests that there ought to be a way of combining the two functions $\sin t$ and $\cos t$ to obtain a function x(t) whose transform is the *product* of *their* transforms. But obviously x(t) is *not* simply the product of $\cos t$ and $\sin t$, because

$$\mathcal{L}\{\cos t \sin t\} = \mathcal{L}\left\{\frac{1}{2}\sin 2t\right\} = \frac{1}{s^2 + 4} \neq \frac{s}{(s^2 + 1)^2}.$$

Thus $\mathcal{L}\{\cos t \sin t\} \neq \mathcal{L}\{\cos t\} \cdot \mathcal{L}\{\sin t\}.$

Theorem 1 of this section will tell us that the function

$$h(t) = \int_0^t f(\tau)g(t-\tau)\,d\tau \tag{1}$$

has the desired property that

$$\mathcal{L}{h(t)} = H(s) = F(s) \cdot G(s).$$
⁽²⁾

The new function of t defined as the integral in (1) depends only on f and g and is called the *convolution* of f and g. It is denoted by f * g, the idea being that it is a new type of product of f and g, so tailored that its transform is the product of the transforms of f and g.

DEFINITION The Convolution of Two Functions

The **convolution** f * g of the piecewise continuous functions f and g is defined for $t \ge 0$ as follows:

$$(f * g)(t) = \int_0^t f(\tau)g(t - \tau) \, d\tau.$$
 (3)

We will also write f(t) * g(t) when convenient. In terms of the convolution product, Theorem 1 of this section says that

$$\mathcal{L}{f \ast g} = \mathcal{L}{f} \cdot \mathcal{L}{g}$$

If we make the substitution $u = t - \tau$ in the integral in (3), we see that

$$f(t) * g(t) = \int_0^t f(\tau)g(t-\tau) d\tau = \int_t^0 f(t-u)g(u)(-du)$$
$$= \int_0^t g(u)f(t-u) du = g(t) * f(t).$$

Thus the convolution is *commutative*: f * g = g * f.

1 The convolution of cos t and sin t is

$$(\cos t) * (\sin t) = \int_0^t \cos \tau \sin(t - \tau) \, d\tau$$

We apply the trigonometric identity

$$\cos A \sin B = \frac{1}{2} \left[\sin(A+B) - \sin(A-B) \right]$$

to obtain

$$(\cos t) * (\sin t) = \int_0^t \frac{1}{2} [\sin t - \sin(2\tau - t)] d\tau$$
$$= \frac{1}{2} \bigg[\tau \sin t + \frac{1}{2} \cos(2\tau - t) \bigg]_{\tau=0}^t;$$

that is,

$$(\cos t) * (\sin t) = \frac{1}{2}t\sin t.$$

And we recall from Example 5 of Section 10.2 that the Laplace transform of $\frac{1}{2}t \sin t$ is indeed $s/(s^2 + 1)^2$.

Theorem 1 is proved at the end of this section.

THEOREM 1 The Convolution Property

Suppose that f(t) and g(t) are piecewise continuous for $t \ge 0$ and that |f(t)| and |g(t)| are bounded by Me^{ct} as $t \to +\infty$. Then the Laplace transform of the convolution f(t) * g(t) exists for s > c; moreover,

$$\mathcal{L}\{f(t) * g(t)\} = \mathcal{L}\{f(t)\} \cdot \mathcal{L}\{g(t)\}$$
(4)

and

 \succ

$$\mathcal{L}^{-1}\{F(s) \cdot G(s)\} = f(t) * g(t).$$
(5)

Thus we can find the inverse transform of the product $F(s) \cdot G(s)$, provided that we can evaluate the integral

$$\mathcal{L}^{-1}\{F(s)\cdot G(s)\} = \int_0^t f(\tau)g(t-\tau)\,d\,\tau.$$
^(5')

Example 2 illustrates the fact that convolution often provides a convenient alternative to the use of partial fractions for finding inverse transforms.

Example 2

With $f(t) = \sin 2t$ and $g(t) = e^t$, convolution yields

$$\mathcal{L}^{-1}\left\{\frac{2}{(s-1)(s^2+4)}\right\} = (\sin 2t) * e^t = \int_0^t e^{t-\tau} \sin 2\tau \, d\tau$$
$$= e^t \int_0^t e^{-\tau} \sin 2\tau \, d\tau = e^t \left[\frac{e^{-\tau}}{5}(-\sin 2\tau - 2\cos 2\tau)\right]_0^t,$$

so

$$\mathcal{L}^{-1}\left\{\frac{2}{(s-1)(s^2+4)}\right\} = \frac{2}{5}e^t - \frac{1}{5}\sin 2t - \frac{2}{5}\cos 2t.$$

Differentiation of Transforms

According to Theorem 1 of Section 10.2, if f(0) = 0 then differentiation of f(t) corresponds to multiplication of its transform by *s*. Theorem 2, proved at the end of this section, tells us that differentiation of the transform F(s) corresponds to multiplication of the original function f(t) by -t.

THEOREM 2 Differentiation of Transforms

If f(t) is piecewise continuous for $t \ge 0$ and $|f(t)| \le Me^{ct}$ as $t \to +\infty$, then

$$\mathcal{L}\{-tf(t)\} = F'(s) \tag{6}$$

for s > c. Equivalently,

$$f(t) = \mathcal{L}^{-1}\{F(s)\} = -\frac{1}{t}\mathcal{L}^{-1}\{F'(s)\}.$$
(7)

Repeated application of Eq. (6) gives

$$\mathcal{L}\{t^{n} f(t)\} = (-1)^{n} F^{(n)}(s)$$
(8)

for $n = 1, 2, 3, \ldots$

Example 3 Find \mathcal{L} { $t^2 \sin kt$ }. Solution Equation (8) gives

>

$$\mathcal{L}\{t^{2}\sin kt\} = (-1)^{2} \frac{d^{2}}{ds^{2}} \left(\frac{k}{s^{2}+k^{2}}\right)$$
$$= \frac{d}{ds} \left[\frac{-2ks}{(s^{2}+k^{2})^{2}}\right] = \frac{6ks^{2}-2k^{3}}{(s^{2}+k^{2})^{3}}.$$
(9)

The form of the differentiation property in Eq. (7) is often helpful in finding an inverse transform when the *derivative* of the transform is easier to work with than the transform itself. Example 4Find \mathcal{L}^{-1} {tan⁻¹(1/s)}.SolutionThe derivative of tan⁻¹(1/s) is a simple rational function, so we apply Eq. (7):

$$\mathcal{L}^{-1}\left\{\tan^{-1}\frac{1}{s}\right\} = -\frac{1}{t}\mathcal{L}^{-1}\left\{\frac{d}{ds}\tan^{-1}\frac{1}{s}\right\}$$
$$= -\frac{1}{t}\mathcal{L}^{-1}\left\{\frac{-1/s^2}{1+(1/s)^2}\right\}$$
$$= -\frac{1}{t}\mathcal{L}^{-1}\left\{\frac{-1}{s^2+1}\right\} = -\frac{1}{t}(-\sin t).$$

Therefore,

$$\mathcal{L}^{-1}\left\{\tan^{-1}\frac{1}{s}\right\} = \frac{\sin t}{t}.$$

Equation (8) can be applied to transform a linear differential equation having polynomial, rather than constant, coefficients. The result will be a differential equation involving the transform; whether this procedure leads to success depends, of course, on whether we can solve the new equation more readily than the old one.

Example 5

Let x(t) be the solution of Bessel's equation of order zero,

$$tx'' + x' + tx = 0,$$

such that x(0) = 1 and x'(0) = 0. This solution of Bessel's equation is customarily denoted by $J_0(t)$. Because

$$\mathcal{L}\{x'(t)\} = sX(s) - 1 \quad \text{and} \quad \mathcal{L}\{x''(t)\} = s^2 X(s) - s,$$

and because x and x'' are each multiplied by t, application of Eq. (6) yields the transformed equation

$$-\frac{d}{ds}\left[s^2X(s) - s\right] + \left[sX(s) - 1\right] - \frac{d}{ds}\left[X(s)\right] = 0.$$

The result of differentiation and simplification is the differential equation

$$(s^2 + 1)X'(s) + sX(s) = 0.$$

This equation is separable—

its general solution is

$$X(s) = \frac{C}{\sqrt{s^2 + 1}}.$$

 $\frac{X'(s)}{X(s)} = -\frac{s}{s^2 + 1};$

In Problem 39 we outline the argument that C = 1. Because $X(s) = \mathcal{L}{J_0(t)}$, it follows that

$$\mathcal{L}\{J_0(t)\} = \frac{1}{\sqrt{s^2 + 1}}.$$
(10)

Integration of Transforms

Differentiation of F(s) corresponds to multiplication of f(t) by t (together with a change of sign). It is therefore natural to expect that integration of F(s) will correspond to division of f(t) by t. Theorem 3, proved at the end of this section, confirms this, provided that the resulting quotient f(t)/t remains well behaved as $t \rightarrow 0$ from the right; that is, provided that

$$\lim_{t \to 0^+} \frac{f(t)}{t} \quad \text{exists and is finite.}$$
(11)

THEOREM 3 Integration of Transforms

Suppose that f(t) is piecewise continuous for $t \ge 0$, that f(t) satisfies the condition in (11), and that $|f(t)| \le Me^{ct}$ as $t \to +\infty$. Then

$$\mathcal{L}\left\{\frac{f(t)}{t}\right\} = \int_{s}^{\infty} F(\sigma) \, d\sigma \tag{12}$$

for s > c. Equivalently,

$$f(t) = \mathcal{L}^{-1}\{F(s)\} = t \mathcal{L}^{-1}\left\{\int_{s}^{\infty} F(\sigma) \, d\sigma\right\}.$$
 (13)

Example 6 Find $\mathcal{L}\{(\sinh t)/t\}$.

Solution

We first verify that the condition in (11) holds:

$$\lim_{t \to 0} \frac{\sinh t}{t} = \lim_{t \to 0} \frac{e^t - e^{-t}}{2t} = \lim_{t \to 0} \frac{e^t + e^{-t}}{2} = 1,$$

with the aid of l'Hôpital's rule. Then Eq. (12), with $f(t) = \sinh t$, yields

$$\mathcal{L}\left\{\frac{\sinh t}{t}\right\} = \int_{s}^{\infty} \mathcal{L}\left\{\sinh t\right\} d\sigma = \int_{s}^{\infty} \frac{d\sigma}{\sigma^{2} - 1}$$
$$= \frac{1}{2} \int_{s}^{\infty} \left(\frac{1}{\sigma - 1} - \frac{1}{\sigma + 1}\right) d\sigma = \frac{1}{2} \left[\ln \frac{\sigma - 1}{\sigma + 1}\right]_{s}^{\infty}$$

Therefore,

$$\mathscr{L}\left\{\frac{\sinh t}{t}\right\} = \frac{1}{2}\ln\frac{s+1}{s-1},$$

because $\ln 1 = 0$.

The form of the integration property in Eq. (13) is often helpful in finding an inverse transform when the indefinite *integral* of the transform is easier to handle than the transform itself.

Example 7 Find
$$\mathcal{L}^{-1}\{2s/(s^2-1)^2\}$$
.

Solution We could use partial fractions, but it is much simpler to apply Eq. (13). This gives

$$\mathcal{L}^{-1}\left\{\frac{2s}{(s^2-1)^2}\right\} = t\mathcal{L}^{-1}\left\{\int_s^\infty \frac{2\sigma}{(\sigma^2-1)^2} d\sigma\right\}$$
$$= t\mathcal{L}^{-1}\left\{\left[\frac{-1}{\sigma^2-1}\right]_s^\infty\right\} = t\mathcal{L}^{-1}\left\{\frac{1}{s^2-1}\right\}$$

and therefore

$$\mathcal{L}^{-1}\left\{\frac{2s}{(s^2-1)^2}\right\} = t \sinh t.$$

*Proofs of Theorems

Proof of Theorem 1: The transforms F(s) and G(s) exist when s > c by Theorem 2 of Section 10.1. For any $\tau > 0$ the definition of the Laplace transform gives

$$G(s) = \int_0^\infty e^{-su}g(u)\,du = \int_\tau^\infty e^{-s(t-\tau)}g(t-\tau)\,dt \quad (u=t-\tau),$$

and therefore

$$G(s) = e^{s\tau} \int_0^\infty e^{-st} g(t-\tau) \, dt,$$

because we may *define* f(t) and g(t) to be zero for t < 0. Then

$$F(s)G(s) = G(s) \int_0^\infty e^{-s\tau} f(\tau) d\tau = \int_0^\infty e^{-s\tau} f(\tau)G(s) d\tau$$
$$= \int_0^\infty e^{-s\tau} f(\tau) \left(e^{s\tau} \int_0^\infty e^{-st} g(t-\tau) dt \right) d\tau$$
$$= \int_0^\infty \left(\int_0^\infty e^{-st} f(\tau) g(t-\tau) dt \right) d\tau.$$

Now our hypotheses on f and g imply that the order of integration may be reversed. (The proof of this requires a discussion of uniform convergence of improper integrals, and can be found in Chapter 2 of Churchill's *Operational Mathematics*, 3rd ed. (New York: McGraw-Hill, 1972).) Hence

$$F(s)G(s) = \int_0^\infty \left(\int_0^\infty e^{-st} f(\tau)g(t-\tau) \, d\tau \right) dt$$
$$= \int_0^\infty e^{-st} \left(\int_0^t f(\tau)g(t-\tau) \, d\tau \right) dt$$
$$= \int_0^\infty e^{-st} \left[f(t) * g(t) \right] \, dt,$$

and therefore,

$$F(s)G(s) = \mathcal{L}\{f(t) * g(t)\}$$

We replace the upper limit of the inner integral with t because $g(t - \tau) = 0$ whenever $\tau > t$. This completes the proof of Theorem 1.

Proof of Theorem 2: Because

$$F(s) = \int_0^\infty e^{-st} f(t) \, dt,$$

differentiation under the integral sign yields

$$F(s) = \frac{d}{ds} \int_0^\infty e^{-st} f(t) dt$$
$$= \int_0^\infty \frac{d}{ds} \left[e^{-st} f(t) \right] dt = \int_0^\infty e^{-st} \left[-tf(t) \right] dt;$$

thus

$$F'(s) = \mathcal{L}\{-tf(t)\},\$$

which is Eq. (6). We obtain Eq. (7) by applying \mathcal{L}^{-1} and then dividing by -t. The validity of differentiation under the integral sign depends on uniform convergence of the resulting integral; this is discussed in Chapter 2 of the book by Churchill just mentioned.

Proof of Theorem 3: By definition,

$$F(\sigma) = \int_0^\infty e^{-\sigma t} f(t) \, dt.$$

So integration of $F(\sigma)$ from *s* to $+\infty$ gives

$$\int_{s}^{\infty} F(\sigma) \, d\sigma = \int_{s}^{\infty} \left(\int_{0}^{\infty} e^{-\sigma t} f(t) \, dt \right) d\sigma.$$

Under the hypotheses of the theorem, the order of integration may be reversed (see Churchill's book once again); it follows that

$$\int_{s}^{\infty} F(\sigma) \, d\sigma = \int_{0}^{\infty} \left(\int_{s}^{\infty} e^{-\sigma t} f(t) \, d\sigma \right) dt$$
$$= \int_{0}^{\infty} \left[\frac{e^{-\sigma t}}{-t} \right]_{\sigma=s}^{\infty} f(t) \, dt$$
$$= \int_{0}^{\infty} e^{-st} \frac{f(t)}{t} \, dt = \mathcal{L} \left\{ \frac{f(t)}{t} \right\}$$

This verifies Eq. (12), and Eq. (13) follows upon first applying \mathcal{L}^{-1} and then multiplying by *t*.

10.4 **Problems**

Find the convolution f(t) * g(t) in Problems 1 through 6.

1. $f(t) = t, g(t) \equiv 1$ **3.** $f(t) = g(t) = \sin t$ **5.** $f(t) = g(t) = e^{at}$ **6.** $f(t) = e^{at}, g(t) = e^{bt}$ ($a \neq b$) **2.** $f(t) = t, g(t) = e^{at}$ **4.** $f(t) = t^2, g(t) = \cos t$

Apply the convolution theorem to find the inverse Laplace transforms of the functions in Problems 7 through 14.

7.
$$F(s) = \frac{1}{s(s-3)}$$

8. $F(s) = \frac{1}{s(s^2+4)}$
9. $F(s) = \frac{1}{(s^2+9)^2}$
10. $F(s) = \frac{1}{s^2(s^2+4)^2}$
11. $F(s) = \frac{s^2}{(s^2+4)^2}$
12. $F(s) = \frac{1}{s(s^2+4s+5)}$
13. $F(s) = \frac{s}{(s-3)(s^2+1)}$
14. $F(s) = \frac{s}{s^4+5s^2+4}$

In Problems 15 through 22, apply either Theorem 2 or Theorem 3 to find the Laplace transform of f(t).

15.
$$f(t) = t \sin 3t$$

16. $f(t) = t^2 \cos 2t$
17. $f(t) = te^{2t} \cos 3t$
18. $f(t) = te^{-t} \sin^2 t$
19. $f(t) = \frac{\sin t}{t}$
20. $f(t) = \frac{1 - \cos 2t}{t}$
21. $f(t) = \frac{e^{3t} - 1}{t}$
22. $f(t) = \frac{e^t - e^{-t}}{t}$

Find the inverse transforms of the functions in Problems 23 through 28.

23.
$$F(s) = \ln \frac{s-2}{s+2}$$

24. $F(s) = \ln \frac{s^2+1}{s^2+4}$
25. $F(s) = \ln \frac{s^2+1}{(s+2)(s-3)}$
26. $F(s) = \tan^{-1} \frac{3}{s+2}$

27. $F(s) = \ln\left(1 + \frac{1}{s^2}\right)$ **28.** $F(s) = \frac{s}{(s^2 + 1)^3}$

In Problems 29 through 34, transform the given differential equation to find a nontrivial solution such that x(0) = 0.

- **29.** tx'' + (t-2)x' + x = 0 **30.** tx'' + (3t-1)x' + 3x = 0 **31.** tx'' - (4t+1)x' + 2(2t+1)x = 0 **32.** tx'' + 2(t-1)x' - 2x = 0 **33.** tx'' - 2x' + tx = 0**34.** tx'' + (4t-2)x' + (13t-4)x = 0
- **35.** Apply the convolution theorem to show that

$$\mathcal{L}^{-1}\left\{\frac{1}{(s-1)\sqrt{s}}\right\} = \frac{2e^t}{\sqrt{\pi}}\int_0^{\sqrt{t}} e^{-u^2} du = e^t \operatorname{erf}\sqrt{t}.$$

(Suggestion: Substitute $u = \sqrt{t}$.)

In Problems 36 through 38, apply the convolution theorem to derive the indicated solution x(t) of the given differential equation with initial conditions x(0) = x'(0) = 0.

36.
$$x'' + 4x = f(t); x(t) = \frac{1}{2} \int_0^t f(t-\tau) \sin 2\tau \, d\tau$$

37. $x'' + 2x' + x = f(t); x(t) = \int_0^t \tau e^{-\tau} f(t-\tau) \, d\tau$
38. $x'' + 4x' + 13x = f(t);$
 $x(t) = \frac{1}{3} \int_0^t f(t-\tau) e^{-2\tau} \sin 3\tau \, d\tau$

594 CHAPTER 10 LAPLACE TRANSFORM METHODS

Termwise Inverse Transformation of Series

In Chapter 2 of Churchill's Operational Mathematics, the following theorem is proved. Suppose that f(t) is continuous for $t \ge 0$, that f(t) is of exponential order as $t \to +\infty$, and that

$$F(s) = \sum_{n=0}^{\infty} \frac{a_n}{s^{n+k+1}}$$

where $0 \leq k < 1$ and the series converges absolutely for s > c. Then

$$f(t) = \sum_{n=0}^{\infty} \frac{a_n t^{n+k}}{\Gamma(n+k+1)}.$$

Apply this result in Problems 39 through 41.

39. In Example 5 it was shown that

$$\mathcal{L}\{J_0(t)\} = \frac{C}{\sqrt{s^2 + 1}} = \frac{C}{s} \left(1 + \frac{1}{s^2}\right)^{-1/2}.$$

Expand with the aid of the binomial series and then compute the inverse transformation term by term to obtain

$$J_0(t) = C \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n}}{2^{2n} (n!)^2}.$$

Finally, note that $J_0(0) = 1$ implies that C = 1.

40. Expand the function $F(s) = s^{-1/2}e^{-1/s}$ in powers of s^{-1} to show that

$$\mathcal{L}^{-1}\left\{\frac{1}{\sqrt{s}}e^{-1/s}\right\} = \frac{1}{\sqrt{\pi t}}\cos 2\sqrt{t}.$$

41. Show that

$$\mathcal{L}^{-1}\left\{\frac{1}{s}e^{-1/s}\right\} = J_0\left(2\sqrt{t}\right).$$

10.5 Periodic and Piecewise Continuous Input Functions



FIGURE 10.5.1. The graph of the unit step function at t = a.

Mathematical models of mechanical or electrical systems often involve functions with discontinuities corresponding to external forces that are turned abruptly on or off. One such simple on–off function is the unit step function that we introduced in Section 10.1. Recall that the *unit step function* at t = a is defined by

$$u_{a}(t) = u(t-a) = \begin{cases} 0 & \text{if } t < a, \\ 1 & \text{if } t \ge a. \end{cases}$$
(1)

The notation $u_a(t)$ indicates succinctly where the unit upward step in value takes place (Fig. 10.5.1), whereas u(t - a) connotes the sometimes useful idea of a "time delay" *a* before the step is made.

In Example 8 of Section 10.1 we saw that if $a \ge 0$, then

$$\mathscr{L}\{u(t-a)\} = \frac{e^{-as}}{s}.$$
(2)

Because $\mathcal{L}{u(t)} = 1/s$, Eq. (2) implies that multiplication of the transform of u(t) by e^{-as} corresponds to the translation $t \to t - a$ in the original independent variable. Theorem 1 tells us that this fact, when properly interpreted, is a general property of the Laplace transformation.

THEOREM 1 Translation on the t-Axis

If $\mathcal{L}{f(t)}$ exists for s > c, then

$$\mathscr{L}\{u(t-a)f(t-a)\} = e^{-as}F(s)$$
(3a)

and

$$\mathcal{L}^{-1}\{e^{-as}F(s)\} = u(t-a)f(t-a)$$
(3b)

for s > c + a.

Note that

$$u(t-a)f(t-a) = \begin{cases} 0 & \text{if } t < a, \\ f(t-a) & \text{if } t \ge a. \end{cases}$$
(4)



FIGURE 10.5.2. Translation of f(t) *a* units to the right.

Thus Theorem 1 implies that $\mathcal{L}^{-1}\{e^{-as}F(s)\}$ is the function whose graph for $t \ge a$ is the translation by *a* units to the right of the graph of f(t) for $t \ge 0$. Note that the part (if any) of the graph of f(t) to the left of t = 0 is "cut off" and is *not* translated (Fig. 10.5.2). In some applications the function f(t) describes an incoming signal that starts arriving at time t = 0. Then u(t-a) f(t-a) denotes a signal of the same "shape" but with a time delay of *a*, so it does not start arriving until time t = a.

Proof of Theorem 1: From the definition of $\mathcal{L}{f(t)}$, we get

$$e^{-as}F(s) = e^{-as}\int_0^\infty e^{-s\tau}f(\tau)\,d\tau = \int_0^\infty e^{-s(\tau+a)}f(\tau)\,d\tau.$$

The substitution $t = \tau + a$ then yields

$$e^{-as}F(s) = \int_a^\infty e^{-st} f(t-a) \, dt.$$

From Eq. (4) we see that this is the same as

$$e^{-as}F(s) = \int_0^\infty e^{-st} u(t-a) f(t-a) dt = \mathcal{L}\{u(t-a)f(t-a)\},\$$

because u(t-a)f(t-a) = 0 for t < a. This completes the proof of Theorem 1.

Example 1 With $f(t) = \frac{1}{2}t^2$, Theorem 1 gives

$$\mathcal{L}^{-1}\left\{\frac{e^{-as}}{s^3}\right\} = u(t-a)\frac{1}{2}(t-a)^2 = \begin{cases} 0 & \text{if } t < a, \\ \frac{1}{2}(t-a)^2 & \text{if } t \ge a \end{cases}$$
(Fig. 10.5.3).

Example 2 Find $\mathcal{L}{g(t)}$ if

$$g(t) = \begin{cases} 0 & \text{if } t < 3, \\ t^2 & \text{if } t \ge 3 \end{cases}$$
 (Fig. 10.5.4).

Solution Before applying Theorem 1, we must first write g(t) in the form u(t-3)f(t-3). The function f(t) whose translation 3 units to the right agrees (for $t \ge 3$) with $g(t) = t^2$ is $f(t) = (t+3)^2$ because $f(t-3) = t^2$. But then

$$F(s) = \mathcal{L}\{t^2 + 6t + 9\} = \frac{2}{s^3} + \frac{6}{s^2} + \frac{9}{s},$$

so now Theorem 1 yields

$$\mathscr{L}\{g(t)\} = \mathscr{L}\{u(t-3)f(t-3)\} = e^{-3s}F(s) = e^{-3s}\left(\frac{2}{s^3} + \frac{6}{s^2} + \frac{9}{s}\right).$$



FIGURE 10.5.3. The graph of the inverse transform of Example 1.



FIGURE 10.5.4. The graph of the function g(t) of Example 2.



FIGURE 10.5.5. The function f(t) of Examples 3 and 4.

Example 3 Find $\mathcal{L}{f(t)}$ if $f(t) = \begin{cases} \cos 2t & \text{if } 0 \leq t < 2\pi, \\ 0 & \text{if } t \geq 2\pi \end{cases}$ (Fig. 10.5.5).

Solution We note first that

$$f(t) = [1 - u(t - 2\pi)]\cos 2t = \cos 2t - u(t - 2\pi)\cos 2(t - 2\pi)$$

because of the periodicity of the cosine function. Hence Theorem 1 gives

$$\mathcal{L}{f(t)} = \mathcal{L}{\cos 2t} - e^{-2\pi s} \mathcal{L}{\cos 2t} = \frac{s(1 - e^{-2\pi s})}{s^2 + 4}.$$

Example 4 Discontinuous forcing A mass that weighs 32 lb (mass m = 1 slug) is attached to the free end of a long light spring that is stretched 1 ft by a force of 4 lb (k = 4 lb/ft). The mass is initially at rest in its equilibrium position. Beginning at time t = 0 (seconds), an external force $F(t) = \cos 2t$ is applied to the mass, but at time $t = 2\pi$ this force is turned off (abruptly discontinued) and the mass is allowed to continue its motion unimpeded. Find the resulting position function x(t) of the mass.

Solution We need to solve the initial value problem

$$x'' + 4x = f(t); \quad x(0) = x'(0) = 0,$$

where f(t) is the function of Example 3. The transformed equation is

$$(s^{2} + 4)X(s) = F(s) = \frac{s(1 - e^{-2\pi s})}{s^{2} + 4}$$

so

Because

$$X(s) = \frac{s}{(s^2 + 4)^2} - e^{-2\pi s} \frac{s}{(s^2 + 4)^2}$$

 $\mathcal{L}^{-1}\left\{\frac{s}{(s^2+4)^2}\right\} = \frac{1}{4}t\sin 2t$

by Eq. (16) of Section 10.3, it follows from Theorem 1 that

$$\begin{aligned} x(t) &= \frac{1}{4}t\sin 2t - u(t-2\pi) \cdot \frac{1}{4}(t-2\pi)\sin 2(t-2\pi) \\ &= \frac{1}{4}\left[t - u(t-2\pi) \cdot (t-2\pi)\right]\sin 2t. \end{aligned}$$

If we separate the cases $t < 2\pi$ and $t \ge 2\pi$, we find that the position function may be written in the form

$$x(t) = \begin{cases} \frac{1}{4}t\sin 2t & \text{if } t < 2\pi, \\ \frac{1}{2}\pi\sin 2t & \text{if } t \ge 2\pi. \end{cases}$$



FIGURE 10.5.6. The graph of the function x(t) of Example 4.

As indicated by the graph of x(t) shown in Fig. 10.5.6, the mass oscillates with circular frequency $\omega = 2$ and with linearly increasing amplitude until the force is removed at time $t = 2\pi$. Thereafter, the mass continues to oscillate with the same frequency but with constant amplitude $\pi/2$. The force $F(t) = \cos 2t$ would produce pure resonance if continued indefinitely, but we see that its effect ceases immediately at the moment it is turned off.

If we were to attack Example 4 with the methods of Chapter 5, we would need to solve one problem for the interval $0 \le t < 2\pi$ and then solve a new problem with different initial conditions for the interval $t \ge 2\pi$. In such a situation the Laplace transform method enjoys the distinct advantage of not requiring the solution of different problems on different intervals.

Transforms of Periodic Functions

Periodic forcing functions in practical mechanical or electrical systems often are more complicated than pure sines or cosines. The nonconstant function f(t) defined for $t \ge 0$ is said to be **periodic** if there is a number p > 0 such that

$$f(t+p) = f(t) \tag{5}$$



FIGURE 10.5.7. The graph of a function with period p.

for all $t \ge 0$. The least positive value of p (if any) for which Eq. (11) holds is called the **period** of f. Such a function is shown in Fig. 10.5.7. Theorem 2 simplifies the computation of the Laplace transform of a periodic function.

THEOREM 2 Transforms of Periodic Functions

Let f(t) be periodic with period p and piecewise continuous for $t \ge 0$. Then the transform $F(s) = \mathcal{L}{f(t)}$ exists for s > 0 and is given by

$$F(s) = \frac{1}{1 - e^{-ps}} \int_0^p e^{-st} f(t) dt.$$
 (6)

Proof: The definition of the Laplace transform gives

$$F(s) = \int_0^\infty e^{-st} f(t) \, dt = \sum_{n=0}^\infty \int_{np}^{(n+1)p} e^{-st} f(t) \, dt.$$

The substitution $t = \tau + np$ in the *n*th integral following the summation sign yields

$$\int_{np}^{(n+1)p} e^{-st} f(t) dt = \int_{0}^{p} e^{-s(\tau+np)} f(\tau+np) d\tau = e^{-nps} \int_{0}^{p} e^{-s\tau} f(\tau) d\tau$$

because $f(\tau + np) = f(\tau)$ by periodicity. Thus

$$F(s) = \sum_{n=0}^{\infty} \left(e^{-nps} \int_0^p e^{-s\tau} f(\tau) \, d\tau \right)$$

= $\left(1 + e^{-ps} + e^{-2ps} + \cdots \right) \int_0^p e^{-s\tau} f(\tau) \, d\tau$

Consequently,

$$F(s) = \frac{1}{1 - e^{-ps}} \int_0^p e^{-s\tau} f(\tau) \, d\tau.$$
We use the geometric series

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots,$$

with $x = e^{-ps} < 1$ (for s > 0) to sum the series in the final step. Thus we have derived Eq. (6).

The principal advantage of Theorem 2 is that it enables us to find the Laplace transform of a periodic function without the necessity of an explicit evaluation of an improper integral.

Example 5



FIGURE 10.5.8. The graph of the square wave function of Example 5.

|g(t)|a

a



$$F(s) = \frac{1}{1 - e^{-2as}} \int_0^{2a} e^{-st} f(t) dt$$

= $\frac{1}{1 - e^{-2as}} \left(\int_0^a e^{-st} dt + \int_a^{2a} (-1)e^{-st} dt \right)$
= $\frac{1}{1 - e^{-2as}} \left(\left[-\frac{1}{s}e^{-st} \right]_0^a - \left[-\frac{1}{s}e^{-st} \right]_a^{2a} \right)$
= $\frac{(1 - e^{-as})^2}{s(1 - e^{-2as})} = \frac{1 - e^{-as}}{s(1 + e^{-as})}.$

$$F(s) = \frac{1 - e^{-as}}{s(1 + e^{-as})}$$
(7a)

$$=\frac{e^{as/2} - e^{-as/2}}{s(e^{as/2} + e^{-as/2})} = \frac{1}{s} \tanh \frac{as}{2}.$$
 (7b)



Example 6

Figure 10.5.9 shows the graph of a triangular wave function g(t) of period p = 2a. Because the derivative g'(t) is the square wave function of Example 6, it follows from the formula in (7b) and Theorem 2 of Section 10.2 that the transform of this triangular wave function is

$$G(s) = \frac{F(s)}{s} = \frac{1}{s^2} \tanh \frac{as}{2}.$$
 (8)

Consider a mass-spring-dashpot system with m = 1, c = 4, and Example 7 Square wave forcing k = 20 in appropriate units. Suppose that the system is initially at rest at equilibrium (x(0) =x'(0) = 0 and that the mass is acted on beginning at time t = 0 by the external force f(t)whose graph is shown in Fig. 10.5.10: the square wave with amplitude 20 and period 2π . Find the position function f(t).

Solution The initial value problem is

Therefore,

$$x'' + 4x' + 20x = f(t); \quad x(0) = x'(0) = 0.$$

The transformed equation is

$$s^{2}X(s) + 4sX(s) + 20X(s) = F(s).$$
(9)

10.5 PERIODIC AND PIECEWISE CONTINUOUS INPUT FUNCTIONS 599





From Example 5 with $a = \pi$ we see that the transform of f(t) is

$$F(s) = \frac{20}{s} \cdot \frac{1 - e^{-\pi s}}{1 + e^{-\pi s}}$$
$$= \frac{20}{s} \left(1 - e^{-\pi s} \right) \left(1 - e^{-\pi s} + e^{-2\pi s} - e^{-3\pi s} + \cdots \right)$$
$$= \frac{20}{s} \left(1 - 2e^{-\pi s} + 2e^{-2\pi s} - 2e^{-3\pi s} + \cdots \right),$$

so that

$$F(s) = \frac{20}{s} + \frac{40}{s} \sum_{n=1}^{\infty} (-1)^n e^{-n\pi s}.$$
 (10)

Substitution of Eq. (10) in Eq. (9) yields

$$X(s) = \frac{F(s)}{s^2 + 4s + 20}$$

= $\frac{20}{s[(s+2)^2 + 16]} + 2\sum_{n=1}^{\infty} (-1)^n \frac{20e^{-n\pi s}}{s[(s+2)^2 + 16]}.$ (11)

From the transform in Eq. (8) of Section 10.3, we get

$$\mathcal{L}^{-1}\left\{\frac{20}{(s+2)^2+16}\right\} = 5e^{-2t}\sin 4t,$$

so by Theorem 2 of Section 10.2 we have

$$g(t) = \mathcal{L}^{-1} \left\{ \frac{20}{s[(s+2)^2 + 16]} \right\} = \int_0^t 5e^{-2\tau} \sin 4\tau \, d\tau.$$

Using a tabulated formula for $\int e^{at} \sin bt \, dt$, we get

$$g(t) = 1 - e^{-2t} \left(\cos 4t + \frac{1}{2} \sin 4t \right) = 1 - h(t), \tag{12}$$

where

$$h(t) = e^{-2t} \left(\cos 4t + \frac{1}{2} \sin 4t \right).$$
(13)

Now we apply Theorem 1 to find the inverse transform of the right-hand term in Eq. (11). The result is

$$x(t) = g(t) + 2\sum_{n=1}^{\infty} (-1)^n u(t - n\pi)g(t - n\pi),$$
(14)

and we note that for any fixed value of t the sum in Eq. (14) is finite. Moreover,

$$g(t - n\pi) = 1 - e^{-2(t - n\pi)} \left[\cos 4(t - n\pi) + \frac{1}{2} \sin 4(t - n\pi) \right]$$
$$= 1 - e^{2n\pi} e^{-2t} \left(\cos 4t + \frac{1}{2} \sin 4t \right).$$

Therefore,

$$g(t - n\pi) = 1 - e^{2n\pi} h(t).$$
(15)

Hence if $0 < t < \pi$, then

$$x(t) = 1 - h(t).$$

If $\pi < t < 2\pi$, then

$$x(t) = [1 - h(t)] - 2\left[1 - e^{2\pi}h(t)\right] = -1 + h(t) - 2h(t)\left[1 - e^{2\pi}\right]$$

600 CHAPTER 10 LAPLACE TRANSFORM METHODS

If $2\pi < t < 3\pi$, then

$$\begin{aligned} x(t) &= [1 - h(t)] - 2 \left[1 - e^{2\pi} h(t) \right] + 2 \left[1 - e^{4\pi} h(t) \right] \\ &= 1 + h(t) - 2h(t) \left[1 - e^{2\pi} + e^{4\pi} \right]. \end{aligned}$$

The general expression for $n\pi < t < (n+1)\pi$ is

$$x(t) = h(t) + (-1)^{n} - 2h(t) \left[1 - e^{2\pi} + \dots + (-1)^{n} e^{2n\pi} \right]$$

= $h(t) + (-1)^{n} - 2h(t) \frac{1 + (-1)^{n} e^{2(n+1)\pi}}{1 + e^{2\pi}},$ (16)

which we obtained with the aid of the familiar formula for the sum of a finite geometric progression. A rearrangement of Eq. (16) finally gives, with the aid of Eq. (13),

$$x(t) = \frac{e^{2\pi} - 1}{e^{2\pi} + 1} e^{-2t} \left(\cos 4t + \frac{1}{2} \sin 4t \right) + (-1)^n - \frac{2 \cdot (-1)^n e^{2\pi}}{e^{2\pi} + 1} e^{-2(t - n\pi)} \left(\cos 4t + \frac{1}{2} \sin 4t \right)$$
(17)

for $n\pi < t < (n + 1)\pi$. The first term in Eq. (17) is the transient solution

$$x_{\rm tr}(t) \approx (0.9963)e^{-2t} \left(\cos 4t + \frac{1}{2}\sin 4t\right) \approx (1.1139)e^{-2t} \cos(4t - 0.4636).$$
 (18)

The last two terms in Eq. (17) give the steady periodic solution x_{sp} . To investigate it, we write $\tau = t - n\pi$ for t in the interval $n\pi < t < (n + 1)\pi$. Then

$$x_{\rm sp}(t) = (-1)^n \left[1 - \frac{2e^{2\pi}}{e^{2\pi} + 1} e^{-2\tau} \left(\cos 4\tau + \frac{1}{2} \sin 4\tau \right) \right]$$

$$\approx (-1)^n \left[1 - (2.2319)e^{-2\tau} \cos(4\tau - 0.4636) \right].$$
 (19)

Figure 10.5.11 shows the graph of $x_{sp}(t)$. Its most interesting feature is the appearance of periodically damped oscillations with a frequency *four times* that of the imposed force f(t).



FIGURE 10.5.11. The graph of the steady periodic solution for Example 7; note the "periodically damped" oscillations with frequency four times that of the imposed force.

10.5 **Problems**

Find the inverse Laplace transform f(t) of each function given in Problems 1 through 10. Then sketch the graph of f.

1. $F(s) = \frac{e^{-3s}}{s^2}$ 2. $F(s) = \frac{e^{-s} - e^{-3s}}{s^2}$ 3. $F(s) = \frac{e^{-s}}{s+2}$ 4. $F(s) = \frac{e^{-s} - e^{2-2s}}{s-1}$ 5. $F(s) = \frac{e^{-\pi s}}{s^2 + 1}$ 6. $F(s) = \frac{se^{-s}}{s^2 + \pi^2}$ 7. $F(s) = \frac{1 - e^{-2\pi s}}{s^2 + 1}$ 8. $F(s) = \frac{s(1 - e^{-2s})}{s^2 + \pi^2}$ 9. $F(s) = \frac{s(1 + e^{-3s})}{s^2 + \pi^2}$ 10. $F(s) = \frac{2s(e^{-\pi s} - e^{-2\pi s})}{s^2 + 4}$

Find the Laplace transforms of the functions given in Problems 11 through 22.

- 11. $f(t) = 2 \text{ if } 0 \le t < 3; f(t) = 0 \text{ if } t \ge 3$ 12. $f(t) = 1 \text{ if } 1 \le t \le 4; f(t) = 0 \text{ if } t < 1 \text{ or if } t > 4$ 13. $f(t) = \sin t \text{ if } 0 \le t \le 2\pi; f(t) = 0 \text{ if } t > 2\pi$ 14. $f(t) = \cos \pi t \text{ if } 0 \le t \le 2; f(t) = 0 \text{ if } t > 2\pi$ 15. $f(t) = \sin t \text{ if } 0 \le t \le 3\pi; f(t) = 0 \text{ if } t > 3\pi$ 16. $f(t) = \sin 2t \text{ if } \pi \le t \le 2\pi; f(t) = 0 \text{ if } t < \pi \text{ or if } t > 2\pi$ 17. $f(t) = \sin \pi t \text{ if } 2 \le t \le 3; f(t) = 0 \text{ if } t < 2 \text{ or if } t > 3$ 18. $f(t) = \cos \frac{1}{2}\pi t \text{ if } 3 \le t \le 5; f(t) = 0 \text{ if } t < 3 \text{ or if } t > 5$ 19. $f(t) = 0 \text{ if } t < 1; f(t) = t \text{ if } t \ge 1$ 20. $f(t) = t \text{ if } t \le 1; f(t) = 1 \text{ if } t > 1$ 21. $f(t) = t \text{ if } t \le 1; f(t) = 2 - t \text{ if } 1 \le t \le 2; f(t) = 0 \text{ if } t > 2$
- **23.** Apply Theorem 2 with p = 1 to verify that $\mathcal{L}\{1\} = 1/s$.
- **24.** Apply Theorem 2 to verify that $\mathcal{L}\{\cos kt\} = s/(s^2 + k^2)$.
- **25.** Apply Theorem 2 to show that the Laplace transform of the square wave function of Fig. 10.5.12 is



FIGURE 10.5.12. The graph of the square wave function of Problem 25.

26. Apply Theorem 2 to show that the Laplace transform of the sawtooth function f(t) of Fig. 10.5.13 is

$$F(s) = \frac{1}{as^2} - \frac{e^{-as}}{s(1 - e^{-as})}$$



FIGURE 10.5.13. The graph of the sawtooth function of Problem 26.

27. Let g(t) be the staircase function of Fig. 10.5.14. Show that g(t) = (t/a) - f(t), where *f* is the sawtooth function of Fig. 10.5.14, and hence deduce that



FIGURE 10.5.14. The graph of the staircase function of Problem 27.

- **28.** Suppose that f(t) is a periodic function of period 2*a* with f(t) = t if $0 \le t < a$ and f(t) = 0 if $a \le t < 2a$. Find $\mathcal{L}{f(t)}$.
- **29.** Suppose that f(t) is the half-wave rectification of $\sin kt$, shown in Fig. 10.5.15. Show that



FIGURE 10.5.15. The half-wave rectification of $\sin kt$.

30. Let $g(t) = u(t - \pi/k) f(t - \pi/k)$, where f(t) is the function of Problem 29 and k > 0. Note that h(t) = f(t) + g(t) is the full-wave rectification of $\sin kt$ shown in Fig. 10.5.16. Hence deduce from Problem 29 that



FIGURE 10.5.16. The full-wave rectification of $\sin kt$.

Discontinuous Forcing

In Problems 31 through 35, the values of mass m, spring constant k, dashpot resistance c, and force f(t) are given for a mass-spring-dashpot system with external forcing function. Solve the initial value problem

$$mx'' + cx' + kx = f(t); \quad x(0) = x'(0) = 0$$

and construct the graph of the position function x(t).

- **31.** m = 1, k = 4, c = 0; f(t) = 1 if $0 \le t < \pi, f(t) = 0$ if $t \ge \pi$
- **32.** m = 1, k = 4, c = 5; f(t) = 1 if $0 \le t < 2, f(t) = 0$ if $t \ge 2$
- **33.** $m = 1, k = 9, c = 0; f(t) = \sin t$ if $0 \le t \le 2\pi, f(t) = 0$ if $t > 2\pi$
- **34.** m = 1, k = 1, c = 0; f(t) = t if $0 \le t < 1, f(t) = 0$ if $t \ge 1$
- **35.** m = 1, k = 4, c = 4; f(t) = t if $0 \le t \le 2, f(t) = 0$ if t > 2

Transient and Steady Periodic Motions

In Problems 36 and 37, a mass-spring-dashpot system with external force f(t) is described. Under the assumption that x(0) = x'(0) = 0, use the method of Example 7 to find the transient and steady periodic motions of the mass. Then construct the graph of the position function x(t). If you would like to check your graph using a numerical DE solver, it may be useful to note that the function

$$f(t) = A[2u((t - \pi)(t - 2\pi)(t - 3\pi) \cdot (t - 4\pi)(t - 5\pi)(t - 6\pi)) - 1]$$

has the value +A if $0 < t < \pi$, the value -A if $\pi < t < 2\pi$, and so forth, and hence agrees on the interval $[0, 6\pi]$ with the square wave function that has amplitude A and period 2π . (See also the definition of a square wave function in terms of sawtooth and triangular wave functions in the application material for this section.)

- **36.** m = 1, k = 4, c = 0; f(t) is a square wave function with amplitude 4 and period 2π .
- **37.** m = 1, k = 10, c = 2; f(t) is a square wave function with amplitude 10 and period 2π .
- **38.** Suppose the function x(t) satisfies the initial value problem

$$mx'' + cx' + kx = F(t), \quad x(a) = b_0, \ x'(a) = b_1$$

for $t \ge a$ and x(t) = 0 for t < a. Then show that $X(s) = \mathcal{L}\{x(t)\}$ satisfies the equation

$$m\left(s^2(e^{as}X) - sb_0 - b_1\right) + c\left(s(e^{as}X) - b_0\right) + k\left(e^{as}X\right)$$
$$= \mathcal{L}\{F(t+a)\}.$$

39. This is an alternate approach to Example 7 that was suggested by Keng C. Wu of Lockheed Martin (*Maritime Systems & Sensors*). Let x(t) denote the *steady periodic solution* of the given differential equation x" + 4x' + 20x = F(t). Suppose we write v(t) for the restriction of x(t) to the first half [0, π] of the fundamental interval [0, 2π], and w(t) for its restriction to the second half-interval [π, 2π]. We may regard v(t) as the solution of the initial value problem

$$v'' + 4v' + 20v = +20, \quad v(0) = b_0, \ v'(0) = b_1$$

and w(t) as the solution of the initial value problem

$$w'' + 4w' + 20w = -20, \quad w(\pi) = c_0, \ w'(\pi) = c_1,$$

where the initial values b_0 , b_1 and c_0 , c_1 are to be determined so that x(t) and x'(t) are continuous.

(a) Transform the first of these initial value problems to show that $V(s) = A(s)b_0 + B(s)b_1 + C(s)$, where

$$A(s) = \frac{s+4}{s^2+4s+20},$$

$$B(s) = \frac{1}{s^2+4s+20},$$

$$C(s) = \frac{20}{s(s^2+4s+20)}$$

- (b) Apply the result of Problem 38 to the second initial value problem above to show that $W(s) = e^{-\pi s} (A(s)c_0 + B(s)c_1 C(s))$, where the coefficient functions A(s), B(s), C(s) are as defined in part (a).
- (c) After finding the inverse transforms

$$a(t) = e^{-2t} \left(\cos 4t + \frac{1}{2} \sin 4t \right),$$

$$b(t) = \frac{1}{4} e^{-2t} \sin 4t,$$

$$c(t) = 1 - e^{-2t} \left(\cos 4t + \frac{1}{2} \sin 4t \right)$$

solve the four continuity equations $v(\pi) = c_0$, $v'(\pi) = c_1$, $w(2\pi) = b_0$, $w'(2\pi) = b_1$ to find the values of the previously undetermined initial values. Conclude that

$$v(t) \approx 1 - 0.9981e^{-2t} (2\cos 4t + \sin 4t),$$

$$w(t) \approx \left[-1 + 0.9981e^{-2(t-\pi)} (2\cos 4t + \sin 4t)\right] u(t - \pi).$$

Finally, use these expressions to verify that the graph of the steady periodic solution x(t) looks as indicated in Fig. 10.5.11.

10.5 Application Engineering Functions



download this application's computing resources including *Maple/Mathematica/*MATLAB. Periodic piecewise linear functions occur so frequently as input functions in engineering applications that they are sometimes called **engineering functions**. Computations with such functions are readily handled by computer algebra systems. In *Mathematica*, for instance, the **SawToothWave**, **TriangleWave**, and **Square-Wave** functions can be used to create the corresponding inputs with specified range, period, etc. Alternatively, we can define our own engineering functions using elementary functions available in any computer algebra system:

```
sawtooth[t_] := t - 2 Floor[t/2] - 1
triangularwave[t_] := 2 Abs[sawtooth[t - 1/2]] - 1
squarewave[t_] := Sign[ triangularwave[t]]
```

Plot each of the functions to verify that it has period 2 and that its name is aptly chosen. For instance, the result of

```
Plot[squarewave[t], {t, 0, 6}]
```

should look like Fig. 10.5.8. If f(t) is one of these engineering functions and p > 0, then the function f(2t/p) will have period p. To illustrate this, try

```
Plot[triangularwave[ 2 t/p ], {t, 0, 3 p}]
```

with various values of p.

Now let's consider the mass-spring-dashpot equation

```
diffEq = m x''[t] + c x'[t] + k x[t] == input
```

with selected parameter values and an input forcing function with period p and amplitude F_0 .

```
m = 4; c = 8; k = 5; p = 1; F0 = 4;
input = F0 squarewave[2 t/p];
```

You can plot this **input** function to verify that it has period 1:

Plot[input, {t, 0, 2}]

Finally, let's suppose that the mass is initially at rest in its equilibrium position and solve numerically the resulting initial value problem.

In the resulting Fig. 10.5.17 we see that after an initial transient dies out, the response function x(t) settles down (as expected?) to a periodic oscillation with the same period as the input.

Investigate this initial value problem with several mass–spring–dashpot parameters—for instance, selected digits of your student ID number—and with input engineering functions having various amplitudes and periods.



FIGURE 10.5.17. Response x(t) to period 1 square wave input.



11.1 Introduction and Review of Power Series

In Section 5.3 we saw that solving a homogeneous linear differential equation with constant coefficients can be reduced to the algebraic problem of finding the roots of its characteristic equation. There is no similar procedure for solving linear differential equations with *variable* coefficients, at least not routinely and in finitely many steps. With the exception of special types, such as the occasional equation that can be solved by inspection, linear equations with variable coefficients generally require the power series techniques of this chapter.

These techniques suffice for many of the nonelementary differential equations that appear most frequently in applications. Perhaps the most important (because of its applications in such areas as acoustics, heat flow, and electromagnetic radiation) is **Bessel's equation** of order n:

$$x^{2}y'' + xy' + (x^{2} - n^{2})y = 0.$$

Legendre's equation of order *n* is important in many applications. It has the form

>
$$(1-x^2)y'' - 2xy' + n(n+1)y = 0.$$

In this section we introduce the **power series method** in its simplest form and, along the way, state (without proof) several theorems that constitute a review of the basic facts about power series. Recall first that a **power series** in (powers of) x - a is an infinite series of the form

$$\sum_{n=0}^{\infty} c_n (x-a)^n = c_0 + c_1 (x-a) + c_2 (x-a)^2 + \dots + c_n (x-a)^n + \dots$$
(1)

If a = 0, this is a power series in x:

>
$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$
 (2)

We will confine our review mainly to power series in x, but every general property of power series in x can be converted to a general property of power series in x - a by replacement of x with x - a.

The power series in (2) converges on the interval I provided that the limit

$$\sum_{n=0}^{\infty} c_n x^n = \lim_{N \to \infty} \sum_{n=0}^{N} c_n x^n \tag{3}$$

exists for all x in I. In this case the sum

$$f(x) = \sum_{n=0}^{\infty} c_n x^n \tag{4}$$

is defined on *I*, and we call the series $\sum c_n x^n$ a **power series representation** of the function *f* on *I*. The following power series representations of elementary functions should be familiar to you from introductory calculus:

$$e^{x} = \sum_{n=0}^{\infty} \frac{x^{n}}{n!} = 1 + x + \frac{x^{2}}{2!} + \frac{x^{3}}{3!} + \dots;$$
 (5)

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots;$$
(6)

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots;$$
(7)

$$\cosh x = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots;$$
(8)

$$\sinh x = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!} = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots;$$
(9)

$$\ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots;$$
(10)

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots;$$
(11)

and

$$(1+x)^{\alpha} = 1 + \alpha x + \frac{\alpha(\alpha-1)x^2}{2!} + \frac{\alpha(\alpha-1)(\alpha-2)x^3}{3!} + \cdots .$$
(12)

In compact summation notation, we observe the usual conventions that 0! = 1 and that $x^0 = 1$ for all x, including x = 0. The series in (5) through (9) converge to the indicated functions for all x. In contrast, the series in (10) and (11) converge if |x| < 1 but diverge if |x| > 1. (What if |x| = 1?) The series in (11) is the **geometric series**. The series in (12), with α an arbitrary real number, is the **binomial series**. If α is a nonnegative integer n, then the series in (12) terminates and the binomial series reduces to a polynomial of degree n which converges for all x. Otherwise, the series is actually infinite and it converges if |x| < 1 and diverges if |x| > 1; its behavior for |x| = 1 depends on the value of α .

Remark Power series such as those listed in (5) through (12) are often derived as Taylor series. The **Taylor series** with **center** x = a of the function f is the power series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \cdots$$
(13)

in powers of x - a, under the hypothesis that f is infinitely differentiable at x = a (so that the coefficients in Eq. (13) are all defined). If a = 0, then the series in (13) is the Maclaurin series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = f(0) + f'(0)x + \frac{f''(0)}{2!} x^2 + \frac{f^{(3)}(0)}{3!} x^3 + \cdots .$$
(13')

For example, suppose that $f(x) = e^x$. Then $f^{(n)}(x) = e^x$, and hence $f^{(n)}(0) = 1$ for all $n \ge 0$. In this case Eq. (13') reduces to the **exponential series** in (5).

Power Series Operations

If the Taylor series of the function f converges to f(x) for all x in some open interval containing a, then we say that the function f is **analytic** at x = a. For example,

- every polynomial function is analytic everywhere;
- every rational function is analytic wherever its denominator is nonzero;
- more generally, if the two functions f and g are both analytic at x = a, then so are their sum f + g and their product $f \cdot g$, as is their quotient f/g if $g(a) \neq 0$.

For instance, the function $h(x) = \tan x = (\sin x)/(\cos x)$ is analytic at x = 0 because $\cos 0 = 1 \neq 0$ and the sine and cosine functions are analytic (by virtue of their convergent power series representations in Eqs. (6) and (7)). It is rather awkward to compute the Taylor series of the tangent function using Eq. (13) because of the way in which its successive derivatives grow in complexity (try it!). Fortunately, power series may be manipulated algebraically in much the same way as polynomials. For example, if

$$f(x) = \sum_{n=0}^{\infty} a_n x^n \quad \text{and} \quad g(x) = \sum_{n=0}^{\infty} b_n x^n, \tag{14}$$

then

$$f(x) + g(x) = \sum_{n=0}^{\infty} (a_n + b_n) x^n$$
(15)

and

$$f(x)g(x) = \sum_{n=0}^{\infty} c_n x^n$$

= $a_0 b_0 + (a_0 b_1 + a_1 b_0) x + (a_0 b_2 + a_1 b_1 + a_2 b_0) x^2 + \cdots$, (16)

where $c_n = a_0b_n + a_1b_{n-1} + \cdots + a_nb_0$. The series in (15) is the result of **termwise** addition, and the series in (16) is the result of **formal multiplication**—multiplying each term of the first series by each term of the second and then collecting coefficients of like powers of x. (Thus the processes strongly resemble addition and multiplication of ordinary polynomials.) The series in (15) and (16) converge to

f(x) + g(x) and f(x)g(x), respectively, on any open interval on which both the series in (14) converge. For example,

$$\sin x \cos x = \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \cdots\right) \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \cdots\right)$$
$$= x + \left(-\frac{1}{6} - \frac{1}{2}\right)x^3 + \left(\frac{1}{24} + \frac{1}{12} + \frac{1}{120}\right)x^5 + \cdots$$
$$= x - \frac{4}{6}x^3 + \frac{16}{120}x^5 - \cdots$$
$$= \frac{1}{2}\left[(2x) - \frac{(2x)^3}{3!} + \frac{(2x)^5}{5!} - \cdots\right] = \frac{1}{2}\sin 2x$$

for all *x*.

Similarly, the quotient of two power series can be computed by long division, as illustrated by the computation shown in Fig. 11.1.1. This division of the Taylor series for $\cos x$ into that for $\sin x$ yields the first few terms of the series

$$\tan x = x + \frac{1}{3}x^3 + \frac{2}{15}x^5 + \frac{17}{315}x^7 + \cdots .$$
 (17)

Division of power series is more treacherous than multiplication; the series thus obtained for f/g may fail to converge at some points where the series for f and g both converge. For example, the sine and cosine series converge for all x, but the tangent series in (17) converges only if $|x| < \pi/2$.

x	+	$\frac{x^3}{3}$	+	$\frac{2x^5}{15}$	+	$\frac{17x^7}{315}$	+	
$1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + \dots \right) x$	_	$\frac{x^3}{6}$	+	$\frac{x^5}{120}$	_	$\frac{x^7}{5040}$	+	
<i>x</i>	_	$\frac{x^3}{2}$	+	$\frac{x^5}{24}$	_	$\frac{x^7}{720}$	+	
		$\frac{x^3}{3}$	_	$\frac{x^5}{30}$	+	$\frac{x^7}{840}$	+	
	_	$\frac{x^3}{3}$	_	$\frac{x^5}{6}$	+	$\frac{x^7}{72}$	_	
				$\frac{2x^5}{15}$	_	$\frac{4x^7}{315}$	+	
			-	$\frac{2x^5}{15}$	_	$\frac{x^7}{15}$	+	••••
						$\frac{17x^7}{315}$	+	
						:		

FIGURE 11.1.1. Obtaining the series for tan x by division of series.

The Power Series Method

The **power series method** for solving a differential equation consists of substituting the power series

in the differential equation and then attempting to determine what the coefficients c_0, c_1, c_2, \ldots must be in order that the power series will satisfy the differential equation. This is reminiscent of the method of undetermined coefficients, but now we have infinitely many coefficients somehow to determine. This method is not always successful, but when it is we obtain an infinite series representation of a solution, in contrast to the "closed form" solutions that our previous methods have yielded.

Before we can substitute the power series in (18) in a differential equation, we must first know what to substitute for the derivatives y', y'', \ldots . The following theorem (stated without proof) tells us that the derivative y' of $y = \sum c_n x^n$ is obtained by the simple procedure of writing the sum of the derivatives of the individual terms in the series for y.

THEOREM 1 Termwise Differentiation of Power Series

If the power series representation

•
$$f(x) = \sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots$$
 (19)

of the function f converges on the open interval I, then f is differentiable on I, and

>
$$f'(x) = \sum_{n=1}^{\infty} nc_n x^{n-1} = c_1 + 2c_2 x + 3c_3 x^2 + \cdots$$
 (20)

at each point of I.

For example, differentiation of the geometric series

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$$
(11)

gives

$$\frac{1}{(1-x)^2} = \sum_{n=1}^{\infty} nx^{n-1} = 1 + 2x + 3x^2 + 4x^3 + \cdots$$

The process of determining the coefficients in the series $y = \sum c_n x^n$ so that it will satisfy a given differential equation depends also on Theorem 2. This theorem—also stated without proof—tells us that if two power series represent the same function, then they are the same series. In particular, the Taylor series in (13) is the only power series (in powers of x - a) that represents the function f.

THEOREM 2 Identity Principle

If

$$\sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} b_n x^n$$

for every point x in some open interval I, then $a_n = b_n$ for all $n \ge 0$.

In particular, if $\sum a_n x^n = 0$ for all x in some open interval, it follows from Theorem 2 that $a_n = 0$ for all $n \ge 0$.

Example 1 So Solution We

Solve the equation y' + 2y = 0. We substitute the series

P



$$y = \sum_{n=0}^{\infty} c_n x^n$$
 and $y' = \sum_{n=1}^{\infty} n c_n x^{n-1}$

and obtain

$$\sum_{n=1}^{\infty} nc_n x^{n-1} + 2\sum_{n=0}^{\infty} c_n x^n = 0.$$
 (21)

To compare coefficients here, we need the general term in each sum to be the term containing x^n . To accomplish this, we shift the index of summation in the first sum. To see how to do this, note that

$$\sum_{n=1}^{\infty} nc_n x^{n-1} = c_1 + 2c_2 x + 3c_3 x^2 + \dots = \sum_{n=0}^{\infty} (n+1)c_{n+1} x^n.$$

Thus we can replace n with n + 1 if, at the same time, we start counting one step lower; that is, at n = 0 rather than at n = 1. This is a shift of +1 in the index of summation. The result of making this shift in Eq. (21) is the identity

$$\sum_{n=0}^{\infty} (n+1)c_{n+1}x^n + 2\sum_{n=0}^{\infty} c_n x^n = 0;$$

that is,

>

$$\sum_{n=0}^{\infty} [(n+1)c_{n+1} + 2c_n]x^n = 0$$

If this equation holds on some interval, then it follows from the identity principle that $(n + 1)c_{n+1} + 2c_n = 0$ for all $n \ge 0$; consequently,

$$c_{n+1} = -\frac{2c_n}{n+1}$$
(22)

for all $n \ge 0$. Equation (22) is a **recurrence relation** from which we can successively compute c_1, c_2, c_3, \ldots in terms of c_0 ; the latter will turn out to be the arbitrary constant that we expect to find in a general solution of a first-order differential equation.

With n = 0, Eq. (22) gives

$$c_1 = -\frac{2c_0}{1}.$$

With n = 1, Eq. (22) gives

$$c_2 = -\frac{2c_1}{2} = +\frac{2^2c_0}{1\cdot 2} = \frac{2^2c_0}{2!}.$$

With n = 2, Eq. (22) gives

$$c_3 = -\frac{2c_2}{3} = -\frac{2^3c_0}{1\cdot 2\cdot 3} = -\frac{2^3c_0}{3!}.$$

By now it should be clear that after *n* such steps, we will have

$$c_n = (-1)^n \frac{2^n c_0}{n!}, \qquad n \ge 1.$$

(This is easy to prove by induction on n.) Consequently, our solution takes the form

$$y(x) = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} (-1)^n \frac{2^n c_0}{n!} x^n = c_0 \sum_{n=0}^{\infty} \frac{(-2x)^n}{n!} = c_0 e^{-2x}.$$

In the final step we have used the familiar exponential series in Eq. (5) to identify our power series solution as the same solution $y(x) = c_0 e^{-2x}$ we could have obtained by the method of separation of variables.

Shift of Index of Summation

In the solution of Example 1 we wrote

$$\sum_{n=1}^{\infty} nc_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)c_{n+1} x^n$$
(23)

by shifting the index of summation by +1 in the series on the left. That is, we simultaneously *increased* the index of summation by 1 (replacing *n* with n + 1, $n \rightarrow n + 1$) and *decreased* the starting point by 1, from n = 1 to n = 0, thereby obtaining the series on the right. This procedure is valid because each infinite series in (23) is simply a compact notation for the single series

$$c_1 + 2c_2x + 3c_3x^2 + 4c_4x^3 + \cdots . (24)$$

More generally, we can shift the index of summation by k in an infinite series by simultaneously *increasing* the summation index by $k (n \rightarrow n + k)$ and *decreasing* the starting point by k. For instance, a shift by $+2 (n \rightarrow n + 2)$ yields

$$\sum_{n=3}^{\infty} a_n x^{n-1} = \sum_{n=1}^{\infty} a_{n+2} x^{n+1}$$

If k is negative, we interpret a "decrease by k" as an increase by -k = |k|. Thus a shift by $-2 (n \rightarrow n - 2)$ in the index of summation yields

$$\sum_{n=1}^{\infty} n c_n x^{n-1} = \sum_{n=3}^{\infty} (n-2) c_{n-2} x^{n-3};$$

we have *decreased* the index of summation by 2 but *increased* the starting point by 2, from n = 1 to n = 3. You should check that the summation on the right is merely another representation of the series in (24).

We know that the power series obtained in Example 1 converges for all x because it is an exponential series. More commonly, a power series solution is not recognizable in terms of the familiar elementary functions. When we get an unfamiliar power series solution, we need a way of finding where it converges. After all, $y = \sum c_n x^n$ is merely an *assumed* form of the solution. The procedure illustrated in Example 1 for finding the coefficients $\{c_n\}$ is merely a formal process and may or may not be valid. Its validity—in applying Theorem 1 to compute y' and applying Theorem 2 to obtain a recurrence relation for the coefficients—depends on the convergence of the initially *unknown* series $y = \sum c_n x^n$. Hence this formal process is justified only if in the end we can show that the power series we obtain converges on some open interval. If so, it then represents a solution of the differential equation on that interval. The following theorem (which we state without proof) may be used for this purpose.

THEOREM 3 Radius of Convergence

Given the power series $\sum c_n x^n$, suppose that the limit

$$\rho = \lim_{n \to \infty} \left| \frac{c_n}{c_{n+1}} \right| \tag{25}$$

exists (ρ is finite) or is infinite (in this case we will write $\rho = \infty$). Then

- (a) If $\rho = 0$, then the series diverges for all $x \neq 0$.
- (**b**) If $0 < \rho < \infty$, then $\sum c_n x^n$ converges if $|x| < \rho$ and diverges if $|x| > \rho$.
- (c) If $\rho = \infty$, then the series converges for all x.

The number ρ in (25) is called the **radius of convergence** of the power series $\sum c_n x^n$. For instance, for the power series obtained in Example 1, we have

$$\rho = \lim_{n \to \infty} \left| \frac{(-1)^n 2^n c_0 / n!}{(-1)^{n+1} 2^{n+1} c_0 / (n+1)!} \right| = \lim_{n \to \infty} \frac{n+1}{2} = \infty,$$

and consequently the series we obtained in Example 1 converges for all x. Even if the limit in (25) fails to exist, there always will be a number ρ such that exactly one of the three alternatives in Theorem 3 holds. This number may be difficult to find, but for the power series we will consider in this chapter, Eq. (25) will be quite sufficient for computing the radius of convergence.

Example 2 Solve the equation
$$(x - 3)y' + 2y = 0$$
.
Solution As before, we substitute

$$y = \sum_{n=0}^{\infty} c_n x^n$$
 and $y' = \sum_{n=1}^{\infty} n c_n x^{n-1}$

to obtain

$$(x-3)\sum_{n=1}^{\infty} nc_n x^{n-1} + 2\sum_{n=0}^{\infty} c_n x^n = 0$$

so that

$$\sum_{n=1}^{\infty} nc_n x^n - 3 \sum_{n=1}^{\infty} nc_n x^{n-1} + 2 \sum_{n=0}^{\infty} c_n x^n = 0$$

In the first sum we can replace n = 1 with n = 0 with no effect on the sum. In the second sum we shift the index of summation by +1. This yields

$$\sum_{n=0}^{\infty} nc_n x^n - 3\sum_{n=0}^{\infty} (n+1)c_{n+1} x^n + 2\sum_{n=0}^{\infty} c_n x^n = 0;$$

that is,

$$\sum_{n=0}^{\infty} \left[nc_n - 3(n+1)c_{n+1} + 2c_n \right] x^n = 0.$$

The identity principle then gives

$$nc_n - 3(n+1)c_{n+1} + 2c_n = 0,$$

from which we obtain the recurrence relation

$$c_{n+1} = \frac{n+2}{3(n+1)}c_n \quad \text{for} \quad n \ge 0$$

We apply this formula with n = 0, n = 1, and n = 2, in turn, and find that

$$c_1 = \frac{2}{3}c_0$$
, $c_2 = \frac{3}{3 \cdot 2}c_1 = \frac{3}{3^2}c_0$, and $c_3 = \frac{4}{3 \cdot 3}c_2 = \frac{4}{3^3}c_0$.

This is almost enough to make the pattern evident; it is not difficult to show by induction on n that

$$c_n = \frac{n+1}{3^n} c_0 \quad \text{if} \quad n \ge 1.$$

Hence our proposed power series solution is

$$y(x) = c_0 \sum_{n=0}^{\infty} \frac{n+1}{3^n} x^n.$$
 (26)

Its radius of convergence is

$$\rho = \lim_{n \to \infty} \left| \frac{c_n}{c_{n+1}} \right| = \lim_{n \to \infty} \frac{3n+3}{n+2} = 3.$$

Thus the series in (26) converges if -3 < x < 3 and diverges if |x| > 3. In this particular example we can explain why. An elementary solution (obtained by separation of variables) of our differential equation is $y = 1/(3 - x)^2$. If we differentiate termwise the geometric series

$$\frac{1}{3-x} = \frac{\frac{1}{3}}{1-\frac{x}{3}} = \frac{1}{3} \sum_{n=0}^{\infty} \frac{x^n}{3^n},$$

we get a constant multiple of the series in (26). Thus this series (with the arbitrary constant c_0 appropriately chosen) represents the solution

$$y(x) = \frac{1}{(3-x)^2}$$

on the interval -3 < x < 3, and the singularity at x = 3 is the reason why the radius of convergence of the power series solution turned out to be $\rho = 3$.

Example 3 Solv Solution We

Solve the equation $x^2 y' = y - x - 1$. We make the usual substitutions $y = \sum c_n x^n$ and $y' = \sum n c_n x^{n-1}$, which yield

$$x^{2} \sum_{n=1}^{\infty} nc_{n} x^{n-1} = -1 - x + \sum_{n=0}^{\infty} c_{n} x^{n}$$

so that

$$\sum_{n=1}^{\infty} nc_n x^{n+1} = -1 - x + \sum_{n=0}^{\infty} c_n x^n.$$

Because of the presence of the two terms -1 and -x on the right-hand side, we need to split off the first two terms, $c_0 + c_1 x$, of the series on the right for comparison. If we also shift the index of summation on the left by -1 (replace n = 1 with n = 2 and n with n - 1), we get

$$\sum_{n=2}^{\infty} (n-1)c_{n-1}x^n = -1 - x + c_0 + c_1x + \sum_{n=2}^{\infty} c_n x^n.$$

Because the left-hand side contains neither a constant term nor a term containing *x* to the first power, the identity principle now yields $c_0 = 1$, $c_1 = 1$, and $c_n = (n-1)c_{n-1}$ for $n \ge 2$. It follows that

$$c_2 = 1 \cdot c_1 = 1!, \quad c_3 = 2 \cdot c_2 = 2!, \quad c_4 = 3 \cdot c_3 = 3!,$$

and, in general, that

$$c_n = (n-1)!$$
 for $n \ge 2$.

Thus we obtain the power series

$$y(x) = 1 + x + \sum_{n=2}^{\infty} (n-1)! x^n$$

But the radius of convergence of this series is

$$\rho = \lim_{n \to \infty} \frac{(n-1)!}{n!} = \lim_{n \to \infty} \frac{1}{n} = 0,$$

so the series converges only for x = 0. What does this mean? Simply that the given differential equation does not have a (convergent) power series solution of the assumed form $y = \sum c_n x^n$. This example serves as a warning that the simple act of writing $y = \sum c_n x^n$ involves an assumption that may be false.

Example 4Solve the equation
$$y'' + y = 0.$$
SolutionIf we assume a solution of the form

$$y = \sum_{n=0}^{\infty} c_n x^n$$

we find that

>
$$y' = \sum_{n=1}^{\infty} nc_n x^{n-1}$$
 and $y'' = \sum_{n=2}^{\infty} n(n-1)c_n x^{n-2}$.

Substitution for y and y'' in the differential equation then yields

$$\sum_{n=2}^{\infty} n(n-1)c_n x^{n-2} + \sum_{n=0}^{\infty} c_n x^n = 0.$$

We shift the index of summation in the first sum by +2 (replace n = 2 with n = 0 and n with n + 2). This gives

$$\sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n + \sum_{n=0}^{\infty} c_n x^n = 0.$$

The identity $(n + 2)(n + 1)c_{n+2} + c_n = 0$ now follows from the identity principle, and thus we obtain the recurrence relation

$$c_{n+2} = -\frac{c_n}{(n+1)(n+2)} \tag{27}$$

for $n \ge 0$. It is evident that this formula will determine the coefficients c_n with even subscript in terms of c_0 and those of odd subscript in terms of c_1 ; c_0 and c_1 are not predetermined and thus will be the two arbitrary constants we expect to find in a general solution of a second-order equation.

When we apply the recurrence relation in (27) with n = 0, 2, and 4 in turn, we get

$$c_2 = -\frac{c_0}{2!}, \quad c_4 = \frac{c_0}{4!}, \quad \text{and} \quad c_6 = -\frac{c_0}{6!}.$$

Taking n = 1, 3, and 5 in turn, we find that

$$c_3 = -\frac{c_1}{3!}, \quad c_5 = \frac{c_1}{5!}, \text{ and } c_7 = -\frac{c_1}{7!}$$

Again, the pattern is clear; we leave it for you to show (by induction) that for $k \ge 1$,

$$c_{2k} = \frac{(-1)^k c_0}{(2k)!} \quad \text{and} \quad c_{2k+1} = \frac{(-1)^k c_1}{(2k+1)!}$$

Thus we get the power series solution

$$y(x) = c_0 \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \right) + c_1 \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots \right);$$

that is, $y(x) = c_0 \cos x + c_1 \sin x$. Note that we have no problem with the radius of convergence here; the Taylor series for the sine and cosine functions converge for all x.

The solution of Example 4 can bear further comment. Suppose that we had never heard of the sine and cosine functions, let alone their Taylor series. We would then have discovered the two power series solutions

$$C(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$
(28)

and

$$S(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$
(29)

of the differential equation y'' + y = 0. Both of these power series converge for all x. For instance, the ratio test in Theorem 3 implies convergence for all z of the series $\sum (-1)^n z^n/(2n)!$ obtained from (28) by writing $z = x^2$. Hence it follows that (28) itself converges for all x, as does (by a similar ploy) the series in (29).

It is clear that C(0) = 1 and S(0) = 0, and termwise differentiation of the two series in (28) and (29) yields

$$C'(x) = -S(x)$$
 and $S'(x) = C(x)$. (30)

Consequently, C'(0) = 0 and S'(0) = 1. Thus with the aid of the power series method (all the while knowing nothing about the sine and cosine functions), we have discovered that y = C(x) is the unique solution of

$$v'' + v = 0$$

that satisfies the initial conditions y(0) = 1 and y'(0) = 0, and that y = S(x) is the unique solution that satisfies the initial conditions y(0) = 0 and y'(0) = 1. It follows that C(x) and S(x) are linearly independent, and—recognizing the importance of the differential equation y'' + y = 0—we can agree to call *C* the *cosine* function and *S* the *sine* function. Indeed, all the usual properties of these two functions can be established, using only their initial values (at x = 0) and the derivatives in (30); there is no need to refer to triangles or even to angles. (Can you use the series in (28) and (29) to show that $[C(x)]^2 + [S(x)]^2 = 1$ for all x?) This demonstrates that

The cosine and sine functions are fully determined by the differential equation y'' + y = 0 of which they are the two natural linearly independent solutions.

Figures 11.1.2 and 11.1.3 show how the geometric character of the graphs of $\cos x$ and $\sin x$ is revealed by the graphs of the Taylor polynomial approximations that we get by truncating the infinite series in (28) and (29).

This is by no means an uncommon situation. Many important special functions of mathematics occur in the first instance as power series solutions of differential equations and thus are in practice *defined* by means of these power series. In the remaining sections of this chapter we will see numerous examples of such functions.



FIGURE 11.1.2. Taylor polynomial approximations to $\cos x$.



In Problems 1 through 10, find a power series solution of the given differential equation. Determine the radius of convergence of the resulting series, and use the series in Eqs. (5) through (12) to identify the series solution in terms of familiar elementary functions. (Of course, no one can prevent you from checking your work by also solving the equations by the methods of earlier chapters!)

1.
$$y' = y$$
2. $y' = 4y$ 3. $2y' + 3y = 0$ 4. $y' + 2xy = 0$ 5. $y' = x^2y$ 6. $(x - 2)y' + y = 0$ 7. $(2x - 1)y' + 2y = 0$ 8. $2(x + 1)y' = y$ 9. $(x - 1)y' + 2y = 0$ 10. $2(x - 1)y' = 3y$

In Problems 11 through 14, use the method of Example 4 to find two linearly independent power series solutions of the given differential equation. Determine the radius of convergence of each series, and identify the general solution in terms of familiar elementary functions.

11.
$$y'' = y$$
12. $y'' = 4y$ **13.** $y'' + 9y = 0$ **14.** $y'' + y = x$

Show (as in Example 3) that the power series method fails to yield a power series solution of the form $y = \sum c_n x^n$ for the differential equations in Problems 15 through 18.

15.
$$xy' + y = 0$$
16. $2xy' = y$ **17.** $x^2y' + y = 0$ **18.** $x^3y' = 2y$

In Problems 19 through 22, first derive a recurrence relation giving c_n for $n \ge 2$ in terms of c_0 or c_1 (or both). Then apply the given initial conditions to find the values of c_0 and c_1 . Next determine c_n (in terms of n, as in the text) and, finally, identify the particular solution in terms of familiar elementary functions.

19.
$$y'' + 4y = 0; y(0) = 0, y'(0) = 3$$

20. $y'' - 4y = 0; y(0) = 2, y'(0) = 0$
21. $y'' - 2y' + y = 0; y(0) = 0, y'(0) = 1$



FIGURE 11.1.3. Taylor polynomial approximations to sin x.

22. y'' + y' - 2y = 0; y(0) = 1, y'(0) = -2

23. Show that the equation

$$x^2y'' + x^2y' + y = 0$$

has no power series solution of the form $y = \sum c_n x^n$.

- 24. Establish the binomial series in (12) by means of the following steps. (a) Show that $y = (1 + x)^{\alpha}$ satisfies the initial value problem $(1 + x)y' = \alpha y$, y(0) = 1. (b) Show that the power series method gives the binomial series in (12) as the solution of the initial value problem in part (a), and that this series converges if |x| < 1. (c) Explain why the validity of the binomial series given in (12) follows from parts (a) and (b).
- **25.** For the initial value problem

$$y'' = y' + y, \quad y(0) = 0, \quad y(1) = 1$$

derive the power series solution

$$y(x) = \sum_{n=1}^{\infty} \frac{F_n}{n!} x^n$$

where $\{F_n\}_{n=0}^{\infty}$ is the sequence 0, 1, 1, 2, 3, 5, 8, 13, ... of *Fibonacci numbers* defined by $F_0 = 0, F_1 = 1, F_n = F_{n-2} + F_{n-1}$ for n > 1.

26. (a) Show that the solution of the initial value problem

$$y' = 1 + y^2$$
, $y(0) = 0$

is $y(x) = \tan x$. (b) Because $y(x) = \tan x$ is an odd function with y'(0) = 1, its Taylor series is of the form

$$y = x + c_3 x^3 + c_5 x^5 + c_7 x^7 + \cdots$$

Substitute this series in $y' = 1 + y^2$ and equate like powers of x to derive the following relations:

$$3c_3 = 1, 5c_5 = 2c_3, 7c_7 = 2c_5 + (c_3)^2, 9c_9 = 2c_7 + 2c_3c_5, 11c_{11} = 2c_9 + 2c_3c_7 + (c_5)^2.$$

(c) Conclude that

$$\tan x = x + \frac{1}{3}x^3 + \frac{2}{15}x^5 + \frac{17}{315}x^7 + \frac{62}{2835}x^9 + \frac{1382}{155925}x^{11} + \cdots$$

(d) Would you prefer to use the Maclaurin series formula in (13) to derive the tangent series in part (c)? Think about it!

27. This section introduces the use of infinite series to solve differential equations. Conversely, differential equations can sometimes be used to sum infinite series. For example, consider the infinite series

$$1 + \frac{1}{1!} - \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} - \frac{1}{5!} + \cdots;$$

note the $+ + - + + - \cdots$ pattern of signs superimposed on the terms of the series for the number e. We could evaluate this series if we could obtain a formula for the function

$$f(x) = 1 + x - \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 - \frac{1}{5!}x^5 + \cdots,$$

because the sum of the numerical series in question is simply f(1). (a) It's possible to show that the power series given here converges for all x and that termwise differentiation is valid. Given these facts, show that f(x) satisfies the initial value problem

$$y^{(3)} = y; \quad y(0) = y'(0) = 1, \quad y''(0) = -1.$$

(b) Solve this initial value problem to show that

$$f(x) = \frac{1}{3}e^x + \frac{2}{3}e^{-x/2}\left(\cos\frac{\sqrt{3}}{2}x + \sqrt{3}\sin\frac{\sqrt{3}}{2}x\right).$$

For a suggestion, see Problem 48 of Section 5.3. (c) Evaluate f(1) to find the sum of the numerical series given here.

11.2 Power Series Solutions

The power series method introduced in Section 11.1 can be applied to linear equations of any order (as well as to certain nonlinear equations), but its most important applications are to homogeneous second-order linear differential equations of the form

$$A(x)y'' + B(x)y' + C(x)y = 0,$$
(1)

where the coefficients A, B, and C are analytic functions of x. Indeed, in most applications these coefficient functions are simple polynomials.

We saw in Example 3 of Section 11.1 that the series method does not always yield a series solution. To discover when it does succeed, we rewrite Eq. (1) in the form

$$y'' + P(x)y' + Q(x)y = 0$$
(2)

with leading coefficient 1, and with P = B/A and Q = C/A. Note that P(x) and Q(x) will generally fail to be analytic at points where A(x) vanishes. For instance, consider the equation

$$xy'' + y' + xy = 0.$$
 (3)

The coefficient functions in (3) are continuous everywhere. But in the form of (2) it is the equation

$$y'' + \frac{1}{x}y' + y = 0 \tag{4}$$

with P(x) = 1/x not analytic at x = 0.

The point x = a is called an **ordinary point** of Eq. (2)—and of the equivalent Eq. (1)—provided that the functions P(x) and Q(x) are both analytic at x = a. Otherwise, x = a is a **singular point**. Thus the only singular point of Eqs. (3) and (4) is x = 0. Recall that a quotient of analytic functions is analytic wherever the denominator is nonzero. It follows that, if $A(a) \neq 0$ in Eq. (1) with analytic coefficients, then x = a is an ordinary point. If A(x), B(x), and C(x) are *polynomials* with no common factors, then x = a is an ordinary point if and only if $A(a) \neq 0$.

Example 1 The point x = 0 is an ordinary point of the equation

$$xy'' + (\sin x)y' + x^2y = 0,$$

despite the fact that A(x) = x vanishes at x = 0. The reason is that

$$P(x) = \frac{\sin x}{x} = \frac{1}{x} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right) = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \dots$$

is nevertheless analytic at x = 0 because the division by x yields a convergent power series.

Example 2The point x = 0 is *not* an ordinary point of the equation $y'' + x^2y' + x^{1/2}y = 0.$ For while $P(x) = x^2$ is analytic at the origin, $Q(x) = x^{1/2}$ is not. The reason is that Q(x)is not differentiable at x = 0 and hence is not analytic there. (Theorem 1 of Section 11.1implies that an analytic function must be differentiable.)Example 3

$$(1 - x^3)y'' + (7x^2 + 3x^5)y' + (5x - 13x^4)y = 0$$

because the coefficient functions A(x), B(x), and C(x) are polynomials with $A(0) \neq 0$.

Theorem 2 of Section 3.1 implies that Eq. (2) has two linearly independent solutions on any open interval where the coefficient functions P(x) and Q(x) are continuous. The basic fact for our present purpose is that near an *ordinary* point *a*, these solutions will be power series in powers of x - a. A proof of the following theorem can be found in Chapter 3 of Coddington, *An Introduction to Ordinary Differential Equations* (Dover Publications, 1989).

THEOREM 1 Solutions Near an Ordinary Point

Suppose that *a* is an ordinary point of the equation

$$A(x)y'' + B(x)y' + C(x)y = 0;$$
(1)

that is, the functions P = B/A and Q = C/A are analytic at x = a. Then Eq. (1) has two linearly independent solutions, each of the form

•
$$y(x) = \sum_{n=0}^{\infty} c_n (x-a)^n.$$
 (5)

The radius of convergence of any such series solution is at least as large as the distance from a to the nearest (real or complex) singular point of Eq. (1). The coefficients in the series in (5) can be determined by its substitution in Eq. (1).

Example 4

)

Determine the radius of convergence guaranteed by Theorem 1 of a series solution of

$$(x2 + 9)y'' + xy' + x2y = 0$$
(6)

in powers of x. Repeat for a series in powers of x - 4.

Solution This example illustrates the fact that we must take into account complex singular points as well as real ones. Because

$$P(x) = \frac{x}{x^2 + 9}$$
 and $Q(x) = \frac{x^2}{x^2 + 9}$

the only singular points of Eq. (6) are $\pm 3i$. The distance (in the complex plane) of each of these from 0 is 3, so a series solution of the form $\sum c_n x^n$ has radius of convergence at least 3. The distance of each singular point from 4 is 5, so a series solution of the form $\sum c_n (x-4)^n$ has radius of convergence at least 5 (see Fig. 11.2.1).

Example 5 Find the general solution in powers of *x* of

$$(x2 - 4)y'' + 3xy' + y = 0.$$
 (7)

 $3i \phi$ Solution

FIGURE 11.2.1. Radius of convergence as distance to nearest singularity.

Then find the particular solution with y(0) = 4, y'(0) = 1.

The only singular points of Eq. (7) are ± 2 , so the series we get will have radius of convergence at least 2. (See Problem 35 for the exact radius of convergence.) Substitution of

$$y = \sum_{n=0}^{\infty} c_n x^n$$
, $y' = \sum_{n=1}^{\infty} n c_n x^{n-1}$, and $y'' = \sum_{n=2}^{\infty} n(n-1)c_n x^{n-2}$

in Eq. (7) yields

$$\sum_{n=2}^{\infty} n(n-1)c_n x^n - 4 \sum_{n=2}^{\infty} n(n-1)c_n x^{n-2} + 3 \sum_{n=1}^{\infty} nc_n x^n + \sum_{n=0}^{\infty} c_n x^n = 0.$$

We can begin the first and third summations at n = 0 as well, because no nonzero terms are thereby introduced. We shift the index of summation in the second sum by +2, replacing nwith n + 2 and using the initial value n = 0. This gives

$$\sum_{n=0}^{\infty} n(n-1)c_n x^n - 4\sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2} x^n + 3\sum_{n=0}^{\infty} nc_n x^n + \sum_{n=0}^{\infty} c_n x^n = 0.$$

After collecting coefficients of c_n and c_{n+2} , we obtain

$$\sum_{n=0}^{\infty} \left[(n^2 + 2n + 1)c_n - 4(n+2)(n+1)c_{n+2} \right] x^n = 0.$$

The identity principle yields

$$(n+1)^2c_n - 4(n+2)(n+1)c_{n+2} = 0,$$

which leads to the recurrence relation

$$c_{n+2} = \frac{(n+1)c_n}{4(n+2)} \tag{8}$$

for $n \ge 0$. With n = 0, 2, and 4 in turn, we get

$$c_2 = \frac{c_0}{4 \cdot 2}, \quad c_4 = \frac{3c_2}{4 \cdot 4} = \frac{3c_0}{4^2 \cdot 2 \cdot 4}, \quad \text{and} \quad c_6 = \frac{5c_4}{4 \cdot 6} = \frac{3 \cdot 5c_0}{4^3 \cdot 2 \cdot 4 \cdot 6}.$$

Continuing in this fashion, we evidently would find that

$$c_{2n} = \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{4^n \cdot 2 \cdot 4 \cdots (2n)} c_0.$$

With the common notation

$$(2n+1)!! = 1 \cdot 3 \cdot 5 \cdots (2n+1) = \frac{(2n+1)!}{2^n \cdot n!}$$

11.2 POWER SERIES SOLUTIONS 619

and the observation that $2 \cdot 4 \cdot 6 \cdots (2n) = 2^n \cdot n!$, we finally obtain

$$c_{2n} = \frac{(2n-1)!!}{2^{3n} \cdot n!} c_0. \tag{9}$$

(We also used the fact that $4^n \cdot 2^n = 2^{3n}$.)

With n = 1, 3, and 5 in Eq. (8), we get

$$c_3 = \frac{2c_1}{4 \cdot 3}, \quad c_5 = \frac{4c_3}{4 \cdot 5} = \frac{2 \cdot 4c_1}{4^2 \cdot 3 \cdot 5}, \quad \text{and} \quad c_7 = \frac{6c_5}{4 \cdot 7} = \frac{2 \cdot 4 \cdot 6c_1}{4^3 \cdot 3 \cdot 5 \cdot 7}.$$

It is apparent that the pattern is

$$c_{2n+1} = \frac{2 \cdot 4 \cdot 6 \cdots (2n)}{4^n \cdot 1 \cdot 3 \cdot 5 \cdots (2n+1)} c_1 = \frac{n!}{2^n \cdot (2n+1)!!} c_1.$$
(10)

The formula in (9) gives the coefficients of even subscript in terms of c_0 ; the formula in (10) gives the coefficients of odd subscript in terms of c_1 . After we separately collect the terms of the series of even and odd degree, we get the general solution

$$y(x) = c_0 \left(1 + \sum_{n=1}^{\infty} \frac{(2n-1)!!}{2^{3n} \cdot n!} x^{2n} \right) + c_1 \left(x + \sum_{n=1}^{\infty} \frac{n!}{2^n \cdot (2n+1)!!} x^{2n+1} \right).$$
(11)

Alternatively,

≻

$$y(x) = c_0 \left(1 + \frac{1}{8}x^2 + \frac{3}{128}x^4 + \frac{5}{1024}x^6 + \cdots \right) + c_1 \left(x + \frac{1}{6}x^3 + \frac{1}{30}x^5 + \frac{1}{140}x^7 + \cdots \right).$$
(11')

Because $y(0) = c_0$ and $y'(0) = c_1$, the given initial conditions imply that $c_0 = 4$ and $c_1 = 1$. Using these values in Eq. (11'), the first few terms of the particular solution satisfying y(0) = 4 and y'(0) = 1 are

$$y(x) = 4 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{3}{32}x^4 + \frac{1}{30}x^5 + \cdots$$
 (12)

Remark As in Example 5, substitution of $y = \sum c_n x^n$ in a linear second-order equation with x = 0 an ordinary point typically leads to a recurrence relation that can be used to express each of the successive coefficients c_2, c_3, c_4, \ldots in terms of the first two, c_0 and c_1 . In this event two linearly independent solutions are obtained as follows. Let $y_0(x)$ be the solution obtained with $c_0 = 1$ and $c_1 = 0$, and let $y_1(x)$ be the solution obtained with $c_0 = 0$ and $c_1 = 1$. Then

$$y_0(0) = 1$$
, $y'_0(0) = 0$ and $y_1(0) = 0$, $y'_1(0) = 1$,

so it is clear that y_0 and y_1 are linearly independent. In Example 5, $y_0(x)$ and $y_1(x)$ are defined by the two series that appear on the right-hand side in Eq. (11), which expresses the general solution in the form $y = c_0 y_0 + c_1 y_1$.

Translated Series Solutions

If in Example 5 we had sought a particular solution with given initial values y(a) and y'(a), we would have needed the general solution in the form

$$y(x) = \sum_{n=0}^{\infty} c_n (x-a)^n;$$
(13)

that is, in powers of x - a rather than in powers of x. For only with a solution of the form in (13) is it true that the initial conditions

$$y(a) = c_0$$
 and $y'(a) = c_1$

determine the arbitrary constants c_0 and c_1 in terms of the initial values of y and y'. Consequently, to solve an initial value problem, we need a series expansion of the general solution centered at the point where the initial conditions are specified.

Example 6 Solve the initial value problem

$$(t^{2} - 2t - 3)\frac{d^{2}y}{dt^{2}} + 3(t - 1)\frac{dy}{dt} + y = 0; \quad y(1) = 4, \quad y'(1) = -1.$$
(14)

Solution

We need a general solution of the form $\sum c_n(t-1)^n$. But instead of substituting this series in (14) to determine the coefficients, it simplifies the computations if we first make the substitution x = t - 1, so that we wind up looking for a series of the form $\sum c_n x^n$ after all. To transform Eq. (14) into one with the new independent variable x, we note that

$$t^{2} - 2t - 3 = (x + 1)^{2} - 2(x + 1) - 3 = x^{2} - 4,$$
$$\frac{dy}{dt} = \frac{dy}{dx}\frac{dx}{dt} = \frac{dy}{dx} = y',$$

and

$$\frac{d^2 y}{dt^2} = \left[\frac{d}{dx}\left(\frac{dy}{dx}\right)\right]\frac{dx}{dt} = \frac{d}{dx}(y') = y'',$$

where primes denote differentiation with respect to x. Hence we transform Eq. (14) into

$$(x^2 - 4)y'' + 3xy' + y = 0$$

with initial conditions y = 4 and y' = 1 at x = 0 (corresponding to t = 1). This is the initial value problem we solved in Example 5, so the particular solution in (12) is available. We substitute t - 1 for x in Eq. (12) and thereby obtain the desired particular solution

$$y(t) = 4 + (t-1) + \frac{1}{2}(t-1)^2 + \frac{1}{6}(t-1)^3 + \frac{3}{32}(t-1)^4 + \frac{1}{30}(t-1)^5 + \cdots$$

This series converges if -1 < t < 3. (Why?) A series such as this can be used to estimate numerical values of the solution. For instance,

$$y(0.8) = 4 + (-0.2) + \frac{1}{2}(-0.2)^2 + \frac{1}{6}(-0.2)^3 + \frac{3}{32}(-0.2)^4 + \frac{1}{30}(-0.2)^5 + \cdots,$$

so that $y(0.8) \approx 3.8188$.

The last computation in Example 6 illustrates the fact that series solutions of differential equations are useful not only for establishing general properties of a solution, but also for numerical computations when an expression of the solution in terms of elementary functions is unavailable.

Types of Recurrence Relation

The formula in Eq. (8) is an example of a **two-term** recurrence relation; it expresses each coefficient in the series in terms of *one* of the preceding coefficients. A **manyterm** recurrence relation expresses each coefficient in the series in terms of two or more preceding coefficients. In the case of a many-term recurrence relation, it is generally inconvenient or even impossible to find a formula that gives the typical coefficient c_n in terms of n. The next example shows what we sometimes can do with a three-term recurrence relation. Find two linearly independent solutions of

$$y'' - xy' - x^2y = 0. (15)$$

Solution

Example 7

We make the usual substitution of the power series $y = \sum c_n x^n$. This results in the equation

$$\sum_{n=2}^{\infty} n(n-1)c_n x^{n-2} - \sum_{n=1}^{\infty} nc_n x^n - \sum_{n=0}^{\infty} c_n x^{n+2} = 0.$$

We can start the second sum at n = 0 without changing anything else. To make each term include x^n in its general term, we shift the index of summation in the first sum by +2 (replace n with n + 2), and we shift it by -2 in the third sum (replace n with n - 2). These shifts yield

$$\sum_{n=0}^{\infty} (n+2)(n+1)c_{n+2}x^n - \sum_{n=0}^{\infty} nc_n x^n - \sum_{n=2}^{\infty} c_{n-2}x^n = 0.$$

The common range of these three summations is $n \ge 2$, so we must separate the terms corresponding to n = 0 and n = 1 in the first two sums before collecting coefficients of x^n . This gives

$$2c_2 + 6c_3x - c_1x + \sum_{n=2}^{\infty} \left[(n+2)(n+1)c_{n+2} - nc_n - c_{n-2} \right] x^n = 0.$$

The identity principle now implies that $2c_2 = 0$, that $c_3 = \frac{1}{6}c_1$, and the three-term recurrence relation

$$c_{n+2} = \frac{nc_n + c_{n-2}}{(n+2)(n+1)} \tag{16}$$

for $n \ge 2$. In particular,

$$c_{4} = \frac{2c_{2} + c_{0}}{12}, \qquad c_{5} = \frac{3c_{3} + c_{1}}{20}, \qquad c_{6} = \frac{4c_{4} + c_{2}}{30},$$

$$c_{7} = \frac{5c_{5} + c_{3}}{42}, \qquad c_{8} = \frac{6c_{6} + c_{4}}{56}.$$
(17)

Thus all values of c_n for $n \ge 4$ are given in terms of the arbitrary constants c_0 and c_1 because $c_2 = 0$ and $c_3 = \frac{1}{6}c_1$.

To get our first solution y_1 of Eq. (15), we choose $c_0 = 1$ and $c_1 = 0$, so that $c_2 = c_3 = 0$. Then the formulas in (17) yield

$$c_4 = \frac{1}{12}, \quad c_5 = 0, \quad c_6 = \frac{1}{90}, \quad c_7 = 0, \quad c_8 = \frac{3}{1120}$$

thus

>

$$y_1(x) = 1 + \frac{1}{12}x^4 + \frac{1}{90}x^6 + \frac{3}{1120}x^8 + \cdots$$
 (18)

Because $c_1 = c_3 = 0$, it is clear from Eq. (16) that this series contains only terms of even degree.

To obtain a second linearly independent solution y_2 of Eq. (15), we take $c_0 = 0$ and $c_1 = 1$, so that $c_2 = 0$ and $c_3 = \frac{1}{6}$. Then the formulas in (17) yield

$$c_4 = 0, \quad c_5 = \frac{3}{40}, \quad c_6 = 0, \quad c_7 = \frac{13}{1008},$$

so that

$$y_2(x) = x + \frac{1}{6}x^3 + \frac{3}{40}x^5 + \frac{13}{1008}x^7 + \cdots$$
 (19)

Because $c_0 = c_2 = 0$, it is clear from Eq. (16) that this series contains only terms of odd degree. The solutions $y_1(x)$ and $y_2(x)$ are linearly independent because $y_1(0) = 1$ and $y'_1(0) = 0$, whereas $y_2(0) = 0$ and $y'_2(0) = 1$. A general solution of Eq. (15) is a linear combination of the power series in (18) and (19). Equation (15) has no singular points, so the power series representing $y_1(x)$ and $y_2(x)$ converge for all x.

The Legendre Equation

The **Legendre equation** of order α is the second-order linear differential equation

$$(1 - x2)y'' - 2xy' + \alpha(\alpha + 1)y = 0,$$
(20)

where the real number α satisfies the inequality $\alpha > -1$. This differential equation has extensive applications, ranging from numerical integration formulas (such as Gaussian quadrature) to the problem of determining the steady-state temperature within a solid spherical ball when the temperatures at points of its boundary are known. The only singular points of the Legendre equation are at +1 and -1, so it has two linearly independent solutions that can be expressed as power series in powers of x with radius of convergence at least 1. The substitution $y = \sum c_m x^m$ in Eq. (20) leads (see Problem 31) to the recurrence relation

$$c_{m+2} = -\frac{(\alpha - m)(\alpha + m + 1)}{(m+1)(m+2)}c_m$$
(21)

for $m \ge 0$. We are using *m* as the index of summation because we have another role for *n* to play.

In terms of the arbitrary constants c_0 and c_1 , Eq. (21) yields

$$c_{2} = -\frac{\alpha(\alpha + 1)}{2!}c_{0},$$

$$c_{3} = -\frac{(\alpha - 1)(\alpha + 2)}{3!}c_{1},$$

$$c_{4} = \frac{\alpha(\alpha - 2)(\alpha + 1)(\alpha + 3)}{4!}c_{0},$$

$$c_{5} = \frac{(\alpha - 1)(\alpha - 3)(\alpha + 2)(\alpha + 4)}{5!}c_{1}.$$

We can show without much trouble that for m > 0,

$$c_{2m} = (-1)^m \frac{\alpha(\alpha - 2)(\alpha - 4)\cdots(\alpha - 2m + 2)(\alpha + 1)(\alpha + 3)\cdots(\alpha + 2m - 1)}{(2m)!} c_0$$
(22)

and

$$c_{2m+1} = (-1)^m \frac{(\alpha - 1)(\alpha - 3)\cdots(\alpha - 2m + 1)(\alpha + 2)(\alpha + 4)\cdots(\alpha + 2m)}{(2m + 1)!}c_1.$$
(23)

Alternatively,

$$c_{2m} = (-1)^m a_{2m} c_0$$
 and $c_{2m+1} = (-1)^m a_{2m+1} c_1$

where a_{2m} and a_{2m+1} denote the fractions in Eqs. (22) and (23), respectively. With this notation, we get two linearly independent power series solutions

$$y_1(x) = c_0 \sum_{m=0}^{\infty} (-1)^m a_{2m} x^{2m}$$
 and $y_2(x) = c_1 \sum_{m=0}^{\infty} (-1)^m a_{2m+1} x^{2m+1}$ (24)

of Legendre's equation of order α .

11.2 POWER SERIES SOLUTIONS 623

Now suppose that $\alpha = n$, a nonnegative *integer*. If $\alpha = n$ is even, we see from Eq. (22) that $a_{2m} = 0$ when 2m > n. In this case, $y_1(x)$ is a *polynomial* of degree n and y_2 is a (nonterminating) infinite series. If $\alpha = n$ is an odd positive integer, we see from Eq. (23) that $a_{2m+1} = 0$ when 2m + 1 > n. In this case, $y_2(x)$ is a *polynomial* of degree n and y_1 is a (nonterminating) infinite series. Thus in either case, one of the two solutions in (24) is a polynomial and the other is a nonterminating series.

With an appropriate choice (made separately for each n) of the arbitrary constants c_0 (n even) or c_1 (n odd), the nth-degree polynomial solution of Legendre's equation of order n,

$$(1 - x^2)y'' - 2xy' + n(n+1)y = 0, (25)$$

is denoted by $P_n(x)$ and is called the **Legendre polynomial** of degree *n*. It is customary (for a reason indicated in Problem 32) to choose the arbitrary constant so that the coefficient of x^n in $P_n(x)$ is $(2n)!/[2^n(n!)^2]$. It then turns out that

$$P_n(x) = \sum_{k=0}^{N} \frac{(-1)^k (2n-2k)!}{2^n k! (n-k)! (n-2k)!} x^{n-2k},$$
(26)

where N = [n/2], the integral part of n/2. The first six Legendre polynomials are

$$P_0(x) \equiv 1, \qquad P_1(x) = x,$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1), \qquad P_3(x) = \frac{1}{2}(5x^3 - 3x),$$

$$P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3), \qquad P_5(x) = \frac{1}{8}(63x^5 - 70x^3 + 15x),$$

and their graphs are shown in Fig. 11.2.2.



FIGURE 11.2.2. Graphs $y = P_n(x)$ of the Legendre polynomials for n = 1, 2, 3, 4, and 5. The graphs are distinguished by the fact that all *n* zeros of $P_n(x)$ lie in the interval -1 < x < 1.

11.2 **Problems**

Find general solutions in powers of x of the differential equations in Problems 1 through 15. State the recurrence relation and the guaranteed radius of convergence in each case.

1. $(x^2 - 1)y'' + 4xy' + 2y = 0$ 2. $(x^2 + 2)y'' + 4xy' + 2y = 0$ 3. y'' + xy' + y = 04. $(x^2 + 1)y'' + 6xy' + 4y = 0$ 5. $(x^2 - 3)y'' + 2xy' = 0$ 6. $(x^2 - 1)y'' - 6xy' + 12y = 0$ 7. $(x^2 + 3)y'' - 7xy' + 16y = 0$ 8. $(2 - x^2)y'' - xy' + 16y = 0$ 9. $(x^2 - 1)y'' + 8xy' + 12y = 0$ 10. 3y'' + xy' - 4y = 011. 5y'' - 2xy' + 10y = 012. $y'' - x^2y' - 3xy = 0$ 13. $y'' + x^2y' + 2xy = 0$ 14. y'' + xy = 0 (an *Airy equation*) 15. $y'' + x^2y = 0$

Use power series to solve the initial value problems in Problems 16 and 17.

16. $(1 + x^2)y'' + 2xy' - 2y = 0; y(0) = 0, y'(0) = 1$ **17.** y'' + xy' - 2y = 0; y(0) = 1, y'(0) = 0

Solve the initial value problems in Problems 18 through 22. First make a substitution of the form t = x - a, then find a solution $\sum c_n t^n$ of the transformed differential equation. State the interval of values of x for which Theorem 1 of this section guarantees convergence.

- **18.** y'' + (x 1)y' + y = 0; y(1) = 2, y'(1) = 0
- **19.** $(2x x^2)y'' 6(x 1)y' 4y = 0; y(1) = 0, y'(1) = 1$
- **20.** $(x^2 6x + 10)y'' 4(x 3)y' + 6y = 0; y(3) = 2, y'(3) = 0$
- **21.** $(4x^2 + 16x + 17)y'' = 8y; y(-2) = 1, y'(-2) = 0$
- **22.** $(x^2 + 6x)y'' + (3x + 9)y' 3y = 0; y(-3) = 0, y'(-3) = 2$

In Problems 23 through 26, find a three-term recurrence relation for solutions of the form $y = \sum c_n x^n$. Then find the first three nonzero terms in each of two linearly independent solutions.

- **23.** y'' + (1+x)y = 0 **24.** $(x^2 - 1)y'' + 2xy' + 2xy = 0$ **25.** $y'' + x^2y' + x^2y = 0$ **26.** $(1 + x^3)y'' + x^4y = 0$
- 27. Solve the initial value problem

$$y'' + xy' + (2x^2 + 1)y = 0;$$
 $y(0) = 1, y'(0) = -1.$

Determine sufficiently many terms to compute y(1/2) accurate to four decimal places.

In Problems 28 through 30, find the first three nonzero terms in each of two linearly independent solutions of the form $y = \sum c_n x^n$. Substitute known Taylor series for the analytic functions and retain enough terms to compute the necessary coefficients.

- **28.** $y'' + e^{-x}y = 0$
- **29.** $(\cos x)y'' + y = 0$ **30.** $xy'' + (\sin x)y' + xy = 0$
- **31.** Derive the recurrence relation in (21) for the Legendre equation.
- Follow the steps outlined in this problem to establish Rodrigues's formula

$$P_n(x) = \frac{1}{n! \, 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n$$

for the *n*th-degree Legendre polynomial. (a) Show that $v = (x^2 - 1)^n$ satisfies the differential equation

$$(1 - x^2)v' + 2nxv = 0.$$

Differentiate each side of this equation to obtain

$$(1 - x2)v'' + 2(n - 1)xv' + 2nv = 0.$$

(b) Differentiate each side of the last equation n times in succession to obtain

$$(1 - x2)v(n+2) - 2xv(n+1) + n(n+1)v(n) = 0.$$

Thus $u = v^{(n)} = D^n (x^2 - 1)^n$ satisfies Legendre's equation of order *n*. (c) Show that the coefficient of x^n in *u* is (2n)!/n!; then state why this proves Rodrigues' formula. (Note that the coefficient of x^n in $P_n(x)$ is $(2n)!/[2^n (n!)^2]$.)

33. The **Hermite equation** of order α is

$$y'' - 2xy' + 2\alpha y = 0.$$

(a) Derive the two power series solutions

$$y_1 = 1 + \sum_{m=1}^{\infty} (-1)^m \frac{2^m \alpha (\alpha - 2) \cdots (\alpha - 2m + 2)}{(2m)!} x^{2m}$$

and

$$y_2 = x + \sum_{m=1}^{\infty} (-1)^m \frac{2^m (\alpha - 1)(\alpha - 3) \cdots (\alpha - 2m + 1)}{(2m + 1)!} x^{2m + 1}.$$

Show that y_1 is a polynomial if α is an even integer, whereas y_2 is a polynomial if α is an odd integer. (b) The **Hermite polynomial** of degree *n* is denoted by $H_n(x)$. It is the *n*th-degree polynomial solution of Hermite's equation, multiplied by a suitable constant so that the coefficient of x^n is 2^n . Show that the first six Hermite polynomials are

$$H_0(x) \equiv 1, \qquad H_1(x) = 2x,$$

$$H_2(x) = 4x^2 - 2, \qquad H_3(x) = 8x^3 - 12x,$$

$$H_4(x) = 16x^4 - 48x^2 + 12,$$

$$H_5(x) = 32x^5 - 160x^3 + 120x.$$

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} \left(e^{-x^2} \right)$$

Verify that this formula does in fact give an *n*th-degree polynomial. It is interesting to use a computer algebra system to investigate the conjecture that (for each *n*) the zeros of the Hermite polynomials H_n and H_{n+1} are "interlaced"—that is, the *n* zeros of H_n lie in the *n* bounded open intervals whose endpoints are successive pairs of zeros of H_{n+1} .

34. The discussion following Example 4 in Section 11.1 suggests that the differential equation y'' + y = 0 could be used to introduce and define the familiar sine and cosine functions. In a similar fashion, the *Airy equation*

$$y'' = xy$$

serves to introduce two new special functions that appear in applications ranging from radio waves to molecular vibrations. Derive the first three or four terms of two different power series solutions of the Airy equation. Then verify that your results agree with the formulas

$$y_1(x) = 1 + \sum_{k=1}^{\infty} \frac{1 \cdot 4 \cdots (3k-2)}{(3k)!} x^{3k}$$

and

$$y_2(x) = x + \sum_{k=1}^{\infty} \frac{2 \cdot 5 \cdots (3k-1)}{(3k+1)!} x^{3k+1}$$

for the solutions that satisfy the initial conditions $y_1(0) = 1$, $y'_1(0) = 0$ and $y_2(0) = 0$, $y'_2(0) = 1$, respectively. The special combinations

$$\operatorname{Ai}(x) = \frac{y_1(x)}{3^{2/3}\Gamma(\frac{2}{3})} - \frac{y_2(x)}{3^{1/3}\Gamma(\frac{1}{3})}$$

and

$$\operatorname{Bi}(x) = \frac{y_1(x)}{3^{1/6}\Gamma(\frac{2}{3})} + \frac{y_2(x)}{3^{-1/6}\Gamma(\frac{1}{3})}$$

define the standard *Airy functions* that appear in mathematical tables and computer algebra systems. Their graphs shown in Fig. 11.2.3 exhibit trigonometric-like oscillatory behavior for x < 0, whereas Ai(x) decreases exponentially and Bi(x) increases exponentially as $x \to +\infty$. It is interesting to use a computer algebra system to investigate how many terms must be retained in the y_1 - and y_2 -series above to produce a figure that is visually indistinguishable from Fig. 11.2.3 (which is based on high-precision approximations to the Airy functions).



FIGURE 11.2.3. The Airy function graphs y = Ai(x) and y = Bi(x).

35. (a) To determine the radius of convergence of the series solution in Example 5, write the series of terms of even degree in Eq. (11) in the form

$$y_0(x) = 1 + \sum_{n=1}^{\infty} c_{2n} x^{2n} = 1 + \sum_{n=1}^{\infty} a_n z^n$$

where $a_n = c_{2n}$ and $z = x^2$. Then apply the recurrence relation in Eq. (8) and Theorem 3 in Section 11.1 to show that the radius of convergence of the series in z is 4. Hence the radius of convergence of the series in x is 2. How does this corroborate Theorem 1 in this section? (b) Write the series of terms of odd degree in Eq. (11) in the form

$$y_1(x) = x \left(1 + \sum_{n=1}^{\infty} c_{2n+1} x^{2n} \right) = x \left(1 + \sum_{n=1}^{\infty} b_n z^n \right)$$

to show similarly that its radius of convergence (as a power series in x) is also 2.

11.2 Application Automatic Computation of Series Coefficients

Go to goo.gl/7BhXBb to download this application's computing resources including *Maple/Mathematica*/MATLAB. Repeated application of a recurrence relation to grind out successive coefficients is—especially in the case of a recurrence relation with three or more terms—a tedious aspect of the infinite series method. Here we illustrate the use of a computer algebra system not only to automate this task, but also to explore interactively the graphical effect of changing the number k of terms we include in a partial-sum approximation to the actual solution given by the full infinite series. In Example 7 we saw that the coefficients in the series solution $y = \sum c_n x^n$ of the differential equation

$$y'' - xy' - x^2 y = 0 (1)$$

are given in terms of the two arbitrary coefficients c_0 and c_1 by

$$c_2 = 0, \quad c_3 = \frac{c_1}{6}, \quad \text{and} \quad c_{n+2} = \frac{nc_n + c_{n-2}}{(n+2)(n+1)} \quad \text{for} \quad n \ge 2.$$
 (2)

It would appear to be a routine matter to implement such a recurrence relation, but a twist results from the fact that a typical computer system array is indexed by the subscripts 1, 2, 3, ..., rather than by the subscripts 0, 1, 2, ... that match the exponents in the successive terms of a power series that begins with a constant term. For this reason we first rewrite our proposed power series solution in the form

$$y = \sum_{n=0}^{\infty} c_n x^n = \sum_{n=1}^{\infty} b_n x^{n-1}$$
(3)

where $b_n = c_{n-1}$ for each $n \ge 1$. Then the first two conditions in (1) imply that $b_3 = 0$ and $b_4 = \frac{1}{6}b_2$; also, the recurrence relation (with *n* replaced with n-1) yields the new recurrence relation

$$b_{n+2} = c_{n+1} = \frac{(n-1)c_{n-1} + c_{n-3}}{(n+1)n} = \frac{(n-1)b_n + b_{n-2}}{n(n+1)}.$$
(4)

Now we are ready to begin. Suppose that we want to calculate the terms through the 10th degree in (2) with the initial conditions $b_1 = b_2 = 1$. Then either the *Maple* commands

or the Mathematica commands

quickly yield the coefficients $\{b_n\}$ corresponding to the solution

$$y(x) = 1 + x + \frac{x^3}{6} + \frac{x^4}{12} + \frac{3x^5}{40} + \frac{x^6}{90} + \frac{13x^7}{1008} + \frac{3x^8}{1120} + \frac{119x^9}{51840} + \frac{41x^{10}}{113400} + \dots$$
(5)

You might note that the even- and odd-degree terms here agree with those shown in Eqs. (18) and (19), respectively, of Example 7.

The MATLAB commands

give the same results, *except that* the coefficient b_{10} of x^9 is shown as 73/31801 rather than the correct value 119/51840 shown in Eq. (4). It happens that

$$\frac{73}{31801} \approx 0.0022955253$$
 while $\frac{119}{51840} \approx 0.0022955247$,

so the two rational fractions agree when rounded to 9 decimal places. The explanation is that (unlike *Mathematica* and *Maple*) MATLAB works internally with decimal rather than exact arithmetic. But at the end its **format** rat algorithm converts a correct 14-place approximation for b_{10} into an incorrect rational fraction that's "close but no cigar."

The MATLAB commands above form the basis for the interactive display shown in Fig. 11.2.4, which graphs the actual solution (blue curve) of the differential equation (1), with initial conditions $b_1 = b_2 = 1$, together with the approximate solution (black curve) consisting of the terms through the fourth degree (k = 5) in (5). The pop-up menu allows the user to vary the number of terms k and thus immediately see the graphical effect of changing the number of terms included in the series expansion. With k = 10, the actual and approximate solutions are indistinguishable in this viewing window.

Finally, you can substitute $b_1 = 1$, $b_2 = 0$ and $b_1 = 0$, $b_2 = 1$ separately (instead of $b_1 = b_2 = 1$) in the commands shown here to derive partial sums of the two linearly independent solutions displayed in Eqs. (18) and (19) of Example 7. This technique can be applied to any of the examples and problems in this section.

11.3 Frobenius Series Solutions

We now investigate the solution of the homogeneous second-order linear equation

$$A(x)y'' + B(x)y' + C(x)y = 0$$
(1)

near a singular point. Recall that if the functions A, B, and C are polynomials having no common factors, then the singular points of Eq. (1) are simply those points where A(x) = 0. For instance, x = 0 is the only singular point of the Bessel equation of order n,

$$x^{2}y'' + xy' + (x^{2} - n^{2})y = 0,$$

whereas the Legendre equation of order n,

$$(1 - x2)y'' - 2xy' + n(n + 1)y = 0,$$

has the two singular points x = -1 and x = 1. It turns out that some of the features of the solutions of such equations of the most importance for applications are largely determined by their behavior near their singular points.



FIGURE 11.2.4. MATLAB interactive display. The blue curve represents the actual solution of the initial value problem $y'' - xy' - x^2y = 0$, y(0) = y'(0) = 1, whereas the black curve shows the partial-sum approximation of the series solution (5) with terms through the fourth degree (k = 5).

We will restrict our attention to the case in which x = 0 is a singular point of Eq. (1). A differential equation having x = a as a singular point is easily transformed by the substitution t = x - a into one having a corresponding singular point at 0. For example, let us substitute t = x - 1 into the Legendre equation of order *n*. Because

$$y' = \frac{dy}{dx} = \frac{dy}{dt}\frac{dt}{dx} = \frac{dy}{dt},$$
$$y'' = \frac{d^2y}{dx^2} = \left[\frac{d}{dt}\left(\frac{dy}{dx}\right)\right]\frac{dt}{dx} = \frac{d^2y}{dt^2}$$

and $1 - x^2 = 1 - (t + 1)^2 = -2t - t^2$, we get the equation

$$-t(t+2)\frac{d^2y}{dt^2} - 2(t+1)\frac{dy}{dt} + n(n+1)y = 0.$$

This new equation has the singular point t = 0 corresponding to x = 1 in the original equation; it also has the singular point t = -2 corresponding to x = -1.

Types of Singular Points

A differential equation having a singular point at 0 ordinarily will *not* have power series solutions of the form $y(x) = \sum c_n x^n$, so the straightforward method of Section 11.2 fails in this case. To investigate the form that a solution of such an equation might take, we assume that Eq. (1) has analytic coefficient functions and rewrite it in the standard form

$$y'' + P(x)y' + Q(x)y = 0,$$
(2)

where P = B/A and Q = C/A. Recall that x = 0 is an ordinary point (as opposed to a singular point) of Eq. (2) if the functions P(x) and Q(x) are analytic at x = 0; that is, if P(x) and Q(x) have convergent power series expansions in powers of x on some open interval containing x = 0. Now it can be proved that each of the functions P(x) and Q(x) either is analytic or approaches $\pm \infty$ as $x \to 0$. Consequently, x = 0is a singular point of Eq. (2) provided that either P(x) or Q(x) (or both) approaches $\pm \infty$ as $x \to 0$. For instance, if we rewrite the Bessel equation of order n in the form

$$y'' + \frac{1}{x}y' + \left(1 - \frac{n^2}{x^2}\right)y = 0,$$

we see that P(x) = 1/x and $Q(x) = 1 - (n/x)^2$ both approach infinity as $x \to 0$.

We will see presently that the power series method can be generalized to apply near the singular point x = 0 of Eq. (2), provided that P(x) approaches infinity no more rapidly than 1/x, and Q(x) no more rapidly than $1/x^2$, as $x \to 0$. This is a way of saying that P(x) and Q(x) have only "weak" singularities at x = 0. To state this more precisely, we rewrite Eq. (2) in the form

$$y'' + \frac{p(x)}{x}y' + \frac{q(x)}{x^2}y = 0,$$
(3)

where

>

>

>

p(x) = xP(x) and $q(x) = x^2Q(x)$. (4)

DEFINITION Regular Singular Point

The singular point x = 0 of Eq. (3) is a **regular singular point** if the functions p(x) and q(x) are both analytic at x = 0. Otherwise it is an **irregular singular point**.

In particular, the singular point x = 0 is a *regular* singular point if p(x) and q(x) are both polynomials. For instance, we see that x = 0 is a regular singular point of Bessel's equation of order *n* by writing that equation in the form

$$y'' + \frac{1}{x}y' + \frac{x^2 - n^2}{x^2}y = 0,$$

noting that $p(x) \equiv 1$ and $q(x) = x^2 - n^2$ are both polynomials in x.

By contrast, consider the equation

$$2x^{3}y'' + (1+x)y' + 3xy = 0,$$

which has the singular point x = 0. If we write this equation in the form of (3), we get

$$y'' + \frac{(1+x)/(2x^2)}{x}y' + \frac{\frac{3}{2}}{x^2}y = 0.$$

Because

$$p(x) = \frac{1+x}{2x^2} = \frac{1}{2x^2} + \frac{1}{2x} \to \infty$$

as $x \to 0$ (although $q(x) \equiv \frac{3}{2}$ is a polynomial), we see that x = 0 is an irregular singular point. We will not discuss the solution of differential equations near irregular singular points; this is a considerably more advanced topic than the solution of differential equations near regular singular points.

Example 1 Consider the differential equation

$$x^{2}(1+x)y'' + x(4-x^{2})y' + (2+3x)y = 0.$$

In the standard form y'' + Py' + Qy = 0 it is

$$y'' + \frac{4 - x^2}{x(1 + x)}y' + \frac{2 + 3x}{x^2(1 + x)}y = 0.$$

Because

$$P(x) = \frac{4 - x^2}{x(1 + x)}$$
 and $Q(x) = \frac{2 + 3x}{x^2(1 + x)}$

both approach ∞ as $x \to 0$, we see that x = 0 is a singular point. To determine the nature of this singular point, we write the differential equation in the form of Eq. (3):

$$y'' + \frac{(4-x^2)/(1+x)}{x}y' + \frac{(2+3x)/(1+x)}{x^2}y = 0.$$

Thus

$$p(x) = \frac{4 - x^2}{1 + x}$$
 and $q(x) = \frac{2 + 3x}{1 + x}$.

Because a quotient of polynomials is analytic wherever the denominator is nonzero, we see that p(x) and q(x) are both analytic at x = 0. Hence x = 0 is a *regular* singular point of the given differential equation.

It may happen that when we begin with a differential equation in the general form in Eq. (1) and rewrite it in the form in (3), the functions p(x) and q(x) as given in (4) are indeterminate forms at x = 0. In this case the situation is determined by the limits

$$p_0 = p(0) = \lim_{x \to 0} p(x) = \lim_{x \to 0} x P(x)$$
(5)

and

$$q_0 = q(0) = \lim_{x \to 0} q(x) = \lim_{x \to 0} x^2 Q(x).$$
(6)

If $p_0 = 0 = q_0$, then x = 0 may be an ordinary point of the differential equation $x^2y'' + xp(x)y' + q(x)y = 0$ in (3). Otherwise:

- If both the limits in (5) and (6) exist and are *finite*, then x = 0 is a regular singular point.
- If either limit fails to exist or is infinite, then x = 0 is an irregular singular point.

Remark The most common case in applications, for the differential equation written in the form

$$y'' + \frac{p(x)}{x}y' + \frac{q(x)}{x^2}y = 0,$$
(3)

is that the functions p(x) and q(x) are *polynomials*. In this case $p_0 = p(0)$ and $q_0 = q(0)$ are simply the constant terms of these polynomials, so there is no need to evaluate the limits in Eqs. (5) and (6).

Example 2 To investigate the nature of the point x = 0 for the differential equation

$$x^{4}y'' + (x^{2}\sin x)y' + (1 - \cos x)y = 0,$$

we first write it in the form in (3):

$$y'' + \frac{(\sin x)/x}{x}y' + \frac{(1 - \cos x)/x^2}{x^2}y = 0$$

Then l'Hôpital's rule gives the values

$$p_0 = \lim_{x \to 0} \frac{\sin x}{x} = \lim_{x \to 0} \frac{\cos x}{1} = 1$$

and

$$q_0 = \lim_{x \to 0} \frac{1 - \cos x}{x^2} = \lim_{x \to 0} \frac{\sin x}{2x} = \frac{1}{2}$$

for the limits in (5) and (6). Since they are not both zero, we see that x = 0 is not an ordinary point. But both limits are finite, so the singular point x = 0 is regular. Alternatively, we could write

$$p(x) = \frac{\sin x}{x} = \frac{1}{x} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots \right) = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \cdots$$

and

$$q(x) = \frac{1 - \cos x}{x^2} = \frac{1}{x^2} \left[1 - \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \right) \right] = \frac{1}{2!} - \frac{x^2}{4!} + \frac{x^4}{6!} - \cdots$$

These (convergent) power series show explicitly that p(x) and q(x) are analytic and moreover that $p_0 = p(0) = 1$ and $q_0 = q(0) = \frac{1}{2}$, thereby verifying directly that x = 0 is a regular singular point.

The Method of Frobenius

We now approach the task of actually finding solutions of a second-order linear differential equation near the regular singular point x = 0. The simplest such equation is the constant-coefficient *equidimensional equation*

$$x^2 y'' + p_0 x y' + q_0 y = 0 (7)$$

to which Eq. (3) reduces when $p(x) \equiv p_0$ and $q(x) \equiv q_0$ are constants. In this case we can verify by direct substitution that the simple power function $y(x) = x^r$ is a solution of Eq. (7) if and only if r is a root of the quadratic equation

$$r(r-1) + p_0 r + q_0 = 0. (8)$$

In the general case, in which p(x) and q(x) are power series rather than constants, it is a reasonable conjecture that our differential equation might have a solution of the form

$$y(x) = x^r \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_n x^{n+r} = c_0 x^r + c_1 x^{r+1} + c_2 x^{r+2} + \dots$$
(9)

—the product of x^r and a power series. This turns out to be a very fruitful conjecture; according to Theorem 1 (soon to be stated formally), every equation of the form in (1) having x = 0 as a regular singular point does, indeed, have at least one such solution. This fact is the basis for the **method of Frobenius**, named for the German mathematician Georg Frobenius (1848–1917), who discovered the method in the 1870s.

An infinite series of the form in (9) is called a **Frobenius series**. Note that a Frobenius series is generally *not* a power series. For instance, with $r = -\frac{1}{2}$ the series in (9) takes the form

$$y = c_0 x^{-1/2} + c_1 x^{1/2} + c_2 x^{3/2} + c_3 x^{5/2} + \cdots;$$

it is not a series in *integral* powers of x.

To investigate the possible existence of Frobenius series solutions, we begin with the equation

≻

 $x^{2}y'' + xp(x)y' + q(x)y = 0,$ (10)

obtained by multiplying the equation in (3) by x^2 . If x = 0 is a regular singular point, then p(x) and q(x) are analytic at x = 0, so

$$p(x) = p_0 + p_1 x + p_2 x^2 + p_3 x^3 + \cdots,$$

$$q(x) = q_0 + q_1 x + q_2 x^2 + q_3 x^3 + \cdots.$$
(11)

Suppose that Eq. (10) has the Frobenius series solution

$$y = \sum_{n=0}^{\infty} c_n x^{n+r}.$$
 (12)

We may (and always do) assume that $c_0 \neq 0$ because the series must have a first nonzero term. Termwise differentiation in Eq. (12) leads to

>
$$y' = \sum_{n=0}^{\infty} c_n (n+r) x^{n+r-1}$$
 (13)

and

>

>

$$y'' = \sum_{n=0}^{\infty} c_n (n+r)(n+r-1)x^{n+r-2}.$$
 (14)

Substitution of the series in Eqs. (11) through (14) in Eq. (10) now yields

$$[r(r-1)c_0x^r + (r+1)rc_1x^{r+1} + \cdots] + [p_0x + p_1x^2 + \cdots] \cdot [rc_0x^{r-1} + (r+1)c_1x^r + \cdots] + [q_0 + q_1x + \cdots] \cdot [c_0x^r + c_1x^{r+1} + \cdots] = 0.$$
(15)

Upon multiplying initial terms of the two products on the left-hand side here and then collecting coefficients of x^r , we see that the lowest-degree term in Eq. (15) is $c_0[r(r-1) + p_0r + q_0]x^r$. If Eq. (15) is to be satisfied identically, then the coefficient of this term (as well as those of the higher-degree terms) must vanish. But we are assuming that $c_0 \neq 0$, so it follows that r must satisfy the quadratic equation

$$r(r-1) + p_0 r + q_0 = 0 \tag{16}$$

of precisely the same form as that obtained with the equidimensional equation in (7). Equation (16) is called the **indicial equation** of the differential equation in (10), and its two roots (possibly equal) are the **exponents** of the differential equation (at the regular singular point x = 0).

Our derivation of Eq. (16) shows that *if* the Frobenius series $y = x^r \sum c_n x^n$ is to be a solution of the differential equation in (10), *then* the exponent *r* must be one of the roots r_1 and r_2 of the indicial equation in (16). If $r_1 \neq r_2$, it follows that there are two possible Frobenius series solutions, whereas if $r_1 = r_2$ there is only one possible Frobenius series solution; the second solution cannot be a Frobenius series. The exponents r_1 and r_2 in the possible Frobenius series solutions are determined (using the indicial equation) by the values $p_0 = p(0)$ and $q_0 = q(0)$ that we have discussed. In practice, particularly when the coefficients in the differential equation in the original form in (1) are polynomials, the simplest way of finding p_0 and q_0 is often to write the equation in the form

$$y'' + \frac{p_0 + p_1 x + p_2 x^2 + \dots}{x} y' + \frac{q_0 + q_1 x + q_2 x^2 + \dots}{x^2} y = 0.$$
 (17)

Then inspection of the series that appear in the two numerators reveals the constants p_0 and q_0 .

Example 3 Find the exponents in the possible Frobenius series solutions of the equation

$$2x^{2}(1+x)y'' + 3x(1+x)^{3}y' - (1-x^{2})y = 0$$

Solution

We divide each term by
$$2x^2(1 + x)$$
 to recast the differential equation in the form

$$y'' + \frac{\frac{3}{2}(1+2x+x^2)}{x}y' + \frac{-\frac{1}{2}(1-x)}{x^2}y = 0$$

and thus see that $p_0 = \frac{3}{2}$ and $q_0 = -\frac{1}{2}$. Hence the indicial equation is

$$r(r-1) + \frac{3}{2}r - \frac{1}{2} = r^2 + \frac{1}{2}r - \frac{1}{2} = (r+1)(r-\frac{1}{2}) = 0$$

with roots $r_1 = \frac{1}{2}$ and $r_2 = -1$. The two possible Frobenius series solutions are then of the forms

$$y_1(x) = x^{1/2} \sum_{n=0}^{\infty} a_n x^n$$
 and $y_2(x) = x^{-1} \sum_{n=0}^{\infty} b_n x^n$.

Frobenius Series Solutions

>

3

Once the exponents r_1 and r_2 are known, the coefficients in a Frobenius series solution are determined by substitution of the series in Eqs. (12) through (14) in the differential equation, essentially the same method as was used to determine coefficients in power series solutions in Section 11.2. If the exponents r_1 and r_2 are complex conjugates, then there always exist two linearly independent Frobenius series solutions. We will restrict our attention here to the case in which r_1 and r_2 are both real. We also will seek solutions only for x > 0. Once such a solution has been found, we need only replace x^{r_1} with $|x|^{r_1}$ to obtain a solution for x < 0. The following theorem is proved in Chapter 4 of Coddington's *An Introduction to Ordinary Differential Equations*.

THEOREM 1 Frobenius Series Solutions

Suppose that x = 0 is a regular singular point of the equation

$$x^{2}y'' + xp(x)y' + q(x)y = 0.$$
 (10)

Let $\rho > 0$ denote the minimum of the radii of convergence of the power series

$$p(x) = \sum_{n=0}^{\infty} p_n x^n$$
 and $q(x) = \sum_{n=0}^{\infty} q_n x^n$.

Let r_1 and r_2 be the (real) roots, with $r_1 \ge r_2$, of the indicial equation $r(r-1) + p_0r + q_0 = 0$. Then

(a) For x > 0, there exists a solution of Eq. (10) of the form

$$y_1(x) = x^{r_1} \sum_{n=0}^{\infty} a_n x^n \qquad (a_0 \neq 0)$$
(18)

corresponding to the larger root r_1 .

(b) If $r_1 - r_2$ is neither zero nor a positive integer, then there exists a second linearly independent solution for x > 0 of the form

$$y_2(x) = x^{r_2} \sum_{n=0}^{\infty} b_n x^n \qquad (b_0 \neq 0)$$
 (19)

corresponding to the smaller root r_2 .

The radii of convergence of the power series in Eqs. (18) and (19) are each at least ρ . The coefficients in these series can be determined by substituting the series in the differential equation

$$x^{2}y'' + xp(x)y' + q(x)y = 0.$$

We have already seen that if $r_1 = r_2$, then there can exist only one Frobenius series solution. It turns out that, if $r_1 - r_2$ is a positive integer, then there may or may not exist a second Frobenius series solution of the form in Eq. (19) corresponding to the smaller root r_2 . Examples 4 through 6 illustrate the process of determining the coefficients in those Frobenius series solutions that are guaranteed by Theorem 1.
Example 4 Find the Frobenius series solutions of

$$2x^{2}y'' + 3xy' - (x^{2} + 1)y = 0.$$
(20)

Solution

on First we divide each term by
$$2x^2$$
 to put the equation in the form in (17):

$$y'' + \frac{\frac{3}{2}}{x}y' + \frac{-\frac{1}{2} - \frac{1}{2}x^2}{x^2}y = 0.$$
 (21)

We now see that x = 0 is a regular singular point, and that $p_0 = \frac{3}{2}$ and $q_0 = -\frac{1}{2}$. Because $p(x) \equiv \frac{3}{2}$ and $q(x) = -\frac{1}{2} - \frac{1}{2}x^2$ are polynomials, the Frobenius series we obtain will converge for all x > 0. The indicial equation is

$$r(r-1) + \frac{3}{2}r - \frac{1}{2} = \left(r - \frac{1}{2}\right)(r+1) = 0,$$

so the exponents are $r_1 = \frac{1}{2}$ and $r_2 = -1$. They do not differ by an integer, so Theorem 1 guarantees the existence of two linearly independent Frobenius series solutions. Rather than separately substituting

$$y_1 = x^{1/2} \sum_{n=0}^{\infty} a_n x^n$$
 and $y_2 = x^{-1} \sum_{n=0}^{\infty} b_n x^n$

in Eq. (20), it is more efficient to begin by substituting $y = x^r \sum c_n x^n$. We will then get a recurrence relation that depends on r. With the value $r_1 = \frac{1}{2}$ it becomes a recurrence relation for the series for y_1 , whereas with $r_2 = -1$ it becomes a recurrence relation for the series for y_2 .

When we substitute

$$y = \sum_{n=0}^{\infty} c_n x^{n+r}, \quad y' = \sum_{n=0}^{\infty} (n+r)c_n x^{n+r-1},$$

and

>

y'' =
$$\sum_{n=0}^{\infty} (n+r)(n+r-1)c_n x^{n+r-2}$$

in Eq. (20)—the original differential equation, rather than Eq. (21)—we get

$$2\sum_{n=0}^{\infty} (n+r)(n+r-1)c_n x^{n+r} + 3\sum_{n=0}^{\infty} (n+r)c_n x^{n+r} - \sum_{n=0}^{\infty} c_n x^{n+r+2} - \sum_{n=0}^{\infty} c_n x^{n+r} = 0.$$
 (22)

At this stage there are several ways to proceed. A good standard practice is to shift indices so that each exponent will be the same as the smallest one present. In this example, we shift the index of summation in the third sum by -2 to reduce its exponent from n + r + 2 to n + r. This gives

$$2\sum_{n=0}^{\infty} (n+r)(n+r-1)c_n x^{n+r} + 3\sum_{n=0}^{\infty} (n+r)c_n x^{n+r} - \sum_{n=2}^{\infty} c_{n-2} x^{n+r} - \sum_{n=0}^{\infty} c_n x^{n+r} = 0.$$
 (23)

The common range of summation is $n \ge 2$, so we must treat n = 0 and n = 1 separately. Following our standard practice, the terms corresponding to n = 0 will always give the indicial equation

$$[2r(r-1) + 3r - 1]c_0 = 2\left(r^2 + \frac{1}{2}r - \frac{1}{2}\right)c_0 = 0.$$

The terms corresponding to n = 1 yield

$$[2(r+1)r + 3(r+1) - 1]c_1 = (2r^2 + 5r + 2)c_1 = 0.$$

Because the coefficient $2r^2 + 5r + 2$ of c_1 is nonzero whether $r = \frac{1}{2}$ or r = -1, it follows that

$$c_1 = 0 \tag{24}$$

in either case.

>

>

≻

The coefficient of x^{n+r} in Eq. (23) is

$$2(n+r)(n+r-1)c_n + 3(n+r)c_n - c_{n-2} - c_n = 0.$$

We solve for c_n and simplify to obtain the recurrence relation

$$c_n = \frac{c_{n-2}}{2(n+r)^2 + (n+r) - 1} \quad \text{for } n \ge 2.$$
(25)

CASE 1: $r_1 = \frac{1}{2}$. We now write a_n in place of c_n and substitute $r = \frac{1}{2}$ in Eq. (25). This gives the recurrence relation

$$a_n = \frac{a_{n-2}}{2n^2 + 3n}$$
 for $n \ge 2$. (26)

With this formula we can determine the coefficients in the first Frobenius solution y_1 . In view of Eq. (24) we see that $a_n = 0$ whenever n is odd. With n = 2, 4, and 6 in Eq. (26), we get

$$a_2 = \frac{a_0}{14}, \quad a_4 = \frac{a_2}{44} = \frac{a_0}{616}, \quad \text{and} \quad a_6 = \frac{a_4}{90} = \frac{a_0}{55,440}.$$

Hence the first Frobenius solution is

$$y_1(x) = a_0 x^{1/2} \left(1 + \frac{x^2}{14} + \frac{x^4}{616} + \frac{x^6}{55,440} + \cdots \right).$$

CASE 2: $r_2 = -1$. We now write b_n in place of c_n and substitute r = -1 in Eq. (25). This gives the recurrence relation

$$b_n = \frac{b_{n-2}}{2b^2 - 3n}$$
 for $n \ge 2$. (27)

Again, Eq. (24) implies that $b_n = 0$ for *n* odd. With n = 2, 4, and 6 in (27), we get

$$b_2 = \frac{b_0}{2}, \quad b_4 = \frac{b_2}{20} = \frac{b_0}{40}, \text{ and } b_6 = \frac{b_4}{54} = \frac{b_0}{2160}.$$

Hence the second Frobenius solution is

$$y_2(x) = b_0 x^{-1} \left(1 + \frac{x^2}{2} + \frac{x^4}{40} + \frac{x^6}{2160} + \cdots \right).$$

Example 5 Find a Frobenius solution of Bessel's equation of order zero,

$$x^2y'' + xy' + x^2y = 0.$$
 (28)

Solution In t

In the form of (17), Eq. (28) becomes

$$y'' + \frac{1}{x}y' + \frac{x^2}{x^2}y = 0$$

Hence x = 0 is a regular singular point with $p(x) \equiv 1$ and $q(x) = x^2$, so our series will converge for all x > 0. Because $p_0 = 1$ and $q_0 = 0$, the indicial equation is

$$r(r-1) + r = r^2 = 0.$$

636 CHAPTER 11 POWER SERIES METHODS

Thus we obtain only the single exponent r = 0, and so there is only one Frobenius series solution

$$y(x) = x^0 \sum_{n=0}^{\infty} c_n x^n$$

of Eq. (28); it is in fact a power series.

Thus we substitute $y = \sum c_n x^n$ in (28); the result is

$$\sum_{n=0}^{\infty} n(n-1)c_n x^n + \sum_{n=0}^{\infty} nc_n x^n + \sum_{n=0}^{\infty} c_n x^{n+2} = 0.$$

We combine the first two sums and shift the index of summation in the third by -2 to obtain

$$\sum_{n=0}^{\infty} n^2 c_n x^n + \sum_{n=2}^{\infty} c_{n-2} x^n = 0.$$

The term corresponding to x^0 gives 0 = 0: no information. The term corresponding to x^1 gives $c_1 = 0$, and the term for x^n yields the recurrence relation

$$c_n = -\frac{c_{n-2}}{n^2} \qquad \text{for } n \ge 2.$$
⁽²⁹⁾

Because $c_1 = 0$, we see that $c_n = 0$ whenever *n* is odd. Substituting n = 2, 4, and 6 in Eq. (29), we get

$$c_2 = -\frac{c_0}{2^2}, \quad c_4 = -\frac{c_2}{4^2} = \frac{c_0}{2^2 \cdot 4^2}, \quad \text{and} \quad c_6 = -\frac{c_4}{6^2} = -\frac{c_0}{2^2 \cdot 4^2 \cdot 6^2}.$$

Evidently, the pattern is

$$c_{2n} = \frac{(-1)^n c_0}{2^2 \cdot 4^2 \cdots (2n)^2} = \frac{(-1)^n c_0}{2^{2n} (n!)^2}$$

The choice $c_0 = 1$ gives us one of the most important special functions in mathematics, the **Bessel function of order zero of the first kind**, denoted by $J_0(x)$. Thus

$$J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2} = 1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + \cdots .$$
 (30)

In this example we have not been able to find a second linearly independent solution of Bessel's equation of order zero. We will derive that solution in Section 11.4; it will not be a Frobenius series.

When $r_1 - r_2$ Is an Integer

Recall that, if $r_1 - r_2$ is a positive integer, then Theorem 1 guarantees only the existence of the Frobenius series solution corresponding to the larger exponent r_1 . Example 6 illustrates the fortunate case in which the series method nevertheless yields a second Frobenius series solution. An example in which the second solution is not a Frobenius series will be discussed in Section 11.4.

Example 6 Find the Frobenius series solutions of

$$xy'' + 2y' + xy = 0. (31)$$

Solution In standard form the equation becomes

$$y'' + \frac{2}{x}y' + \frac{x^2}{x^2}y = 0$$

so we see that x = 0 is a regular singular point with $p_0 = 2$ and $q_0 = 0$. The indicial equation

$$r(r-1) + 2r = r(r+1) = 0$$

has roots $r_1 = 0$ and $r_2 = -1$, which differ by an integer. In this case when $r_1 - r_2$ is an integer, it is better to depart from the standard procedure of Example 4 and begin our work with the *smaller* exponent. As you will see, the recurrence relation will then tell us whether or not a second Frobenius series solution exists. If it does exist, our computations will simultaneously yield *both* Frobenius series solutions. If the second solution does not exist, we begin anew with the larger exponent $r = r_1$ to obtain the one Frobenius series solution guaranteed by Theorem 1.

Hence we begin by substituting

$$y = x^{-1} \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_n x^{n-1}$$

in Eq. (31). This gives

$$\sum_{n=0}^{\infty} (n-1)(n-2)c_n x^{n-2} + 2\sum_{n=0}^{\infty} (n-1)c_n x^{n-2} + \sum_{n=0}^{\infty} c_n x^n = 0.$$

We combine the first two sums and shift the index by -2 in the third to obtain

$$\sum_{n=0}^{\infty} n(n-1)c_n x^{n-2} + \sum_{n=2}^{\infty} c_{n-2} x^{n-2} = 0.$$
(32)

The cases n = 0 and n = 1 reduce to

$$0 \cdot c_0 = 0$$
 and $0 \cdot c_1 = 0$.

Hence we have *two* arbitrary constants c_0 and c_1 and therefore can expect to find a general solution incorporating two linearly independent Frobenius series solutions. If, for n = 1, we had obtained an equation such as $0 \cdot c_1 = 3$, which can be satisfied for *no* choice of c_1 , this would have told us that no second Frobenius series solution could exist.

Now knowing that all is well, from (32) we read the recurrence relation

•
$$c_n = -\frac{c_{n-2}}{n(n-1)}$$
 for $n \ge 2$. (33)

The first few values of n give

$$c_{2} = -\frac{1}{2 \cdot 1}c_{0}, \qquad c_{3} = -\frac{1}{3 \cdot 2}c_{1},$$

$$c_{4} = -\frac{1}{4 \cdot 3}c_{2} = \frac{c_{0}}{4!}, \qquad c_{5} = -\frac{1}{5 \cdot 4}c_{3} = \frac{c_{1}}{5!},$$

$$c_{6} = -\frac{1}{6 \cdot 5}c_{4} = -\frac{c_{0}}{6!}, \qquad c_{7} = -\frac{1}{7 \cdot 6}c_{6} = -\frac{c_{1}}{7!};$$

evidently the pattern is

$$c_{2n} = \frac{(-1)^n c_0}{(2n)!}, \qquad c_{2n+1} = \frac{(-1)^n c_1}{(2n+1)!}$$

for $n \ge 1$. Therefore, a general solution of Eq. (31) is

$$y(x) = x^{-1} \sum_{n=0}^{\infty} c_n x^n$$

= $\frac{c_0}{x} \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots \right) + \frac{c_1}{x} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots \right)$
= $\frac{c_0}{x} \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} + \frac{c_1}{x} \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}.$

y y_1 y_2 2π 4π x

FIGURE 11.3.1. The solutions $y_1(x) = \frac{\cos x}{x}$ and $y_2(x) = \frac{\sin x}{x}$ in Example 6.

Thus

$$y(x) = \frac{1}{x}(c_0 \cos x + c_1 \sin x).$$

We have thus found a general solution expressed as a linear combination of the two Frobenius series solutions

$$y_1(x) = \frac{\cos x}{x}$$
 and $y_2(x) = \frac{\sin x}{x}$. (34)

As indicated in Fig. 11.3.1, one of these Frobenius series solutions is bounded but the other is unbounded near the regular singular point x = 0—a common occurrence in the case of exponents differing by an integer.

Summary

>

When confronted with a linear second-order differential equation

$$A(x)y'' + B(x)y' + C(x)y = 0$$

with analytic coefficient functions, in order to investigate the possible existence of series solutions we first write the equation in the standard form

$$y'' + P(x)y' + Q(x)y = 0.$$

If P(x) and Q(x) are both analytic at x = 0, then x = 0 is an ordinary point, and the equation has two linearly independent power series solutions.

Otherwise, x = 0 is a singular point, and we next write the differential equation in the form

>
$$y'' + \frac{p(x)}{x}y' + \frac{q(x)}{x^2}y = 0.$$

If p(x) and q(x) are both analytic at x = 0, then x = 0 is a regular singular point. In this case we find the two exponents r_1 and r_2 (assumed real, and with $r_1 \ge r_2$) by solving the indicial equation

 $\mathbf{>}$

$$r(r-1) + p_0 r + q_0 = 0,$$

where $p_0 = p(0)$ and $q_0 = q(0)$. There always exists a Frobenius series solution $y = x^{r_1} \sum a_n x^n$ associated with the larger exponent r_1 , and if $r_1 - r_2$ is not an integer, the existence of a second Frobenius series solution $y_2 = x^{r_2} \sum b_n x^n$ is also guaranteed.

11.3 **Problems**

In Problems 1 through 8, determine whether x = 0 is an ordinary point, a regular singular point, or an irregular singular point. If it is a regular singular point, find the exponents of the differential equation at x = 0.

- 1. $xy'' + (x x^3)y' + (\sin x)y = 0$ 2. $xy'' + x^2y' + (e^x - 1)y = 0$ 3. $x^2y'' + (\cos x)y' + xy = 0$ 4. $3x^3y'' + 2x^2y' + (1 - x^2)y = 0$
- **5.** x(1+x)y'' + 2y' + 3xy = 0
- 6. $x^2(1-x^2)y'' + 2xy' 2y = 0$

7. $x^2 y'' + (6 \sin x)y' + 6y = 0$ **8.** $(6x^2 + 2x^3)y'' + 21xy' + 9(x^2 - 1)y = 0$

If $x = a \neq 0$ is a singular point of a second-order linear differential equation, then the substitution t = x - a transforms it into a differential equation having t = 0 as a singular point. We then attribute to the original equation at x = a the behavior of the new equation at t = 0. Classify (as regular or irregular) the singular points of the differential equations in Problems 9 through 16. 9. $(1-x)y'' + xy' + x^2y = 0$ 10. $(1-x)^2y'' + (2x-2)y' + y = 0$ 11. $(1-x^2)y'' - 2xy' + 12y = 0$ 12. $(x-2)^3y'' + 3(x-2)^2y' + x^3y = 0$ 13. $(x^2-4)y'' + (x-2)y' + (x+2)y = 0$ 14. $(x^2-9)^2y'' + (x^2+9)y' + (x^2+4)y = 0$ 15. $(x-2)^2y'' - (x^2-4)y' + (x+2)y = 0$ 16. $x^3(1-x)y'' + (3x+2)y' + xy = 0$

Find two linearly independent Frobenius series solutions (for x > 0) of each of the differential equations in Problems 17 through 26.

17. 4xy'' + 2y' + y = 018. 2xy'' + 3y' - y = 019. 2xy'' - y' - y = 020. 3xy'' + 2y' + 2y = 021. $2x^2y'' + xy' - (1 + 2x^2)y = 0$ 22. $2x^2y'' + xy' - (3 - 2x^2)y = 0$ 23. $6x^2y'' + 7xy' - (x^2 + 2)y = 0$ 24. $3x^2y'' + 2xy' + x^2y = 0$ 25. 2xy'' + (1 + x)y' + y = 026. $2xy'' + (1 - 2x^2)y' - 4xy = 0$

Use the method of Example 6 to find two linearly independent Frobenius series solutions of the differential equations in Problems 27 through 31. Then construct a graph showing their graphs for x > 0.

27. xy'' + 2y' + 9xy = 0 **28.** xy'' + 2y' - 4xy = 0 **29.** 4xy'' + 8y' + xy = 0 **30.** $xy'' - y' + 4x^3y = 0$ **31.** $4x^2y'' - 4xy' + (3 - 4x^2)y = 0$

In Problems 32 through 34, find the first three nonzero terms of each of two linearly independent Frobenius series solutions.

- **32.** $2x^2y'' + x(x+1)y' (2x+1)y = 0$ **33.** $(2x^2 + 5x^3)y'' + (3x - x^2)y' - (1+x)y = 0$ **34.** $2x^2y'' + (\sin x)y' - (\cos x)y = 0$
- **35.** Note that x = 0 is an irregular point of the equation

$$x^{2}y'' + (3x - 1)y' + y = 0$$

(a) Show that $y = x^r \sum c_n x^n$ can satisfy this equation only if r = 0. (b) Substitute $y = \sum c_n x^n$ to derive the "formal" solution $y = \sum n! x^n$. What is the radius of convergence of this series?

- 36. (a) Suppose that A and B are nonzero constants. Show that the equation x²y" + Ay' + By = 0 has at most one solution of the form y = x^r ∑ c_nxⁿ. (b) Repeat part (a) with the equation x³y" + Axy' + By = 0. (c) Show that the equation x³y" + Ax²y' + By = 0 has no Frobenius series solution. (Suggestion: In each case substitute y = x^r ∑ c_nxⁿ in the given equation to determine the possible values of r.)
- **37.** (a) Use the method of Frobenius to derive the solution $y_1 = x$ of the equation $x^3y'' xy' + y = 0$. (b) Verify by substitution the second solution $y_2 = xe^{-1/x}$. Does y_2 have a Frobenius series representation?

38. Apply the method of Frobenius to Bessel's equation of order $\frac{1}{2}$,

$$x^{2}y'' + xy' + \left(x^{2} - \frac{1}{4}\right)y = 0,$$

to derive its general solution for x > 0,

$$y(x) = c_0 \frac{\cos x}{\sqrt{x}} + c_1 \frac{\sin x}{\sqrt{x}}$$

Figure 11.3.2 shows the graphs of the two indicated solutions.



FIGURE 11.3.2. The solutions

$$y_1(x) = \frac{\cos x}{\sqrt{x}}$$
 and $y_2(x) = \frac{\sin x}{\sqrt{x}}$ in

Problem 38.

39. (a) Show that Bessel's equation of order 1,

$$x^{2}y'' + xy' + (x^{2} - 1)y = 0,$$

has exponents $r_1 = 1$ and $r_2 = -1$ at x = 0, and that the Frobenius series corresponding to $r_1 = 1$ is

$$J_1(x) = \frac{x}{2} \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{n! (n+1)! 2^{2n}}.$$

(b) Show that there is no Frobenius solution corresponding to the smaller exponent $r_2 = -1$; that is, show that it is impossible to determine the coefficients in

$$y_2(x) = x^{-1} \sum_{n=0}^{\infty} c_n x^n$$

40. Consider the equation $x^2y'' + xy' + (1 - x)y = 0$. (a) Show that its exponents are $\pm i$, so it has complex-valued Frobenius series solutions

$$y_{+} = x^{i} \sum_{n=0}^{\infty} p_{n} x^{n}$$
 and $y_{-} = x^{-i} \sum_{n=0}^{\infty} q_{n} x^{n}$

with $p_0 = q_0 = 1$. (b) Show that the recursion formula is

$$c_n = \frac{c_{n-1}}{n^2 + 2rn}.$$

Apply this formula with r = i to obtain $p_n = c_n$, then with r = -i to obtain $q_n = c_n$. Conclude that p_n and q_n are complex conjugates: $p_n = a_n + ib_n$ and $q_n = a_n - ib_n$, where the numbers $\{a_n\}$ and $\{b_n\}$ are real. (c) Deduce

from part (b) that the differential equation given in this problem has real-valued solutions of the form

$$y_1(x) = A(x)\cos(\ln x) - B(x)\sin(\ln x),$$

$$y_2(x) = A(x)\sin(\ln x) + B(x)\cos(\ln x),$$

where $A(x) = \sum a_n x^n$ and $B(x) = \sum b_n x^n$. **41.** Consider the differential equation

$$x(x-1)(x+1)^{2}y'' + 2x(x-3)(x+1)y' - 2(x-1)y = 0$$

that appeared in an advertisement for a symbolic algebra program in the March 1984 issue of the *American Mathematical Monthly*. (a) Show that x = 0 is a regular singular point with exponents $r_1 = 1$ and $r_2 = 0$. (b) It follows from Theorem 1 that this differential equation has a power series solution of the form

$$y_1(x) = x + c_2 x^2 + c_3 x^3 + \cdots$$

Substitute this series (with $c_1 = 1$) in the differential equation to show that $c_2 = -2$, $c_3 = 3$, and

$$c_{n+2} = \frac{(n^2 - n)c_{n-1} + (n^2 - 5n - 2)c_n - (n^2 + 7n + 4)c_{n+1}}{(n+1)(n+2)}$$

for $n \ge 2$. (c) Use the recurrence relation in part (b) to prove by induction that $c_n = (-1)^{n+1}n$ for $n \ge 1$ (!). Hence deduce (using the geometric series) that

$$y_1(x) = \frac{x}{(1+x)^2}$$

for 0 < x < 1.

42. This problem is a brief introduction to Gauss's **hypergeometric equation**

$$x(1-x)y'' + [\gamma - (\alpha + \beta + 1)x]y' - \alpha\beta y = 0, \quad (35)$$

where α , β , and γ are constants. This famous equation has wide-ranging applications in mathematics and physics. (a) Show that x = 0 is a regular singular point of Eq. (35), with exponents 0 and $1 - \gamma$. (b) If γ is not zero or a negative integer, it follows (why?) that Eq. (35) has a power series solution

$$y(x) = x^0 \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_n x^n$$

with $c_0 \neq 0$. Show that the recurrence relation for this series is

$$c_{n+1} = \frac{(\alpha+n)(\beta+n)}{(\gamma+n)(1+n)}c_n$$

for $n \ge 0$. (c) Conclude that with $c_0 = 1$ the series in part (b) is

$$y(x) = 1 + \sum_{n=0}^{\infty} \frac{\alpha_n \beta_n}{n! \gamma_n} x^n$$
(36)

where $\alpha_n = \alpha(\alpha + 1)(\alpha + 2)\cdots(\alpha + n - 1)$ for $n \ge 1$, and β_n and γ_n are defined similarly. (d) The series in (36) is known as the **hypergeometric series** and is commonly denoted by $F(\alpha, \beta, \gamma, x)$. Show that

(i)
$$F(1, 1, 1, x) = \frac{1}{1-x}$$
 (the geometric series);
(ii) $xF(1, 1, 2, -x) = \ln(1+x)$;
(iii) $xF\left(\frac{1}{2}, 1, \frac{3}{2}, -x^2\right) = \tan^{-1}x$;
(iv) $F(-k, 1, 1, -x) = (1+x)^k$ (the binomial series).

11.3 Application Automating the Frobenius Series Method

Go to goo.gl/e4wzfl to download this application's computing resources including *Maple/Mathematica*/MATLAB. Here we illustrate the use of a computer algebra system such as *Maple* to apply the method of Frobenius. More complete versions of this application—illustrating the use of *Maple*, *Mathematica*, and MATLAB—can be downloaded from our Expanded Applications site using the URL indicated in the margin. We consider the differential equation

$$2x^{2}y'' + 3xy' - (x^{2} + 1)y = 0$$
⁽¹⁾

of Example 4 in this section, where we found the two indicial roots $r_1 = \frac{1}{2}$ and $r_2 = -1$.

Beginning with the indicial root $r_1 = \frac{1}{2}$, we first write the initial seven terms of a proposed Frobenius series solution:

a := array(0..6):
y :=
$$x^{(1/2)}$$
sum(a[n] $x^{(n)}$, n = 0..6);
y := $\sqrt{x} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6)$

Then we substitute this series (actually, partial sum) into the left-hand side of Eq. (1).

Evidently $x^{3/2}$ will factor out upon simplification, so we multiply by $x^{-3/2}$ and then collect coefficients of like powers of x.

deq2 := collect(
$$x^{(-3/2)}$$
*simplify(deq1), x);
 $deq2 := -x^7a_6 - x^6a_5 + (90a_6 - a_4)x^5 + (-a_3 + 65a_5)x^4$
 $+ (-a_2 + 44a_4)x^3 + (-a_1 + 27a_3)x^2 + (14a_2 - a_0)x + 5a_5$

We see here the equations that the successive coefficients must satisfy. We can select them automatically by defining an array, then filling the elements of this array by equating to zero (in turn) each of the coefficients in the series.

```
eqs := array(0..5):
for n from 0 to 5 do
eqs[n] := coeff(deq1,x,n) = 0: od:
coeffEqs := convert(eqs, set);
coeffEqs := \{5a_1 = 0, -a_2 + 44a_4 = 0, -a_3 + 65a_5 = 0, 90a_6 - a_4 = 0, 14a_2 - a_0 = 0, -a_1 + 27a_3 = 0\}
```

We now have a collection of six linear equations relating the seven coefficients $(a_0 \text{ through } a_6)$. Hence we can proceed to solve for the successive coefficients in terms of a_0 .

succCoeffs := convert([seq(a[n], n=1..6)], set); ourCoeffs := solve(coeffEqs, succCoeffs); $ourCoeffs := \left\{ a_1 = 0, \ a_6 = \frac{1}{55440} a_0, \ a_4 = \frac{1}{616} a_0, \ a_2 = \frac{1}{14} a_0, \ a_5 = 0, \ a_3 = 0 \right\}$

Thus we get the first particular solution

$$y_1(x) = a_0 x^{1/2} \left(1 + \frac{x^2}{14} + \frac{x^4}{616} + \frac{x^6}{55440} + \dots \right)$$

found in Example 4. You can now repeat this process, beginning with the indicial root $r_2 = -1$, to derive similarly the second particular solution.

In the following problems, use this method to derive Frobenius series solutions that can be checked against the given known general solutions.

1. $xy'' - y' + 4x^3y = 0$, $y(x) = A\cos x^2 + B\sin x^2$ 2. $xy'' - 2y' + 9x^5y = 0$, $y(x) = A\cos x^3 + B\sin x^3$ 3. 4xy'' - 2y' + y = 0, $y(x) = A\cos \sqrt{x} + B\sin \sqrt{x}$ 4. xy'' + 2y' + xy = 0, $y(x) = \frac{1}{x}(A\cos x + B\sin x)$ 5. 4xy'' + 6y' + y = 0, $y(x) = \frac{1}{\sqrt{x}}(A\cos \sqrt{x} + B\sin \sqrt{x})$ 6. $x^2y'' + xy' + (4x^4 - 1)y = 0$, $y(x) = \frac{1}{x}(A\cos x^2 + B\sin x^2)$ 7. $xy'' + 3y' + 4x^3y = 0$, $y(x) = \frac{1}{x^2}(A\cos x^2 + B\sin x^2)$ 8. $x^2y'' + x^2y' - 2y = 0$, $y(x) = \frac{1}{x}[A(2 - x) + B(2 + x)e^{-x}]$

Problems 9 through 11 involve the arctangent series

$$\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots$$

9.
$$(x + x^3)y'' + (2 + 4x^2)y' + 2xy = 0$$
, $y(x) = \frac{1}{x}(A + B\tan^{-1}x)$
10. $(2x + 2x^2)y'' + (3 + 5x)y' + y = 0$, $y(x) = \frac{1}{\sqrt{x}}(A + B\tan^{-1}\sqrt{x})$
11. $(x + x^5)y'' + (3 + 7x^4)y' + 8x^3y = 0$, $y(x) = \frac{1}{x^2}(A + B\tan^{-1}x^2)$

11.4 Bessel Functions

We have already seen several cases of Bessel's equation of order $p \ge 0$,

>
$$x^2y'' + xy' + (x^2 - p^2)y = 0.$$
 (1)

Its solutions are now called Bessel functions of order *p*. Such functions first appeared in the 1730s in the work of Daniel Bernoulli and Euler on the oscillations of a vertically suspended chain. The equation itself appears in a 1764 article by Euler on the vibrations of a circular drumhead, and Fourier used Bessel functions in his classical treatise on heat (1822). But their general properties were first studied systematically in an 1824 memoir by the German astronomer and mathematician Friedrich W. Bessel (1784–1846), who was investigating the motion of planets. The standard source of information on Bessel functions is G. N. Watson's *A Treatise on the Theory of Bessel Functions*, 2nd ed. (Cambridge: Cambridge University Press, 1944). Its 36 pages of references, which cover only the period up to 1922, give some idea of the vast literature of this subject.

Bessel's equation in (1) has indicial equation $r^2 - p^2 = 0$, with roots $r = \pm p$. If we substitute $y = \sum c_m x^{m+r}$ in Eq. (1), we find in the usual manner that $c_1 = 0$ and that

$$\left[(m+r)^2 - p^2\right]c_m + c_{m-2} = 0 \tag{2}$$

for $m \ge 2$. The verification of Eq. (2) is left to the reader (Problem 6).

The Case r = p > 0

If we use r = p and write a_m in place of c_m , then Eq. (2) yields the recursion formula

$$a_m = -\frac{a_{m-2}}{m(2p+m)}.$$
 (3)

Because $a_1 = 0$, it follows that $a_m = 0$ for all odd values of *m*. The first few even coefficients are

$$a_{2} = -\frac{a_{0}}{2(2p+2)} = -\frac{a_{0}}{2^{2}(p+1)},$$

$$a_{4} = -\frac{a_{2}}{4(2p+4)} = \frac{a_{0}}{2^{4} \cdot 2(p+1)(p+2)},$$

$$a_{6} = -\frac{a_{4}}{6(2p+6)} = -\frac{a_{0}}{2^{6} \cdot 2 \cdot 3(p+1)(p+2)(p+3)}.$$

The general pattern is

$$a_{2m} = \frac{(-1)^m a_0}{2^{2m} m! (p+1)(p+2) \cdots (p+m)}$$

so with the larger root r = p we get the solution

$$y_1(x) = a_0 \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m+p}}{2^{2m} m! (p+1)(p+2) \cdots (p+m)}.$$
(4)

If p = 0 this is the only Frobenius series solution; with $a_0 = 1$ as well, it is the function $J_0(x)$ we have seen before.

The Case r = -p < 0

>

If we use r = -p and write b_m in place of c_m , Eq. (2) takes the form

$$m(m-2p)b_m + b_{m-2} = 0 (5)$$

for $m \ge 2$, whereas $b_1 = 0$. We see that there is a potential difficulty if it happens that 2p is a positive integer—that is, if p is either a positive integer or an odd positive integral multiple of $\frac{1}{2}$. For then when m = 2p, Eq. (5) is simply $0 \cdot b_m + b_{m-2} = 0$. Thus if $b_{m-2} \ne 0$, then no value of b_m can satisfy this equation.

But if p is an odd positive integral multiple of $\frac{1}{2}$, we can circumvent this difficulty. For suppose that p = k/2, where k is an odd positive integer. Then we need only choose $b_m = 0$ for all odd values of m. The crucial step is the kth step,

$$k(k-k)b_k + b_{k-2} = 0;$$

and this equation will hold because $b_k = b_{k-2} = 0$.

Hence if p is not a positive integer, we take $b_m = 0$ for m odd and define the coefficients of even subscript in terms of b_0 by means of the recursion formula

$$b_m = -\frac{b_{m-2}}{m(m-2p)}, \qquad m \ge 2.$$
 (6)

In comparing (6) with (3), we see that (6) will lead to the same result as that in (4), except with p replaced with -p. Thus in this case we obtain the second solution

>
$$y_2(x) = b_0 \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m-p}}{2^{2m} m! (-p+1)(-p+2) \cdots (-p+m)}.$$
 (7)

The series in (4) and (7) converge for all x > 0 because x = 0 is the only singular point of Bessel's equation. If p > 0, then the leading term in y_1 is a_0x^p , whereas the leading term in y_2 is b_0x^{-p} . Hence $y_1(0) = 0$, but $y_2(x) \to \pm \infty$ as $x \to 0$, so it is clear that y_1 and y_2 are linearly independent solutions of Bessel's equation of order p > 0.

The Gamma Function

The formulas in (4) and (7) can be simplified by use of the **gamma function** $\Gamma(x)$, which (as in Section 10.1) is defined for x > 0 by

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} \, dt.$$
 (8)

It is not difficult to show that this improper integral converges for each x > 0. The gamma function is a generalization for x > 0 of the factorial function n!, which

644 CHAPTER 11 POWER SERIES METHODS

is defined only if *n* is a nonnegative integer. To see the way in which $\Gamma(x)$ is a generalization of *n*!, we note first that

$$\Gamma(1) = \int_0^\infty e^{-t} dt = \lim_{b \to \infty} \left[-e^{-t} \right]_0^b = 1.$$
(9)

Then we integrate by parts with $u = t^x$ and $dv = e^{-t} dt$:

$$\Gamma(x+1) = \lim_{b \to \infty} \int_0^b e^{-t} t^x dt = \lim_{b \to \infty} \left(\left[-e^{-t} t^x \right]_0^b + \int_0^b x e^{-t} t^{x-1} dt \right)$$
$$= x \left(\lim_{b \to \infty} \int_0^b e^{-t} t^{x-1} dt \right);$$

that is,

 $\mathbf{>}$

$$\Gamma(x+1) = x\Gamma(x). \tag{10}$$

This is the most important property of the gamma function.

If we combine Eqs. (9) and (10), we see that

$$\Gamma(2) = 1 \cdot \Gamma(1) = 1!, \quad \Gamma(3) = 2 \cdot \Gamma(2) = 2!, \quad \Gamma(4) = 3 \cdot \Gamma(3) = 3!,$$

and in general that

$$\Gamma(n+1) = n!$$
 for $n \ge 0$ an integer. (11)

An important special value of the gamma function is

$$\Gamma\left(\frac{1}{2}\right) = \int_0^\infty e^{-t} t^{-1/2} \, dt = 2 \int_0^\infty e^{-u^2} \, du = \sqrt{\pi},\tag{12}$$

where we have substituted u^2 for t in the first integral; the fact that

$$\int_0^\infty e^{-u^2} \, du = \frac{\sqrt{\pi}}{2}$$

is known, but is far from obvious. (See, for instance, Example 5 in Section 13.4 of Edwards and Penney, *Calculus: Early Transcendentals*, 7th edition, Hoboken, NJ: Pearson, 2008.)

Although $\Gamma(x)$ is defined in (8) only for x > 0, we can use the recursion formula in (10) to define $\Gamma(x)$ whenever x is neither zero nor a negative integer. If -1 < x < 0, then

$$\Gamma(x) = \frac{\Gamma(x+1)}{x};$$

the right-hand side is defined because 0 < x + 1 < 1. The same formula may then be used to extend the definition of $\Gamma(x)$ to the open interval (-2, -1), then to the open interval (-3, -2), and so on. The graph of the gamma function thus extended is shown in Fig. 11.4.1. The student who would like to pursue this fascinating topic further should consult Artin's *The Gamma Function* (New York: Holt, Rinehart and Winston, 1964). In only 39 pages, this is one of the finest expositions in the entire literature of mathematics.



FIGURE 11.4.1. The graph of the extended gamma function.

Bessel Functions of the First Kind

If we choose $a_0 = 1/[2^p \Gamma(p+1)]$ in (4), where p > 0, and note that

$$\Gamma(p+m+1) = (p+m)(p+m-1)\cdots(p+2)(p+1)\Gamma(p+1)$$

by repeated application of Eq. (10), we can write the **Bessel function of the first** kind of order p very concisely with the aid of the gamma function:

$$J_p(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(p+m+1)} \left(\frac{x}{2}\right)^{2m+p}.$$
 (13)

Similarly, if p > 0 is not an integer, we choose $b_0 = 1/[2^{-p}\Gamma(-p+1)]$ in (7) to obtain the linearly independent second solution

>
$$J_{-p}(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \, \Gamma(-p+m+1)} \left(\frac{x}{2}\right)^{2m-p}$$
 (14)

of Bessel's equation of order p. If p is not an integer, we have the general solution

$$y(x) = c_1 J_p(x) + c_2 J_{-p}(x)$$
(15)

for x > 0; x^p must be replaced with $|x|^p$ in Eqs. (13) through (15) to get the correct solutions for x < 0.

If p = n, a nonnegative integer, then Eq. (13) gives

$$J_n(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! (m+n)!} \left(\frac{x}{2}\right)^{2m+n}$$
(16)

for the Bessel functions of the first kind of integral order. Thus

$$J_0(x) = \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m}}{2^{2m} (m!)^2} = 1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots$$
(17)

and

$$J_1(x) = \sum_{m=0}^{\infty} \frac{(-1)^m 2^{2m+1}}{2^{2m+1} m! (m+1)!} = \frac{x}{2} - \frac{1}{2!} \left(\frac{x}{2}\right)^3 + \frac{1}{2! \cdot 3!} \left(\frac{x}{2}\right)^5 - \dots$$
(18)

The graphs of $J_0(x)$ and $J_1(x)$ are shown in Fig. 11.4.2. In a general way they resemble damped cosine and sine oscillations, respectively. Indeed, if you examine the series in (17), you can see part of the reason why $J_0(x)$ and $\cos x$ might be similar—only minor changes in the denominators in (17) are needed to produce the Taylor series for $\cos x$. As suggested by Fig. 11.4.2, the zeros of the functions $J_0(x)$ and $J_1(x)$ are *interlaced*—between any two consecutive zeros of $J_0(x)$ there is precisely one zero of $J_1(x)$, and vice versa. The first four zeros of $J_0(x)$ are approximately 2.4048, 5.5201, 8.6537, and 11.7915. For n large, the nth zero of $J_0(x)$ is approximately $(n - \frac{1}{4}) \pi$; the nth zero of $J_1(x)$ is approximately $(n + \frac{1}{4}) \pi$. Thus the interval between consecutive zeros of either $J_0(x)$ or $J_1(x)$ is approximately π another similarity with $\cos x$ and $\sin x$. You can see the way the accuracy of these approximations increases with increasing n by rounding the entries in the table in Fig. 11.4.3 to two decimal places.

It turns out that $J_p(x)$ is an elementary function if the order p is half an odd integer. For instance, on substitution of $p = \frac{1}{2}$ in Eqs. (13) and (14), respectively, the results can be recognized (Problem 2) as

$$J_{1/2}(x) = \sqrt{\frac{2}{\pi x}} \sin x$$
 and $J_{-1/2}(x) = \sqrt{\frac{2}{\pi x}} \cos x.$ (19)



n	<i>n</i> th zero of $J_0(x)$	$\left(n-\frac{1}{4}\right)\pi$	<i>n</i> th zero of $J_1(x)$	$\left(n+\frac{1}{4}\right)\pi$
1	2.4048	2.3562	3.8317	3.9270
2	5.5201	5.4978	7.0156	7.0686
3	8.6537	8.6394	10.1735	10.2102
4	11.7915	11.7810	13.3237	13.3518
5	14.9309	14.9226	16.4706	16.4934

FIGURE 11.4.2. The graphs of the Bessel functions $J_0(x)$ and $J_1(x)$.

FIGURE 11.4.3. Zeros of $J_0(x)$ and $J_1(x)$.

Bessel Functions of the Second Kind

If *n* is not an integer, then Bessel's equation of order *n* has *no* second Frobenius series solution independent of $J_n(x)$. For Bessel's equation of order 0, Example 4 in Section 8.4 of Edwards and Penney, *Differential Equations and Boundary Value Problems: Computing and Modeling* (5th edition, Hoboken, NJ: Pearson, 2014) gives the second solution

$$Y_0(x) = \frac{2}{\pi} \left[\left(\gamma + \ln \frac{x}{2} \right) J_0(x) + \sum_{m=1}^{\infty} \frac{(-1)^{m+1} H_m}{(m!)^2} \left(\frac{x}{2} \right)^{2m} \right],$$
(20)

where

$$H_m = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{m}$$

denotes the *m*th partial sum of the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ and
 $\gamma = \lim_{n \to \infty} (H_n - \ln n) \approx 0.57722$

is *Euler's constant*. The logarithmic term in (20) implies that this **Bessel function** $Y_0(x)$ of the second kind is not a Frobenius series, and it is typical of the case in which a second Frobenius series solution does not exist. It also implies that $Y_0(x) \rightarrow -\infty$ as $x \rightarrow 0$ (Fig. 11.4.4), so $Y_0(x)$ is not continuous at x = 0. These properties are shared by the general **Bessel function** $Y_n(x)$ of the second kind (with *n* a positive integer), which is defined by a complicated generalization of the formula in (20).

A general solution of Bessel's equation of integral order *n* is given by

$$y(x) = c_1 J_n(x) + c_2 Y_n(x).$$
(21)

But if y(x) is continuous at x = 0, the fact that $Y_n(x) \to -\infty$ as $x \to 0$ implies that $c_2 = 0$. It follows that any *continuous* solution of Bessel's equation of integral order *n* must be a constant multiple of the Bessel function $J_n(x)$ of the first kind. Numerous physical applications of this fact—to heat flow in circular plates or cylinders and to vibrations of circular membranes, for instance—are discussed in Section 10.4 of the reference cited previously.

Figure 11.4.5 illustrates the fact that for n > 1 the graphs of $J_n(x)$ and $Y_n(x)$ look generally like those of $J_1(x)$ and $Y_1(x)$. In particular, $J_n(0) = 0$ while $Y_n(x) \rightarrow -\infty$ as $x \rightarrow 0^+$, and both functions undergo damped oscillation as $x \rightarrow +\infty$.





FIGURE 11.4.4. The graphs of the Bessel functions $Y_0(x)$ and $Y_1(x)$ of the second kind.

FIGURE 11.4.5. The graphs of the Bessel functions $J_2(x)$ and $Y_2(x)$.

Bessel Function Identities

Bessel functions are analogous to trigonometric functions in that they satisfy a large number of standard identities of frequent utility, especially in the evaluation of integrals involving Bessel functions. Differentiation of

$$J_p(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \, \Gamma(p+m+1)} \left(\frac{x}{2}\right)^{2m+p} \tag{13}$$

in the case that p is a nonnegative integer gives

$$\frac{d}{dx} \left[x^p J_p(x) \right] = \frac{d}{dx} \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m+2p}}{2^{2m+p} m! (p+m)!}$$
$$= \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m+2p-1}}{2^{2m+p-1} m! (p+m-1)!}$$
$$= x^p \sum_{m=0}^{\infty} \frac{(-1)^m x^{2m+p-1}}{2^{2m+p-1} m! (p+m-1)!},$$

and thus we have shown that

$$\frac{d}{dx}\left[x^p J_p(x)\right] = x^p J_{p-1}(x).$$
(22)

Similarly,

$$\frac{d}{dx} \left[x^{-p} J_p(x) \right] = -x^{-p} J_{p+1}(x).$$
(23)

If we carry out the differentiations in Eqs. (22) and (23) and then divide the resulting identities by x^p and x^{-p} , respectively, we obtain (Problem 8) the identities

$$J'_{p}(x) = J_{p-1}(x) - \frac{p}{x}J_{p}(x)$$
(24)

and

$$J'_{p}(x) = \frac{p}{x}J_{p}(x) - J_{p+1}(x).$$
(25)

Thus we may express the derivatives of Bessel functions in terms of Bessel functions themselves. Subtraction of Eq. (25) from Eq. (24) gives the recursion formula

$$J_{p+1}(x) = \frac{2p}{x} J_p(x) - J_{p-1}(x),$$
(26)

which can be used to express Bessel functions of higher order in terms of Bessel functions of lower orders. In the form

$$J_{p-1}(x) = \frac{2p}{x} J_p(x) - J_{p+1}(x),$$
(27)

it can be used to express Bessel functions of large negative order in terms of Bessel functions of numerically smaller negative orders.

The identities in Eqs. (22) through (27) hold wherever they are meaningful that is, whenever no Bessel functions of negative integral order appear. In particular, they hold for all nonintegral values of p.

Example 1 With p = 0, E

/ith
$$p = 0$$
, Eq. (22) gives

$$\int x J_0(x) \, dx = x J_1(x) + C.$$

Similarly, with p = 0, Eq. (23) gives

$$\int J_1(x) \, dx = -J_0(x) + C.$$

Example 2

Using first p = 2 and then p = 1 in Eq. (26), we get

$$J_3(x) = \frac{4}{x}J_2(x) - J_1(x) = \frac{4}{x}\left[\frac{2}{x}J_1(x) - J_0(x)\right] - J_1(x),$$

so that

$$J_3(x) = -\frac{4}{x}J_0(x) + \left(\frac{8}{x^2} - 1\right)J_1(x).$$

With similar manipulations every Bessel function of positive integral order can be expressed in terms of $J_0(x)$ and $J_1(x)$.

Example 3

To antidifferentiate $xJ_2(x)$, we first note that

$$\int x^{-1} J_2(x) \, dx = -x^{-1} J_1(x) + C$$

by Eq. (23) with p = 1. We therefore write

$$\int x J_2(x) \, dx = \int x^2 \left[x^{-1} J_2(x) \right] \, dx$$

and integrate by parts with

$$u = x^2$$
, $dv = x^{-1}J_2(x) dx$,
 $du = 2x dx$, and $v = -x^{-1}J_1(x)$.

This gives

$$\int x J_2(x) \, dx = -x J_1(x) + 2 \int J_1(x) \, dx = -x J_1(x) - 2J_0(x) + C,$$

with the aid of the second result of Example 1.

Applications of Bessel Functions

The importance of Bessel functions stems not only from the frequent appearance of Bessel's equation in applications, but also from the fact that the solutions of many other second-order linear differential equations can be expressed in terms of Bessel functions. To see how this comes about, we begin with Bessel's equation of order p in the form

$$z^{2}\frac{d^{2}w}{dz^{2}} + z\frac{dw}{dz} + (z^{2} - p^{2})w = 0$$
(28)

and substitute

$$w = x^{-\alpha}y, \qquad z = kx^{\beta}.$$
 (29)

Then a routine (but lengthy) transformation of Eq. (28) yields

$$x^{2}y'' + (1 - 2\alpha)xy' + (\alpha^{2} - \beta^{2}p^{2} + \beta^{2}k^{2}x^{2\beta})y = 0;$$

that is,

$$x^{2}y'' + Axy' + (B + Cx^{q})y = 0,$$
(30)

where the constants A, B, C, and q are given by

$$A = 1 - 2\alpha, \quad B = \alpha^2 - \beta^2 p^2, \quad C = \beta^2 k^2, \text{ and } q = 2\beta.$$
 (31)

It is a simple matter to solve the equations in (31) for

$$\alpha = \frac{1-A}{2}, \qquad \beta = \frac{q}{2},$$

$$k = \frac{2\sqrt{C}}{q}, \qquad \text{and} \qquad p = \frac{\sqrt{(1-A)^2 - 4B}}{q}.$$
(32)

Under the assumption that the square roots in (32) are real, it follows that the general solution of Eq. (30) is

$$y(x) = x^{\alpha}w(z) = x^{\alpha}w(kx^{\beta}),$$

where

$$w(z) = c_1 J_p(z) + c_2 J_{-p}(z)$$

(assuming that p is not an integer) is the general solution of the Bessel equation in (28). This establishes the following result.

THEOREM 1 Solutions in Bessel Functions

If C > 0, $q \neq 0$, and $(1 - A)^2 \ge 4B$, then the general solution (for x > 0) of Eq. (30) is

$$y(x) = x^{\alpha} \left[c_1 J_p(kx^{\beta}) + c_2 J_{-p}(kx^{\beta}) \right],$$
 (33)

where α , β , k, and p are given by the equations in (32). If p is an integer, then J_{-p} is to be replaced with Y_p .

$$4x^{2}y'' + 8xy' + (x^{4} - 3)y = 0.$$
(34)

Solution To compare Eq. (34) with Eq. (30), we rewrite the former as

$$x^{2}y'' + 2xy' + \left(-\frac{3}{4} + \frac{1}{4}x^{4}\right)y = 0$$

and see that A = 2, $B = -\frac{3}{4}$, $C = \frac{1}{4}$, and q = 4. Then the equations in (32) give $\alpha = -\frac{1}{2}$, $\beta = 2$, $k = \frac{1}{4}$, and $p = \frac{1}{2}$. Thus the general solution in (33) of Eq. (34) is

$$y(x) = x^{-1/2} \left[c_1 J_{1/2} \left(\frac{1}{4} x^2 \right) + c_2 J_{-1/2} \left(\frac{1}{4} x^2 \right) \right]$$

If we recall from Eq. (19) that

$$J_{1/2}(z) = \sqrt{\frac{2}{\pi z}} \sin z$$
 and $J_{-1/2}(z) = \sqrt{\frac{2}{\pi z}} \cos z$,

we see that a general solution of Eq. (34) can be written in the elementary form

$$y(x) = x^{-3/2} \left(A \cos \frac{x^2}{4} + B \sin \frac{x^2}{4} \right).$$

Example 5 Solve the Airy equation

y'' + 9xy = 0. (35)

Solution First we rewrite the given equation in the form

$$x^2 y'' + 9x^3 y = 0.$$

This is the special case of Eq. (30) with A = B = 0, C = 9, and q = 3. It follows from the equations in (32) that $\alpha = \frac{1}{2}$, $\beta = \frac{3}{2}$, k = 2, and $p = \frac{1}{3}$. Thus the general solution of Eq. (35) is

$$y(x) = x^{1/2} \left[c_1 J_{1/3}(2x^{3/2}) + c_2 J_{-1/3}(2x^{3/2}) \right].$$

11.4 Problems

- 1. Differentiate termwise the series for $J_0(x)$ to show directly that $J'_0(x) = -J_1(x)$ (another analogy with the cosine and sine functions).
- **2.** (a) Deduce from Eqs. (10) and (12) that

$$\Gamma\left(n+\frac{1}{2}\right) = \frac{1\cdot 3\cdot 5\cdot (2n-1)}{2^n}\sqrt{\pi}.$$

- (b) Use the result of part (a) to verify the formulas in Eq. (19) for $J_{1/2}(x)$ and $J_{-1/2}(x)$.
- 3. (a) Suppose that m is a positive integer. Show that

$$\Gamma\left(m+\frac{2}{3}\right) = \frac{2\cdot 5\cdot 8\cdots (3m-1)}{3^m}\Gamma\left(\frac{2}{3}\right)$$

(b) Conclude from part (a) and Eq. (13) that

$$J_{-1/3}(x) = \frac{(x/2)^{-1/3}}{\Gamma\left(\frac{2}{3}\right)} \left(1 + \sum_{m=1}^{\infty} \frac{(-1)^m 3^m x^{2m}}{2^{2m} m! \cdot 2 \cdot 5 \cdots (3m-1)}\right)$$

4. Apply Eqs. (19), (26), and (27) to show that

$$J_{3/2}(x) = \sqrt{\frac{2}{\pi x^3}} (\sin x - x \cos x)$$

and

$$J_{-3/2}(x) = -\sqrt{\frac{2}{\pi x^3}} (\cos x + x \sin x).$$

- 5. Express $J_4(x)$ in terms of $J_0(x)$ and $J_1(x)$.
- **6.** Derive the recursion formula in Eq. (2) for Bessel's equation.
- 7. Verify the identity in (23) by termwise differentiation.
- **8.** Deduce the identities in Eqs. (24) and (25) from those in Eqs. (22) and (23).
- 9. Use the relation $\Gamma(x + 1) = x\Gamma(x)$ to deduce from Eqs. (13) and (14) that if p is not a negative integer, then

$$J_p(x) = \frac{(x/2)^p}{\Gamma(p+1)} \left[1 + \sum_{m=1}^{\infty} \frac{(-1)^m (x/2)^{2m}}{m! (p+1)(p+2) \cdots (p+m)} \right].$$

This form is more convenient for the computation of $J_p(x)$ because only the single value $\Gamma(p + 1)$ of the gamma function is required.

10. Use the series of Problem 9 to find $y(0) = \lim_{x \to 0} y(x)$ if

$$y(x) = x^{2} \left[\frac{J_{5/2}(x) + J_{-5/2}(x)}{J_{1/2}(x) + J_{-1/2}(x)} \right]$$

Any integral of the form $\int x^m J_n(x) dx$ can be evaluated in terms of Bessel functions and the indefinite integral $\int J_0(x) dx$. The latter integral cannot be simplified further, but the function $\int_0^x J_0(t) dt$ is tabulated in Table 11.1 of Abramowitz and Stegun. Use the identities in Eqs. (22) and (23) to evaluate the integrals in Problems 11 through 18.

11.
$$\int x^2 J_0(x) dx$$
 12. $\int x^3 J_0(x) dx$

 13. $\int x^4 J_0(x) dx$
 14. $\int x J_1(x) dx$

 15. $\int x^2 J_1(x) dx$
 16. $\int x^3 J_1(x) dx$

 17. $\int x^4 J_1(x) dx$
 18. $\int J_2(x) dx$

In Problems 19 through 30, express the general solution of the given differential equation in terms of Bessel functions.

19.
$$x^{2}y'' - xy' + (1 + x^{2})y = 0$$

20. $xy'' + 3y' + xy = 0$
21. $xy'' - y' + 36x^{3}y = 0$
22. $x^{2}y'' - 5xy' + (8 + x)y = 0$
23. $36x^{2}y'' + 60xy' + (9x^{3} - 5)y = 0$
24. $16x^{2}y'' + 24xy' + (1 + 144x^{3})y = 0$
25. $x^{2}y'' + 3xy' + (1 + x^{2})y = 0$
26. $4x^{2}y'' - 12xy' + (15 + 16x)y = 0$
27. $16x^{2}y'' - (5 - 144x^{3})y = 0$
28. $2x^{2}y'' - 3xy' - 2(14 - x^{5})y = 0$
29. $y'' + x^{4}y = 0$
30. $y'' + 4x^{3}y = 0$

31. Apply Theorem 1 to show that the general solution of

$$xy'' + 2y' + xy = 0$$

is
$$y(x) = x^{-1}(A\cos x + B\sin x)$$
.

- **32.** Verify that the substitutions in (2) in Bessel's equation [Eq. (1)] yield Eq. (3).
- 33. (a) Show that the substitution

$$y = -\frac{1}{u}\frac{du}{dx}$$

transforms the Riccati equation $dy/dx = x^2 + y^2$ into $u'' + x^2u = 0$. (b) Show that the general solution of $dy/dx = x^2 + y^2$ is

$$y(x) = x \frac{J_{3/4}\left(\frac{1}{2}x^2\right) - cJ_{-3/4}\left(\frac{1}{2}x^2\right)}{cJ_{1/4}\left(\frac{1}{2}x^2\right) + J_{-1/4}\left(\frac{1}{2}x^2\right)},$$

Suggestion: Apply the identities in Eqs. (22) and (23).

34. (a) Substitute the series of Problem 9 in the result of Problem 33 to show that the solution of the initial value problem $\frac{dy}{dx} = 2 - 2$

is

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0) = 0$$

$$y(x) = x \frac{J_{3/4}\left(\frac{1}{2}x^2\right)}{J_{-1/4}\left(\frac{1}{2}x^2\right)}$$

(b) Deduce similarly that the solution of the initial value problem

$$\frac{dy}{dx} = x^2 + y^2, \quad y(0) = 1$$

is

$$y(x) = x \frac{2\Gamma\left(\frac{3}{4}\right) J_{3/4}\left(\frac{1}{2}x^{2}\right) + \Gamma\left(\frac{1}{4}\right) J_{-3/4}\left(\frac{1}{2}x^{2}\right)}{2\Gamma\left(\frac{3}{4}\right) J_{-1/4}\left(\frac{1}{2}x^{2}\right) - \Gamma\left(\frac{1}{4}\right) J_{1/4}\left(\frac{1}{2}x^{2}\right)}.$$

Some solution curves of the equation $dy/dx = x^2 + y^2$ are shown in Fig. 11.4.6. The location of the asymptotes where $y(x) \rightarrow +\infty$ can be found by using Newton's method to find the zeros of the denominators in the formulas for the solutions as listed here.



REFERENCES FOR FURTHER STUDY

The literature of the theory and applications of differential equations is vast. The following list includes a selection of books that might be useful to readers who wish to pursue further the topics introduced in this book.

- 1. ABRAMOWITZ, M. and I. A. STEGUN, *Handbook of Mathematical Functions*. New York: Dover, 1965. The comprehensive collection of tables to which frequent reference is made in the text.
- 2. BIRKHOFF, G. and G.-C. ROTA, *Ordinary Differential Equations* (4th ed.). New York: John Wiley, 1989. An intermediate-level text that includes a more complete treatment of existence and uniqueness theorems, Sturm-Liouville problems, and eigenfunction expansions.
- 3. BRAUN, M., *Differential Equations and Their Applications* (3rd ed.). New York: Springer-Verlag, 1983. An introductory text at a slightly higher level than this book; it includes several interesting "case study" applications.
- 4. CHURCHILL, R. V., *Operational Mathematics* (3rd ed.). New York: McGraw-Hill, 1972. The standard reference for theory and applications of Laplace transforms, starting at about the same level as Chapter 10 of this book.
- CODDINGTON, E. A., An Introduction to Ordinary Differential Equations. Hoboken, NJ: Pearson, 1961. An intermediate-level introduction; Chapters 3 and 4 include proofs of the theorems on power series and Frobenius series solutions stated in Chapter 11 of this book.
- CODDINGTON, E. A. and N. LEVINSON, *Theory of Ordinary Differential Equations*. New York: McGraw-Hill, 1955. An advanced theoretical text; Chapter 5 discusses solutions near an irregular singular point.
- 7. DORMAND, J. R., *Numerical Methods for Differential Equations*. Boca Raton: CRC Press, 1996. More complete coverage of modern computational methods for approximate solution of differential equations.
- 8. HUBBARD, J. H. and B. H. WEST, *Differential Equations: A Dynamical Systems Approach*. New York: Springer-Verlag, 1992 (part I) and 1995 (Higher-Dimensional Systems). Detailed treatment of qualitative phenomena, with a balanced combination of computational and theoretical viewpoints.
- 9. INCE, E. L., *Ordinary Differential Equations*. New York: Dover, 1956. First published in 1926, this is the classic older reference work on the subject.
- 10. MCLACHLAN, N. W., Ordinary Non-Linear Differential Equations in Engineering and Physical Sciences. London: Oxford University Press, 1956. A concrete introduction to the effects of nonlinearity in physical systems.

- 11. NOBLE, B. and J.W. DANIEL, *Applied Linear Algebra* (3rd ed.). Hoboken, NJ: Pearson, 1988. An especially concrete introduction to linear algebra, with significant applications included.
- 12. POLKING, J. C. and D. ARNOLD, *Ordinary Differential Equations Using MAT-LAB* (3rd ed.). Hoboken, NJ: Pearson, 2003. A manual for using MATLAB in an elementary differential equations course; based on the MATLAB programs **dfield** and **pplane** that are used and referenced in this text.
- 13. PRESS, W. H., B. P. FLANNERY, S. A. TEUKOLSKY, and W. T. VETTERLING, *Numerical Recipes: The Art of Scientific Computing*, 3rd ed. Cambridge: Cambridge University Press, 2007. Chapter 17 discusses modern techniques for the numerical solution of ordinary differential equations. Programs in C⁺⁺, as well as C, FORTRAN, Pascal, and other languages can be downloaded from the accompanying web site numerical.recipes.
- 14. SIMMONS, G. F., *Differential Equations* (2nd ed.). New York: McGraw-Hill, 1991. An introductory text with interesting historical notes and fascinating applications and with the most eloquent preface in any mathematics book currently in print.
- 15. STRANG, W. G., *Linear Algebra and Its Applications*. (4th ed.). New York: Brooks Cole, 2006. An introductory treatment of linear algebra with motivating applications. Appendix B contains a concrete derivation of the Jordan normal form for square matrices.

APPENDIX A

Existence and Uniqueness of Solutions

In Chapter 1 we saw that an initial value problem of the form

$$\frac{dy}{dx} = f(x, y), \quad y(a) = b \tag{1}$$

can fail (on a given interval containing the point x = a) to have a unique solution. For instance, in Example 4 of Section 1.3, we saw that the initial value problem

$$x^{2}\frac{dy}{dx} + y^{2} = 0, \quad y(0) = b$$
⁽²⁾

has no solutions at all unless b = 0, in which case there are infinitely many solutions. According to Problem 31 of Section 1.3, the initial value problem

$$\frac{dy}{dx} = -\sqrt{1 - y^2}, \quad y(0) = 1 \tag{3}$$

has the two distinct solutions $y_1(x) \equiv 1$ and $y_2(x) = \cos x$ on the interval $0 \leq x \leq \pi$. In this appendix we investigate conditions on the function f(x, y) that suffice to guarantee that the initial value problem in (1) has one and only one solution, and we then proceed to establish appropriate versions of the existence-uniqueness theorems that were stated without proof in Sections 1.3, 5.1, 5.2, and 7.1.

A.1 Existence of Solutions

The approach we employ is the **method of successive approximations**, which was developed by the French mathematician Emile Picard (1856–1941). This method is based on the fact that the function y(x) satisfies the initial value problem in (1) on the open interval *I* containing x = a if and only if it satisfies the integral equation

$$y(x) = b + \int_{a}^{x} f(t, y(t)) dt$$
 (4)

for all x in I. In particular, if y(x) satisfies Eq. (4), then clearly y(a) = b, and differentiation of both sides in (4)—using the fundamental theorem of calculus—yields the differential equation y'(x) = f(x, y(x)).

To attempt to solve Eq. (4), we begin with the initial function

$$y_0(x) \equiv b \tag{5}$$

and then define iteratively a sequence y_1, y_2, y_3, \ldots of functions that we hope will converge to the solution. Specifically, we let

$$y_1(x) = b + \int_a^x f(t, y_0(t)) dt$$
 and $y_2(x) = b + \int_a^x f(t, y_1(t)) dt$. (6)

In general, y_{n+1} is obtained by substitution of y_n for y in the right-hand side in Eq. (4):

$$y_{n+1}(x) = b + \int_{a}^{x} f(t, y_n(t)) dt.$$
 (7)

Suppose we know that each of these functions $\{y_n(x)\}_0^\infty$ is defined on some open interval (the same for each *n*) containing x = a and that the limit

$$y(x) = \lim_{n \to \infty} y_n(x) \tag{8}$$

exists at each point of this interval. Then it will follow that

$$y(x) = \lim_{n \to \infty} y_{n+1}(x) = \lim_{n \to \infty} \left[b + \int_a^x f(t, y_n(t)) dt \right]$$
$$= b + \lim_{n \to \infty} \int_a^x f(t, y_n(t)) dt$$
(9)

$$= b + \int_{a}^{x} f\left(t, \lim_{n \to \infty} y_n(t)\right) dt \tag{10}$$

and, hence, that

$$y(x) = b + \int_a^x f(t, y(t)) dt,$$

provided that we can validate the interchange of limit operations involved in passing from (9) to (10). It is therefore reasonable to expect that, under favorable conditions, the sequence $\{y_n(x)\}$ defined iteratively in Eqs. (5) and (7) will converge to a solution y(x) of the integral equation in (4), and hence to a solution of the original initial value problem in (1).

Example 1 To apply the method of successive approximations to the initial value problem

$$\frac{dy}{dx} = y, \quad y(0) = 1, \tag{11}$$

we write Eqs. (5) and (7), thereby obtaining

$$y_0(x) \equiv 1, \quad y_{n+1}(x) = 1 + \int_0^x y_n(t) dt.$$
 (12)

The iteration formula in (12) yields

$$y_1(x) = 1 + \int_0^x 1 \, dt = 1 + x,$$

$$y_2(x) = 1 + \int_0^x (1+t) \, dt = 1 + x + \frac{1}{2}x^2,$$

$$y_3(x) = 1 + \int_0^x \left(1 + t + \frac{1}{2}t^2\right) dt = 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3,$$

and

$$y_4(x) = 1 + \int_0^x \left(1 + t + \frac{1}{2}t^2 + \frac{1}{6}t^3\right) dt$$
$$= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4.$$

It is clear that we are generating the sequence of partial sums of a power series solution; indeed, we immediately recognize the series as that of $y(x) = e^x$. There is no difficulty in demonstrating that the exponential function is indeed the solution of the initial value problem in (11); moreover, a diligent student can verify (by using a proof by induction on *n*) that $y_n(x)$, obtained in the aforementioned manner, is indeed the *n*th partial sum for the Taylor series with center zero for $y(x) = e^x$.

Example 2 To apply the method of successive approximations to the initial value problem

$$\frac{dy}{dx} = 4xy, \qquad y(0) = 3,$$
 (13)

we write Eqs. (5) and (7) as in Example 1. Now we obtain

$$y_0(x) \equiv 3, \quad y_{n+1}(x) = 3 + \int_0^x 4t y_n(t) dt.$$
 (14)

The iteration formula in (14) yields

$$y_1(x) = 3 + \int_0^x (4t)(3) dt = 3 + 6x^2,$$

$$y_2(x) = 3 + \int_0^x (4t)(3 + 6t^2) dt = 3 + 6x^2 + 6x^4,$$

$$y_3(x) = 3 + \int_0^x (4t)(3 + 6t^2 + 6t^4) dt = 3 + 6x^2 + 6x^4 + 4x^6,$$

and

$$y_4(x) = 3 + \int_0^x (4t)(3 + 6t^2 + 6t^4 + 4t^6) dt$$
$$= 3 + 6x^2 + 6x^4 + 4x^6 + 2x^8.$$

It is again clear that we are generating partial sums of a power series solution. It is not quite so obvious what function has such a power series representation, but the initial value problem in (13) is readily solved by separation of variables:

$$y(x) = 3 \exp(2x^2) = 3 \sum_{n=0}^{\infty} \frac{(2x^2)^n}{n!}$$

= 3 + 6x^2 + 6x^4 + 4x^6 + 2x^8 + $\frac{4}{5}x^{10} + \cdots$.

In some cases, it may be necessary to compute a much larger number of terms, either in order to identify the solution or to use a partial sum of its series with large subscript to approximate the solution accurately for *x* near its initial value. Fortunately, computer algebra systems such as *Maple* and *Mathematica* can perform the symbolic integrations (as opposed to numerical integrations) of the sort in Examples 1 and 2. If necessary, you could generate the first hundred terms in Example 2 in a matter of minutes.

In general, of course, we apply Picard's method because we cannot find a solution by elementary methods. Suppose that we have produced a large number of terms of what we believe to be the correct power series expansion of the solution. We *must* have conditions under which the sequence $\{y_n(x)\}$ provided by the method of successive approximations is guaranteed in advance to converge to a solution. It is just as convenient to discuss the initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t), \quad \mathbf{x}(a) = \mathbf{b}$$
(15)

for a system of *m* first-order equations, where

_

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_m \end{bmatrix}, \quad \mathbf{f} = \begin{bmatrix} f_1 \\ f_2 \\ f_3 \\ \vdots \\ f_m \end{bmatrix}, \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_m \end{bmatrix}.$$

It turns out that with the aid of this vector notation (which we introduced in Section 7.1), most results concerning a single [scalar] equation x' = f(x,t) can be generalized readily to analogous results for a system of *m* first-order equations, as abbreviated in (15). Consequently, the effort of using vector notation is amply justified by the generality it provides.

The method of successive approximations for the system in (15) calls for us to compute the sequence $\{\mathbf{x}_n(t)\}_0^\infty$ of vector-valued functions of *t*,

$$\mathbf{x}_n(t) = \begin{bmatrix} x_{1n}(t) \\ x_{2n}(t) \\ x_{3n}(t) \\ \vdots \\ x_{mn}(t) \end{bmatrix},$$

defined iteratively by

$$\mathbf{x}_0(a) \equiv \mathbf{b}, \quad \mathbf{x}_{n+1}(t) = \mathbf{b} + \int_a^t \mathbf{f}(\mathbf{x}_n(s), s) \, ds.$$
 (16)

Recall that vector-valued functions are integrated componentwise.

nple 3 Consider the *m*-dimensional initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{A}\mathbf{x}, \quad \mathbf{x}(0) = \mathbf{b} \tag{17}$$

for a homogeneous linear system with $m \times m$ constant coefficient matrix **A**. The equations in (16) take the form

$$\mathbf{x}_0(t) = \mathbf{b}, \quad \mathbf{x}_{n+1} = \mathbf{b} + \int_0^x \mathbf{A} \mathbf{x}_n(s) \, ds. \tag{18}$$

Thus

$$\mathbf{x}_1(t) = \mathbf{b} + \int_0^t \mathbf{A}\mathbf{b} \, ds = \mathbf{b} + \mathbf{A}\mathbf{b}t = (\mathbf{I} + \mathbf{A}t)\mathbf{b},$$

$$\mathbf{x}_2(t) = \mathbf{b} + \int_0^t \mathbf{A}(\mathbf{b} + \mathbf{A}\mathbf{b}s) \, ds = \mathbf{b} + \mathbf{A}\mathbf{b}t + \frac{1}{2}\mathbf{A}^2\mathbf{b}t^2 = (\mathbf{I} + \mathbf{A}t + \frac{1}{2}\mathbf{A}^2t^2)\mathbf{b},$$

and

$$\mathbf{x}_{3}(t) = \mathbf{b} + \int_{0}^{t} \mathbf{A}(\mathbf{b} + \mathbf{A}\mathbf{b}s + \frac{1}{2}\mathbf{A}^{2}\mathbf{b}s^{2}) \, ds = (\mathbf{I} + \mathbf{A}t + \frac{1}{2}\mathbf{A}^{2}t^{2} + \frac{1}{6}\mathbf{A}^{3}t^{3})\mathbf{b}.$$

We have therefore obtained the first several partial sums of the exponential series solution

$$\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{b} = \left(\sum_{n=0}^{\infty} \frac{(\mathbf{A}t)^n}{n!}\right)\mathbf{b}$$
(19)

of (17), which was derived earlier in Section 8.1.

The key to establishing convergence in the method of successive approximations is an appropriate condition on the rate at which $\mathbf{f}(\mathbf{x}, t)$ changes when \mathbf{x} varies but *t* is held fixed. If *R* is a region in (m + 1)-dimensional (\mathbf{x}, t) -space, then the function $\mathbf{f}(\mathbf{x}, t)$ is said to be **Lipschitz continuous** on *R* if there exists a constant k > 0 such that

$$|\mathbf{f}(\mathbf{x}_1, t) - \mathbf{f}(\mathbf{x}_2, t)| \le k |\mathbf{x}_1 - \mathbf{x}_2|$$
(20)

if (\mathbf{x}_1, t) and (\mathbf{x}_2, t) are points of *R*. Recall that the **norm** of an *m*-dimensional point or vector **x** is defined to be

$$\mathbf{x}| = \sqrt{x_1^2 + x_2^2 + x_3^2 + \dots + x_m^2}.$$
(21)

Then $|\mathbf{x}_1 - \mathbf{x}_2|$ is simply the Euclidean distance between the points \mathbf{x}_1 and \mathbf{x}_2 .

Example 4

Let $f(x,t) = x^2 \exp(-t^2) \sin t$ and let R be the strip $0 \le x \le 2$ in the xy-plane. If (x_1,t) and (x_2,t) are both points of R, then

$$|f(x_1,t) - f(x_2,t)| = |\exp(-t^2)\sin t| \cdot |x_1 + x_2| \cdot |x_1 - x_2| \le 4|x_1 - x_2|,$$

because $|\exp(-t^2)\sin t| \le 1$ for all t and $|x_1 + x_2| \le 4$ if x_1 and x_2 are both in the interval [0, 2]. Thus f satisfies the Lipschitz condition in (20) with k = 4 and is therefore Lipschitz continuous in the strip R.

Example 5

Let $f(x, t) = t\sqrt{x}$ on the rectangle *R* consisting of the points (x, t) in the *xt*-plane for which $0 \le x \le 1$ and $0 \le t \le 1$. Then, taking $x_1 = x$, $x_2 = 0$, and t = 1, we find that

$$|f(x, 1) - f(0, 1)| = \sqrt{x} = \frac{1}{\sqrt{x}}|x - 0|.$$

Because $x^{-1/2} \to +\infty$ as $x \to 0^+$, we see that the Lipschitz condition in (20) cannot be satisfied by any (finite) constant k > 0. Thus the function f, though obviously continuous on R, is *not* Lipschitz continuous on R.

Suppose, however, that the function f(x, t) has a continuous partial derivative $f_x(x, t)$ on the closed rectangle R in the xt-plane, and denote by k the maximum value of $|f_x(x, t)|$ on R. Then the mean-value theorem of differential calculus yields

$$|f(x_1,t) - f(x_2,t)| = |f_x(\overline{x},t) \cdot (x_1 - x_2)|$$

for some \overline{x} in (x_1, x_2) , so it follows that

$$|f(x_1,t) - f(x_2,t)| \le k|x_1 - x_2|$$

because $|f_x(\overline{x}, t)| \leq k$. Thus a continuously differentiable function f(x, t) defined on a closed rectangle *is* Lipschitz continuous there. More generally, the multivariable mean value theorem of advanced calculus can be used similarly to prove that a *vector-valued function* $\mathbf{f}(\mathbf{x}, t)$ with continuously differentiable component functions on a closed rectangular region R in (\mathbf{x}, t) -space is Lipschitz continuous on R.

Example 6

The function $f(x,t) = x^2$ is Lipschitz continuous on any closed [bounded] region in the *xt*-plane. But consider this function on the infinite strip *R* consisting of the points (x, t) for which $0 \le t \le 1$ and *x* is arbitrary. Then

$$|f(x_1,t) - f(x_2,t)| = |x_1^2 - x_2^2| = |x_1 + x_2| \cdot |x_1 - x_2|.$$

Because $|x_1 + x_2|$ can be made arbitrarily large, it follows that f is *not* Lipschitz continuous on the infinite strip R.



FIGURE A.1. An infinite slab in (m + 1)-space.

If *I* is an interval on the *t*-axis, then the set of all points (\mathbf{x}, t) with *t* in *I* is an infinite strip or slab in (m + 1)-space (as indicated in Fig. A.1). Example 6 shows that Lipschitz continuity of $\mathbf{f}(\mathbf{x}, t)$ on such an infinite slab is a very strong condition. Nevertheless, the existence of a solution of the initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t), \quad \mathbf{x}(a) = \mathbf{b}$$
(15)

under the hypothesis of Lipschitz continuity of \mathbf{f} in such a slab is of considerable importance.

THEOREM 1 Global Existence of Solutions

Let **f** be a vector-valued function (with *m* components) of m + 1 real variables, and let *I* be a (bounded or unbounded) open interval containing t = a. If $\mathbf{f}(\mathbf{x}, t)$ is continuous and satisfies the Lipschitz condition in (20) for all *t* in *I* and for all \mathbf{x}_1 and \mathbf{x}_2 , then the initial value problem in (15) has a solution on the entire interval *I*.

Proof: We want to show that the sequence $\{\mathbf{x}_n(t)\}_0^\infty$ of successive approximations determined iteratively by

$$\mathbf{x}_0(a) = \mathbf{b}, \quad \mathbf{x}_{n+1}(t) = \mathbf{b} + \int_0^t \mathbf{f}(\mathbf{x}_n(s), s) \, ds \tag{16}$$

converges to a solution $\mathbf{x}(t)$ of (15). We see that each of these functions in turn is continuous on *I*, as each is an (indefinite) integral of a continuous function.

We may assume that a = 0, because the transformation $t \rightarrow t + a$ converts (15) into an equivalent problem with initial point t = 0. Also, we will consider only the portion $t \ge 0$ of the interval *I*; the details for the case $t \le 0$ are very similar.

The main part of the proof consists in showing that if [0, T] is a closed (and bounded) interval contained in *I*, then the sequence $\{\mathbf{x}_n(t)\}$ converges *uniformly* on [0, T] to a limit function $\mathbf{x}(t)$. This means that, given $\epsilon > 0$, there exists an integer *N* such that

$$|\mathbf{x}_n(t) - \mathbf{x}(t)| < \epsilon \tag{22}$$

for all $n \ge N$ and all *t* in [0, T]. For ordinary (perhaps nonuniform) convergence the integer *N*, for which (22) holds for all $n \ge N$, might depend on *t*, with no single value of *N* working for all *t* in *I*. Once this uniform convergence of the sequence $\{\mathbf{x}_n(t)\}$ has been established, the following conclusions will follow from standard theorems of advanced calculus [see pages 620–622 of A. E. Taylor and W. R. Mann, *Advanced Calculus*, 3rd ed. (New York: John Wiley, 1983)].

- **1.** The limit function $\mathbf{x}(t)$ is continuous on [0, T].
- **2.** If *N* is so chosen that the inequality in (22) holds for $n \ge N$, then the Lipschitz continuity of **f** implies that

$$|\mathbf{f}(\mathbf{x}_n(t), t) - \mathbf{f}(\mathbf{x}(t), t)| \leq k |\mathbf{x}_n(t) - \mathbf{x}(t)| < k\epsilon$$

for all t in [0, T] and $n \ge N$, so it follows that the sequence $\{\mathbf{f}(\mathbf{x}_n(t), t)\}_0^\infty$ converges uniformly to $\mathbf{f}(\mathbf{x}(t), t)$ on [0, T].

3. But a uniformly convergent sequence or series can be integrated termwise, so it follows that, on taking limits in the iterative formula in (16),

$$\mathbf{x}(t) = \lim_{n \to \infty} \mathbf{x}_{n+1}(t) = \mathbf{b} + \lim_{n \to \infty} \int_0^t \mathbf{f}(\mathbf{x}_n(s), s) \, ds$$
$$= \mathbf{b} + \int_0^t \lim_{n \to \infty} \mathbf{f}(\mathbf{x}_n(s), s) \, ds;$$

thus

$$\mathbf{x}(t) = \mathbf{b} + \int_0^t \mathbf{f}(\mathbf{x}(s), s) \, ds.$$
(23)

4. Because the function $\mathbf{x}(t)$ is continuous on [0, T], the integral equation in (23) [analogous to the 1-dimensional case in (4)] implies that $\mathbf{x}'(t) = \mathbf{f}(\mathbf{x}(t), t)$ on [0, T]. If this is true on every closed subinterval of the open interval *I*, however, then it is true on the entire interval *I* as well.

It therefore remains only to prove that the sequence $\{\mathbf{x}_n(t)\}_0^\infty$ converges uniformly on the closed interval [0, T]. Let *M* be the maximum value of $|\mathbf{f}(\mathbf{b}, t)|$ for *t* in [0, T]. Then

$$|\mathbf{x}_1(t) - \mathbf{x}_0(t)| = \left| \int_0^t \mathbf{f}(\mathbf{x}_0(s), s) \, ds \right| \le \int_0^t |\mathbf{f}(\mathbf{b}, s)| \, ds \le Mt.$$
(24)

Next,

$$|\mathbf{x}_{2}(t) - \mathbf{x}_{1}(t)| = \left| \int_{0}^{t} [\mathbf{f}(\mathbf{x}_{1}(s), s) - \mathbf{f}(\mathbf{x}_{0}(s), s)] \, ds \right| \leq k \int_{0}^{t} |\mathbf{x}_{1}(s) - \mathbf{x}_{0}(s)| \, ds,$$

and hence

$$|\mathbf{x}_{2}(t) - \mathbf{x}_{1}(t)| \leq k \int_{0}^{t} Ms \, ds = \frac{1}{2}kMt^{2}.$$
(25)

We now proceed by induction. Assume that

$$|\mathbf{x}_n(t) - \mathbf{x}_{n-1}(t)| \le \frac{M}{k} \cdot \frac{(kt)^n}{n!}.$$
(26)

It then follows that

$$|\mathbf{x}_{n+1}(t) - \mathbf{x}_n(t)| = \left| \int_0^t \mathbf{f}(\mathbf{x}_n(s), s) - \mathbf{f}(\mathbf{x}_{n-1}(s), s) \right] ds$$
$$\leq k \int_0^t |\mathbf{x}_n(s) - \mathbf{x}_{n-1}(s)| ds;$$

consequently,

$$|\mathbf{x}_{n+1}(t) - \mathbf{x}_n(t)| \leq k \int_0^t \frac{M}{k} \cdot \frac{(ks)^n}{n!} \, ds.$$

It follows, upon evaluation of this integral, that

$$|\mathbf{x}_{n+1}(t) - \mathbf{x}_n(t)| \leq \frac{M}{k} \cdot \frac{(kt)^{n+1}}{(n+1)!}.$$

Thus (26) holds on [0, T] for all $n \ge 1$.

Hence the terms of the infinite series

$$\mathbf{x}_{0}(t) + \sum_{n=1}^{\infty} [\mathbf{x}_{n}(t) - \mathbf{x}_{n-1}(t)]$$
(27)

are dominated (in magnitude on the interval [0, T]) by the terms of the convergent series

$$\sum_{n=1}^{\infty} \frac{M}{k} \cdot \frac{(kT)^{n+1}}{(n+1)!} = \frac{M}{k} (e^{kT} - 1),$$
(28)

which is a series of positive constants. It therefore follows (from the Weierstrass *M*-test on pages 618–619 of Taylor and Mann) that the series in (27) converges uniformly on [0, T]. But the sequence of partial sums of this series is, however, simply our original sequence $\{\mathbf{x}_n(t)\}_0^\infty$ of successive approximations, so the proof of Theorem 1 is finally complete.

A.2 Linear Systems

An important application of the global existence theorem just given is to the initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{A}(t)\mathbf{x} + \mathbf{g}(t), \quad \mathbf{x}(a) = \mathbf{b}$$
(29)

for a linear system, where the $m \times m$ matrix-valued function $\mathbf{A}(t)$ and the vectorvalued function $\mathbf{g}(t)$ are continuous on a (bounded or unbounded) open interval Icontaining the point t = a. In order to apply Theorem 1 to the linear system in (29), we note first that the proof of Theorem 1 requires only that, for each closed and bounded subinterval J of I, there exists a Lipschitz constant k such that

$$|\mathbf{f}(\mathbf{x}_1, t) - \mathbf{f}(\mathbf{x}_2, t)| \le k |\mathbf{x}_1 - \mathbf{x}_2|$$
(20)

for all t in J (and all \mathbf{x}_1 and \mathbf{x}_2). Thus we do not need a single Lipschitz constant for the entire open interval I.

In (29) we have $\mathbf{f}(\mathbf{x}, t) = \mathbf{A}(t)\mathbf{x} + \mathbf{g}$, so

$$\mathbf{f}(\mathbf{x}_1, t) - \mathbf{f}(\mathbf{x}_2, t) = \mathbf{A}(t)(\mathbf{x}_1 - \mathbf{x}_2).$$
(30)

It therefore suffices to show that, if A(t) is continuous on the closed and bounded interval J, then there is a constant k such that

$$|\mathbf{A}(t)\mathbf{x}| \le k|\mathbf{x}| \tag{31}$$

for all t in J. But this follows from the fact (Problem 17) that

$$|\mathbf{A}\mathbf{x}| \le \|\mathbf{A}\| \cdot |\mathbf{x}|,\tag{32}$$

where the **norm** $||\mathbf{A}||$ of the matrix \mathbf{A} is defined to be

$$\|\mathbf{A}\| = \left(\sum_{i,j=1}^{m} (a_{ij})^2\right)^{1/2}.$$
(33)

Because $\mathbf{A}(t)$ is continuous on the closed and bounded interval J, the norm $\|\mathbf{A}\|$ is bounded on J, so Eq. (31) follows, as desired. Thus we have the following global existence theorem for the linear initial value problem in (29).

THEOREM 1 Existence for Linear Systems

Let the $m \times m$ matrix-valued function $\mathbf{A}(t)$ and the vector-valued function $\mathbf{g}(t)$ be continuous on the (bounded or unbounded) open interval I containing the point t = a. Then the initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{A}(t)\mathbf{x} + \mathbf{g}(t), \quad \mathbf{x}(a) = \mathbf{b}$$
(29)

has a solution on the (entire) interval I.

As we saw in Section 4.1, the *m*th-order initial value problem

$$x^{(m)} + a_1(t)x^{(m-1)} + \dots + a_{m-1}(t)x' + a_m(t)x = p(t),$$

$$x(a) = b_0, \quad x'(a) = b_1, \quad \dots, \quad x^{(m-1)}(a) = b_{m-1}$$
(34)

is readily transformed into an equivalent $m \times m$ system of the form in (29). It therefore follows from Theorem 2 that if the functions $a_1(t), a_2(t), \ldots, a_m(t)$ and p(t) in (34) are all continuous on the (bounded or unbounded) open interval *I* containing t = a, then the initial value problem in (34) has a solution on the (entire) interval *I*.

A.3 Local Existence

In the case of a nonlinear initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t), \quad \mathbf{x}(a) = \mathbf{b},$$
(35)

the hypothesis in Theorem 1 that **f** satisfies a Lipschitz condition on a slab (\mathbf{x}, t) (t in I, all \mathbf{x}) is unrealistic and rarely satisfied. This is illustrated by the following simple example.

Example 1 Consider the initial value problem

$$\frac{dy}{dx} = x^2, \quad x(0) = b > 0.$$
 (36)

As we saw in Example 6, the equation $x' = x^2$ does not satisfy a "strip Lipschitz condition." When we solve (36) by separation of variables, we get

X

$$x(t) = \frac{b}{1 - bt}.$$
(37)

Because the denominator vanishes for t = 1/b, Eq. (37) provides a solution of the initial value problem in (36) only for t < 1/b, despite the fact that the differential equation $x' = x^2$ "looks nice" on the entire real line—certainly the function appearing on the right-hand side of the equation is continuous everywhere. In particular, if *b* is large, then we have a solution only on a very small interval to the right of t = 0.

Although Theorem 2 assures us that *linear* equations have global solutions, Example 7 shows that, in general, even a "nice" nonlinear differential equation can be expected to have a solution only on a small interval around the initial point t = a, and it also shows that the length of this interval of existence can depend on the initial value $\mathbf{x}(a) = \mathbf{b}$, as well as on the differential equation itself. The reason is this: If $\mathbf{f}(\mathbf{x}, t)$ is continuously differentiable in a neighborhood of the point (\mathbf{b}, a) in (m + 1)-dimensional space, then—as indicated in the discussion preceding Example 6—we can conclude that $\mathbf{f}(\mathbf{x}, t)$ satisfies a Lipschitz condition on some rectangular region *R* centered at (\mathbf{b}, a) , of the form

$$|t - a| < A, \quad |x_i - b_i| < B_i$$
 (38)

(i = 1, 2, ..., m). In the proof of Theorem 1, we need to apply the Lipschitz condition on the function **f** in analyzing the iterative formula

$$\mathbf{x}_{n+1}(t) = \mathbf{b} + \int_{a}^{t} \mathbf{f}(\mathbf{x}_{n}(s), s) \, ds.$$
(39)

The potential difficulty is that, unless the values of t are suitably restricted, the point $(\mathbf{x}_n(t), t)$ appearing in the integrand in (39) may not lie in the region R where \mathbf{f} is known to satisfy a Lipschitz condition. On the other hand, it can be shown that—on a sufficiently small open interval J containing the point t = a—the graphs of the functions $\{\mathbf{x}_n(t)\}$ given iteratively by the formula in (39) remain within the region R, so the proof of convergence can then be carried out as in the proof of Theorem 1. A proof of the following *local* existence theorem can be found in Chapter 6 of G. Birkhoff and G.-C. Rota, *Ordinary Differential Equations*, 4th ed. (New York: John Wiley, 1989).

THEOREM 1 Local Existence of Solutions

Let **f** be a vector-valued function (with *m* components) of the m + 1 real variables $x_1, x_2, ..., x_m$, and *t*. If the first-order partial derivatives of **f** all exist and are continuous in some neighborhood of the point $\mathbf{x} = \mathbf{b}, t = a$, then the initial value problem

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t), \quad \mathbf{x}(a) = \mathbf{b},$$
(35)

has a solution on some open interval containing the point t = a.

A.4 Uniqueness of Solutions

It is possible to establish the existence of solutions of the initial value problem in (35) under the much weaker hypothesis that $\mathbf{f}(\mathbf{x}, t)$ is merely continuous; techniques other than those used in this section are required. By contrast, the Lipschitz condition that we used in proving Theorem 1 is the key to *uniqueness* of solutions. In particular, the solution provided by Theorem 3 is unique near the point t = a.

THEOREM 1 Uniqueness of Solutions

Suppose that on some region R in (m + 1)-space, the function **f** in (35) is continuous and satisfies the Lipschitz condition

$$|\mathbf{f}(\mathbf{x}_1, t) - \mathbf{f}(\mathbf{x}_2, t)| \le k \cdot |\mathbf{x}_1 - \mathbf{x}_2|.$$
(20)

If $\mathbf{x}_1(t)$ and $\mathbf{x}_2(t)$ are two solutions of the initial problem in (35) on some open interval *I* containing x = a, such that the solution curves $(\mathbf{x}_1(t), t)$ and $(\mathbf{x}_2(t), t)$ both lie in *R* for all *t* in *I*, then $\mathbf{x}_1(t) = \mathbf{x}_2(t)$ for all *t* in *I*.

We will outline the proof of Theorem 4 for the 1-dimensional case in which *x* is a real variable. A generalization of this proof to the multivariable case can be found in Chapter 6 of Birkhoff and Rota.

Let us consider the function

$$\phi(t) = [x_1(t) - x_2(t)]^2 \tag{40}$$

for which $\phi(a) = 0$, because $x_1(a) = x_2(a) = b$. We want to show that $\phi(t) \equiv 0$, so that $x_1(t) \equiv x_2(t)$. We will consider only the case $t \ge a$; the details are similar for the case $t \le a$.

If we differentiate each side in Eq. (40), we find that

$$\begin{aligned} \phi'(t) &| = |2[x_1(t) - x_2(t)] \cdot [x_1'(t) - x_2'(t)]| \\ &= |2[x_1(t) - x_2(t)] \cdot [f(x_1(t), t) - f(x_2(t), t)]| \\ &\leq 2k |x_1(t) - x_2(t)|^2 = 2k\phi(t), \end{aligned}$$

by using the Lipschitz condition on f. Hence

$$\phi'(t) \le 2k\phi(t). \tag{41}$$

Now let us temporarily ignore the fact that $\phi(a) = 0$ and compare $\phi(t)$ with the solution of the differential equation

$$\Phi'(t) = 2k\Phi(t) \tag{42}$$

such that $\Phi(a) = \phi(a)$; clearly

$$\Phi(t) = \Phi(a)e^{2k(t-a)}.$$
(43)

In comparing (41) with (42), it seems inevitable that

 $\phi(t) \leq \Phi(t) \qquad \text{for } t \geq a, \tag{44}$

and this is easily proved (Problem 18). Hence

$$0 \leq [x_1(t) - x_2(t)]^2 \leq [x_1(a) - x_2(a)]^2 e^{2k(t-a)}.$$

On taking square roots, we get

$$0 \leq |x_1(t) - x_2(t)| \leq |x_1(a) - x_2(a)|e^{k(t-a)}.$$
(45)

But $x_1(a) - x_2(a) = 0$, so (45) implies that $x_1(t) \equiv x_2(t)$.

Example

The initial value problem

$$\frac{dx}{dt} = 3x^{2/3}, \quad x(0) = 0 \tag{46}$$

has both the obvious solution $x_1(t) \equiv 0$ and the solution $x_2(t) = t^3$ (which is readily found by separation of variables). Hence the function f(x, t) must *fail* to satisfy a Lipschitz condition near (0, 0). Indeed, the mean value theorem yields

$$|f(x,0) - f(0,0)| = |f_x(\overline{x},0)| \cdot |x-0|$$

for some \overline{x} between 0 and x. But $f_x(x, 0) = 2x^{-1/3}$ is unbounded as $x \to 0$, so no Lipschitz condition can be satisfied.

A.5 Well-Posed Problems and Mathematical Models

In addition to uniqueness, another consequence of the inequality in (45) is the fact that solutions of the differential equation

$$\frac{dx}{dt} = f(x,t) \tag{47}$$

depend *continuously* on the initial value x(a); that is, if $x_1(t)$ and $x_2(t)$ are two solutions of (47) on the interval $a \leq t \leq T$ such that the initial values $x_1(a)$ and $x_2(a)$ are sufficiently close to one another, then the values of $x_1(t)$ and $x_2(t)$ remain close to one another. In particular, if $|x_1(a) - x_2(a)| \leq \delta$, then (45) implies that

$$|x_1(t) - x_2(t)| \le \delta e^{k(T-a)} = \epsilon \tag{48}$$

for all *t* with $a \leq t \leq T$. Obviously, we can make ϵ as small as we wish by choosing δ sufficiently close to zero.

This continuity of solutions of (47) with respect to initial values is important in practical applications where we are unlikely to know the initial value $x_0 = x(a)$ with absolute precision. For example, suppose that the initial value problem

$$\frac{dx}{dt} = f(x,t), \quad x(a) = x_0 \tag{49}$$

models a population for which we know only that the initial population is within $\delta > 0$ of the assumed value x_0 . Then even if the function f(x, t) is accurate, the solution x(t) of (49) will be only an approximation to the actual population. But (45) implies that the actual population at time t will be within $\delta e^{k(T-a)}$ of the approximate population x(t). Thus, on a given closed interval [a, T], x(t) will be a close approximation to the actual population provided that $\delta > 0$ is sufficiently small.

An initial value problem is usually considered *well posed* as a mathematical model for a real-world situation only if the differential equation has unique solutions that are continuous with respect to initial values. Otherwise it is unlikely that the initial value problem adequately mirrors the real-world situation.

An even stronger "continuous dependence" of solutions is often desirable. In addition to possible inaccuracy in the initial value, the function f(x,t) may not model precisely the physical situation. For instance, it may involve physical parameters (such as resistance coefficients) whose values cannot be measured with absolute precision. Birkhoff and Rota generalize the proof of Theorem 4 to establish the following result.

THEOREM 1 Continuous Dependence of Solutions

Let $\mathbf{x}(t)$ and $\mathbf{y}(t)$ be solutions of the equations

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}, t) \text{ and } \frac{d\mathbf{y}}{dt} = \mathbf{g}(\mathbf{y}, t)$$
 (50)

on the closed interval [a, T]. Let **f** and **g** be continuous for $a \le t \le T$ and for **x** and **y** in a common region *D* of *n*-space and assume that **f** satisfies the Lipschitz condition in (20) on the region *D*. If

$$|\mathbf{f}(\mathbf{z},t) - \mathbf{g}(\mathbf{z},t)| \le \mu \tag{51}$$

for all t in the interval [a, T] and all \mathbf{z} in D, it then follows that

$$|\mathbf{x}(t) - \mathbf{y}(t)| \leq |\mathbf{x}(a) - \mathbf{y}(a)| \cdot e^{k(t-a)} + \frac{\mu}{k} \left[e^{k(t-a)} - 1 \right]$$
(52)

on the interval [a, T].

If $\mu > 0$ is small, then (51) implies that the functions **f** and **g** appearing in the two differential equations, though different, are "close" to each other. If $\epsilon > 0$ is

given, then it is apparent from (52) that

$$|\mathbf{x}(t) - \mathbf{y}(t)| \le \epsilon \tag{53}$$

for all t in [a, T] if both $|\mathbf{x}(a) - \mathbf{y}(a)|$ and μ are sufficiently small. Thus Theorem 5 says (roughly) that if both the two initial values and the two differential equations in (50) are close to each other, then the two solutions remain close to each other for $a \leq t \leq T$.

For example, suppose that a falling body is subject both to constant gravitational acceleration g and to resistance proportional to some power of its velocity, so (with the positive axis directed downward) its velocity v satisfies the differential equation

$$\frac{dv}{dt} = g - cv^{\rho}.$$
(54)

Assume, however, that only an approximation \overline{c} to the actual resistance c and an approximation $\overline{\rho}$ to the actual exponent ρ are known. Then our mathematical model is based on the differential equation

$$\frac{du}{dt} = g - \overline{c}u^{\overline{\rho}} \tag{55}$$

instead of the actual equation in (54). Thus if we solve Eq. (55), we obtain only an approximation u(t) to the actual velocity v(t). But if the parameters \overline{c} and $\overline{\rho}$ are sufficiently close to the actual values c and ρ , then the right-hand sides in (54) and (55) will be close to each other. If so, then Theorem 5 implies that the actual and approximate velocity functions v(t) and u(t) are close to each other. In this case, the approximation in (55) will be a good model of the actual physical situation.

Problems

In Problems 1 through 8, apply the successive approximation formula to compute $y_n(x)$ for $n \leq 4$. Then write the exponential series for which these approximations are partial sums (perhaps with the first term or two missing; for example,

$$e^{x} - 1 = x + \frac{1}{2}x^{2} + \frac{1}{6}x^{3} + \frac{1}{24}x^{4} + \cdots).$$

1.
$$\frac{dy}{dx} = y, y(0) = 3$$
 2. $\frac{dy}{dx} = -2y, y(0) = 4$

3.
$$\frac{dy}{dx} = -2xy, y(0) = 1$$
 4. $\frac{dy}{dx} = 3x^2y, y(0) = 2$

5.
$$\frac{dy}{dx} = 2y + 2$$
, $y(0) = 0$
6. $\frac{dy}{dx} = x + y$, $y(0) = 0$

7.
$$\frac{dy}{dx} = 2x(1+y), y(0) = 0$$

8. $\frac{dy}{dx} = 4x(y+2x^2), y(0) = 0$

1.

In Problems 9 through 12, compute the successive approximations $y_n(x)$ for $n \leq 3$; then compare them with the appropriate partial sums of the Taylor series of the exact solution.

9.
$$\frac{dy}{dx} = x + y, y(0) = 1$$
 10. $\frac{dy}{dx} = y + e^x, y(0) = 0$

11.
$$\frac{dy}{dx} = y^2$$
, $y(0) = 1$ **12.** $\frac{dy}{dx} = \frac{1}{2}y^3$, $y(0) = 1$

13. Apply the iterative formula in (16) to compute the first three successive approximations to the solution of the initial value problem

$$\frac{dx}{dt} = 2x - y,$$
 $x(0) = 1;$
 $\frac{dy}{dt} = 3x - 2y,$ $y(0) = -1.$

14. Apply the matrix exponential series in (19) to solve (in closed form) the initial value problem

$$\mathbf{x}'(t) = \begin{bmatrix} 1 & 1\\ 0 & 1 \end{bmatrix} \mathbf{x}, \quad \mathbf{x}(0) = \begin{bmatrix} 1\\ 1 \end{bmatrix}$$

(Suggestion: Show first that

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$$

for each positive integer *n*.)

15. For the initial value problem $dy/dx = 1 + y^3$, y(1) = 1, show that the second Picard approximation is

$$y_2(x) = 1 + 2(x-1) + 3(x-1)^2 + 4(x-1)^3 + 2(x-1)^4.$$

Then compute $y_2(1.1)$ and $y_2(1.2)$. The fourth-order Runge–Kutta method with step size h = 0.005 yields $y(1.1) \approx 1.2391$ and $y(1.2) \approx 1.6269$.

16. For the initial value problem $dy/dx = x^2 + y^2$, y(0) = 0, show that the third Picard approximation is

$$y_3(x) = \frac{1}{3}x^3 + \frac{1}{63}x^7 + \frac{2}{2079}x^{11} + \frac{1}{59535}x^{15}.$$

Compute $y_3(1)$. The fourth-order Runge–Kutta method yields $y(1) \approx 0.350232$, both with step size h = 0.05 and with step size h = 0.025.

17. Prove as follows the inequality $|\mathbf{A}\mathbf{x}| \leq ||\mathbf{A}|| \cdot |\mathbf{x}|$, where **A** is an $m \times m$ matrix with row vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m$, and **x** is an *m*-dimensional vector. First note that the components of the vector $\mathbf{A}\mathbf{x}$ are $\mathbf{a}_1 \cdot \mathbf{x}, \mathbf{a}_2 \cdot \mathbf{x}, \dots, \mathbf{a}_m \cdot \mathbf{x}$, so

$$|\mathbf{A}\mathbf{x}| = \left[\sum_{n=1}^{m} (\mathbf{a}_i \cdot \mathbf{x})^2\right]^{1/2}.$$

Then use the Cauchy–Schwarz inequality $(a \cdot x)^2 \leqq |a|^2 |x|^2$ for the dot product.

18. Suppose that $\phi(t)$ is a differentiable function with

$$\phi'(t) \le k\phi(t) \qquad (k > 0)$$

for $t \ge a$. Multiply both sides by e^{-kt} , then transpose to show that

$$\frac{d}{dt} \left[\phi(t) e^{-kt} \right] \leq 0$$

for $t \ge a$. Then apply the mean value theorem to conclude that

$$\phi(t) \le \phi(a)e^{k(t-a)}$$

for $t \geq a$.

APPENDIX B

Theory of Determinants

The proof of the theorem on row and column cofactor expansions (Theorem 1 in Section 3.6) involves an alternative interpretation of determinants that in more advanced treatments is usually taken as the original definition. Consider the following elementary scheme for evaluating a 3×3 determinant:

 $\begin{array}{c} a_{11} & a_{12} & a_{13} & a_{11} & a_{12} \\ a_{21} & a_{22} & a_{23} & a_{21} & a_{22} \\ a_{31} & a_{32} & a_{33} & a_{31} & a_{32} \end{array}$

The first two columns of $\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}$ are duplicated to its right. It is readily verified that det \mathbf{A} is equal to the sum of the three products along the indicated diagonals that point down and to the *right*, minus the sum of the products along the three diagonals that point down and to the *left*:

$$\det \mathbf{A} = + a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33}.$$
(1)

Remark This scheme works **only** in dimensions 2 and 3. Neither the analogous scheme nor anything resembling it gives the correct value of an $n \times n$ determinant with $n \ge 4$.

Note that each of the six terms on the right in (1) is of the form $\pm a_{1i}a_{2j}a_{3k}$, where $(i \ j \ k)$ is a **permutation** of (1 2 3). That is, the triple $(i \ j \ k)$ consists of the three distinct numbers 1, 2, and 3 written in some specific order. The six terms in (1) correspond to the six possible permutations of (1 2 3):

The + and - signs in (1) can also be explained in terms of these permutations. A **transposition** of an ordered sequence of objects (such as the numbers 1, 2, 3) is the operation of *interchanging* some single pair of them. For instance, the operation $(1 \ 2 \ 3) \rightarrow (1 \ 3 \ 2)$ is a transposition that consists of interchanging 2 and 3. But it requires two transpositions to change $(1 \ 2 \ 3)$ into the permutation $(3 \ 1 \ 2)$:

$$(1 \ 2 \ 3) \rightarrow (3 \ 2 \ 1) \rightarrow (3 \ 1 \ 2).$$

Given a permutation $P = (i \ j \ k)$ of $(1 \ 2 \ 3)$, we denote by s(P) the *minimum* number of transpositions required to change $(1 \ 2 \ 3)$ into $(i \ j \ k)$. For the six permutations in (2) it is easy to verify that

$$s(1 \ 2 \ 3) = 0, \qquad s(2 \ 1 \ 3) = 1, \qquad s(3 \ 1 \ 2) = 2, s(1 \ 3 \ 2) = 1, \qquad s(2 \ 3 \ 1) = 2, \qquad s(3 \ 2 \ 1) = 1.$$
(3)

Now for the point to all this: If you check each of the six terms in (1) you will find that the *sign* of the term $a_{1i}a_{2j}a_{3k}$ is $(-1)^{s(P)}$, where $P = (i \ j \ k)$. Therefore, we can rewrite (1) more concisely as

$$\det \mathbf{A} = \sum_{P} (-1)^{s(P)} a_{1i} a_{2j} a_{3k}, \tag{4}$$

where there is one term on the right for each possible permutation $P = (i \ j \ k)$ of $(1 \ 2 \ 3)$.

The formula in (4) generalizes to determinants of higher order. If **A** is an $n \times n$ matrix, then

$$\det \mathbf{A} = \sum_{P} (-1)^{s(P)} a_{1i_1} a_{2i_2} \cdots a_{ni_n},$$
(5)

where there is one term on the right for each possible permutation $P = (i_1 \ i_2 \ \cdots \ i_n)$ of $(1 \ 2 \ \cdots \ n)$, and s(P) denotes the minimum number of transpositions required to change $(1 \ 2 \ \cdots \ n)$ into $(i_1 \ i_2 \ \cdots \ i_n)$. The formula in (5) can be established by induction on n. Assuming its validity for $(n-1) \times (n-1)$ determinants, the formula can be verified for an $n \times n$ matrix **A** by expanding along its first row. It is fairly easy to see that this gives terms of the form $\pm a_{1i_1}a_{2i_2}\cdots a_{ni_n}$, but somewhat more difficult to check the signs.

Finally, the proof of the cofactor expansion theorem consists of verifying similarly that the cofactor expansion of det \mathbf{A} along *any* row *or* column of \mathbf{A} agrees with the formula in (5). The details are more lengthy than instructive, and therefore we omit them.

Determinants and Elementary Row Operations

As we described in Section 3.6, the determinant of any square matrix \mathbf{A} can be evaluated directly by carrying out a cofactor expansion of det \mathbf{A} along any row or column of \mathbf{A} . Because this approach involves so much computational labor, it is more efficient instead to reduce \mathbf{A} to an echelon matrix \mathbf{R} . Because any square echelon matrix is (upper) triangular, the determinant of the echelon matrix \mathbf{R} is simply the product of its diagonal elements. But because we have altered the matrix \mathbf{A} by transforming it into \mathbf{R} , the question is this: What effects do elementary row operations have on the determinant of \mathbf{A} ?

The following theorem summarizes Properties 1, 2, and 5 of Section 3.6 and tells us how to keep track of the effect each elementary row operation will have on det \mathbf{A} as we reduce \mathbf{A} to echelon form. We use here the concise notation

$$|\mathbf{A}| = \det \mathbf{A}$$

for the determinant of the matrix A.

THEOREM 1 Effect of Elementary Row Operations

Let A be a square matrix and let B be the matrix obtained by performing one of the three types of elementary row operations on A.

(a) Type 1: If **B** is obtained by *dividing* some row of **A** by k, then

$$|\mathbf{A}| = k|B|. \tag{6}$$

(b) Type 2: If **B** is obtained by interchanging two rows of **A**, then

$$\mathbf{A}| = -|\mathbf{B}|.\tag{7}$$
(c) Type 3: If **B** is obtained by adding a multiple of some row of **A** to another row, then

$$|\mathbf{A}| = |\mathbf{B}|.\tag{8}$$

The following example shows how to lay out the computation of $\det A$ by using row operations to reduce A to echelon form.

Example 1	$\begin{vmatrix} 2 & 2 & -2 & 0 \\ -4 & 2 & 3 & 1 \\ 0 & 4 & -1 & -4 \\ 3 & 1 & 3 & -1 \end{vmatrix} = 2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ -4 & 2 & 3 & 1 \\ 0 & 4 & -1 & -4 \\ 3 & 1 & 3 & -1 \end{vmatrix}$	We have divided row 1 by 2.
	$= 2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & 6 & -1 & 1 \\ 0 & 4 & -1 & -4 \\ 0 & -2 & 6 & -1 \end{vmatrix}$	We have added 4 times row 1 to row 2, and -3 times row 1 to row 4.
	$= +2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & 0 & 17 & -2 \\ 0 & 0 & 11 & -6 \\ 0 & -2 & 6 & -1 \end{vmatrix}$	We have added 3 times row 4 to row 2, and 2 times row 4 to row 3.
	$= -2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 6 & -1 \\ 0 & 0 & 11 & -6 \\ 0 & 0 & 17 & -2 \end{vmatrix}$	We have interchanged rows 2 and 4.
	$= -22 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 6 & -1 \\ 0 & 0 & 1 & -\frac{6}{11} \\ 0 & 0 & 17 & -2 \end{vmatrix}$	We have divided row 3 by 11.
	$= -22 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 6 & -1 \\ 0 & 0 & 1 & -\frac{6}{11} \\ 0 & 0 & 0 & \frac{80}{11} \end{vmatrix}$	We have added -17 times row 3 to row 4.
	$= (-22)(1)(-2)(1)(\frac{80}{11}) = 320.$	•

If the square matrix **A** has only *integer* entries, it is always possible to carry out the reduction to echelon form without introducing any fractions. In the computation of determinants (as opposed to the solution of linear systems), we have the additional flexibility of using elementary *column* operations; their effects are precisely analogous to those of elementary row operations, as described in Theorem 1. In Example 2, we illustrate this by finishing differently the evaluation in Example 1.

Example 2	$\begin{vmatrix} 2 & 2 & -2 & 0 \\ -4 & 2 & 3 & 1 \\ 0 & 4 & -1 & -4 \\ 3 & 1 & 3 & -1 \end{vmatrix} = -2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 6 & -1 \\ 0 & 0 & 11 & -6 \\ 0 & 0 & 17 & -2 \end{vmatrix}$	(As in Example 1.)
	$= -2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 5 & -1 \\ 0 & 0 & 5 & -6 \\ 0 & 0 & 15 & -2 \end{vmatrix}$	We have added column 4 to column 3.

$$= -2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 5 & -1 \\ 0 & 0 & 5 & -6 \\ 0 & 0 & 0 & 16 \end{vmatrix}$$
 We have added (-3) times row 3 to row 4.

$$= (-2)(1)(-2)(5)(16) = 320.$$

Yet another alternative is to reduce only the first n - 2 rows of the $n \times n$ matrix **A**. We then directly calculate the 2×2 determinant that remains in the lower right-hand corner. We get det **A** by multiplying this 2×2 determinant by the preceding factors and diagonal elements. Thus

$$\begin{vmatrix} 2 & 2 & -2 & 0 \\ -4 & 2 & 3 & 1 \\ 0 & 4 & -1 & -4 \\ 3 & 1 & 3 & -1 \end{vmatrix} = -2 \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & -2 & 6 & -1 \\ 0 & 0 & 11 & -6 \\ 0 & 0 & 17 & -2 \end{vmatrix}$$
$$= (-2)(1)(-2) \begin{vmatrix} 11 & -6 \\ 17 & -2 \end{vmatrix} = (4)(-22 + 102) = 320.$$

Determinants and Invertibility

We began Section 3.6 with the remark that a 2×2 matrix **A** is invertible if and only if its determinant is nonzero: $|\mathbf{A}| \neq 0$. Now we want to show that this result also holds for $n \times n$ matrices. This connection between determinants and invertibility is closely related to the fact that the determinant function "respects" matrix multiplication in the sense that

$$\mathbf{AB}| = |\mathbf{A}| \, |\mathbf{B}| \tag{9}$$

if **A** and **B** are $n \times n$ matrices. Our first step is to show that Eq. (9) holds if **A** is an *elementary matrix* obtained from the $n \times n$ identity matrix **I** by performing a single elementary row operation.

LEMMA Multiplication by an Elementary Matrix

If **B** is an $n \times n$ matrix and **E** is an $n \times n$ elementary matrix then

$$|\mathbf{EB}| = |\mathbf{E}| |\mathbf{B}|. \tag{10}$$

Proof: This is really just a restatement of Theorem 1. For instance, suppose that **E** is obtained from **I** by multiplying the *p*th row by *c*, so that $|\mathbf{E}| = c$. Then Theorem 5 in Section 3.5 tells us that the product **EB** is the result of multiplying the *p*th row of **B** by *c*. Therefore,

$$|\mathbf{E}\mathbf{B}| = c|\mathbf{B}| = |\mathbf{E}||\mathbf{B}|,$$

and so we have verified Eq. (10) for the first of the three types of elementary matrices. The verifications for the other two types are similar.

Now, let **A** be an $n \times n$ matrix whose invertibility we want to discuss, and let **R** be the *reduced* echelon form of **A**. If the elementary matrices $\mathbf{F}_1, \mathbf{F}_2, \ldots, \mathbf{F}_k$ correspond to the elementary row operations that reduce **A** to **R**, then

$$\mathbf{F}_k \cdots \mathbf{F}_2 \mathbf{F}_1 \mathbf{A} = \mathbf{R} \tag{11}$$

by Theorem 5 in Section 3.5. Recalling that every elementary matrix is invertible (Section 3.5), we can rewrite Eq. (11) as

$$\mathbf{A} = \mathbf{E}_1 \mathbf{E}_2 \cdots \mathbf{E}_k \mathbf{R},\tag{12}$$

where each $\mathbf{E}_i = (\mathbf{F}_i)^{-1}$ is an elementary matrix. It now follows, by *k* applications of the lemma, that

$$\mathbf{A} = |\mathbf{E}_1| |\mathbf{E}_2| \cdots |\mathbf{E}_k| |\mathbf{R}|.$$
(13)

This relation is the key both to the proof of (9) and to the proof of the following theorem.

THEOREM 2 Determinants and Invertibility

The $n \times n$ matrix **A** is invertible if and only if det $\mathbf{A} \neq 0$.

Proof: If (as previously) **R** is the reduced echelon form of **A**, then Theorem 6 in Section 3.5 implies that

A is invertible if and only if $\mathbf{R} = \mathbf{I}$. (14)

Because **R** is a *square* reduced echelon matrix, we see that *either* **R** is the identity matrix **I** and $|\mathbf{R}| = 1$, or **R** has an all-zero row and, consequently, $|\mathbf{R}| = 0$. Therefore, $\mathbf{R} = \mathbf{I}$ if and only if $|\mathbf{R}| \neq 0$. (15)

Finally, because $|\mathbf{E}| \neq 0$ if \mathbf{E} is an elementary matrix, it follows immediately from Eq. (13) that

$$|\mathbf{R}| \neq 0$$
 if and only if $|\mathbf{A}| \neq 0$. (16)

Combining the statements in (14), (15), and (16), we see that **A** is invertible if and only if $|\mathbf{A}| \neq 0$.

So now we can add the statement det $\mathbf{A} \neq 0$ to the list of equivalent properties of nonsingular matrices stated in Theorem 7 of Section 3.5. Indeed, some texts *define* the square matrix \mathbf{A} to be nonsingular if and only if det $\mathbf{A} \neq 0$.

THEOREM 3 Determinants and Matrix Products

If **A** and **B** are $n \times n$ matrices, then

$$\mathbf{AB}| = |\mathbf{A}| \, |\mathbf{B}|. \tag{9}$$

Proof: If **R** is the reduced echelon form of **A**, then we see that

$$\mathbf{A} = \mathbf{E}_1 \mathbf{E}_2 \cdots \mathbf{E}_k \mathbf{R},\tag{12}$$

where $\mathbf{E}_1, \mathbf{E}_2, \ldots$, and \mathbf{E}_k are elementary matrices. Hence

$$\mathbf{AB} = \mathbf{E}_1 \mathbf{E}_2 \cdots \mathbf{E}_k \mathbf{RB}.$$

We now take the determinant of both sides, using the lemma stated earlier to "split off" the elementary matrices:

$$|\mathbf{AB}| = |\mathbf{E}_1| |\mathbf{E}_2 \mathbf{E}_3 \cdots \mathbf{E}_k \mathbf{RB}| \qquad (\text{lemma once})$$
$$= |\mathbf{E}_1| |\mathbf{E}_2| |\mathbf{E}_3 \cdots \mathbf{E}_k \mathbf{RB}| \qquad (\text{lemma twice})$$
$$= |\mathbf{E}_1 \mathbf{E}_2| |\mathbf{E}_3 \cdots \mathbf{E}_k \mathbf{RB}|. \qquad (\text{lemma thrice})$$

After 2k - 1 steps, we get

$$|\mathbf{AB}| = |\mathbf{E}_1 \mathbf{E}_2 \cdots \mathbf{E}_k| |\mathbf{RB}|. \tag{17}$$

The remainder of the proof depends on whether or not **A** is invertible.

If **A** *is* invertible, then $\mathbf{R} = \mathbf{I}$, so Eq. (12) yields

$$\mathbf{A} = \mathbf{E}_1 \mathbf{E}_2 \cdots \mathbf{E}_k$$

and also that $\mathbf{RB} = \mathbf{IB} = \mathbf{B}$. In this case the meaning of Eq. (17) is precisely that $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$.

If **A** is *not* invertible, then $|\mathbf{A}| = 0$ by Theorem 2. Also, as we noted previously, the reduced echelon form **R** of **A** has an all-zero row in this case. Hence it follows from the definition of matrix multiplication that the product **RB** has an all-zero row and, therefore, that $|\mathbf{RB}| = 0$. In this case Eq. (17) implies that $|\mathbf{AB}| = 0$. Because both $|\mathbf{A}| = 0$ and $|\mathbf{AB}| = 0$, the equation $|\mathbf{AB}| = |\mathbf{A}| |\mathbf{B}|$ holds, and the proof is complete.

Cramer's Rule and Inverse Matrices

Suppose that we need to solve the $n \times n$ linear system

$$\mathbf{A}\mathbf{x} = \mathbf{b},\tag{18}$$

where

$$\mathbf{A} = \begin{bmatrix} a_{ij} \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}. \tag{19}$$

We assume that the coefficient matrix **A** is invertible, so we know in advance that a unique solution **x** of (18) exists. The question is how to write **x** explicitly in terms of the coefficients a_{ij} and the constants b_i . In the following discussion, we think of **x** as a fixed (though as yet unknown) vector.

If we denote by $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ the column vectors of the $n \times n$ matrix **A**, then

$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}. \tag{20}$$

By Fact 1 in Section 3.5, we can rewrite Eq. (18) as

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \dots + x_n\mathbf{a}_n = \mathbf{b}.$$

Thus the constant vector **b** is expressed in terms of the entries $x_1, x_2, ..., x_n$ of the solution vector **x** and the column vectors of **A** by

$$\mathbf{b} = \sum_{j=1}^{n} x_j \mathbf{a}_j. \tag{21}$$

The trick for finding the *i*th unknown x_i is to compute the determinant of the matrix

$$\begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{b} & \cdots & \mathbf{a}_n \end{bmatrix} = \begin{bmatrix} a_{11} & \cdots & b_1 & \cdots & a_{1n} \\ a_{21} & \cdots & b_2 & \cdots & a_{2n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & \cdots & b_n & \cdots & a_{nn} \end{bmatrix}$$
(22)

that we obtain by replacing the *i* th column \mathbf{a}_i of **A** with the constant vector **b**. Using

Eq. (21) to substitute for **b**, we find that

$$\mathbf{a}_{1} \cdots \mathbf{b} \cdots \mathbf{a}_{n} \mid = \begin{vmatrix} \mathbf{a}_{1} & \cdots & \sum_{j=1}^{n} x_{j} \mathbf{a}_{j} & \cdots & \mathbf{a}_{n} \end{vmatrix}$$
$$= \sum_{j=1}^{n} \mid \mathbf{a}_{1} & \cdots & x_{j} \mathbf{a}_{j} & \cdots & \mathbf{a}_{n} \mid \qquad \text{(by Property 4 of determinants)}$$
$$= \sum_{j=1}^{n} x_{j} \mid \mathbf{a}_{1} & \cdots & \mathbf{a}_{j} & \cdots & \mathbf{a}_{n} \mid \qquad \text{(by Property 1 of determinants)}.$$

Note that, in the *j* th term of this summation, the vector \mathbf{a}_j appears in the *i*th position. Thus we have found that

$$|\mathbf{a}_{1} \cdots \mathbf{b} \cdots \mathbf{a}_{n}| = x_{1} |\mathbf{a}_{1} \mathbf{a}_{2} \cdots \mathbf{a}_{1} \cdots \mathbf{a}_{n}|$$

$$+ x_{2} |\mathbf{a}_{1} \mathbf{a}_{2} \cdots \mathbf{a}_{2} \cdots \mathbf{a}_{n}|$$

$$\vdots$$

$$+ x_{i} |\mathbf{a}_{1} \mathbf{a}_{2} \cdots \mathbf{a}_{i} \cdots \mathbf{a}_{n}|$$

$$\vdots$$

$$+ x_{n} |\mathbf{a}_{1} \mathbf{a}_{2} \cdots \mathbf{a}_{n} \cdots \mathbf{a}_{n}|.$$

Of the *n* determinants on the right-hand side here, all but the *i*th one have two identical columns and therefore are equal to zero. The coefficient of x_i in the *i*th term is simply

$$|\mathbf{A}| = |\mathbf{a}_1 \ \mathbf{a}_2 \ \cdots \ \mathbf{a}_i \ \cdots \ \mathbf{a}_n |$$

Consequently, a result of our computation is that

$$|\mathbf{a}_1 \cdots \mathbf{b} \cdots \mathbf{a}_n| = x_i |\mathbf{A}|.$$
 (23)

We get the desired simple formula for x_i after we divide each side by $|\mathbf{A}| \neq 0$.

THEOREM 4 Cramer's Rule

Consider the $n \times n$ linear system Ax = b with

$$\mathbf{A} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}.$$

If $|\mathbf{A}| \neq 0$, then the *i*th entry of the unique solution $\mathbf{x} = (x_1, x_2, \dots, x_n)$ is given by

$$x_{i} = \frac{\begin{vmatrix} \mathbf{a}_{1} & \cdots & \mathbf{b} & \cdots & \mathbf{a}_{n} \end{vmatrix}}{|\mathbf{A}|}$$
$$= \frac{1}{|\mathbf{A}|} \begin{vmatrix} a_{11} & \cdots & b_{1} & \cdots & a_{1n} \\ a_{21} & \cdots & b_{2} & \cdots & a_{2n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & \cdots & b_{n} & \cdots & a_{nn} \end{vmatrix},$$
(24)

where (in the last expression) the constant vector **b** replaces the *i* th column vector \mathbf{a}_i of **A**.

Inverses and the Adjoint Matrix

We now use Cramer's rule to develop an explicit formula for the inverse A^{-1} of the invertible matrix **A**. First, we need to rewrite Cramer's rule more concisely. Expansion of the determinant in the numerator in Eq. (24) along its *i*th column yields

$$x_i = \frac{1}{|\mathbf{A}|} (b_1 A_{1i} + b_2 A_{2i} + \dots + b_n A_{ni}),$$
(25)

because the cofactor of b_p is simply the cofactor A_{pi} of a_{pi} in $|\mathbf{A}|$. The formula in Eq. (25) gives the solution vector

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \frac{1}{|\mathbf{A}|} \begin{bmatrix} b_1 A_{11} + b_2 A_{21} + \dots + b_n A_{n1} \\ b_1 A_{12} + b_2 A_{22} + \dots + b_n A_{n2} \\ \vdots \\ b_1 A_{1n} + b_2 A_{2n} + \dots + b_n A_{nn} \end{bmatrix}.$$

Then the definition of matrix multiplication yields

$$\mathbf{x} = \frac{1}{|\mathbf{A}|} \begin{bmatrix} A_{11} & A_{21} & \cdots & A_{n1} \\ A_{12} & A_{22} & \cdots & A_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{1n} & A_{2n} & \cdots & A_{nn} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$
(26)

for the solution \mathbf{x} of $\mathbf{A}\mathbf{x} = \mathbf{b}$.

Observe that the double subscripts in (26) are reversed from their usual order; the element in the *i*th row and *j*th column is A_{ji} (rather than A_{ij}). We therefore see in (26) the *transpose* of the **cofactor matrix** $[A_{ij}]$ of the $n \times n$ matrix **A**. The transpose of the cofactor matrix of **A** is called the **adjoint matrix** of **A** and is denoted by

$$\operatorname{adj} \mathbf{A} = \left[A_{ij} \right]^T = \left[A_{ji} \right].$$
(27)

With the aid of this notation, Cramer's rule as expressed in Eq. (26) can be written in the especially simple form

$$\mathbf{x} = \frac{\left[A_{ji}\right]\mathbf{b}}{|\mathbf{A}|} = \frac{(\mathrm{adj}\,\mathbf{A})\mathbf{b}}{|\mathbf{A}|}.$$
(28)

The fact that the formula in (28) gives the unique solution \mathbf{x} of $\mathbf{A}\mathbf{x} = \mathbf{b}$ implies that

$$\mathbf{A}\frac{(\mathrm{adj}\,\mathbf{A})\mathbf{b}}{|\mathbf{A}|} = \mathbf{b} \tag{29}$$

for every *n*-vector **b**. If we write

$$\mathbf{C} = \frac{\mathrm{adj}\,\mathbf{A}}{|\mathbf{A}|} \tag{30}$$

for brevity, then

$$\mathbf{ACb} = \mathbf{b} \tag{31}$$

for every *n*-vector **b**. From this it follows (one column at a time, as we use Fact 2 in Section 3.5) that

$$\mathbf{ACB} = \mathbf{B} \tag{32}$$

for every matrix **B** having *n* rows. In particular, with $\mathbf{B} = \mathbf{I}$ (the $n \times n$ identity matrix), we see that

$$\mathbf{AC} = \mathbf{I}.\tag{33}$$

Therefore, we have discovered that the matrix C as defined in Eq. (30) is the inverse matrix A^{-1} of A.

THEOREM 5 The Inverse Matrix

The inverse of the invertible matrix A is given by the formula

$$\mathbf{A}^{-1} = \frac{\operatorname{adj} \mathbf{A}}{|\mathbf{A}|} = \frac{\left[a_{ij}\right]^{T}}{|\mathbf{A}|},$$
(34)

where, as usual, A_{ij} is the *ij*th cofactor of **A**—that is, A_{ij} is the product of $(-1)^{i+j}$ and the *ij*th minor of **A**.

ANSWERS TO SELECTED PROBLEMS

Chapter 1

Section 1.1

- 11. If $y = y_1 = x^{-2}$, then $y'(x) = -2x^{-3}$ and $y''(x) = 6x^{-4}$, so $x^2y'' + 5xy' + 4y = x^2(6x^{-4}) + 5x(-2x^{-3}) + 4(x^{-2}) = 6x^{-2} 10x^{-2} + 4x^{-2} = 0$. If $y = y_2 = x^{-2} \ln x$, then $y'(x) = x^{-3} 2x^{-3} \ln x$ and $y''(x) = -5x^{-4} + 6x^{-4} \ln x$, so $x^2y'' + 5xy' + 4y = x^2(-5x^{-4} + 6x^{-4} \ln x) + 5x(x^{-3} 2x^{-3} \ln x) + 4(x^{-2} \ln x) = 0$.
- **13.** $r = \frac{2}{3}$ **14.** $r = \pm \frac{1}{2}$ **15.** r = -2, 1

16.
$$r = \frac{1}{6}(-3 \pm \sqrt{57})$$

17.
$$C = 2$$







19. C = 6







21. C = 7





23. C = -56

22. C = 1



24. C = 17



25. $C = \pi/4$







(b) The graphs (figure below) of typical solutions with $k = -\frac{1}{2}$ suggest that now the value x(t) approaches 0 as t increases without bound.



- **45.** P(t) = 100/(50 t); P = 100 when t = 49, and P = 1000 when t = 49.9. Thus it appears that P(t) grows without bound as t approaches 50.
- **46.** v(t) = 50/(5+2t); v = 1 when t = 22.5, and $v = \frac{1}{10}$ when t = 247.5. Thus it appears that v(t) approaches 0 as t increases without bound.

47. (a) C = 10.1; (b) No such C, but the constant function $y(x) \equiv 0$ satisfies the conditions $y' = y^2$ and y(0) = 0.

Section 1.2

1. $y(x) = x^2 + x + 3$ 3. $y(x) = \frac{1}{3}(2x^{3/2} - 16)$ 5. $y(x) = 2\sqrt{x+2} - 5$ 6. $y(x) = \frac{1}{3}[(x^2 + 9)^{3/2} - 125]$ 7. $y(x) = 10 \tan^{-1} x$ 8. $y(x) = \frac{1}{2} \sin 2x + 1$ 9. $y(x) = \sin^{-1} x$ 10. $y(x) = -(x + 1)e^{-x} + 2$ 11. $x(t) = 25t^2 + 10t + 20$ 12. $x(t) = -10t^2 - 15t + 5$ 13. $x(t) = \frac{1}{2}t^3 + 5t$ 14. $x(t) = \frac{1}{3}t^3 + \frac{1}{2}t^2 - 7t + 4$ 15. $x(t) = \frac{1}{3}(t + 3)^4 - 37t - 26$ 16. $x(t) = \frac{4}{3}(t + 4)^{3/2} - 5t - \frac{29}{3}$ 17. $x(t) = \frac{1}{2}[(t + 1)^{-1} + t - 1]$ 19. $x(t) = \begin{cases} 5t & \text{if } 0 \le t \le 5, \\ 10t - \frac{1}{2}t^2 - \frac{25}{2} & \text{if } 5 \le t \le 10. \end{cases}$



20.
$$x(t) = \begin{cases} \frac{1}{2}t^2 & \text{if } 0 \le t \le 5, \\ 5t - \frac{25}{2} & \text{if } 5 \le t \le 10. \end{cases}$$





- 23. v(t) = -(9.8)t + 49, so the ball reaches its maximum height (v = 0) after t = 5 seconds. Its maximum height then is y(5) = 122.5 (m).
- **24.** v(5) = -160 ft/s
- **25.** The car stops when $t \approx 2.78$ (s), so the distance traveled before stopping is approximately $x(2.78) \approx 38.58$ (m).
- **26.** (a) $y \approx 530$ m (b) $t \approx 20.41$ s (c) $t \approx 20.61$ s
- **27.** $y_0 \approx 178.57$ (m)
- **28.** $v(4.77) \approx -192.64$ ft/s
- **29.** After 10 seconds the car has traveled 200 ft and is traveling at 70 ft/s.
- **30.** a = 22 ft/s²; it skids for 4 seconds.
- **31.** $v_0 = 10\sqrt{30}$ (m/s), about 197.18 km/h
- **32.** 60 m **33.** $20\sqrt{10} \approx 63.25$ (ft/s)
- **34.** 460.8 ft **36.** About 13.6 ft
- **37.** 25 (mi) **38.** 1:10 pm
- **39.** 6 mph **40.** 2.4 mi
- **41.** $\frac{544}{3} \approx 181.33$ ft/s **42.** 25 mi
- **43.** Time: 6.12245×10^9 s ≈ 194 years; Distance: 1.8367×10^{17} m ≈ 19.4 light-years
- **44.** About 54 mi/h



















- 11. A unique solution exists in some neighborhood of x = 1.
- 12. A unique solution exists in some neighborhood of x = 1.
- **13.** A unique solution exists in some neighborhood of x = 0.
- 14. Existence but not uniqueness is guaranteed in some neighborhood of x = 0.
- **15.** Neither existence nor uniqueness is guaranteed in any neighborhood of x = 2.
- 16. A unique solution exists in some neighborhood of x = 2.
- 17. A unique solution exists in some neighborhood of x = 0.
- 18. Neither existence nor uniqueness is guaranteed.
- **19.** A unique solution exists in some neighborhood of x = 0.
- **20.** A unique solution exists in some neighborhood of x = 0.
- **21.** Your figure should suggest that $y(-4) \approx 3$; an exact solution of the differential equation gives $y(-4) = 3 + e^{-4} \approx 3.0183$.



22.
$$y(-4) \approx -3$$

ANSWERS TO SELECTED PROBLEM 681

23. Your figure should suggest that $y(2) \approx 1$; the actual value is closer to 1.004.



24. $y(2) \approx 1.5$

25. Your figure should suggest that the limiting velocity is about 20 ft/sec (quite survivable) and that the time required to reach 19 ft/sec is a little less than 2 seconds. An exact solution gives v(t) = 19 when $t = \frac{5}{8} \ln 20 \approx 1.8723$.



- **26.** A figure suggests that there are 40 deer after about 60 months; a more accurate value is $t \approx 61.61$. The limiting population is 75 deer.
- 27. The initial value problem $y' = 2\sqrt{y}$, y(0) = b has no solution if b < 0; a unique solution if b > 0; infinitely many solutions if b = 0.



- **28.** The initial value problem xy' = y, y(a) = b has a unique solution if $a \neq 0$; infinitely many solutions if a = b = 0; no solution if a = 0 but $b \neq 0$.
- **29.** The initial value problem $y' = 3y^{2/3}$, y(a) = b always has infinitely many solutions defined for all *x*. However, if $b \neq 0$

then it has a unique solution near x = a.



- **30.** The initial value problem $y' = -\sqrt{1-y^2}$, y(a) = b has a unique solution if |b| < 1; no solution if |b| > 1, and infinitely many solutions (defined for all x) if $b = \pm 1$.
- **31.** The initial value problem $y' = \sqrt{1 y^2}$, y(a) = b has a unique solution if |b| < 1; no solution if |b| > 1, and infinitely many solutions (defined for all x) if $b = \pm 1$.



- **32.** The initial value problem $y' = 4x\sqrt{y}$, y(a) = b has infinitely many solutions (defined for all x) if $b \ge 0$; no solutions if b < 0. However, if b > 0 then it has a unique solution near x = a.
- **33.** The initial value problem $x^2y' + y^2 = 0$, y(a) = b has a unique solution with initial point (a, b) if $a \neq 0$, no solution if a = 0 but $b \neq 0$, infinitely many solutions if a = b = 0.



34. (a) If y(-1) = -1.2 then $y(1) \approx -0.48$. If y(-1) = -0.8 then $y(1) \approx 2.48$. (b) If y(-3) = -3.01 then $y(3) \approx -1.0343$.

If
$$y(-3) = -2.99$$
 then $y(3) \approx 7.0343$.

35. (a) If y(-3) = -0.2 then $y(2) \approx 2.019$. If y(-3) = +0.2 then $y(2) \approx 2.022$. In either case, $y(2) \approx 2.02$. (b) If $y(-3) \approx -0.5$ then $y(2) \approx 2.017$. If $y(-3) \approx +0.5$ then $y(2) \approx 2.024$. In either case, $y(2) \approx 2.02$.

Section 1.4

1.
$$y(x) = C \exp(-x^2)$$

3. $y(x) = C \exp(-\cos x)$
5. $y(x) = \sin(C + \sqrt{x})$
7. $y(x) = (2x^{4/3} + C)^{3/2}$
9. $y(x) = C(1 + x)/(1 - x)$
10. $y(x) = (1 + x)/[1 + C(1 + x)] - 1$
11. $y(x) = (C - x^2)^{-1/2}$
12. $y^2 + 1 = Ce^{x^2}$
13. $\ln(y^4 + 1) = C + 4 \sin x$
14. $3y + 2y^{3/2} = 3x + 2x^{3/2} + C$
15. $1/(3y^3) - 2/y = 1/x + \ln |x| + C$
16. $y(x) = \sec^{-1}(C\sqrt{1 + x^2})$
17. $\ln |1 + y| = x + \frac{1}{2}x^2 + C$
18. $y(x) = \tan\left(C - \frac{1}{x} - x\right)$
19. $y(x) = 2\exp(e^x)$
20. $y(x) = \tan(x^3 + \pi/4)$
21. $y^2 = 1 + \sqrt{x^2 - 16}$
22. $y(x) = -3\exp(x^4 - x)$
23. $y(x) = \frac{1}{2}(1 + e^{2x-2})$
24. $y(x) = \frac{\pi}{2}\sin x$
25. $y(x) = x\exp(x^2 - 1)$
26. $y(x) = \tan^{-1}(\sqrt{x} - 1)$

29. (a) General solution y(x) = -1/(x - C); (b) The singular solution $y(x) \equiv 0$. (c) In the following figure we see that there is a unique solution through every point of the xy-plane.



- 30. General solution y(x) = (x C)²; singular solution y(x) ≡ 0.
 (a) No solution if b < 0; (b) Infinitely many solutions (for all x) if b ≥ 0; (c) Two solutions near (a, b) if b > 0.
- **31.** Separation of variables gives the same general solution $y = (x C)^2$ as in Problem 30, but the restriction that $y' = 2\sqrt{y} \ge 0$ implies that only the *right halves* of the parabolas qualify as solution curves. In the figure below we see that through the point (a, b) there passes (a) No solution curve if b < 0, (b) a unique solution curve if b > 0, (c) Infinitely many

solution curves if b = 0.



32. General solution $y(x) = \pm \sec(x - C)$; singular solutions $y(x) \equiv \pm 1$.

(a) No solution if |b| < 1;
(b) A unique solution if |b| > 1;
(c) Infinitely many solutions if b = ±1.

36. Age about 686 years

40. About 35 years

38. \$44.52

- **33.** About 51840 persons **34.** $t \approx 3.87$ hr
- **35.** About 14735 years

37. \$21103.48

- **39.** 2585 mg
- **41.** About 4.86×10^9 years ago
- 42. About 1.25 billion years
- 43. After a total of about 63 min have elapsed
- 44. About 2.41 minutes
- **45.** (a) 0.495 m; (b $(8.32 \times 10^{-7})I_0$; (c) 3.29 m
- 46. (a) About 9.60 inches; (b) About 18,200 ft
- **47.** After about 46 days
- **48.** About 6 billion years
- **49.** After about 66 min 40 s
- **50.** (a) $A(t) = 10 \cdot 3^{2t/15}$; (b) About 20.80 pu; (c) About 15.72 years

55. 972 s

- **51.** (a) $A(t) = 15 \cdot (\frac{2}{3})^{t/5}$; (b) approximately 7.84 su; (c) After about 33.4 months
- **52.** About 120 thousand years ago
- **53.** About 74 thousand years ago
- **54.** 3 hours
- **56.** At time $t = 2048/1562 \approx 1.31$ (in hours)
- 58. 1:20 P.M.
- **59.** (a) $y(t) = (8-7t)^{2/3}$; (b) at 1:08:34 P.M.; (c) $r = \frac{1}{60}\sqrt{\frac{7}{12}} \approx 0.15$ (in.)
- 60. About 6 min 3 sec
- 61. Approximately 14 min 29 s
- 62. The tank is empty about 14 seconds after 2:00 P.M.
- **63.** (a) 1:53:34 P.M.; (b) $r \approx 0.04442$ ft ≈ 0.53 in.
- 64. $r = \frac{1}{720}\sqrt{3}$ ft, about $\frac{1}{35}$ in.
- 65. At approximately 10:29 A.M.

Section 1.5

1.
$$y(x) = 2(1 - e^{-x})$$
 2. $y(x) = (3x + C)e^{2x}$

 3. $y(x) = e^{-3x}(x^2 + C)$
 4. $y(x) = (x + C)e^{x^2}$

 5. $y(x) = x + 4x^{-2}$
 6. $y(x) = x^2 + 32/x^5$

 7. $y(x) = 5x^{1/2} + Cx^{-1/2}$
 8. $y(x) = 3x + Cx^{-1/3}$

 9. $y(x) = x(7 + \ln x)$
 10. $y(x) = 3x^3 + Cx^{-3/2}$

 11. $y(x) \equiv 0$
 12. $y(x) = \frac{1}{4}x^5 - 56x^{-3}$

ANSWERS TO SELECTED PROBLEM 683

13.	$y(x) = (e^x + e^{-x})/2$ 14. $y(x) = x^3 \ln x + 10x^3$
15.	$y(x) = [1 - 5\exp(-x^2)]/2$
16.	$y(x) = 1 + e^{-\sin x}$
17.	$y(x) = (1 + \sin x)/(1 + x)$
18.	$y(x) = x^2(\sin x + C)$
19.	$y(x) = \frac{1}{2}\sin x + C\csc x$
20.	$y(x) = -1 + \exp\left(x + \frac{1}{2}x^2\right)$
21.	$y(x) = x^3 \sin x$
22.	$y(x) = (x^3 + 5)e^{x^2}$
23.	$y(x) = x^3(2 + Ce^{-2x})$
24.	$y(x) = \frac{1}{3} [1 + 16(x^2 + 4)^{-3/2}]$
25.	$y(x) = \left[\exp\left(-\frac{3}{2}x^2\right)\right] \left[3(x^2+1)^{3/2}-2\right]$
26.	$x(y) = 1/2y^2 + C/y^4$
27.	$x(y) = e^{y} \left(C + \frac{1}{2} y^{2} \right)$
28.	$x(y) = \frac{1}{2}[y + (1 + y^2)(\tan^{-1} y + C)]$
30.	$y(x) = x^{1/2} \int_{1}^{x} t^{-1/2} \cos t dt$
29.	$y(x) = \left[\exp(x^2)\right] \left[C + \frac{1}{2}\sqrt{\pi}\operatorname{erf}(x)\right]$
32.	(a) $y(x) = \sin x - \cos x$; (b) $y(x) = Ce^{-x} + \sin x - \cos x$;
	(c) $y(x) = 2e^{-x} + \sin x - \cos x$
33.	After about 7 min 41 s
34.	About 22.2 days
35.	About 5.5452 years
36.	(a) $x(t) = (60 - t) - (60 - t)^3/3600$; (b) About 23.09 lb
37.	393.75 lb
38.	(a) $x(t) = 50e^{-t/20}$; (b) $y(t) = 150e^{-t/40} - 100e^{-t/20}$;
	(c) 56.25 lb
39.	(b) $y_{\text{max}} = 100e^{-1} \approx 36.79$ (gal)

- **41.** (b) Approximately \$1,308,283
- 43. -50.0529, -28.0265, -6.0000, 16.0265, 38.0529
- 44. 3.99982, 4.00005, 4.00027, 4.00050, 4.00073
- **45.** $x(t) = 20(1 e^{-t/10}); x = 10$ after $t = 10 \ln 2 \approx 6.93$ months.
- **46.** $x(t) = \frac{20}{101}(101 102e^{-t/10} + \cos t + 10\sin t); x = 10$ after t = 6.47 months.

Section 1.6

2. $y^2 = x^2(\ln x + C)$ 1. $x^2 - 2xy - y^2 = C$ 3. $y(x) = x (C + \ln |x|)^2$ 4. $2\tan^{-1}(y/x) - \ln(y^2/x^2 + 1) = 2\ln x + C$ 5. $\ln |xy| = C + xy^{-1}$ 6. $2y \ln y = x + Cy$ 7. $y^3 = 3x^3 (C + \ln |x|)$ 8. $y = -x \ln(C - \ln x)$ 9. $y(x) = x/(C - \ln |x|)$ 10. $x^2 + 2y^2 = Cx^6$ 11. $y = C(x^2 + y^2)$ 12. $4x^2 + y^2 = x^2(\ln x + C)^2$ 13. $y + \sqrt{x^2 + y^2} = Cx^2$ 14. $x - \sqrt{x^2 + y^2} = C$ 15. $x^2(2xy + y^2) = C$ 16. $x = 2\sqrt{x+y+1} - 2\ln(1+\sqrt{x+y+1}) + C$ 17. $y(x) = -4x + 2\tan(2x + C)$ 18. $y = \ln(x + y + 1) + C$ 19. $y^2 = x/(2 + Cx^5)$ 20. $y^3 = 3 + Ce^{-3x^2}$ 21. $y^2 = 1/(Ce^{-2x} - 1)$ 22. $y^3 = 7x/(7Cx^7 + 15)$ 23. $y(x) = (x + Cx^2)^{-3}$ 24. $y^2 = e^{2x}/(C + \ln x)$ 25. $2x^3y^3 = 3\sqrt{1+x^4} + C$ 26. $y^3 = e^{-x}(x+C)$ 27. $y(x) = (x^4 + Cx)^{1/3}$ 28. $y = \ln(Cx^2 + x^2e^{2x})$ **29.** $\sin^2 y = 4x^2 + Cx$ **30.** $x^2 - 2xe^y - e^{2y} = C$ 32. $2x^2 - xy + 3y^2 = C$ 31. $x^2 + 3xy + y^2 = C$ 33. $x^3 + 2xy^2 + 2y^3 = C$ 34. $x^3 + x^2y^2 + y^4 = C$ 35. $3x^4 + 4y^3 + 12y \ln x = C$ 36. $x + e^{xy} + y^2 = C$ **37.** $\sin x + x \ln y + e^y = C$ **38.** $x^2 + 2x \tan^{-1} y + \ln(1 + y^2) = C$

39. $5x^3y^3 + 5xy^4 + y^5 = C$ **40.** $e^x \sin y + x \tan y = C$ 41. $x^2y^{-1} + y^2x^{-3} + 2y^{1/2} = C$ 42. $xy^{-2/3} + x^{-3/2}y = C$ 44. $x(y) = Ay^2 + B$ 43. $y(x) = Ax^2 + B$ 46. $y(x) = x^2 + A \ln x + B$ 45. $y(x) = A \cos 2x + B \sin 2x$ 47. $y(x) = A - \ln|x + B|$ 48. $y(x) = \ln x + Ax^{-2} + B$ **49.** $y(x) = \pm (A + Be^x)^{1/2}$ 50. $y(x) = \ln|\sec(x+A)| - \frac{1}{2}x^2 + B$ 51. $x(y) = -\frac{1}{3}(y^3 + Ay + B)$ 52. $Ay^2 - (Ax + B)^2 = 1$ 54. $Av^2(B-x) = 1$ 53. $y(x) = A \tan(Ax + B)$ 58. $y = \exp(x^2 + C/x^2)$ **59.** $x^2 - 2xy - y^2 - 2x - 6y = C$ 60. $(x + 3y + 3)^5 = C(y - x - 5)$ 61. $x = \tan(x - y) + \sec(x - y) + C$ 64. $y(x) = x + e^{-x^2} \left[C + \frac{1}{2} \sqrt{\pi} \operatorname{erf}(x) \right]^{-1}$ 65. $y(x) = x + (C - x)^{-1}$ 69. Approximately 3.68 mi

Chapter 1 Review Problems

1. Linear: $y(x) = x^3(C + \ln x)$ 2. Separable: $y(x) = x/(3 - Cx - x \ln x)$ 3. Homogeneous: $y(x) = x/(C - \ln x)$ 4. Exact: $x^2 y^3 + e^x - \cos y = C$ 5. Separable: $y(x) = C \exp(x^{-3} - x^{-2})$ 6. Separable: $y(x) = x/(1 + Cx + 2x \ln x)$ 7. Linear: $y(x) = x^{-2}(C + \ln x)$ 8. Homogeneous: $y(x) = 3Cx/(C - x^3) = 3x/(1 + Kx^3)$ 9. Bernoulli: $y(x) = (x^2 + Cx^{-1})^2$ 10. Separable: $y(x) = \tan(C + x + \frac{1}{3}x^3)$ 11. Homogeneous: $y(x) = x/(C - 3\ln x)$ 12. Exact: $3x^2y^3 + 2xy^4 = C$ 13. Separable: $y(x) = 1/(C + 2x^2 - x^5)$ 14. Homogeneous: $y^2 = x^2/(C + 2 \ln x)$ 15. Linear: $y(x) = (x^3 + C)e^{-3x}$ **16.** Substitution: v = y - x; solution: $y - x - 1 = Ce^{2\tilde{x}}(y - x + 1)$ 17. Exact: $e^{x} + e^{y} + e^{xy} = C$ **18.** Homogeneous: $y^2 = Cx^2(x^2 - y^2)$ **19.** Separable: $y(x) = x^2/(x^5 + Cx^2 + 1)$ **20.** Linear: $y(x) = 2x^{-3/2} + Cx^{-3}$ **21.** Linear: $y(x) = [C + \ln(x - 1)]/(x + 1)$ 22. Bernoulli: $y(x) = (2x^4 + Cx^2)^3$ **23.** Exact: $xe^{y} + y\sin x = C$ 24. Separable: $y(x) = \frac{x^{1/2}}{(6x^2 + Cx^{1/2} + 2)}$ **25.** Linear: $y(x) = (x + 1)^{-2}(x^3 + 3x^2 + 3x + C)$ $= x + 1 + K(x + 1)^{-2}$ **26.** Exact: $3x^{3/2}y^{4/3} - 5x^{6/5}y^{3/2} = C$ 27. Bernoulli: $y(x) = x^{-1}(C + \ln x)^{-1/3}$ **28.** Linear: $y(x) = x^{-1}(C + e^{2x})$ **29.** Linear: $y(x) = (x^2 + x + C)(2x + 1)^{-1/2}$ **30.** Substitution: v = x + y; solution: $x = 2(x + y)^{1/2} - 2\ln[1 + (x + y)^{1/2}] + C$ 31. Separable and linear 32. Separable and Bernoulli 33. Exact and homogeneous 34. Exact and homogeneous 35. Separable and linear 36. Separable and Bernoulli

Chapter 2

Section 2.1

1.
$$x(t) = \frac{2}{2 - e^{-t}}$$



$$2. \ x(t) = \frac{10}{1 + 9e^{-10t}}$$



3.
$$x(t) = \frac{2 + e^{-2t}}{2 - e^{-2t}}$$





ANSWERS TO SELECTED PROBLEM685



- **26.** $50 \ln \frac{9}{8} \approx 5.89$ months
- 27. (a) $100 \ln \frac{9}{5} \approx 58.78$ months; (b) $100 \ln 2 \approx 69.31$ months
- **28.** (a) The alligators eventually die out. (b) Doomsday occurs after about 9 years 2 months.
- (a) P(140) ≈ 127.008 million; (b) About 210.544 million;
 (c) In 2000 we get P ≈ 196.169, whereas the actual 2000 population was about 281.422 million.
- **31.** $\alpha \approx 0.3915$; 2.15 × 10⁶ cells
- **37.** $k \approx 0.0000668717$, $M \approx 338.027$
- **38.** $k \approx 0.000146679$, $M \approx 208.250$
- **39.** $P(t) = P_0 \exp\left(kt + \frac{b}{2\pi}\sin 2\pi t\right)$; the colored curve in the figure below shows the graph with $P_0 = 100$, k = 0.03, and b = 0.06. It oscillates about the black curve which represents natural growth with $P_0 = 100$ and k = 0.03. We see that the two agree at the end of each full year.



Section 2.2

1. Unstable critical point: x = 4;

$$x(t) = 4 + (x_0 - 4)e^t$$



4. Stable critical point: x = 3; unstable critical point: x = 0; $3x_0$



5. Stable critical point: x = -2; unstable critical point: x = 2;



6. Stable critical point: x = 3; unstable critical point: x = -3;







3. Stable critical point: x = 0; unstable critical point: x = 4;



ANSWERS TO SELECTED PROBLEM 687

7. Semi-stable (see Problem 18) critical point: x = 2;



8. Semi-stable critical point: x = 3;



9. Stable critical point: x = 1; unstable critical point: x = 4;



10. Stable critical point: x = 5; unstable critical point: x = 2;



11. Unstable critical point: x = 1;





12. Stable critical point: x = 2;

$$\frac{1}{(2-x(t))^2} = \frac{1}{(2-x_0)^2} + 2t$$



For each of Problems 13 through 18 we show a plot of slope field and typical solution curves. The equilibrium solutions of the given differential equation are labeled, and the stability or instability of each should be clear from the picture.







- **19.** There are two critical points if $h < 2\frac{1}{2}$, one critical point if $h = 2\frac{1}{2}$, and no critical points if $h > 2\frac{1}{2}$. The bifurcation diagram is the parabola $(c-5)^2 = 25 - 10h$ in the *hc*-plane.
- **20.** There are two critical points if $s < \frac{1}{16}$, one critical point if $s = \frac{1}{16}$, and no critical points if $s > \frac{1}{16}$. The bifurcation diagram is the parabola $(2c - 5)^2 = 25(1 - 16s)$ in the *sc*-plane.

Section 2.3

- 1. Approximately 31.5 s
- 3. $400/(\ln 2) \approx 577$ ft
- 5. 400 ln 7 \approx 778 ft
- 7. (a) 100 ft/sec; (b) about 23 sec and 1403 ft to reach 90 ft/sec
- 8. (a) 100 ft/sec; (b) about 14.7 sec and 830 ft to reach 90 ft/sec
- 9. 50 ft/s
- 10. About 5 min 47 s
- 11. Time of fall: about 12.5 s
- 12. Approximately 648 ft
- 19. Approximately 30.46 ft/s; exactly 40 ft/s
- 20. Approximately 277.26 ft
- 22. Approximately 20.67 ft/s; about 484.57 s
- 23. Approximately 259.304 s
- 24. (a) About 0.88 cm; (b) about 2.91 km
- 25. (b) About 1.389 km/sec; (c) $r_{\text{max}} = 100R/19 \approx 5.26R$
- 26. Yes
- **28.** (b) After about $8\frac{1}{2}$ minutes it hits the surface at about 4.116 km/sec.
- 29. About 51.427 km
- 30. Approximately 11.11 km/sec (as compared with the earth's escape velocity of about 11.18 km/sec).

ANSWERS TO SELECTED PROBLEM 689

Section 2.4

In Problems 1 through 10 we round off the indicated values to 3 decimal places.

- 1. Approximate values 1.125 and 1.181; true value 1.213
- 2. Approximate values 1.125 and 1.244; true value 1.359
- 3. Approximate values 2.125 and 2.221; true value 2.297
- 4. Approximate values 0.625 and 0.681; true value 0.713
- 5. Approximate values 0.938 and 0.889; true value 0.851
- **6.** Approximate values 1.750 and 1.627; true value 1.558
- Approximate values 2.859 and 2.737; true value 2.647
 Approximate values 0.445 and 0.420; true value 0.405
- **9.** Approximate values 1.267 and 1.278; true value 1.287
- **10.** Approximate values 1.125 and 1.231; true value 1.333

Problems 11 through 24 call for tables of values that would occupy too much space for inclusion here. In Problems 11 through 16 we give first the final *x*-value, next the corresponding approximate *y*-values obtained with step sizes h = 0.01 and h = 0.005, and then the final true *y*-value. (All *y*-values rounded off accurate to 4 decimal places.)

- **11.** 1.0, -0.7048, -0.7115, -0.7183 **12.** 1.0, 2.9864, 2.9931, 3.0000 **13.** 2.0, 4.8890, 4.8940, 4.8990 **14.** 2.0, 3.2031, 3.2304, 3.2589 **15.** 3.0, 3.4422, 3.4433, 3.4444
- **16.** 3.0, 8.8440, 8.8445, 8.8451

In Problems 17 through 24 we give first the final *x*-value and then the corresponding approximate *y*-values obtained with step sizes h = 0.1, h = 0.02, h = 0.004, and h = 0.0008 respectively. (All *y*-values rounded off accurate to 4 decimal places.)

- 17. 1.0, 0.2925, 0.3379, 0.3477, 0.3497
- **18.** 2.0, 1.6680, 1.6771, 1.6790, 1.6794
- 19. 2.0, 6.1831, 6.3653, 6.4022, 6.4096
- 20. 2.0, -1.3792, -1.2843, -1.2649, -1.2610
- 21. 2.0, 2.8508, 2.8681, 2.8716, 2.8723
- **22.** 2.0, 6.9879, 7.2601, 7.3154, 7.3264
- **23.** 1.0, 1.2262, 1.2300, 1.2306, 1.2307
- **24.** 1.0, 0.9585, 0.9918, 0.9984, 0.9997
- **25.** With both step sizes h = 0.01 and h = 0.005, the approximate velocity after 1 second is 16.0 ft/sec (80% of the limiting velocity of 20 ft/sec); after 2 seconds it is 19.2 ft/sec (96% of the limiting velocity).
- **26.** With both step sizes h = 1 and h = 0.5, the approximate population after 5 years is 49 deer (65% of the limiting population of 75 deer); after 10 years it is 66 deer (88% of the limiting population).
- 27. With successive step sizes $h = 0.1, 0.01, 0.001, \ldots$ the first four approximations to y(2) we obtain are 0.7772, 0.9777, 1.0017, and 1.0042. It therefore seems likely that $y(2) \approx 1.00$.
- **28.** With successive step sizes h = 0.1, 0.01, 0.001, ... the first four approximations to y(2) we obtain are 1.2900, 1.4435, 1.4(21) with the formula of the state of t
- 1.4613, and 1.4631. It therefore seems likely that $y(2) \approx 1.46$.

	h = 0.15	h = 0.03	h = 0.006
x	У	У	У
-1.0	1.0000	1.0000	1.0000
-0.7	1.0472	1.0512	1.0521
-0.4	1.1213	1.1358	1.1390
-0.1	1.2826	1.3612	1.3835
0.2	0.8900	1.4711	0.8210
0.5	0.7460	1.2808	0.7192

30.			
	x	$\begin{array}{c} h = 0.1 \\ y \end{array}$	h = 0.01
	1.8	2.8200	4.3308
	1.9	3.9393	7.9425
	2.0	5.8521	28.3926
31.			
31.		h = 0.1	h = 0.01
31.	x	h = 0.1	h = 0.01
31.	x 0.7	<i>h</i> = 0.1 <i>y</i> 4.3460	h = 0.01 y 6.4643
31.	x 0.7 0.8	<i>h</i> = 0.1 <i>y</i> 4.3460 5.8670	h = 0.01 y 6.4643 11.8425
31.	x 0.7 0.8 0.9	h = 0.1 y 4.3460 5.8670 8.3349	<i>h</i> = 0.01 <i>y</i> 6.4643 11.8425 39.5010

Section 2.5

1

x	Improved Euler y	Actual y
0.1	1.8100	1.8097
0.2	1.6381	1.6375
0.3	1.4824	1.4816
0.4	1.3416	1.3406
0.5	1.2142	1.2131

Note: In Problems 2 through 10, we give the value of x, the corresponding improved Euler value of y, and the true value of y.

2.	0.5, 1.3514, 1.3191	3. 0.5, 2.2949, 2.2974
4.	0.5, 0.7142, 0.7131	5. 0.5, 0.8526, 0.8513
6.	0.5, 1.5575, 1.5576	7. 0.5, 2.6405, 2.6475
8.	0.5, 0.4053, 0.4055	9. 0.5, 1.2873, 1.2874
10.	0.5, 1.3309, 1.3333	

In Problems 11 through 16 we give the final value of x, the corresponding values of y with h = 0.01 and with h = 0.005, and the true value of y.

10, -0.71824, -0.71827, -0.71828
 10, 2.99995, 2.99999, 3.00000
 2.0, 4.89901, 4.89899, 4.89898
 2.0, 3.25847, 3.25878, 3.25889
 3.0, 3.44445, 3.44445, 3.44444

16. 3.0, 8.84511, 8.84509, 8.84509

In Problems 17 through 24 we give the final value of x and the corresponding values of y for h = 0.1, 0.02, 0.004, and 0.0008.

- 17. 1.0, 0.35183, 0.35030, 0.35023, 0.35023
- **18.** 2.0, 1.68043, 1.67949, 1.67946, 1.67946
- 19. 2.0, 6.40834, 6.41134, 6.41147, 6.41147
- $\textbf{20. } 2.0, \ -1.26092, \ -1.26003, \ -1.25999, \ -1.25999$
- 21. 2.0, 2.87204, 2.87245, 2.87247, 2.87247
- **22.** 2.0, 7.31578, 7.32841, 7.32916, 7.32920
- **23.** 1.0, 1.22967, 1.23069, 1.23073, 1.23073
- 24. 1.0, 1.00006, 1.00000, 1.00000, 1.00000
- **25.** With both step sizes h = 0.01 and h = 0.005 the approximate velocity after 1 second is 15.962 ft/sec (80% of the limiting velocity of 20 ft/sec); after 2 seconds it is 19.185 ft/sec (96% of the limiting velocity).

690 ANSWERS TO SELECTED PROBLEMS

- **26.** With both step sizes h = 1 and h = 0.5 the approximate population after 5 years is 49.391 deer (65% of the limiting population of 75 deer); after 10 years it is 66.113 deer (88% of the limiting population).
- 27. With successive step sizes h = 0.1, 0.01, 0.001, ... the first three approximations to y(2) we obtain are 1.0109, 1.0045, and 1.0045. It therefore seems likely that $y(2) \approx 1.0045$.
- **28.** With successive step sizes h = 0.1, 0.01, 0.001, ... the first four approximations to y(2) we obtain are 1.4662, 1.4634, 1.4633, and 1.4633. It therefore seems likely that $y(2) \approx 1.4633$.
- 29. Impact speed approximately 43.22 m/s
- 30. Impact speed approximately 43.48 m/s

Section 2.6

1. $y(0.25) \approx 1.55762$; y(0.25) = 1.55760. $y(0.5) \approx 1.21309$; y(0.5) = 1.21306. Solution: $y = 2e^{-x}$

In Problems 2 through 10 we give the approximation to y(0.5), its true value, and the solution.

- 2. 1.35867, 1.35914; $y = \frac{1}{2}e^{2x}$ 3. 2.29740, 2.29744; $y = 2e^x - 1$ 4. 0.71309, 0.71306; $y = 2e^{-x} + x - 1$ 5. 0.85130, 0.85128; $y = -e^x + x + 2$ 6. 1.55759, 1.55760; $u = 2\exp(-x^2)$ 7. 2.64745, 2.64749; $y = 3\exp(-x^3)$ 8. 0.40547, 0.40547; $y = \ln(x + 1)$ 9. 1.28743, 1.28743; $y = \tan \frac{1}{4}(x + \pi)$ 10. 1.33337, 1.33333; $y = (1 - x^2)^{-1}$
- 11. Solution: $y(x) = 2 e^x$.

	h = 0.2	h = 0.1	Exact
x	У	У	У
0.0	1.00000	1.00000	1.00000
0.2	0.77860	0.77860	0.77860
0.4	0.50818	0.50818	0.50818
0.6	0.17789	0.17788	0.17788
0.8	-0.22552	-0.22554	-0.22554
1.0	-0.71825	-0.71828	-0.71828

In Problems 12 through 16 we give the final value of x, the corresponding Runge-Kutta approximations with h = 0.2 and with h = 0.1, the exact value of y, and the solution.

12. 1.0, 2.99996, 3.00000, 3.00000;

- y = 1 + 2/(2 x)
- **13.** 2.0, 4.89900, 4.89898, 4.89898; $y = \sqrt{8 + x^4}$
- **14.** 2.0, 3.25795, 3.25882, 3.25889; $y = 1/(1 \ln x)$
- **15.** 3.0, 3.44445, 3.44444, 3.44444; $y = x + 4x^{-2}$
- **16.** 3.0, 8.84515, 8.84509, 8.84509; $y = (x^6 - 37)^{1/3}$

In Problems 17 through 24 we give the final value of x and the corresponding values of y with h = 0.2, 0.1, 0.05, and 0.025.

- **17.** 1.0, 0.350258, 0.350234, 0.350232, 0.350232
- **18.** 2.0, 1.679513, 1.679461, 1.679459, 1.679459
- 19. 2.0, 6.411464, 6.411474, 6.411474, 6.411474

- **20.** 2.0, -1.259990, -1.259992, -1.259993, -1.259993
- **21.** 2.0, 2.872467, 2.872468, 2.872468, 2.872468
- 22. 2.0, 7.326761, 7.328452, 7.328971, 7.329134
- 23. 1.0, 1.230735, 1.230731, 1.230731, 1.230731
- 24. 1.0, 1.000000, 1.000000, 1.000000, 1.000000
- **25.** With both step sizes h = 0.1 and h = 0.05, the approximate velocity after 1 second is 15.962 ft/sec (80% of the limiting velocity of 20 ft/sec); after 2 seconds it is 19.185 ft/sec (96% of the limiting velocity).
- **26.** With both step sizes h = 6 and h = 3, the approximate population after 5 years is 49.3915 deer (65% of the limiting population of 75 deer); after 10 years it is 66.1136 deer (88% of the limiting population).
- 27. With successive step sizes h = 1, 0.1, 0.01, ... the first four approximations to y(2) we obtain are 1.05722, 1.00447, 1.00445 and 1.00445. Thus it seems likely that $y(2) \approx 1.00445$ accurate to 5 decimal places.
- **28.** With successive step sizes h = 1, 0.1, 0.01, ... the first four approximations to y(2) we obtain are 1.48990, 1.46332, 1.46331, and 1.46331. Thus it seems likely that $y(2) \approx 1.4633$ accurate to 5 decimal places.
- 29. Time aloft: approximately 9.41 seconds
- 30. Time aloft: approximately 9.41 seconds

Chapter 3

Section 3.1

- 1. x = 3, y = 22. x = 5, y = -33. x = -4, y = 34. x = 5, y = 45. Inconsistent-no solution 6. Inconsistent-no solution 7. x = -10 + 4t, y = t (infinitely many solutions) 8. x = 4 + 2t, y = t (infinitely many solutions) 9. x = 4, y = -1, z = 310. x = 3, y = 1, z = -211. x = 1, y = 3, z = -412. x = 1, y = 3, z = 513. x = y = z = 014. x = 5, y = 3, z = -415. Inconsistent-no solution 16. Inconsistent-no solution 17. Inconsistent-no solution
 - **18.** Inconsistent—no solution
 - 19. x = 8 + 3t, y = 3 + 2t, z = t (infinitely many solutions)
 - **20.** x = -5t, y = 5 + t, z = t (infinitely many solutions)
 - **21.** x = 3 2t, y = 2 + 3t, z = t (infinitely many solutions)
 - 22. x = -4t, y = -5t, z = t (infinitely many solutions)
 - 23. $y(x) = 3\cos 2x + 4\sin 2x$
 - 24. $y(x) = 5 \cosh 3x + 4 \sinh 3x$
 - 25. $y(x) = 7e^{5x} + 3e^{-5x}$
 - **26.** $y(x) = 23e^{11x} + 21e^{-11x}$
 - **27.** $y(x) = 23e^{3x} + 17e^{-5x}$
 - **28.** $y(x) = 23e^{3x} 8e^{7x}$
 - **29.** $y(x) = 52e^{x/2} 45e^{x/3}$
 - **30.** $y(x) = 81e^{4x/3} 40e^{-7x/5}$
 - **31.** *Suggestion*: The two lines both pass through the origin.
 - **32.** Suggestion: Two distinct planes in space either are parallel or intersect in a straight line.

ANSWERS TO SELECTED PROBLEM 691

- **33.** (a) No solution (b) A unique solution (c) No solution (d) No solution (e) A unique solution (f) Infinitely many solutions
- 34. (a) No solution (b) Infinitely many solutions (c) No solution (d) No solution (e) Infinitely many solutions (f) A unique solution

Section 3.2

1. $x_1 = 1, x_2 = 0, x_3 = 2$ 2. $x_1 = 5, x_2 = 1, x_3 = -3$ 3. $x_1 = 13 + 11t$, $x_2 = 2 + 5t$, $x_3 = t$ 4. $x_1 = 35 + 33t$, $x_2 = 5 + 7t$, $x_3 = t$ 5. $x_1 = 13 + 4t$, $x_2 = 6 + t$, $x_3 = 5 + 3t$, $x_4 = t$ 6. $x_1 = 17 + t$, $x_2 = 11 + 3t$, $x_3 = t$, $x_4 = -4$ 7. $x_1 = 3 - 8s + 19t$, $x_2 = 7 + 2s - 7t$, $x_3 = s$, $x_4 = t$ 8. $x_1 = -25 + 10s + 22t$, $x_2 = s$, $x_3 = 10 - 3t$, $x_4 = t$ 9. $x_1 = 1, x_2 = 3, x_3 = -5, x_4 = 6$ **10.** $x_1 = 63s - 16t$, $x_2 = 13s - 8t$, $x_3 = s$, $x_4 = 5t$, $x_5 = t$ 11. $x_1 = 3, x_2 = -2, x_3 = 4$ 12. $x_1 = 5, x_2 = -3, x_3 = 2$ 13. $x_1 = 4 + 3t, x_2 = 3 - 2t, x_3 = t$ 14. $x_1 = 5 + 2t, x_2 = t, x_3 = 7$ 15. Inconsistent-no solution 16. Inconsistent—no solution 17. $x_1 = 3 - 2t, x_2 = -4 + t, x_3 = 5 - 3t, x_4 = t$ **18.** $x_1 = 4 + 2s - 3t$, $x_2 = s$, $x_3 = 3 - 4t$, $x_4 = t$ **19.** $x_1 = 3 - s - t$, $x_2 = 5 + 2s - 3t$, $x_3 = s$, $x_4 = t$ **20.** $x_1 = 2 + 3t$, $x_2 = 1 + s - 2t$, $x_3 = 2 + 2s$, $x_4 = s$, $x_5 = t$ **21.** $x_1 = 2, x_2 = 1, x_3 = 3, x_4 = 4$ **22.** $x_1 = 3, x_2 = -2, x_3 = 4, x_4 = -1$ **23.** (a) None (b) $k \neq 2$ (c) k = 2**24.** (a) $k \neq 2$ (b) None (c) k = 4**25.** (a) $k \neq 4$ (b) k = 4 (c) None **26.** (a) All k (b) None (c) None **27.** (a) None (b) $k \neq 11$ (c) k = 11

28. No solution unless c = 2a + 3b, in which case it has infinitely many solutions.

Section 3.3



	$\begin{bmatrix} 1 & 0 & 3 & -2 & 3 \end{bmatrix}$
18.	0 1 4 5 1
	0 0 0 0 0
	$\overline{1}$ 0 2 1 3 $\overline{1}$
19.	$\begin{bmatrix} 0 & 1 & -2 & -3 & 1 \end{bmatrix}$
	0 0 0 0 0
20	
20.	
21.	$x_1 = 3, x_2 = -2, x_3 = 4$
22.	$x_1 = 5, x_2 = -3, x_3 = 2$
23.	$x_1 = 4 + 3t$, $x_2 = 3 - 2t$, $x_3 = t$
24.	$x_1 = 5 + 2t$ $x_2 = t$ $x_3 = 7$
25	Inconsistent no solution
25.	Inconsistent no solution
20.	
27.	$x_1 = 3 - 2t, x_2 = -4 + t, x_3 = 5 - 3t, x_4 = t$
28.	$x_1 = 4 + 2s - 3t, x_2 = s, x_3 = 3 - 4t, x_4 = t$
29.	$x_1 = 3 - s - t, x_2 = 5 + 2s - 3t, x_3 = s, x_4 = t$
30.	$x_1 = 2 + 3t, x_2 = 1 + s - 2t, x_3 = 2 + 2s, x_4 = s, x_5 = t$
31.	The sequence $\frac{1}{6}R_3$, $R_2 - 5R_3$, $\frac{1}{4}R_2$, $R_1 - 2R_2$, $R_1 - 3R_3$ of
	row operations transforms the first matrix to the identity matrix.
22	$\begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & * \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 \end{bmatrix}$
33.	

Section 3.4

1.
$$\begin{bmatrix} 5 & -15 \\ 18 & 5 \end{bmatrix}$$

3. $\begin{bmatrix} -26 & 20 \\ 12 & -6 \\ 22 & 18 \end{bmatrix}$
5. $AB = \begin{bmatrix} -9 & 1 \\ -10 & 12 \end{bmatrix}, BA = \begin{bmatrix} -2 & 8 \\ 11 & 5 \end{bmatrix}$
6. $AB = \begin{bmatrix} 7 & -13 & -24 \\ 23 & 10 & 41 \\ 11 & -8 & 57 \end{bmatrix}, BA = \begin{bmatrix} 1 & -17 & -22 \\ 12 & 16 & 7 \\ 27 & -21 & 57 \end{bmatrix}$
7. $AB = \begin{bmatrix} 26 \end{bmatrix}, BA = \begin{bmatrix} 3 & 6 & 9 \\ 4 & 8 & 12 \\ 5 & 10 & 15 \end{bmatrix}$
8. $AB = \begin{bmatrix} 21 & 15 \\ 35 & 0 \end{bmatrix}, BA = \begin{bmatrix} 3 & 0 & 9 \\ 7 & -20 & 13 \\ 16 & -25 & 38 \end{bmatrix}$
9. $BA = \begin{bmatrix} 4 \\ 7 \\ -22 \end{bmatrix}$ but AB is not defined.
10. $AB = \begin{bmatrix} 1 & -2 & 13 \\ 5 & -6 & 31 \end{bmatrix}$ but BA is not defined.
11. $AB = \begin{bmatrix} 11 & 1 & 5 & 3 \end{bmatrix}$ but BA is not defined.
12. Neither product matrix AB nor BA is defined.
13. $A(BC) = (AB)C = \begin{bmatrix} 32 & 51 \\ -2 & -17 \end{bmatrix}$
14. $A(BC) = (AB)C = \begin{bmatrix} -3 \\ 12 & 15 \\ 8 & 10 \end{bmatrix}$
15. $A(BC) = (AB)C = \begin{bmatrix} -4 & -4 & -2 & 2 \\ -9 & -12 & -9 & 12 \\ -14 & -18 & -13 & 17 \end{bmatrix}$
16. $A(BC) = (AB)C = \begin{bmatrix} -4 & -4 & -2 & 2 \\ -9 & -12 & -9 & 12 \\ -14 & -18 & -13 & 17 \end{bmatrix}$
17. $x = s(5, -2, 1, 0) + t(-4, 7, 0, 1)$
18. $x = s(3, 1, 0, 0) + t(-6, 0, -9, 1)$
19. $x = s(-3, 2, -1, 1, 0) + t(1, -6, 8, 0, 1)$
20. $x = r(1, -2, 1, 0, 0) + s(-2, 3, 0, 1, 0) + t(-7, -4, 0, 0, 1)$
21. $x = r(1, -2, 1, 0, 0) + s(-7, 0, 2, 10, 0)$
21. $x = r(1, -2, -1, 0, 0) + s(-7, 0, 1, 1, 0) + t(-7, -4, 0, 0, 1)$

23.
$$\mathbf{B} = \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix}$$
24.
$$\mathbf{B} = \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix}$$
25.
$$\mathbf{B} = \begin{bmatrix} 3 & -7 \\ -2 & 5 \end{bmatrix}$$
29.
$$\mathbf{A}^{2} = \begin{bmatrix} a^{2} + bc & ab + bd \\ ac + dc & bc + d^{2} \end{bmatrix},$$

$$\mathbf{A}^{2} + (ad - bc)\mathbf{I} = \begin{bmatrix} a^{2} + ad & ab + bd \\ ac + dc & d^{2} + ad \end{bmatrix} =$$

$$\begin{bmatrix} a(a + d) & b(a + d) \\ c(a + d) & d(a + d) \end{bmatrix} = (a + d)\mathbf{A}$$
31. (a)
$$(\mathbf{A} + \mathbf{B})(\mathbf{A} - \mathbf{B}) = \begin{bmatrix} -25 & -34 \\ -71 & -34 \end{bmatrix} \neq \begin{bmatrix} -8 & -45 \\ -44 & -51 \end{bmatrix}$$

$$= \mathbf{A}^{2} - \mathbf{B}^{2}$$
33.
$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} +1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & +1 \end{bmatrix}$$
35.
$$\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix}$$
37.
$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$
39.
$$\mathbf{A}(c_{1}\mathbf{x}_{1} + c_{2}\mathbf{x}_{2}) = c_{1}\mathbf{A}\mathbf{x}_{1} + c_{2}\mathbf{A}\mathbf{x}_{2} = c_{1} \cdot \mathbf{0} + c_{2} \cdot \mathbf{0} = \mathbf{0}$$
43.
$$\mathbf{A}^{2} = \begin{bmatrix} 6 & -3 & -3 \\ -3 & 6 & -3 \\ -3 & -3 & 6 \end{bmatrix} = 3\mathbf{A}$$

Section 3.5

1.
$$A^{-1} = \begin{bmatrix} 3 & -2 \\ -4 & 3 \end{bmatrix}, x = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$$

2. $A^{-1} = \begin{bmatrix} 5 & -7 \\ -2 & 3 \end{bmatrix}, x = \begin{bmatrix} -26 \\ 11 \end{bmatrix}$
3. $A^{-1} = \begin{bmatrix} 6 & -7 \\ -5 & 6 \end{bmatrix}, x = \begin{bmatrix} 33 \\ -28 \end{bmatrix}$
4. $A^{-1} = \begin{bmatrix} 17 & -12 \\ -7 & 5 \end{bmatrix}, x = \begin{bmatrix} 25 \\ -10 \end{bmatrix}$
5. $A^{-1} = \frac{1}{2} \begin{bmatrix} 4 & -2 \\ -5 & 3 \end{bmatrix}, x = \frac{1}{2} \begin{bmatrix} 8 \\ -7 \end{bmatrix}$
6. $A^{-1} = \frac{1}{3} \begin{bmatrix} 6 & -7 \\ -3 & 4 \end{bmatrix}, x = \frac{1}{3} \begin{bmatrix} 25 \\ -10 \end{bmatrix}$
7. $A^{-1} = \frac{1}{4} \begin{bmatrix} 7 & -9 \\ -5 & 7 \end{bmatrix}, x = \frac{1}{4} \begin{bmatrix} 3 \\ -1 \end{bmatrix}$
8. $A^{-1} = \frac{1}{5} \begin{bmatrix} 10 & -15 \\ -5 & 8 \end{bmatrix}, x = \frac{1}{5} \begin{bmatrix} 25 \\ -11 \end{bmatrix}$
9. $A^{-1} = \begin{bmatrix} 5 & -6 \\ -4 & 5 \end{bmatrix}$
10. $A^{-1} = \frac{1}{2} \begin{bmatrix} 6 & -7 \\ -4 & -5 \end{bmatrix}$
11. $A^{-1} = \begin{bmatrix} -5 & -2 & 5 \\ 2 & 1 & -2 \\ -4 & -3 & 5 \end{bmatrix}$
12. $A^{-1} = \begin{bmatrix} 18 & 2 & -7 \\ -3 & 0 & 1 \\ -4 & -1 & 2 \end{bmatrix}$
13. $A^{-1} = \begin{bmatrix} -13 & 42 & -5 \\ 3 & -9 & 1 \\ 2 & -7 & 1 \end{bmatrix}$
14. $A^{-1} = \begin{bmatrix} 11 & -7 & -9 \\ -4 & 3 & 3 \\ -2 & 1 & 2 \end{bmatrix}$
15. $A^{-1} = \begin{bmatrix} -22 & 2 & 7 \\ -7 & 3 & 8 \\ 10 & -1 & -3 \end{bmatrix}$
16. $A^{-1} = \frac{1}{3} \begin{bmatrix} -3 & 0 & 3 \\ -1 & -3 & -1 \\ -1 & 3 & 2 \end{bmatrix}$
17. $A^{-1} = \frac{1}{4} \begin{bmatrix} -2 & -6 & -3 \\ -2 & -2 & -1 \\ -2 & -2 & 1 \end{bmatrix}$

18.
$$A^{-1} = \frac{1}{5} \begin{bmatrix} 1 & 2 & -2 \\ -5 & 0 & 5 \\ -3 & -1 & 6 \end{bmatrix}$$

19. $A^{-1} = \frac{1}{6} \begin{bmatrix} -21 & 11 & 8 \\ 9 & -5 & -2 \\ -3 & 3 & 0 \end{bmatrix}$
20. $A^{-1} = \frac{1}{7} \begin{bmatrix} 3 & 1 & 0 \\ -2 & -3 & 7 \\ -1 & 2 & 0 \end{bmatrix}$
21. $A^{-1} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -2 & 0 & 1 & 0 \\ 0 & -2 & -1 & 2 \\ 0 & 1 & 1 & -1 \\ -3 & 3 & -5 & 1 \end{bmatrix}$
23. $A^{-1} = \begin{bmatrix} 4 & -3 \\ -5 & 4 \end{bmatrix}, X = \begin{bmatrix} 7 & 18 & -35 \\ -9 & -23 & 45 \end{bmatrix}$
24. $A^{-1} = \begin{bmatrix} 7 & -6 & 4 \\ -8 & 7 \end{bmatrix}, X = \begin{bmatrix} 7 & 14 & -30 & 46 \\ -16 & 35 & -53 \end{bmatrix}$
25. $A^{-1} = \begin{bmatrix} -16 & 3 & 11 \\ 6 & -8 & 7 \end{bmatrix}, X = \begin{bmatrix} 7 & -14 & 15 \\ -1 & 3 & -2 \\ -2 & 1 & 0 \end{bmatrix}, X = \begin{bmatrix} -21 & 9 & 6 \\ 8 & -3 & -2 \\ -17 & 6 & 5 \end{bmatrix}$
26. $A^{-1} = \begin{bmatrix} -16 & 3 & 11 \\ 6 & -1 & -4 \\ -13 & 2 & 9 \end{bmatrix}, X = \begin{bmatrix} -21 & 9 & 6 \\ 8 & -3 & -2 \\ -17 & 6 & 5 \end{bmatrix}$
27. $A^{-1} = \begin{bmatrix} 7 & -20 & 17 \\ 0 & -1 & 1 \\ -2 & 6 & -5 \end{bmatrix}, X = \begin{bmatrix} 17 & -20 & 24 & -13 \\ 1 & -1 & 1 & -1 \\ -5 & 6 & -7 & 4 \end{bmatrix}$
28. $A^{-1} = \begin{bmatrix} -5 & 5 & 10 \\ -8 & 8 & 15 \\ 24 & -23 & -45 \end{bmatrix}, X = \begin{bmatrix} -5 & 5 & 10 \\ -8 & 8 & 15 \\ 24 & -23 & -45 & -13 \end{bmatrix}$

Section 3.6

1.	60 2. 4
3.	-210 4. -72
5.	120 6. 60
7.	0 8. 25
9.	30 10. 7
11.	40 12. 10
13.	78 1422
15.	-74 16. 84
17.	8 18. 135
19.	3 9 20. 79
21.	$\Delta = 1, x = 10, y = -7$
22.	$\Delta = 1, x = -1, y = 1$
23.	$\Delta = 1, x = 2, y = -4$
24.	$\Delta = 1, x = 5, y = -3$
25.	$\Delta = 2, x = 6, y = -3$
26.	$\Delta = -2, x = \frac{1}{2}, y = 0$
27.	$\Delta = 96, x_1 = \frac{1}{3}, x_2 = -\frac{2}{3}, x_3 = -\frac{1}{3}$
28.	$\Delta = 35, x_1 = \frac{4}{7}, x_2 = \frac{3}{7}, x_3 = \frac{2}{7}$
29.	$\Delta = 23, x_1 = 2, x_2 = 3, x_3 = 0$
30.	$\Delta = 56, x_1 = -\frac{1}{7}, x_2 = \frac{9}{14}, x_3 = \frac{2}{7}$
31.	$\Delta = 14, x_1 = -\frac{8}{7}, x_2 = -\frac{10}{7}, x_3 = \frac{1}{7}$

32.
$$\Delta = 6, x_1 = -\frac{7}{3}, x_2 = 9, x_3 = 8$$

33. det $A = -4, A^{-1} = \frac{1}{4} \begin{bmatrix} 4 & 4 & 4 \\ 16 & 15 & 13 \\ 28 & 25 & 23 \end{bmatrix}$
34. det $A = 35, A^{-1} = \frac{1}{4} \begin{bmatrix} -2 & -3 & 12 \\ 9 & -4 & -19 \\ 13 & 2 & -8 \end{bmatrix}$
35. det $A = 35, A^{-1} = \frac{1}{35} \begin{bmatrix} -15 & 25 & -26 \\ 10 & -5 & 8 \\ 15 & -25 & 19 \end{bmatrix}$
36. det $A = 23, A^{-1} = \frac{1}{23} \begin{bmatrix} 5 & 20 & -17 \\ 10 & 17 & -11 \\ 1 & 4 & -8 \end{bmatrix}$
37. det $A = 29, A^{-1} = \frac{1}{29} \begin{bmatrix} -11 & -14 & -15 \\ -17 & 19 & 10 \\ 18 & -15 & -14 \end{bmatrix}$
38. det $A = 6, A^{-1} = \frac{1}{6} \begin{bmatrix} -6 & 10 & 2 \\ 15 & -21 & -6 \\ 12 & -18 & -6 \end{bmatrix}$
39. det $A = 37, A^{-1} = \frac{1}{37} \begin{bmatrix} -21 & -1 & -13 \\ 4 & 9 & 6 \\ -6 & 5 & -9 \end{bmatrix}$
40. det $A = 107, A^{-1} = \frac{1}{107} \begin{bmatrix} 9 & 12 & -13 \\ 11 & -21 & -4 \\ -15 & -20 & -14 \end{bmatrix}$

Section 3.7

1. y(x) = -2 + 3x2. y(x) = 4 - 7x4. $y(x) = 2x + 3x^2$ 3. $y(x) = 3 - 2x^2$ 5. $y(x) = 5 - 3x + x^2$ 6. $y(x) = -10 - 7x + 2x^2$ 7. $y(x) = \frac{1}{3}(4x + 3x^2 - 4x^3)$ 8. $y(x) = 5 + 3x - x^3$ 9. $y(x) = 4 + 3x + 2x^2 + x^3$ 10. $y(x) = 17 - 5x + 3x^2 - 2x^3$ 11. $x^2 + y^2 - 6x - 4y - 12 = 0$, center (3, 2) and radius 5 12. $x^2 + y^2 + 6x - 8y - 75 = 0$, center (-3, 4) and radius 10 13. $x^2 + y^2 + 4x + 4y - 5 = 0$, center (-2, -2) and radius $\sqrt{13}$ **14.** $x^2 + y^2 - 10x - 24y = 0$, center (5, 12) and radius 13 **15.** $x^2 - xy + y^2 = 25$ 16. $4x^2 - 7xy + 4y^2 = 100$ 17. $100x^2 - 199xy + 100y^2 = 100$ **18.** $400x^2 - 481xy + 225y^2 = 3600$ 20. $y = 10x + \frac{8}{x} - \frac{16}{x^2}$ **19.** $y = 3 + \frac{2}{3}$ **21.** $x^2 + y^2 + z^2 - 2x - 4y - 6z - 155 = 0$, center (1, 2, 3) and radius 13 22. $x^2 + y^2 + z^2 - 10x + 14y - 18z - 521 = 0$, center (5, -7, 9) and radius 26 **23.** $P(t) = 49.061 - 0.0722t + 0.00798t^2$ **24.** $P(t) = 56.590 + 0.30145t - 0.007375t^2$ **25.** $P(t) = 62.813 + 1.37915t - 0.012375t^2$ **26.** $P(t) = 34.838 + 0.7692t + 0.0064t^2$ 27. $P(t) = 44.678 + 0.850417t - 0.05105t^2 + 0.000983833t^3$ **28.** $P(t) = 51.619 + 0.672433t - 0.019565t^2 + 0.000203167t^3$ **29.** $P(t) = 54.973 + 0.308667t + 0.059515t^2 - 0.00119817t^3$ **30.** $P(t) = 28.053 + 0.592233t + 0.00907t^2 - 0.0000443333t^3$ **31.** $P(t) = 39.478 + 0.209692t + 0.0564163t^2 - 0.0564163t^2 - 0.0564163t^2$ $0.00292992t^3 + 0.0000391375t^4$ **32.** $P(t) = 44.461 + 0.7651t - 0.000489167t^2 - 0.000489167t^2$ $0.000516t^3 + 7.19167 \times 10^{-6}t^4$ **33.** $P(t) = 47.197 + 1.22537t - 0.077192t^2 + 1.2257t - 0.07719t - 0.077t - 0.077t$ $0.00373475t^3 - 0.0000493292t^4$

ANSWERS TO SELECTED PROBLEM 693

34. P(t) = 20.190 + 1.00003t - 0.031775t² + 0.00116067t³ - 0.000041205t⁴
36. y = 2x² - 6x + 7
38. x² + y² + 6x - 8y - 75 = 0, center (-3, 4) and radius 10
40. 400x² - 481xy + 225y² - 3600 = 0

Chapter 4

Section 4.1

1.	$\sqrt{51}$, (5, 8, -11), (2, 23, 0)		
2.	9, (1, 4, -1), (-15, -16, 26)		
3.	$3\sqrt{21}, 9i - 3j + 3k, -14i - 2$	21 j -	- 43 k
4.	$\sqrt{17}$, 4 i - j - 3 k , 6 i - 7 j + 12	2 k	
5.	Linearly dependent	6.	Linearly independent
7.	Linearly independent	8.	Linearly dependent
9.	$\mathbf{w} = 3\mathbf{u} + 2\mathbf{v}$	10.	$\mathbf{w} = 2\mathbf{u} - 3\mathbf{v}$
11.	$\mathbf{w} = \mathbf{u} - 2\mathbf{v}$	12.	$\mathbf{w} = 3\mathbf{u} + 5\mathbf{v}$
13.	$\mathbf{w} = 2\mathbf{u} - 3\mathbf{v}$	14.	$\mathbf{w} = 7\mathbf{u} + 5\mathbf{v}$
15.	Linearly dependent	16.	Linearly dependent
17.	Linearly independent	18.	Linearly independent
19.	Linearly dependent; $a = 3, b =$	= 2,	c = 1
20.	Linearly dependent; $a = -2, b$	= 3	B, c = 1
21.	Linearly dependent; $a = 11, b$	= 4	, c = -1
22.	Linearly independent	23.	Linearly independent
24.	Linearly independent		
25.	$\mathbf{t} = 2\mathbf{u} - \mathbf{v} + 3\mathbf{w}$	26.	$\mathbf{t} = \mathbf{u} + 5\mathbf{v} - \mathbf{w}$
27.	$\mathbf{t} = 2\mathbf{u} + 6\mathbf{v} + \mathbf{w}$	28.	$\mathbf{t} = \mathbf{u} + \mathbf{v} + \mathbf{w}$

Section 4.2

- **1.** W is a subspace of \mathbb{R}^3 .
- 2. W is a subspace of \mathbb{R}^3 .
- **3.** *W* is not a subspace of \mathbb{R}^3 .
- 4. W is not a subspace of \mathbb{R}^3 .
- 5. W is a subspace of \mathbb{R}^4 .
- 6. W is a subspace of \mathbb{R}^4 .
- 7. W is not a subspace of \mathbb{R}^2 .
- 8. W is a subspace of \mathbb{R}^2 .
- 9. W is not a subspace of \mathbb{R}^2 .
- 10. W is not a subspace of \mathbb{R}^2 .
- 11. W is a subspace of \mathbb{R}^4 .
- 12. W is not a subspace of \mathbb{R}^4 .
- 13. *W* is not a subspace of \mathbb{R}^4 .
- 14. W is not a subspace of \mathbb{R}^4 .
- **15.** $\mathbf{u} = (-1, 0, 1, 0)$ and $\mathbf{v} = (-4, -2, 0, 1)$
- **16.** $\mathbf{u} = (-1, -1, 1, 0)$ and $\mathbf{v} = (-5, -3, 0, 1)$
- **17.** $\mathbf{u} = (1, -3, 1, 0)$ and $\mathbf{v} = (-2, 1, 0, 1)$
- **18.** $\mathbf{u} = (2, -1, -2, 1, 0)$ and $\mathbf{v} = (3, -4, 5, 0, 1)$
- **19.** $\mathbf{u} = (1, 2, -1, 0)$
- **20.** $\mathbf{u} = (-5, 3, -2, 1)$
- **21.** $\mathbf{u} = (-3, 2, -4, 1)$
- 22. $\mathbf{u} = (-6, 4, -3, 1)$

2. Linearly dependent

4. Linearly dependent

8. Linearly dependent

12. $w = 2v_1 + 5v_2$

6. Linearly independent

Section 4.3

- **1.** Linearly dependent
- 3. Linearly dependent
- 5. Linearly independent
- 7. Linearly independent
- 9. $w = 2v_1 3v_2$ 10. $w = 7v_1 + 4v_2$
- 11. $w = v_1 2v_2$
 - $v_1 = 2v_2$
- 13. w cannot be expressed as a linear combination of v1 and v2.
 14. w cannot be expressed as a linear combination of v1, v2, and v3
- **14.** We cannot be expressed as a linear co
- **15.** $\mathbf{w} = 3\mathbf{v}_1 2\mathbf{v}_2 + 4\mathbf{v}_3$ **16.** $\mathbf{w} = 6\mathbf{v}_1 - 2\mathbf{v}_2 + 3\mathbf{v}_3$
- **17.** The vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 are linearly independent.
- **18.** $3\mathbf{v}_1 + \mathbf{v}_2 + 5\mathbf{v}_3 = \mathbf{0}$
- **19.** The vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 are linearly independent.
- 20. The vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 are linearly independent.

21. $\mathbf{v}_1 - 2\mathbf{v}_2 - \mathbf{v}_3 = \mathbf{0}$ **22.** $7\mathbf{v}_1 + 5\mathbf{v}_2 - 9\mathbf{v}_3 = \mathbf{0}$

Section 4.4

- **1.** The vectors \mathbf{v}_1 and \mathbf{v}_2 form a basis for \mathbf{R}^2 .
- **2.** The vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 do not form a basis for \mathbf{R}^3 .
- **3.** The given vectors do not form a basis for \mathbf{R}^3 .
- **4.** The given vectors do not form a basis for \mathbf{R}^4 .
- **5.** The three vectors \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 do not form a basis for \mathbf{R}^3 .
- **6.** The three given vectors form a basis for \mathbf{R}^3 .
- **7.** The three given vectors form a basis for \mathbf{R}^3 .
- **8.** The four given vectors form a basis for \mathbf{R}^4 .
- 9. The plane x 2y + 5z = 0 is a 2-dimensional subspace of \mathbb{R}^3 with basis consisting of the vectors $\mathbf{v}_1 = (2, 1, 0)$ and $\mathbf{v}_2 = (-5, 0, 1)$.
- 10. The plane y z = 0 is a 2-dimensional subspace of \mathbb{R}^3 with basis consisting of the vectors $\mathbf{v}_1 = (1, 0, 0)$ and $\mathbf{v}_2 = (0, 1, 1)$.
- 11. The line is a 1-dimensional subspace of \mathbb{R}^3 with basis consisting of the vector $\mathbf{v} = (-3, 1, 1)$.
- 12. Hence the subspace consisting of all such vectors is 3-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (1, 1, 0, 0), \mathbf{v}_2 = (1, 0, 1, 0), \text{ and } \mathbf{v}_3 = (1, 0, 0, 1).$
- 13. The subspace consisting of all such vectors is 2-dimensional with basis consisting of the vectors v₁ = (3, 0, 1, 0) and v₂ = (0, 4, 0, 1).
- 14. The subspace consisting of all such vectors is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-2, 1, 0, 0)$ and $\mathbf{v}_2 = (0, 0, -3, 1)$.
- **15.** The solution space of the given system is 1-dimensional with basis consisting of the vector $\mathbf{v}_1 = (11, 7, 1)$.
- 16. The solution space of the given system is 1-dimensional with basis consisting of the vector $\mathbf{v}_1 = (11, -5, 1)$.
- 17. The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-11, -3, 1, 0)$ and $\mathbf{v}_2(-11, -5, 0, 1)$.
- 18. The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-3, 1, 0, 0)$ and $\mathbf{v}_2(-25, 0, 5, 1)$.
- 19. The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (3, -2, 1, 0)$ and $\mathbf{v}_2 = (-4, -3, 0, 1)$.
- **20.** The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (1, -3, 1, 0)$ and $\mathbf{v}_2 = (-2, 1, 0, 1)$.
- **21.** The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-1, -1, 1, 0)$ and $\mathbf{v}_2 = (-5, -3, 0, 1)$.

- **22.** The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (2, 1, 0, 0)$ and $\mathbf{v}_2 = (-5, 0, -7, 1)$.
- **23.** The solution space of the given system is 1-dimensional with basis consisting of the vector $\mathbf{v}_1 = (2, -3, 1, 0)$.
- 24. The solution space of the given system is 3-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-2, 2, 1, 0, 0)$, $\mathbf{v}_2 = (-1, 3, 0, 1, 0)$, and $\mathbf{v}_3 = (-3, -1, 0, 0, 1)$.
- **25.** The solution space of the given system is 3-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-2, 1, 0, 0, 0)$, $\mathbf{v}_2 = (2, 0, 1, 1, 0)$, and $\mathbf{v}_3 = (-3, 0, -4, 0, 1)$.
- **26.** The solution space of the given system is 2-dimensional with basis consisting of the vectors $\mathbf{v}_1 = (-2, 1, 2, 1, 0)$ and $\mathbf{v}_2 = (3, -4, 5, 0, 1)$.

Section 4.5

- Row basis: The first and second row vectors of E. Column basis: The first and second column vectors of A.
- Row basis: The first and second row vectors of E. Column basis: The first and second column vectors of A.
- **3.** Row basis: The first and second row vectors of **E**. Column basis: The first and second column vectors of **A**.
- **4.** Row basis: The three row vectors of **E**. Column basis: The first three column vectors of **A**.
- Row basis: The three row vectors of E. Column basis: The first, second, and fourth column vectors of A.
- Row basis: The three row vectors of E. Column basis: The first, second, and fourth column vectors of A.
- 7. Row basis: The first two row vectors of **E**. Column basis: The first two column vectors of **A**.
- **8.** Row basis: The first three row vectors of **E**. Column basis: The first, second, and fourth column vectors of **A**.
- **9.** Row basis: The first three row vectors of **E**. Column basis: The first three column vectors of **A**.
- Row basis: The first three row vectors of E. Column basis: The first, second, and fourth column vectors of A.
- Row basis: The first three row vectors of E. Column basis: The first, second, and fifth column vectors of A.
- **12.** Row basis: The first three row vectors of **E**. Column basis: The first, second, and fifth column vectors of **A**.
- **13.** Linearly independent: \mathbf{v}_1 and \mathbf{v}_2
- 14. Linearly independent: \mathbf{v}_1 and \mathbf{v}_2
- 15. Linearly independent: v_1 , v_2 , and v_4
- 16. Linearly independent: $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_4$, and \mathbf{v}_5
- **17.** Basis vectors: $\mathbf{v}_1, \mathbf{v}_2, \mathbf{e}_2$
- **18.** Basis vectors: $\mathbf{v}_1, \mathbf{v}_2, \mathbf{e}_2$
- **19.** Basis vectors: **v**₁, **v**₂, **e**₂, **e**₃
- **20.** Basis vectors: **v**₁, **v**₂, **e**₁, **e**₃
- 21. The first and second equations are irredundant.
- 22. The first and second equations are irredundant.
- 23. The first, second, and fourth equations are irredundant.
- 24. The first, second, and fifth equations are irredundant.

Section 4.6

- 1. Yes, the three vectors are mutually orthogonal.
- 2. Yes, the three vectors are mutually orthogonal. 3. Yes, the three vectors are mutually orthogonal.
- 4. Yes, the three vectors are mutually orthogonal.

5. $a^2 = 7, b^2 = 7, c^2 = 14$ 6. $a^2 = 18, b^2 = 18, c^2 = 36$ 7. $a^2 = 19, b^2 = 25, c^2 = 44$ 8. $a^2 = 103, b^2 = 112, c^2 = 215$ 9. $\angle A = \angle B = 45^{\circ}$ 10. $\angle A = \angle B = 45^{\circ}$ 11. $\angle A = 41.08^{\circ}, \angle B = 48.92^{\circ}$ 12. $\angle A = 43.80^{\circ}, \angle B = 46.20^{\circ}$ **13.** $\mathbf{u}_1 = (2, 1, 0), \mathbf{u}_2 = (-3, 0, 1)$ 14. $\mathbf{u}_1 = (-5, 1, 0), \mathbf{u}_2 = (3, 0, 1)$ **15.** $\mathbf{u}_1 = (2, 1, 0, 0), \mathbf{u}_2 = (3, 0, 1, 0), \mathbf{u}_3 = (-5, 0, 0, 1)$ **16.** $\mathbf{u}_1 = (-7, 1, 0, 0), \mathbf{u}_2 = (6, 0, 1, 0), \mathbf{u}_3 = (9, 0, 0, 1)$ 17. $\mathbf{u}_1 = (7, -3, 1, 0), \mathbf{u}_2 = (-19, 5, 0, 1)$ **18.** $\mathbf{u}_1 = (-12, -3, 1, 0), \mathbf{u}_2 = (16, 7, 0, 1)$ **19.** $\mathbf{u}_1 = (-13, 4, 1, 0, 0), \mathbf{u}_2 = (4, -3, 0, 1, 0),$ $\mathbf{u}_3 = (-11, 4, 0, 0, 1)$ **20.** $\mathbf{u}_1 = (-5, 1, 1, 0, 0), \mathbf{u}_2 = (-12, 4, 0, 1, 0),$ $\mathbf{u}_3 = (-19, 7, 0, 0, 1)$ **21.** $\mathbf{u}_1 = (-1, -1, 1, 0, 0), \mathbf{u}_2 = (0, -1, 0, -1, 1)$

22. $\mathbf{u}_1 = (-2, 1, 1, 0, 0), \mathbf{u}_2 = (1, -2, 0, 1, 0)$

Section 4.7

1.	It is a subspace.	2.	It is a subspace.
3.	It is not a subspace.	4.	It is not a subspace.
5.	It is a subspace.	6.	It is not a subspace.
7.	It is not a subspace.	8.	It is a subspace.
9.	It is not a subspace.	10.	It is a subspace.
11.	It is a subspace.	12.	It is not a subspace.

- 13. The functions $\sin x$ and $\cos x$ are linearly independent.
- 14. The functions e^x and xe^x are linearly independent.
- 15. The functions (1 + x), (1 x), and $(1 x^2)$ are linearly independent.
- 16. The three given polynomials are linearly dependent.
- 17. The three given trigonometric functions are linearly dependent.
- **18.** The two given linear combinations of $\sin x$ and $\cos x$ are linearly independent.
- **19.** A = 3 and B = -2.
- **20.** A = -2 and B = C = 1.
- **21.** A = 2, B = -2, and C = 0.
- **22.** A = -1, B = 4, and C = -3.
- **23.** The solution space is 3-dimensional with basis $\{1, x, x^2\}$.
- 24. The solution space is 4-dimensional with basis $\{1, x, x^2, x^3\}$.
- **25.** The solution space is 2-dimensional with basis $\{1, e^{5x}\}$.
- **26.** The solution space is 2-dimensional with basis $\{1, e^{-10x}\}$.

Chapter 5

Section 5.1

1. $y(x) = \frac{5}{2}e^x - \frac{5}{2}e^{-x}$ 2. $y(x) = 2e^{3x} - 3e^{-3x}$ 3. $y(x) = 3\cos 2x + 4\sin 2x$ 4. $y(x) = 10\cos 5x - 2\sin 5x$ 5. $y(x) = 2e^x - e^{2x}$ 6. $y(x) = 4e^{2x} + 3e^{-3x}$ 7. $y(x) = 6 - 8e^{-x}$ 8. $y(x) = \frac{1}{3}(14 - 2e^{3x})$ 9. $y(x) = 2e^{-x} + xe^{-x}$ 10. $y(x) = 3e^{5x} - 2xe^{5x}$ 11. $y(x) = 5e^x \sin x$ 12. $y(x) = e^{-3x}(2\cos 2x + 3\sin 2x)$ 13. $y(x) = 5x - 2x^2$ 14. $y(x) = 3x^2 - 16/x^3$ 15. $y(x) = 7x - 5x \ln x$ 16. $y(x) = 2\cos(\ln x) + 3\sin(\ln x)$ 21. Linearly independent 22. Linearly independent **23.** Linearly independent 24. Linearly dependent **25.** Linearly independent 26. Linearly independent 28. $y(x) = 1 - 2\cos x - \sin x$ 29. There is no contradiction because if the given differential equation is divided by x^2 to get the form in Eq. (8), then the resulting coefficient functions p(x) = -4/x and $q(x) = 6/x^2$ are not continuous at x = 0. 34. $y(x) = c_1 e^{-5x} + c_2 e^{3x}$ **33.** $y(x) = c_1 e^x + c_2 e^{2x}$ **36.** $y(x) = c_1 + c_2 e^{-3x/2}$ 35. $y(x) = c_1 + c_2 e^{-5x}$ 37. $y(x) = c_1 e^{-x/2} + c_2 e^x$ **38.** $y(x) = c_1 e^{-x/2} + c_2 e^{-3x/2}$ **39.** $y(x) = (c_1 + c_2 x)e^{-x/2}$ **40.** $y(x) = (c_1 + c_2 x)e^{2x/3}$ **41.** $y(x) = c_1 e^{-4x/3} + c_2 e^{5x/2}$ 42. $y(x) = c_1 e^{-4x/7} + c_2 e^{3x/5}$ 43. y'' + 10y' = 044. y'' - 100y = 0**46.** y'' - 110y' + 1000y = 045. y'' + 20y' + 100y = 047. y'' = 048. y'' - 2y' - y = 049. The high point is $\left(\ln \frac{7}{4}, \frac{16}{7}\right)$. 50. $(-\ln 2, -2)$ 52. $y(x) = c_1 x + c_2 / x$ 53. $y(x) = c_1 x^{-4} + c_2 x^3$ 54. $y(x) = c_1 x^{-3/2} + c_2 x^{1/2}$ 55. $y(x) = c_1 + c_2 \ln x$ 56. $y(x) = x^2(c_1 + c_2 \ln x)$ Section 5.2

1. $15 \cdot (2x) - 16 \cdot (3x^2) - 6 \cdot (5x - 8x^2) \equiv 0$ 2. $(-4)(5) + (5)(2 - 3x^2) + (1)(10 + 15x^2) \equiv 0$ 3. $1 \cdot 0 + 0 \cdot \sin x + 0 \cdot e^x \equiv 0$ 4. $(6)(17) + (-51)(2\sin^2 x) + (-34)(3\cos^2 x) \equiv 0$ 5. $1 \cdot 17 - 34 \cdot \cos^2 x + 17 \cdot \cos 2x \equiv 0$ 6. $(-1)(e^x) + (1)(\cosh x) + (1)(\sinh x) \equiv 0$ 13. $y(x) = \frac{4}{3}e^x - \frac{1}{3}e^{-2x}$ 14. $y(x) = \frac{1}{2}(3e^x - 6e^{2x} + 3e^{3x})$ 15. $y(x) = (2 - 2x + x^2)e^x$ 16. $y(x) = -12e^x + 13e^{2x} - 10xe^{2x}$ 17. $y(x) = \frac{1}{9}(29 - 2\cos 3x - 3\sin 3x)$ 18. $y(x) = e^x (2 - \cos x - \sin x)$ 19. $y(x) = x + 2x^2 + 3x^3$ **20.** $y(x) = 2x - x^{-2} + x^{-2} \ln x$ 21. $y(x) = 2\cos x - 5\sin x + 3x$ 22. $v(x) = 4e^{2x} - e^{-2x} - 3$

23.
$$y(x) = e^{-x} + 4e^{3x} - 2$$

24. $y(x) = e^{x}(3\cos x + 4\sin x) + x + 1$
38. $y_{2}(x) = \frac{1}{x^{3}}$
39. $y_{2}(x) = xe^{x/2}$
40. $y_{2}(x) = xe^{x}$
41. $y_{2}(x) = x + 2$
42. $y_{2}(x) = 1 + x^{2}$

Section 5.3

1. $y(x) = c_1 e^{2x} + c_2 e^{-2x}$ 2. $y(x) = c_1 + c_2 e^{3x/2}$ 3. $y(x) = c_1 e^{2x} + c_2 e^{-5x}$ 4. $y(x) = c_1 e^{x/2} + c_2 e^{3x}$ 5. $y(x) = c_1 e^{-3x} + c_2 x e^{-3x}$ 6. $y(x) = e^{-5x/2} \left[c_1 \exp\left(\frac{1}{2}x\sqrt{5}\right) + c_2 \exp\left(-\frac{1}{2}x\sqrt{5}\right) \right]$ 7. $y(x) = c_1 e^{3x/2} + c_2 x e^{3x/2}$ 8. $y(x) = e^{3x}(c_1 \cos 2x + c_2 \sin 2x)$ 9. $y(x) = e^{-4x}(c_1 \cos 3x + c_2 \sin 3x)$ **10.** $y(x) = c_1 + c_2 x + c_3 x^2 + c_4 e^{-3x/5}$ 11. $y(x) = c_1 + c_2 x + c_3 e^{4x} + c_4 x e^{4x}$ 12. $y(x) = c_1 + c_2 e^x + c_3 x e^x + c_4 x^2 e^x$ 13. $y(x) = c_1 + c_2 e^{-2x/3} + c_3 x e^{-2x/3}$ 14. $y(x) = c_1 e^x + c_2 e^{-x} + c_3 \cos 2x + c_4 \sin 2x$ 15. $y(x) = c_1 e^{2x} + c_2 x e^{2x} + c_3 e^{-2x} + c_4 x e^{-2x}$ 16. $y(x) = (c_1 + c_2 x) \cos 3x + (c_3 + c_4 x) \sin 3x$ 17. $y(x) = c_1 \cos\left(x/\sqrt{2}\right) + c_2 \sin\left(x/\sqrt{2}\right) +$ $c_3 \cos\left(\frac{2x}{\sqrt{3}}\right) + c_4 \sin\left(\frac{2x}{\sqrt{3}}\right)$ 18. $y(x) = c_1 e^{2x} + c_2 e^{-2x} + c_3 \cos 2x + c_4 \sin 2x$ $19. \ y(x) = c_1 e^x + c_2 e^{-x} + c_3 x e^{-x}$ **20.** y(x) = $e^{-x/2}\left[(c_1+c_2x)\cos\left(\frac{1}{2}x\sqrt{3}\right)+(c_3+c_3x)\sin\left(\frac{1}{2}x\sqrt{3}\right)\right]$ 21. $y(x) = 5e^x + 2e^{3x}$ 22. $y(x) = e^{-x/3} \left[3\cos\left(x/\sqrt{3}\right) + 5\sqrt{3}\sin\left(x/\sqrt{3}\right) \right]$ 23. $y(x) = e^{3x} (3\cos 4x - 2\sin 4x)$ 24. $y(x) = \frac{1}{2}(-7 + e^{2x} + 8e^{-x/2})$ 25. $y(x) = \frac{1}{4}(-13 + 6x + 9e^{-2x/3})$ **26.** $y(x) = \frac{1}{5}(24 - 9e^{-5x} - 25xe^{-5x})$ 27. $y(x) = c_1 e^x + c_2 e^{-2x} + c_3 x e^{-2x}$ **28.** $y(x) = c_1 e^{2x} + c_2 e^{-x} + c_3 e^{-x/2}$ **29.** $y(x) = c_1 e^{-3x} + e^{3x/2} \left[c_2 \cos\left(\frac{3}{2}x\sqrt{3}\right) + c_3 \sin\left(\frac{3}{2}x\sqrt{3}\right) \right]$ **30.** $y(x) = c_1 e^{-x} + c_2 e^{2x} + c_3 \cos(x\sqrt{3}) + c_4 \sin(x\sqrt{3})$ 31. $y(x) = c_1 e^x + e^{-2x} (c_2 \cos 2x + c_3 \sin 2x)$ 32. $y(x) = c_1 e^{2x} + (c_2 + c_3 x + c_4 x^2) e^{-x}$ 33. $y(x) = c_1 e^{3x} + e^{-3x} (c_2 \cos 3x + c_3 \sin 3x)$ 34. $y(x) = c_1 e^{2x/3} + c_2 \cos 2x + c_3 \sin 2x$ 35. $y(x) = c_1 e^{-x/2} + c_2 e^{-x/3} + c_3 \cos 2x + c_4 \sin 2x$ **36.** $y(x) = c_1 e^{7x/9} + e^{-x} (c_2 \cos x + c_3 \sin x)$ 37. $y(x) = 11 + 5x + 3x^2 + 7e^x$ 38. $y(x) = 2e^{5x} - 2\cos 10x$ **39.** $y^{(3)} - 6y'' + 12y' - 8y = 0$ **40.** $y^{(3)} - 2y'' + 4y' - 8y = 0$ **41.** $y^{(4)} - 16y = 0$ **42.** $y^{(6)} + 12y^{(4)} + 48y'' + 64y = 0$ **44.** (a) x = i, -2i (b) x = -i, 3i $45. \ y(x) = c_1 e^{-ix} + c_2 e^{3ix}$ **46.** $y(x) = c_1 e^{3ix} + c_2 e^{-2ix}$ **47.** $y(x) = c_1 \exp\left(\left[1 + i\sqrt{3}\right]x\right) + c_2 \exp\left(-\left[1 + i\sqrt{3}\right]x\right)$

48.
$$y(x) = \frac{1}{3} \left(e^x + \exp\left[\frac{1}{2} \left(-1 + i\sqrt{3}\right)x\right] + \exp\left[\frac{1}{2} \left(-1 - i\sqrt{3}\right)x\right] \right)$$

49. $y(x) = 2e^{2x} - 5e^{-x} + 3\cos x - 9\sin x$
52. $y(x) = c_1\cos(3\ln x) + c_2\sin(3\ln x)$
53. $y(x) = x^{-3}[c_1\cos(4\ln x) + c_2\sin(4\ln x)]$
54. $y(x) = c_1 + c_2\ln x + c_3x^{-3}$
55. $y(x) = c_1 + x^2(c_2 + c_3\ln x)$
56. $y(x) = c_1 + x^3\left(c_2x^{-\sqrt{3}} + c_3x^{+\sqrt{3}}\right)$
57. $y(x) = c_1 + x^3\left(c_2x^{-\sqrt{3}} + c_3x^{+\sqrt{3}}\right)$
58. $y(x) = x^{-1}[c_1 + c_2\ln x + c_3(\ln x)^2]$
Section 5.4

- **1.** Frequency: 2 rad/s $(1/\pi \text{ Hz})$; period: π s
- 2. Frequency: 8 rad/sec (4/ π Hz); period: $\pi/4$ sec
- Amplitude: 2 m; frequency: 5 rad/s; period: 2π/5 s
- 4. (a) $x(t) = \frac{13}{12}\cos(12t \alpha)$ with $\alpha = 2\pi - \tan^{-1}(5/12) \approx 5.8884;$ (b) Amplitude: $\frac{13}{12}$ m; period: $\pi/6$ sec
- 6. About 7.33 mi
- 7. About 10450 ft
- 8. 29.59 in.
- **10.** Amplitude: 100 cm; period: about 2.01 sec
- **11.** About 3.8 in.
- **13.** (a) $x(t) = 50(e^{-2t/5} e^{-t/2})$; (b) 4.096 exactly
- 14. (a) $x(t) = 25e^{-t/5}\cos(3t \alpha)$ with $\alpha = \tan^{-1}(3/4) \approx 0.6435$; (b) envelope curves $x = \pm 25e^{-t/5}$; pseudoperiod $2\pi/3$
- 15. $x(t) = 4e^{-2t} 2e^{-4t}$, overdamped; $u(t) = 2\cos(2\sqrt{2}t)$



16. $x(t) = 4e^{-3t} - 2e^{-7t}$, overdamped; $u(t) \approx 2\sqrt{\frac{22}{21}}\cos(\sqrt{21}t - 0.2149)$



17. $x(t) = 5e^{-4t}(2t+1)$, critically damped; $u(t) \approx \frac{5}{2}\sqrt{5}\cos(4t-5.8195)$



18. $x(t) = 2e^{-3t} \cos\left(4t - \frac{3\pi}{2}\right)$, underdamped; $u(t) = \frac{8}{5} \cos\left(5t - \frac{3\pi}{2}\right)$



19. $x(t) \approx \frac{1}{3}\sqrt{313} e^{-5t/2} \cos(6t - 0.8254)$, underdamped; $u(t) \approx \frac{4}{13}\sqrt{233} \cos(\frac{13}{2}t - 0.5517)$



20. $x(t) \approx 13e^{-4t} \cos(2t - 1.1760)$, underdamped; $u(t) \approx \sqrt{\frac{129}{5}} \cos(2\sqrt{5}t - 0.1770)$



21. $x(t) \approx 10e^{-5t} \cos(10t - 0.9273)$, underdamped; $u(t) \approx 2\sqrt{14} \cos(5\sqrt{5}t - 0.6405)$



- 22. (b) The time-varying amplitude is $\frac{2}{3}\sqrt{3}$, the frequency is $4\sqrt{3}$ rad/s, and the phase angle is $\pi/6$.
- **23.** (a) $k \approx 7018 \text{ lb/ft}$; (b) After about 2.47 s
- **34.** Damping constant: $c \approx 11.51$ lb/ft/s; spring constant: $k \approx 189.68$ lb/ft

Section 5.5

1. $y_p(x) = \frac{1}{25}e^{3x}$ 3. $y_p(x) = \frac{1}{39}(\cos 3x - 5\sin 3x)$ 4. $y_p(x) = \frac{1}{9}(-4e^x + 3xe^x)$ 5. $y_p(x) = \frac{1}{26}(13 + 3\cos 2x - 2\sin 2x)$ 6. $y_p(x) = \frac{1}{243}(4 - 56x + 49x^2)$ 7. $y_p(x) = -\frac{1}{6}(e^x - e^{-x}) = -\frac{1}{3}\sinh x$ 8. $y_p(x) = \frac{1}{4}x\sinh 2x$ 9. $y_p(x) = -\frac{1}{3} + \frac{1}{16}(2x^2 - x)e^x$ 10. $y_p(x) = \frac{1}{6}(2x\sin 3x - 3x\cos 3x)$

ANSWERS TO SELECTED PROBLEM 697

11.	$y_p(x) = \frac{1}{8}(3x^2 - 2x)$ 12. $y_p(x) = 2x + \frac{1}{2}x\sin x$
13.	$y_p(x) = \frac{1}{65}e^x(7\sin x - 4\cos x)$
14.	$y_p(x) = \frac{1}{24}(-3x^2e^x + x^3e^x)$ 15. $y_p(x) \equiv -17$
16.	$y_p(x) = \frac{1}{81}(45 + e^{3x} - 6xe^{3x} + 9x^2e^{3x})$
17.	$y_p(x) = \frac{1}{4}(x^2 \sin x - x \cos x)$
18.	$y_p(x) = -\frac{1}{144}(24xe^x - 19xe^{2x} + 6x^2e^{2x})$
19.	$y_p(x) = \frac{1}{8}(10x^2 - 4x^3 + x^4)$
20.	$y_p(x) = -7 + \frac{1}{2}xe^x$
21.	$y_p(x) = xe^x (A\cos x + B\sin x)$
22.	$y_p(x) = Ax^3 + Bx^4 + Cx^5 + Dxe^x$
23.	$y_p(x) = Ax \cos 2x + Bx \sin 2x + Cx^2 \cos 2x + Dx^2 \sin 2x$
24.	$y_p(x) = Ax + Bx^2 + (Cx + Dx^2)e^{-3x}$
25. 26	$y_p(x) = Axe^{-x} + Bx^2e^{-x} + Cxe^{-2x} + Dx^2e^{-2x}$ $y_n(x) = (Ax + Bx^2)e^{3x}\cos 2x + (Cx + Dx^2)e^{3x}\sin 2x$
20. 27.	$y_p(x) = (Ax + Bx)e^{-1}\cos 2x + (Cx + Dx)e^{-1}\sin 2x$ $y_p(x) = Ax\cos x + Bx\sin x + Cx\cos 2x + Dx\sin 2x$
28.	$y_p(x) =$
	$(Ax + Bx^2 + Cx^3)\cos 3x + (Dx + Ex^2 + Fx^3)\sin 3x$
29.	$y_p(x) = Ax^3e^x + Bx^4e^x + Cxe^{2x} + Dxe^{-2x}$
30.	$y_p(x) = (A + Bx + Cx^2)\cos x + (D + Ex + Fx^2)\sin x$
31.	$y(x) = \cos 2x + \frac{2}{4} \sin 2x + \frac{1}{2}x$
32.	$y(x) = \frac{1}{6}(15e^{-x} - 16e^{-2x} + e^{x})$
33.	$y(x) = \cos 3x - \frac{2}{15} \sin 3x + \frac{1}{5} \sin 2x$
34.	$y(x) = \cos x - \sin x + \frac{1}{2}x \sin x$
35.	$y(x) = e^{x} (2\cos x - \frac{1}{2}\sin x) + \frac{1}{2}x + 1$
36.	$y(x) = \frac{1}{192}(234 + 240x - 9e^{-2x} - 33e^{2x} - 12x^2 - 4x^4)$
37.	$y(x) = 4 - 4e^{x} + 3xe^{x} + x - \frac{1}{2}x^{2}e^{x} + \frac{1}{6}x^{3}e^{x}$
38.	$y(x) = \frac{1}{85} \left[e^{-x} \left(176 \cos x + 197 \sin x \right) - \left(6 \cos 3x + 7 \sin 3x \right) \right]$
39.	$y(x) = -3 + 3x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + 4e^{-x} + xe^{-x}$
40.	$y(x) = \frac{1}{4}(5e^{-x} + 5e^{x} + 10\cos x - 20)$
41.	$y_p(x) = 255 - 450x + 30x^2 + 20x^3 + 10x^4 - 4x^3$ $y(x) = 10x^{-x} + 25x^{2x} + 210x^{-3} + 10x^{-4} - 4x^{-5}$
42.	$y(x) = 10e^{-x} + 35e^{-x} + 210\cos x + 390\sin x + y_p(x)$ where $y_p(x)$ is the particular solution of Problem 41.
43.	(b) $y(x) = c_1 \cos 2x + c_2 \sin 2x$
	$+\frac{1}{4}\cos x - \frac{1}{20}\cos 3x$
44.	$y(x) = e^{-x/2} \left[c_1 \cos\left(\frac{1}{2}x\sqrt{3}\right) + c_2 \sin\left(\frac{1}{2}x\sqrt{3}\right) \right] +$
	$\frac{1}{26}(-3\cos 2x + 2\sin 2x) + \frac{1}{482}(15\cos 4x + 4\sin 4x)$
45.	$y(x) = c_1 \cos 3x + c_2 \sin 3x + \frac{1}{24} - \frac{1}{10} \cos 2x - \frac{1}{56} \cos 4x$
46.	$y(x) = c_1 \cos x + c_2 \sin x + \frac{1}{16} (3x \cos x + 3x^2 \sin x) + \frac{1}{16} (3x \cos x) + \frac{1}{16} (3x \cos x) + \frac{1}{16} (3x \cos x) + \frac{1}$
	$\frac{1}{128}(3\sin 3x - 4x\cos 3x)$
47.	$y_p(x) = \frac{2}{3}e^x$ 48. $y_p(x) = -\frac{1}{12}(6x+1)e^{-2x}$
49.	$y_p(x) = x^2 e^{2x}$
50.	$y_p(x) = \frac{1}{16}(4x \cosh 2x - \sinh 2x)$
51.	$y_p(x) = -\frac{1}{4}(\cos 2x \cos x - \sin 2x \sin x) +$
	$\frac{1}{20}(\cos 5x \cos 2x + \sin 5x \sin 2x) = -\frac{1}{5}\cos 3x (!)$
52.	$y_p(x) = -\frac{1}{6}x\cos 3x$
53.	$y_p(x) = \frac{2}{3}x\sin 3x + \frac{2}{9}(\cos 3x)\ln \cos 3x $
54.	$y_p(x) = -1 - (\cos x) \ln \csc x - \cot x $
55.	$y_p(x) = \frac{1}{8}(1 - x \sin 2x)$ 56. $y_p(x) = -\frac{1}{9}e^x(3x + 2)$
58.	$y_p(x) = x^3(\ln x - 1)$ 59. $y_p(x) = \frac{1}{4}x^4$
60. 61	$y_p(x) = -\frac{1}{5}x^{4/5}$
UI.	$y_p(x) = mx$ 1+x = 1
62.	$y_p(x) = -x^2 + x \ln \left \frac{1 + x}{1 - x} \right + \frac{1}{2} (1 + x^2) \ln 1 - x^2 $

Section 5.6



2. $x(t) = \frac{3}{2}\sin 2t - \sin 3t$



3. $x(t) = \sqrt{138388} \cos(10t - \alpha) + 5\cos(5t - \beta)$ with $\alpha = 2\pi - \tan^{-1}(1/186) \approx 6.2778$ and $\beta = \tan^{-1}(4/3) \approx 0.9273$.



4. $x(t) = 2\sqrt{106}\cos(5t - \alpha) + 10\cos 4t$ with $\alpha = \pi - \tan^{-1}(9/5) \approx 2.0779$



- 5. $x(t) = (x_0 C) \cos \omega_0 t + C \cos \omega t$, where $C = F_0/(k - m\omega^2)$
- 7. $x_{sp}(t) = \frac{10}{13}\cos(3t \alpha)$ with $\alpha = \pi \tan^{-1}(12/5) \approx 1.9656$



8. $x_{\rm sp}(t) = \frac{4}{25}\cos(5t-\alpha)$ with $\alpha = 2\pi - \tan^{-1}(3/4) \approx 5.6397$





13. $x_{\rm sp}(t) = \frac{300}{\sqrt{1469}} \cos(10t - \alpha)$ with $\alpha = \pi - \tan^{-1}(10/37) \approx 2.8776$



14. $x_{sp}(t) = \sqrt{485} \cos(t-\alpha)$ with $\alpha = \tan^{-1}(22) \approx 1.5254$ $x_{tr}(t) = \sqrt{3665} e^{-4t} \cos(3t-\beta)$ with $\beta = \pi + \tan^{-1}(52/31) \approx 4.1748$



15. $C(\omega) = 2/\sqrt{4} + \omega^4$; there is no practical resonance frequency.



16. $C(\omega) = 10/\sqrt{25 + 6\omega^2 + \omega^4}$; there is no practical resonance frequency.



17. $C(\omega) = 50/\sqrt{2025 - 54\omega^2 + \omega^4}$; there is practical resonance at frequency $\omega = 3\sqrt{3}$.



18. $C(\omega) = 100/\sqrt{422500 - 1200\omega^2 + \omega^4}$; there is practical resonance at frequency $\omega = 10\sqrt{6}$.



- **19.** $\omega = \sqrt{384}$ rad/sec (approximately 3.12 Hz)
- **20.** $\omega \approx 44.27$ rad/sec (approximately 422.75 rpm)
- **21.** $\omega_0 = \sqrt{(g/L) + (k/m)}$
- 22. $\omega_0 = \sqrt{k/(m+I/a^2)}$
- **23.** (a) Natural frequency: $\sqrt{10}$ rad/s (approximately 0.50 Hz); (b) amplitude: approximately 10.625 in.

Chapter 6

Section 6.1

- **1.** $\lambda_1 = 2$: $\mathbf{v}_1 = (1, 1)$; $\lambda_2 = 3$: $\mathbf{v}_2 = (2, 1)$ **2.** $\lambda_1 = -1$: $\mathbf{v}_1 = (1, 1)$; $\lambda_2 = 2$: $\mathbf{v}_2 = (2, 1)$ **3.** $\lambda_1 = 2$: $\mathbf{v}_2 = (1, 1)$; $\lambda_2 = 5$: $\mathbf{v}_1 = (2, 1)$ **4.** $\lambda_1 = 1$: $\mathbf{v}_2 = (1, 1)$; $\lambda_2 = 2$: $\mathbf{v}_1 = (3, 2)$ 5. $\lambda_1 = 1$: $\mathbf{v}_2 = (1, 1)$; $\lambda_2 = 4$: $\mathbf{v}_1 = (3, 2)$ **6.** $\lambda_1 = 2$: $\mathbf{v}_2 = (1, 1)$; $\lambda_2 = 3$: $\mathbf{v}_1 = (4, 3)$ 7. $\lambda_1 = 2$: $\mathbf{v}_2 = (1, 1)$; $\lambda_2 = 4$: $\mathbf{v}_1 = (4, 3)$ 8. $\lambda_1 = -2$: $\mathbf{v}_2 = (2, 3)$; $\lambda_2 = -1$: $\mathbf{v}_1 = (3, 4)$ **9.** $\lambda_1 = 3$: $\mathbf{v}_1 = (2, 1)$; $\lambda_2 = 4$: $\mathbf{v}_2 = (5, 2)$ **10.** $\lambda_1 = 4$: $\mathbf{v}_1 = (2, 1)$; $\lambda_2 = 5$: $\mathbf{v}_2 = (5, 2)$ **11.** $\lambda_1 = 4$: $\mathbf{v}_2 = (2, 3)$; $\lambda_2 = 5$: $\mathbf{v}_1 = (5, 7)$ **12.** $\lambda_1 = 3$: $\mathbf{v}_2 = (3, 2)$; $\lambda_2 = 4$: $\mathbf{v}_1 = (5, 3)$ **13.** $\lambda_1 = 0$: $\mathbf{v}_1 = (0, -1, 2)$; $\lambda_2 = 1$: $\mathbf{v}_2 = (0, -1, 3)$; $\lambda_3 = 2$: $\mathbf{v}_3 = (1, 0, 2)$ 14. $\lambda_1 = 0$: $\mathbf{v}_1 = (0, -1, 2)$; $\lambda_2 = 2$: $\mathbf{v}_2 = (0, -1, 3)$; $\lambda_3 = 5$: $\mathbf{v}_3 = (1, 0, 2)$ **15.** $\lambda_1 = 0$: $\mathbf{v}_1 = (1, 1, 0)$; $\lambda_2 = 1$: $\mathbf{v}_2 = (2, 1, 1)$; $\lambda_3 = 2$: $\mathbf{v}_3 = (1, 0, 2)$ **16.** $\lambda_1 = 0$: $\mathbf{v}_1 = (1, 1, 1)$; $\lambda_2 = 1$: $\mathbf{v}_2 = (1, 1, 0)$; $\lambda_3 = 3$: $\mathbf{v}_3 = (-1, 0, 2)$ 17. $\lambda_1 = 1$: $\mathbf{v}_1 = (1, 0, 1)$; $\lambda_2 = 2$: $\mathbf{v}_2 = (-1, 1, 2)$; $\lambda_3 = 3$: $\mathbf{v}_3 = (1, 0, 0)$ **18.** $\lambda_1 = 1$: $\mathbf{v}_1 = (1, 0, 3)$; $\lambda_2 = 2$: $\mathbf{v}_2 = (0, -1, 3)$; $\lambda_3 = 3$: $\mathbf{v}_3 = (0, -2, 5)$ **19.** $\lambda_1 = \lambda_2 = 1$: $\mathbf{v}_1 = (1, 0, 1), \mathbf{v}_2 = (-3, 1, 0);$ $\lambda_3 = 3, \mathbf{v}_3 = (1, 0, 0)$ **20.** $\lambda_1 = \lambda_2 = 1$: $\mathbf{v}_1 = (1, 0, 2), \mathbf{v}_2 = (3, 2, 0);$ $\lambda_3 = 2, \mathbf{v}_3 = (0, -2, 5)$ **21.** $\lambda_1 = 1$: $\mathbf{v}_1 = (1, 1, 0)$; $\lambda_2 = \lambda_3 = 2$: $\mathbf{v}_2 = (3, 2, 0)$, $\mathbf{v}_3 = (-1, 0, 2)$ **22.** $\lambda_1 = \lambda_2 = -1$: $\mathbf{v}_1 = (1, 1, 0), \mathbf{v}_2 = (-1, 0, 2);$ $\lambda_3 = 2, \mathbf{v}_3 = (1, 1, 1)$
- **23.** $\lambda_1 = 1: \mathbf{v}_1 = (1, 0, 0, 0); \lambda_2 = 2: \mathbf{v}_2 = (2, 1, 0, 0); \lambda_3 = 3: \mathbf{v}_3 = (3, 2, 1, 0); \lambda_4 = 4: \mathbf{v}_4 = (4, 3, 2, 1)$
- **24.** $\lambda_1 = \lambda_2 = 1$: $\mathbf{v}_1 = (1, 0, 0, 0), \mathbf{v}_2 = (0, 1, 0, 0);$ $\lambda_3 = \lambda_4 = 3$: $\mathbf{v}_3 = (0, 0, 0, 1), \mathbf{v}_4 = (2, 2, 1, 0)$
- **25.** $\lambda_1 = \lambda_2 = 1$: $\mathbf{v}_1 = (1, 0, 0, 0), \mathbf{v}_2 = (0, 1, 0, 0);$ $\lambda_3 = \lambda_4 = 2$: $\mathbf{v}_3 = (0, 0, 0, 1), \mathbf{v}_4 = (1, 1, 1, 0)$
- **26.** $\lambda_1 = -2$: $\mathbf{v}_1 = (1, 0, 0, 2); \lambda_2 = -1$: $\mathbf{v}_2 = (0, 0, 1, 0); \lambda_3 = 1$: $\mathbf{v}_3 = (1, 0, 0, 1); \lambda_4 = 2$: $\mathbf{v}_4 = (0, 1, 0, 0)$
- **27.** $\lambda_1 = -i$: $\mathbf{v}_1 = (+i, 1)$; $\lambda_2 = +i$: $\mathbf{v}_2 = (-i, 1)$
- **28.** $\lambda_1 = -6i$: $\mathbf{v}_1 = (-i, 1)$; $\lambda_2 = +6i$: $\mathbf{v}_2 = (+i, 1)$
- **29.** $\lambda_1 = -6i$: $\mathbf{v}_1 = (-i, 2)$; $\lambda_2 = +6i$: $\mathbf{v}_2 = (+i, 2)$

700 ANSWERS TO SELECTED PROBLEMS

- **30.** $\lambda_1 = -12i$: $\mathbf{v}_1 = (-i, 1)$; $\lambda_2 = +12i$: $\mathbf{v}_2 = (+i, 1)$
- **31.** $\lambda_1 = -12i$: $\mathbf{v}_1 = (+2i, 1)$; $\lambda_2 = +12i$: $\mathbf{v}_2 = (-2i, 1)$
- **32.** $\lambda_1 = -12i$: $\mathbf{v}_1 = (-i, 3)$; $\lambda_2 = +12i$: $\mathbf{v}_2 = (+i, 3)$
- **40.** We find that Tr **A** = 12 and det **A** = 60, so the characteristic polynomial of the given matrix **A** is $p(\lambda) = -\lambda^3 + 12\lambda^2 + c_1\lambda + 60$. Eigenvalues and eigenvectors: $\lambda_1 = 3$: $\mathbf{v}_1 = (3, 2, 1)$; $\lambda_2 = 4$: $\mathbf{v}_2 = (5, 7, 7)$; $\lambda_3 = 5$: $\mathbf{v}_3 = (-1, 1, 2)$
- **41.** We find that Tr A = 8 and det A = -60, so the characteristic polynomial of the given matrix A is $p(\lambda) = \lambda^4 8\lambda^3 + c_2\lambda^2 + c_1\lambda 60$. Eigenvalues and eigenvectors: $\lambda_1 = -2$: $\mathbf{v}_1 = (1, 0, 1, 2)$; $\lambda_2 = 2$: $\mathbf{v}_2 = (3, 4, 0, 3)$; $\lambda_3 = 3$: $\mathbf{v}_3 = (3, 1, 2, 4)$; $\lambda_4 = 5$: $\mathbf{v}_4 = (1, 1, 0, 1)$

Section 6.2

1.
$$\lambda_1 = 1, \lambda_2 = 3; \mathbf{P} = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$$

2. $\lambda_1 = 0, \lambda_2 = 2; \mathbf{P} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 0 & 0 \\ 0 & 2 \end{bmatrix}$
3. $\lambda_1 = 2, \lambda_2 = 3; \mathbf{P} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$
4. $\lambda_1 = 1, \lambda_2 = 2; \mathbf{P} = \begin{bmatrix} 1 & 4 \\ 1 & 3 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$
5. $\lambda_1 = 1, \lambda_2 = 3; \mathbf{P} = \begin{bmatrix} 1 & 4 \\ 1 & 3 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$
6. $\lambda_1 = 1, \lambda_2 = 2; \mathbf{P} = \begin{bmatrix} 2 & 3 \\ 3 & 4 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$
7. $\lambda_1 = 1, \lambda_2 = 2; \mathbf{P} = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$
8. $\lambda_1 = 1, \lambda_2 = 2; \mathbf{P} = \begin{bmatrix} 3 & 5 \\ 2 & 3 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$

- The double eigenvalue λ₁ = λ₂ = 1 has only the single associated eigenvector v₁ = (2, 1), so the matrix A is not diagonalizable.
- 10. The double eigenvalue $\lambda_1 = \lambda_2 = 2$ has only the single associated eigenvector $\mathbf{v}_1 = (1, 1)$, so the matrix **A** is not diagonalizable.
- 11. The double eigenvalue $\lambda_1 = \lambda_2 = 2$ has only the single associated eigenvector $\mathbf{v}_1 = (-1, 3)$, so the matrix **A** is not diagonalizable.
- 12. The double eigenvalue $\lambda_1 = \lambda_2 = -1$ has only the single associated eigenvector $\mathbf{v}_1 = (-3, 4)$, so the matrix **A** is not diagonalizable.

13.
$$\lambda_1 = 1, \lambda_2 = \lambda_3 = 2; \mathbf{P} = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

14. $\lambda_1 = \lambda_2 = 0, \lambda_3 = 1; \mathbf{P} = \begin{bmatrix} -1 & 1 & 1 \\ 0 & 1 & 1 \\ 2 & 0 & 1 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
15. $\lambda_1 = 0, \lambda_2 = \lambda_3 = 1; \mathbf{P} = \begin{bmatrix} 1 & -1 & 3 \\ 1 & 0 & 2 \\ 0 & 2 & 0 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

16.
$$\lambda_{1} = \lambda_{2} = 1, \lambda_{3} = 3; \mathbf{P} = \begin{bmatrix} 0 & 1 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 2 \end{bmatrix},$$

 $\mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{bmatrix}$
17. $\lambda_{1} = -1, \lambda_{2} = 1, \lambda_{3} = 2; \mathbf{P} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix},$
 $\mathbf{D} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$
18. $\lambda_{1} = 1, \lambda_{2} = 2, \lambda_{3} = 3; \mathbf{P} = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix},$
 $\mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$
19. $\lambda_{1} = 1, \lambda_{2} = 2, \lambda_{3} = 3; \mathbf{P} = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 2 \end{bmatrix},$
 $\mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$
20. $\lambda_{1} = 2, \lambda_{2} = 5, \lambda_{3} = 6; \mathbf{P} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -2 \\ 3 & 3 & 5 \end{bmatrix},$
 $\mathbf{D} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 6 \end{bmatrix}$
21. The triple eigenvalue $\lambda_{1} = \lambda_{2} = \lambda_{3} = 1$ has only the

- The triple eigenvalue λ₁ = λ₂ = λ₃ = 1 has only the two associated eigenvectors v₁ = (0, 0, 1) and v₂ = (1, 1, 0), so the matrix A is not diagonalizable.
- 22. The triple eigenvalue $\lambda_1 = \lambda_2 = \lambda_3 = 1$ has only the single associated eigenvector $\mathbf{v}_1 = (1, 1, 1)$, so the matrix **A** is not diagonalizable.
- 23. The eigenvalues $\lambda_1 = \lambda_2 = 1$ and $\lambda_3 = 2$ have only the two associated eigenvectors $\mathbf{v}_1 = (1, 1, 1)$ and $\mathbf{v}_3 = (1, 1, 0)$, so the matrix **A** is not diagonalizable.
- 24. The eigenvalues $\lambda_1 = 1$ and $\lambda_2 = \lambda_3 = 2$ have only the two associated eigenvectors $\mathbf{v}_1 = (1, 1, 0)$ and $\mathbf{v}_2 = (1, 1, 1)$, so the matrix **A** is not diagonalizable.

25.
$$\lambda_1 = \lambda_2 = -1, \lambda_3 = \lambda_4 = 1; \mathbf{P} = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix},$$

 $\mathbf{D} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$
26. $\lambda_1 = \lambda_2 = \lambda_3 = 1, \lambda_4 = 2; \mathbf{P} = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix},$
 $\mathbf{D} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$

- 27. The eigenvalues $\lambda_1 = \lambda_2 = \lambda_3 = 1$ and $\lambda_4 = 2$ have only the two associated eigenvectors $\mathbf{v}_1 = (1, 0, 0, 0)$ and $\mathbf{v}_4 = (1, 1, 1, 1)$, so the matrix **A** is not diagonalizable.
- **28.** The eigenvalues $\lambda_1 = \lambda_2 = 1$ and $\lambda_3 = \lambda_4 = 2$ have only the two associated eigenvectors $\mathbf{v}_1 = (1, 0, 0, 0)$ and $\mathbf{v}_3 = (1, 1, 1, 0)$, so the matrix **A** is not diagonalizable.

Section 6.3

1.
$$\mathbf{P} = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 63 & -62 \\ 31 & -30 \end{bmatrix}$$

2. $\mathbf{P} = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 96 & -96 \\ 64 & -64 \end{bmatrix}$
3. $\mathbf{P} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 96 & -96 \\ 64 & -64 \end{bmatrix}$
4. $\mathbf{P} = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 125 & -124 \\ 93 & -92 \end{bmatrix}$
6. $\mathbf{P} = \begin{bmatrix} 2 & 5 \\ 1 & 2 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 125 & -310 \\ 62 & -123 \end{bmatrix}$
7. $\mathbf{P} = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$
8. $\mathbf{P} = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 32 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$
4. $A^{5} = \begin{bmatrix} 1 & -62 & 31 \\ 0 & 1 & 0 \\ 0 & -62 & 32 \end{bmatrix}$
9. $\mathbf{P} = \begin{bmatrix} 1 & 1 & -3 \\ 1 & 0 & 2 & 0 \\ 0 & 0 & 32 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 94 & -93 & 31 \\ 0 & 32 & 0 \\ 0 & 0 & 32 \end{bmatrix}$
10. $\mathbf{P} = \begin{bmatrix} 1 & -73 \\ 1 & 0 & 32 & 0 \\ 0 & 0 & 32 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 94 & -93 & 31 \\ 0 & 32 & 0 \\ 0 & 0 & 32 \end{bmatrix}$
11. $\mathbf{P} = \begin{bmatrix} 1 & -93 & 31 \\ 0 & 0 & 32 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}, A^{5} = \begin{bmatrix} 94 & -93 & 31 \\ 62 & -61 & 31 \\ 0 & 0 & 32 \end{bmatrix}$
12. $\mathbf{P} = \begin{bmatrix} 1 & 1 & -3 \\ 1 & 0 & 0 \\ 2 & 0 & 5 & 0 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A^{10} = \begin{bmatrix} 1 & 0 & 0 \\ -78 & -5 & -2 \\ -195 & 15 & 6 \end{bmatrix}$
12. $\mathbf{P} = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 0 & 5 \\ 0 & 5 & 0 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A^{10} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A^{10} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A^{10} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A^{10} = \begin{bmatrix} 3 & -3 & -1 \\ 2 & -2 & 1 \\ 0 & 0 & 1 \end{bmatrix}, \mathbf{D} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A^{10} = \begin{bmatrix} 3 & -3 & -1 \\ 2 & -2 & 1 \\ 0 & 0 & 1 \end{bmatrix}$

ANSWERS TO SELECTED PROBLEM \$01

15.
$$A^2 - 3A + 2I = 0, A^3 = \begin{bmatrix} 29 & -28 \\ 21 & -20 \end{bmatrix}, A^4 = \begin{bmatrix} 61 & -60 \\ 45 & -44 \end{bmatrix}, A^{-1} = \frac{1}{2} \begin{bmatrix} -2 & 4 \\ -3 & 5 \end{bmatrix}$$

16. $A^2 - 3A + 2I = 0, A^3 = \begin{bmatrix} 36 & -70 \\ 14 & -27 \\ -2 & 6 \end{bmatrix}$
17. $-A^3 + 5A^2 - 8A + 4I = 0, A^3 = \begin{bmatrix} 1 & 21 & 0 \\ 0 & 0 & 8 \\ 0 & 0 & 8 \end{bmatrix}, A^4 = \begin{bmatrix} 1 & 45 & 0 \\ 0 & 16 & 0 \\ 0 & 0 & 16 \end{bmatrix}, A^{-1} = \frac{1}{2} \begin{bmatrix} 2 & -3 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$
18. $-A^3 + 4A^2 - 5A + 2I = 0, A^3 = \begin{bmatrix} 1 & -14 & 7 \\ 0 & 1 & 0 \\ 0 & -14 & 8 \end{bmatrix}, A^4 = \begin{bmatrix} 1 & -30 & 15 \\ 0 & 1 & 0 \\ 0 & -30 & 16 \end{bmatrix} A^{-1} = \frac{1}{2} \begin{bmatrix} 2 & 2 & -1 \\ 0 & 2 & 0 \\ 0 & 2 & 1 \end{bmatrix}$
19. $-A^3 + 5A^2 - 8A + 4I = 0, A^3 = \begin{bmatrix} 1 & -21 & 7 \\ 0 & 1 & 0 \\ 0 & -30 & 16 \end{bmatrix}, A^{-1} = \frac{1}{2} \begin{bmatrix} 2 & 3 & -1 \\ 0 & 2 & 0 \\ 0 & 2 & 1 \end{bmatrix}$
19. $-A^3 + 5A^2 - 8A + 4I = 0, A^3 = \begin{bmatrix} 22 & -21 & 7 \\ 0 & 8 & 0 \\ 0 & 0 & 16 \end{bmatrix}, A^{-1} = \frac{1}{2} \begin{bmatrix} 2 & 3 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
20. $-A^3 + 5A^2 - 8A + 4I = 0, A^3 = \begin{bmatrix} 22 & -21 & 7 \\ 14 & -13 & 7 \\ 0 & 0 & 8 \end{bmatrix}, A^4 = \begin{bmatrix} 46 & -45 & 15 \\ 30 & -29 & 15 \\ 0 & 0 & 16 \end{bmatrix}, A^{-1} = \frac{1}{2} \begin{bmatrix} -1 & 3 & -1 \\ -2 & 4 & -1 \\ 0 & 0 & 1 \end{bmatrix}$
21. $-A^3 + A = 0, A^3 = A = \begin{bmatrix} 1 & 0 & 0 \\ 6 & 5 & 2 \\ 21 & -15 & -6 \end{bmatrix}, A^4 = A^2 = \begin{bmatrix} 1 & 0 & 0 \\ -195 & 15 & -6 \end{bmatrix}$. Because $\lambda = 0$ is an eigenvalue, A is singular and A^{-1} does not exist.
22. $-A^3 + A^2 + A - I = 0, A^3 = A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 1 \\ 0 & 0 & 1 \end{bmatrix} = I, A^{-1} = A$
23. $-A^3 + A = 0, A^3 = A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 1 \\ 0 & 0 & 1 \end{bmatrix}$. Because $\lambda = 0$ is an eigenvalue, A is singular and A^{-1} does not exist.
24. $-A^3 + A = 0, A^3 = A = \begin{bmatrix} 2 & -2 & -3 \\ 2 & -2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$. Because $\lambda = 0$ is an eigenvalue, A is singular and A^{-1} does not exist.
24. $-A^3 + A = 0, A^3 = A = \begin{bmatrix} 5 & -5 & -3 \\ 2 & -2 & -1 \\ 4 & -4 & -3 \end{bmatrix}, A^4 = A^2 = \begin{bmatrix} 3 & -3 & 1 \\ 2 & -2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$. Because $\lambda = 0$ is an eigenvalue, A is singular and A^{-1} does not exist.
25. $x_k = A^k x_0 = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 4/5 \end{bmatrix}^k \frac{1}{2} \begin{bmatrix} -1 & 1 \\ -1 & 1 \end{bmatrix} x_0 \rightarrow = (C_0 + S_0) \begin{bmatrix} 1/2 \\ 1/2 \end{bmatrix}$ as $k \rightarrow \infty$. The long-term distribution of

702 ANSWERS TO SELECTED PROBLEMS

population is 50% city, 50% suburban.

26.
$$\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 1 & -1 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 4/5 \end{bmatrix}^k \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -3 & 1 \end{bmatrix} \mathbf{x}_0 \rightarrow = (C_0 + S_0) \begin{bmatrix} 1/4 \\ 3/4 \\ 3/4 \end{bmatrix} \text{ as } k \rightarrow \infty.$$
 The long-term distribution of population is 25% city, 75% suburban.
27. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 3 & -1 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 3/5 \end{bmatrix}^k \frac{1}{8} \begin{bmatrix} -1 & 1 \\ -5 & 3 \end{bmatrix} \mathbf{x}_0 \rightarrow = (C_0 + S_0) \begin{bmatrix} 3/8 \\ 5/8 \\ 5/8 \end{bmatrix} \text{ as } k \rightarrow \infty.$ The long-term distribution of population is 3/8 city, 5/8 suburban.
28. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 7/10 \end{bmatrix}^k \frac{1}{3} \begin{bmatrix} -1 & 1 \\ -2 & 1 \end{bmatrix} \mathbf{x}_0 \rightarrow = (C_0 + S_0) \begin{bmatrix} 1/3 \\ 2/3 \end{bmatrix} \text{ as } k \rightarrow \infty.$ The long-term distribution of population is 1/3 city, 2/3 suburban.
29. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 1 & 0 \\ 1/2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1/2 \end{bmatrix}^k \frac{1}{3} \begin{bmatrix} 1 & 1 \\ -2 & 1 \end{bmatrix} \mathbf{x}_0 \rightarrow = (C_0 + S_0) \begin{bmatrix} 1/3 \\ 1/3 \\ 2/3 \end{bmatrix} \text{ as } k \rightarrow \infty.$ The long-term distribution of population is 1/3 city, 2/3 suburban.
30. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 3 & -1 \\ 0 & 13/20 \end{bmatrix}^k \frac{1}{7} \begin{bmatrix} 1 & 1 \\ -4 & 3 \end{bmatrix} \mathbf{x}_0 \rightarrow = (C_0 + S_0) \begin{bmatrix} 3/7 \\ 4/7 \\ 4/7 \end{bmatrix} \text{ as } k \rightarrow \infty.$ The long-term distribution of population is 1/3 city, 2/3 suburban.
31. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 5 & 5 \\ 5 & 5 \\ 2.5 R_0 - F_0 \\ 2.6 & 0.8 F_0 \end{bmatrix} \text{ as } k \rightarrow \infty.$ The long-term distribution of population is 3/7 city, 4/7 suburban.
32. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 1 & 0 \\ 0 & 3/4 \end{bmatrix}^k \frac{1}{60} \begin{bmatrix} -2 & 5 \\ 4 & -5 \end{bmatrix} \mathbf{x}_0 \rightarrow = \begin{bmatrix} 2.5 R_0 - F_0 \\ 0 \\ 3 \end{bmatrix} \mathbf{x}_k \rightarrow \infty.$ The fox and rabbit population approaches a stable situation with 2.5 R_0 - F_0 foxes and 2R_0 - 0.8 F_0 rabbits.
32. $\mathbf{x}_k = \mathbf{A}^k \mathbf{x}_0 = \begin{bmatrix} 1 & 0 \\ 0 & 3/4 \end{bmatrix}^k \frac{1}{60} \begin{bmatrix} -3 & 10 \\ 9 & -10 \end{bmatrix} \mathbf{x}_0 \Rightarrow = \begin{bmatrix} 10 & 0 \\ 0 & 3/4 \end{bmatrix} \begin{bmatrix} 21/20 & 0 \\ 0 & 3/4 \end{bmatrix}^k \frac{1}{60} \begin{bmatrix} -3 & 10 \\ 9 & -10 \end{bmatrix} \mathbf{x}_0 \approx \frac{1}{60} (10.5)^k (10R_0 - 3F_0) \begin{bmatrix} 1 \\ 9 \end{bmatrix}$ as when k is sufficiently large. The fox and rabbit populations are both increasing at 5% per year, with 10 foxes for each 9 rabbits.
34. $\mathbf{A} = PDP^{-1} = I = PP^{-1} = I.$ If *n* is even, then $D^n = I$ so $A^n = PD^nP^{-1} = I = PP^{-1} = I.$ If *n* is even, then $D^n = I$ so $A^n = PD^n$

Chapter 7

Section 7.1

1. $x'_1 = x_2, x'_2 = -7x_1 - 3x_2 + t^2$ 2. $x'_1 = x_2, x'_2 = x_3, x'_3 = x_4, x'_4 = -x_1 + 3x_2 - 6x_3 + \cos 3t$ 3. $x'_1 = x_2, t^2 x'_2 = (1 - t^2)x_1 - tx_2$ 4. $x'_1 = x_2, x'_2 = x_3, t^3 x'_3 = -5x_1 - 3tx_2 + 2t^2 x_3 + \ln t$ 5. $x'_1 = x_2, x'_2 = x_3, x'_3 = x_2^2 + \cos x_1$ 6. $x'_1 = x_2, x'_2 = 5x_1 - 4y_1, y'_1 = y_2, y'_2 = -4x_1 + 5y_1$ 7. $x'_1 = x_2, y'_1 = y_2, x'_2 = -kx_1 \cdot (x_1^2 + y_1^2)^{-3/2}, y'_2 = -ky_1 \cdot (x_1^2 + y_1^2)^{-3/2}$ 8. $x'_1 = x_2, x'_2 = 3x_1 - y_1 - 2y_2 + \cos t$ 9. $x'_1 = x_2, y'_1 = y_2, z'_1 = z_2, x'_2 = 3x_1 - y_1 + 2z_1, y'_2 = x_1 + y_1 - 4z_1, z'_2 = 5x_1 - y_1 - z_1$ 10. $x'_1 = x_2, x'_2 = x_1(1 - y_1) y'_1 = y_2, y'_2 = y_1(1 - x_1)$

11.
$$x(t) = A\cos t + B\sin t$$
, $y(t) = B\cos t - A\sin t$



12.
$$x(t) = Ae^{t} + Be^{-t}$$
, $y(t) = Ae^{t} - Be^{-t}$



13. $x(t) = A\cos 2t + B\sin 2t$, $y(t) = -B\cos 2t + A\sin 2t$; $x(t) = \cos 2t$, $y(t) = \sin 2t$

ANSWERS TO SELECTED PROBLEM 303

17.
$$x(t) = Ae^{-3t} + Be^{2t}$$
, $y(t) = -3Ae^{-3t} + 2Be^{2t}$;
 $x(t) = e^{2t}$, $y(t) = 2e^{2t}$



18. $x(t) = Ae^{-2t} + Be^{-5t}$, $y(t) = 2Ae^{-2t} + 5Be^{-5t}$; $A = \frac{17}{3}$ and $B = -\frac{11}{3}$ in the particular solution.



19. $x(t) = -e^{-2t} \sin 3t$, $y(t) = e^{-2t} (3 \cos 3t + 2 \sin 3t)$





14. $x(t) = A\cos 10t + B\sin 10t, y(t) = B\cos 10t - A\sin 10t;$ $x(t) = 3\cos 10t + 4\sin 10t, y(t) = 4\cos 10t - 3\sin 10t$



15. $x(t) = A \cos 2t + B \sin 2t$, $y(t) = 4B \cos 2t - 4A \sin 2t$



16. $x(t) = A\cos 4t + B\sin 4t$, $y(t) = \frac{1}{2}B\cos 4t - \frac{1}{2}A\sin 4t$







Section 7.2

- **1.** $(\mathbf{AB})' = \begin{bmatrix} 1 8t + 18t^2 & 1 + 2t 12t^2 + 32t^3 \\ 3 + 3t^2 4t^3 & 8t + 3t^2 + 4t^3 \end{bmatrix}$ 3. $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}, \mathbf{P}(t) = \begin{bmatrix} 0 & -3 \\ 3 & 0 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ 5. $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}, \mathbf{P}(t) = \begin{bmatrix} 2 & 4 \\ 5 & -1 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 3e^t \\ -t^2 \end{bmatrix}$ 7. $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $\mathbf{P}(t) = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$, $\mathbf{f}(t) = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ **9.** $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \mathbf{P}(t) = \begin{bmatrix} 3 & -4 & 1 \\ 1 & 0 & -3 \\ 0 & 6 & -7 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} t \\ t^2 \\ t^3 \end{bmatrix}$ **11.** $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}, \mathbf{P}(t) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 4 & 0 & 0 & 0 \end{bmatrix}, \mathbf{f}(t) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$ **13.** $W(t) = e^{3t}$; $\mathbf{x}(t) = \begin{bmatrix} 2c_1e^t + c_2e^{2t} \\ -3c_1e^t - c_2e^{2t} \end{bmatrix}$ **15.** $W(t) \equiv 4$; $\mathbf{x}(t) = \begin{bmatrix} c_1 e^{2t} + c_2 e^{-2t} \\ c_1 e^{2t} + 5c_2 e^{-2t} \end{bmatrix}$ **17.** $W(t) = 7e^{-3t}$; $\mathbf{x}(t) = \begin{bmatrix} 3c_1e^{2t} + c_2e^{-5t} \\ 2c_1e^{2t} + 3c_2e^{-5t} \end{bmatrix}$ **19.** $W(t) \equiv 3$; $\mathbf{x}(t) = \begin{bmatrix} c_1 e^{2t} + c_2 e^{-t} \\ c_1 e^{2t} + c_3 e^{-t} \\ c_1 e^{2t} - (c_2 + c_3) e^{-t} \end{bmatrix}$ 21. $W(t) = e^{2t}$; $\mathbf{x}(t) = \begin{bmatrix} 3c_1e^{-2t} + c_2e^t + c_3e^{3t} \\ -2c_1e^{-2t} - c_2e^t - c_3e^{3t} \\ 2c_1e^{-2t} + c_2e^t \end{bmatrix}$ **23.** $\mathbf{x} = 2\mathbf{x}_1 - \mathbf{x}_2$ 25. $x = 15x_1 - 4x_2$ 27. $\mathbf{x} = \mathbf{x}_1 + 2\mathbf{x}_2 + \mathbf{x}_3$ **29.** $\mathbf{x} = 3\mathbf{x}_1 - 3\mathbf{x}_2 - 5\mathbf{x}_3$ **31.** $\mathbf{x} = 3\mathbf{x}_1 + 7\mathbf{x}_2 + \mathbf{x}_3 - 2\mathbf{x}_4$ **32.** $\mathbf{x} = 13\mathbf{x}_1 + 41\mathbf{x}_2 + 3\mathbf{x}_3 - 2\mathbf{x}_4$ $12x_4$
- 33. (a) x₂ = tx₁, so neither is a constant multiple of the other.
 (b) W(x₁, x₂) ≡ 0, whereas Theorem 2 implies that W ≠ 0 if x₁ and x₂ were independent solutions of a system of the indicated form.

Section 7.3

1.
$$x_1(t) = c_1 e^{-t} + c_2 e^{3t}, x_2(t) = -c_1 e^{-t} + c_2 e^{3t}$$



- 2. $x_1(t) = c_1 e^{-t} + 3c_2 e^{4t}, x_2(t) = -c_1 e^{-t} + 2c_2 e^{4t}$
- 3. General solution $x_1(t) = c_1 e^{-t} + 4c_2 e^{6t}$, $x_2(t) = -c_1 e^{-t} + 3c_2 e^{6t}$ Particular solution $x_1(t) = \frac{1}{7}(-e^{-t} + 8e^{6t})$, $x_2(t) = \frac{1}{7}(e^{-t} + 6e^{6t})$.
- 4. $x_1(t) = c_1 e^{-2t} + c_2 e^{5t}$, $x_2(t) = -6c_1 e^{-2t} + c_2 e^{5t}$



5. $x_1(t) = c_1 e^{-t} + 7c_2 e^{5t}, x_2(t) = c_1 e^{-t} + c_2 e^{5t}$



6. General solution $x_1(t) = 5c_1e^{3t} + c_2e^{4t}$, $x_2(t) = -6c_1e^{3t} - c_2e^{4t}$ Particular solution $x_1(t) = -5e^{3t} + 6e^{4t}$,

ANSWERS TO SELECTED PROBLEM \$05

$$x_2(t) = 6e^{3t} - 6e^{4t}$$



7. $x_1(t) = c_1 e^t + 2c_2 e^{-9t}, \ x_2(t) = c_1 e^t - 3c_2 e^{-9t}$



8. $x_1(t) = 5c_1 \cos 2t + 5c_2 \sin 2t$, $x_2(t) = (c_1 - 2c_2) \cos 2t + (2c_1 + c_2) \sin 2t$



- 9. General solution $x_1(t) = 5c_1 \cos 4t + 5c_2 \sin 4t$, $x_2(t) = c_1(2\cos 4t + 4\sin 4t) + c_2(2\sin 4t - 4\cos 4t)$. Particular solution $x_1(t) = 2\cos 4t - \frac{11}{4}\sin 4t$, $x_2(t) = 3\cos 4t + \frac{1}{2}\sin 4t$
- **10.** $x_1(t) = -2c_1 \cos 3t 2c_2 \sin 3t$, $x_2(t) = (3c_1 + 3c_2) \cos 3t + (3c_2 - 3c_1) \sin 3t$
- 11. General solution $x_1(t) = e^t (c_1 \cos 2t c_2 \sin 2t)$, $x_2(t) = e^t (c_1 \sin 2t + c_2 \cos 2t)$ Particular solution $x_1(t) = -4e^t \sin 2t$, $x_2(t) = 4e^t \cos 2t$



- 12. $x_1(t) = e^{2t} (-5c_1 \cos 2t 5c_2 \sin 2t),$ $x_2(t) = e^{2t} [(c_1 + 2c_2) \cos 2t + (-2c_1 + c_2) \sin 2t]$
- 13. $x_1(t) = 3e^{2t}(c_1 \cos 3t c_2 \sin 3t),$ $x_2(t) = e^{2t}[(c_1 + c_2)\cos 3t + (c_1 - c_2)\sin 3t]$
- **14.** $x_1(t) = e^{3t}(c_1\cos 4t + c_2\sin 4t),$ $x_2(t) = e^{3t}(c_1\sin 4t - c_2\cos 4t)$
- **15.** $x_1(t) = 5e^{5t}(c_1\cos 4t c_2\sin 4t),$ $x_2(t) = e^{5t}[(2c_1 + 4c_2)\cos 4t + (4c_1 - 2c_2)\sin 4t]$
- **16.** $x_1(t) = c_1 e^{-10t} + 2c_2 e^{-100t},$ $x_2(t) = 2c_1 e^{-10t} - 5c_2 e^{-100t}$



- **17.** $x_1(t) = c_1 e^{9t} + c_2 e^{6t} + c_3,$ $x_2(t) = c_1 e^{9t} - 2c_2 e^{6t},$ $x_3(t) = c_1 e^{9t} + c_2 e^{6t} - c_3$
- **18.** $x_1(t) = c_1 e^{9t} + 4c_3,$ $x_2(t) = 2c_1 e^{9t} + c_2 e^{6t} - c_3,$ $x_3(t) = 2c_1 e^{9t} - c_2 e^{6t} - c_3$
- **19.** $x_1(t) = c_1 e^{6t} + c_2 e^{3t} + c_3 e^{3t},$ $x_2(t) = c_1 e^{6t} - 2c_2 e^{3t},$
- $x_3(t) = c_1 e^{6t} + c_2 e^{3t} c_3 e^{3t}$ 20. $x_1(t) = c_1 e^{9t} + c_2 e^{6t} + c_3 e^{2t}$,
- $x_{2}(t) = c_{1}e^{9t} 2c_{2}e^{6t},$ $x_{3}(t) = c_{1}e^{9t} + c_{2}e^{6t} - c_{3}e^{2t}$
- **21.** $x_1(t) = 6c_1 + 3c_2e^t + 2c_3e^{-t},$ $x_2(t) = 2c_1 + c_2e^t + c_3e^{-t},$ $x_3(t) = 5c_1 + 2c_2e^t + 2c_3e^{-t}$
- 22. $x_1(t) = c_2 e^t + c_3 e^{3t}$, $x_2(t) = c_1 e^{-2t} - c_2 e^t - c_3 e^{3t}$, $x_3(t) = -c_1 e^{-2t} + c_3 e^{3t}$
- 23. $x_1(t) = c_1 e^{2t} + c_3 e^{3t}$, $x_2(t) = -c_1 e^{2t} + c_2 e^{-2t} - c_3 e^{3t}$, $x_3(t) = -c_2 e^{-2t} + c_3 e^{3t}$
- 24. $x_1(t) = c_1 e^t + c_2(2\cos 2t \sin 2t) + c_3(\cos 2t + 2\sin 2t)$ $x_2(t) = -c_1 e^t - c_2(3\cos 2t + \sin 2t) + c_3(\cos 2t - 3\sin 2t)$ $x_3(t) = c_2(3\cos 2t + \sin 2t) + c_3(3\sin 2t - \cos 2t)$
706 ANSWERS TO SELECTED PROBLEMS

- 25. $x_1(t) = c_1 + e^{2t} [(c_2 + c_3) \cos 3t + (-c_2 + c_3) \sin 3t],$ $x_2(t) = -c_1 + 2e^{2t} (-c_2 \cos 3t - c_3 \sin 3t),$ $x_3(t) = 2e^{2t} (c_2 \cos 3t + c_3 \sin 3t)$
- **26.** $x_1(t) = 4e^{3t} e^{-t}(4\cos t \sin t),$ $x_2(t) = 9e^{3t} - e^{-t}(9\cos t + 2\sin t),$ $x_3(t) = 17e^{-t}\cos t$
- **27.** $x_1(t) = 15e^{-0.2t}, x_2(t) = 15(e^{-0.2t} e^{-0.4t}).$ The maximum amount ever in tank 2 is $x_2(5 \ln 2) = 3.75$ lb.



28. $x_1(t) = 15e^{-0.4t}, x_2(t) = 40(-e^{-0.4t} + e^{-0.25t}).$ The maximum amount ever in tank 2 is about 6.85 lb.



29. $x_1(t) = 10 + 5e^{-0.6t}, x_2(t) = 5 - 5e^{-0.6t}$



30. $x_1(t) = 5c_1 + c_2e^{-0.65t}, x_2(t) = 8c_1 - c_2e^{-0.65t}.$



31. $x_1(t) = 27e^{-t}$, $x_2(t) = 27e^{-t} - 27e^{-2t}$, $x_3(t) = 27e^{-t} - 54e^{-2t} + 27e^{-3t}$. The maximum amount of salt ever in tank 3 is $x_3(\ln 3) = 4$ pounds.



32. $x_1(t) = 45e^{-3t}$, $x_2(t) = -135e^{-3t} + 135e^{-2t}$, $x_3(t) = 135e^{-3t} - 270e^{-2t} + 135e^{-t}$. The maximum amount of salt ever in tank 3 is $x_3(\ln 3) = 20$ pounds.



33. $x_1(t) = 45e^{-4t}$, $x_2(t) = 90e^{-4t} - 90e^{-6t}$, $x_3(t) = -270e^{-4t} + 135e^{-6t} + 135e^{-2t}$. The maximum amount of salt ever in tank 3 is $x_3(\frac{1}{2}\ln 3) = 20$ pounds.



34. $x_1(t) = 40e^{-3t}$, $x_2(t) = 60e^{-3t} - 60e^{-5t}$, $x_3(t) = -150e^{-3t} + 75e^{-5t} + 75e^{-t}$. The maximum amount of salt ever in tank 3 is $x_3(\frac{1}{2} \ln 5) \approx 21.4663$ pounds.



35. $x_1(t) = 10 - \frac{1}{7} (55e^{-18t} - 216e^{-11t}),$ $x_2(t) = 3 - \frac{1}{7} (165e^{-18t} - 144e^{-11t}),$ $x_3(t) = 20 + \frac{1}{7} (220e^{-18t} - 360e^{-11t}).$ The limiting amounts of salt in tanks 1, 2, and 3 are 10 lb, 3 lb, and 20 lb.



36. $x_1(t) = 4 + e^{-3t/5} [14\cos(3t/10) - 2\sin(3t/10)],$ $x_2(t) = 10 - e^{-3t/5} [10\cos(3t/10) - 10\sin(3t/10)],$ $x_3(t) = 4 - e^{-3t/5} [4\cos(3t/10) + 8\sin(3t/10)].$ The limiting amounts of salt in tanks 1, 2, and 3 are 4 lb, 10 lb, and 4 lb.



37. $x_1(t) = 30 + e^{-3t} [25 \cos(t\sqrt{2}) + 10\sqrt{2} \sin(t\sqrt{2})],$ $x_2(t) = 10 - e^{-3t} [10 \cos(t\sqrt{2}) - \frac{25}{2}\sqrt{2} \sin(t\sqrt{2})],$ $x_3(t) = 15 - e^{-3t} [15 \cos(t\sqrt{2}) + \frac{45}{2}\sqrt{2} \sin(t\sqrt{2})].$ The limiting amounts of salt in tanks 1, 2, and 3 are 30 lb, 10 lb, and 15 lb.





Section 7.4

Note that phase portraits for Problems 1–16 are found in the answers for Section 7.2.

- 1. Saddle point (real eigenvalues of opposite sign)
- 2. Saddle point (real eigenvalues of opposite sign)
- 3. Saddle point (real eigenvalues of opposite sign)
- 4. Saddle point (real eigenvalues of opposite sign)
- 5. Saddle point (real eigenvalues of opposite sign)
- 6. Improper nodal source (distinct positive real eigenvalues)
- 7. Saddle point (real eigenvalues of opposite sign)
- 8. Center (pure imaginary eigenvalues)
- 9. Center (pure imaginary eigenvalues)
- **10.** Center (pure imaginary eigenvalues)
- **11.** Spiral source (complex conjugate eigenvalues with positive real part)

- **12.** Spiral source (complex conjugate eigenvalues with positive real part)
- **13.** Spiral source (complex conjugate eigenvalues with positive real part)
- **14.** Spiral source (complex conjugate eigenvalues with positive real part)
- **15.** Spiral source (complex conjugate eigenvalues with positive real part)
- 16. Improper nodal sink (distinct negative real eigenvalues)
- 17. Center; pure imaginary eigenvalues
- **18.** Improper nodal source; distinct positive real eigenvalues; $\mathbf{v}_1 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$, $\mathbf{v}_2 = \begin{bmatrix} -1 & 1 \end{bmatrix}^T$
- **19.** Saddle point; real eigenvalues of opposite sign; $\mathbf{v}_1 = \begin{bmatrix} 0 & 1 \end{bmatrix}^T$ corresponds to the negative eigenvalue and $\mathbf{v}_2 = \begin{bmatrix} -1 & 1 \end{bmatrix}^T$ to the positive one.
- **20.** Spiral source; complex conjugate eigenvalues with positive real part
- Proper nodal source; repeated positive real eigenvalue with linearly independent eigenvectors
- **22.** Parallel lines; one zero and one negative real eigenvalue
- **23.** Spiral sink; complex conjugate eigenvalues with negative real part
- **24.** Improper nodal sink; distinct negative real eigenvalues; $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 \end{bmatrix}^T$, $\mathbf{v}_2 = \begin{bmatrix} -1 & 4 \end{bmatrix}^T$
- **25.** Saddle point; real eigenvalues of opposite sign; $\mathbf{v}_1 = \begin{bmatrix} 1 & 1 \end{bmatrix}^T$ corresponds to the positive eigenvalue and $\mathbf{v}_2 = \begin{bmatrix} 4 & -1 \end{bmatrix}^T$ to the negative one.
- **26.** Center; pure imaginary eigenvalues
- 27. Improper nodal source; distinct positive real eigenvalues; $\mathbf{v}_1 = \begin{bmatrix} 2 & 3 \end{bmatrix}^T$, $\mathbf{v}_2 = \begin{bmatrix} 2 & -1 \end{bmatrix}^T$.
- **28.** Spiral sink; complex conjugate eigenvalues with negative real part

Section 7.5

- The natural frequencies are ω₀ = 0 and ω₁ = 2. In the degenerate natural mode with "frequency" ω₀ = 0 the two masses move linearly with x₁(t) = x₂(t) = a₀ + b₀t. At frequency ω₁ = 2 they oscillate in opposite directions with equal amplitudes.
- 2. The natural frequencies are $\omega_1 = 1$ and $\omega_2 = 3$. In the natural mode with frequency ω_1 , the two masses m_1 and m_2 move in the same direction with equal amplitudes of oscillation. At frequency ω_2 they move in opposite directions with equal amplitudes.
- 3. The natural frequencies are $\omega_1 = 1$ and $\omega_2 = 2$. In the natural mode with frequency ω_1 , the two masses m_1 and m_2 move in the same direction with equal amplitudes of oscillation. In the natural mode with frequency ω_2 they move in opposite directions with the amplitude of oscillation of m_1 twice that of m_2 .
- 4. The natural frequencies are $\omega_1 = 1$ and $\omega_2 = \sqrt{5}$. In the natural mode with frequency ω_1 , the two masses m_1 and m_2 move in the same direction with equal amplitudes of oscillation. At frequency ω_2 they move in opposite directions with equal amplitudes.
- 5. The natural frequencies are $\omega_1 = \sqrt{2}$ and $\omega_2 = 2$. In the natural mode with frequency ω_1 , the two masses m_1 and m_2 move in the same direction with equal amplitudes of oscillation. At frequency ω_2 they move in opposite directions with equal amplitudes.

- 6. The natural frequencies are ω₁ = √2 and ω₂ = √8. In the natural mode with frequency ω₁, the two masses m₁ and m₂ move in the same direction with equal amplitudes of oscillation. In the natural mode with frequency ω₂ they move in opposite directions with the amplitude of oscillation of m₁ twice that of m₂.
- 7. The natural frequencies are $\omega_1 = 2$ and $\omega_2 = 4$. In the natural mode with frequency ω_1 , the two masses m_1 and m_2 move in the same direction with equal amplitudes of oscillation. At frequency ω_2 they move in opposite directions with equal amplitudes.
- 8. $x_1(t) = 2\cos t + 3\cos 3t 5\cos 5t$, $x_2(t) = 2\cos t - 3\cos 3t + \cos 5t$. We have a superposition of three oscillations, in which the two masses move (1) in the same direction with frequency $\omega_1 = 1$ and equal amplitudes; (2) in opposite directions with frequency $\omega_2 = 3$ and equal amplitudes; (3) in opposite directions with frequency $\omega_3 = 5$ and with the amplitude of motion of m_1 being 5 times that of m_2 .
- 9. x₁(t) = 5 cos t 8 cos 2t + 3 cos 3t, x₂(t) = 5 cos t + 4 cos 2t - 9 cos 3t. We have a superposition of three oscillations, in which the two masses move (1) in the same direction with frequency ω₁ = 1 and equal amplitudes; (2) in opposite directions with frequency ω₂ = 2 and with the amplitude of motion of m₁ being twice that of m₂; (3) in opposite directions with frequency ω₃ = 3 and with the amplitude of motion of m₂ being 3 times that of m₁.
- **10.** $x_1(t) = -15 \cos 2t + \cos 4t + 14 \cos t$, $x_2(t) = -15 \cos 2t - \cos 4t + 16 \cos t$. We have a superposition of three oscillations, in which the two masses move (1) in the same direction with frequency $\omega_1 = 1$ and with the amplitude of motion of m_2 being 8/7 times that of m_1 ; (2) in the same direction with frequency $\omega_2 = 2$ and equal amplitudes; (3) in opposite directions with frequency $\omega_3 = 4$ and equal amplitudes.
- (a) The natural frequencies are ω₁ = 6 and ω₂ = 8. In mode 1 the two masses oscillate in the same direction with frequency ω₁ = 6 and with the amplitude of motion of m₁ being twice that of m₂. In mode 2 the two masses oscillate in opposite directions with frequency ω₂ = 8 and with the amplitude of motion of m₂ being 3 times that of m₁.
 (b) x(t) = 2 sin 6t + 19 cos 7t, y(t) = sin 6t + 3 cos 7t We have a superposition of (only two) oscillations, in which the

two masses move (1) in the same direction with frequency $\omega_1 = 6$ and with the amplitude of motion of m_1 being twice that of m_2 ; (2) in the same direction with frequency $\omega_3 = 7$ and with the amplitude of motion of m_1 being 19/3 times that of m_2 .

- 12. The system's three natural modes of oscillation have (1) natural frequency ω₁ = √2 with amplitude ratios 1:0:-1; (2) natural frequency ω₂ = √2 + √2 with amplitude ratios 1:-√2:1; (3) natural frequency ω₃ = √2 √2 with amplitude ratios
- 1: √2:1.
 13. The system's three natural modes of oscillation have (1) natural frequency ω₁ = 2 with amplitude ratios 1:0:-1; (2) natural frequency ω₂ = √4 + 2√2 with amplitude ratios 1:-√2:1; (3) natural frequency ω₃ = √4 2√2 with amplitude ratios 1:
- $\sqrt{2}$: 1. **15.** $x_1(t) = \frac{2}{3}\cos 5t - 2\cos 5\sqrt{3}t + \frac{4}{3}\cos 10t$, $x_2(t) = \frac{4}{3}\cos 5t + 4\cos 5\sqrt{3}t + \frac{16}{3}\cos 10t$. We have a superposition of two oscillations with the natural frequencies $\omega_1 = 5$ and $\omega_2 = 5\sqrt{3}$ and a forced oscillation with frequency $\omega = 10$. In each of the two natural oscillations

the amplitude of motion of m_2 is twice that of m_1 , while in the forced oscillation the amplitude of motion of m_2 is four times that of m_1 .

20.
$$x'_1(t) = -v_0, x'_2(t) = 0, x'_1(t) = v_0$$
 for $t > \pi/2$

21.
$$x'_1(t) = -v_0, x'_2(t) = 0, x'_1(t) = 2v_0$$
 for $t > \pi/2$

22.
$$x'_1(t) = -2v_0, x'_2(t) = v_0, x'_1(t) = v_0$$
 for $t > \pi/2$

- **23.** $x'_1(t) = 2v_0, x'_2(t) = 2v_0, x'_1(t) = 3v_0$ for $t > \pi/2$
- **24.** (a) $\omega_1 \approx 1.0293$ Hz; $\omega_2 \approx 1.7971$ Hz. (b) $v_1 \approx 28$ mi/h; $v_2 \approx 49$ mi/h
- **27.** $\omega_1 = 2\sqrt{10}, v_1 \approx 40.26$ (ft/s (about 27 mi/h), $\omega_2 = 5\sqrt{5}, v_2 \approx 71.18$ ft/s (about 49 mi/h)
- **28.** $\omega_1 \approx 6.1311, v_1 \approx 39.03$ ft/s (about 27 mi/h) $\omega_2 \approx 10.3155, v_2 \approx 65.67$ ft/s (about 45 mi/h)
- **29.** $\omega_1 \approx 5.0424, v_1 \approx 32.10$ ft/s (about 22 mi/h), $\omega_2 \approx 9.9158, v_2 \approx 63.13$ ft/s (about 43 mi/h)

Section 7.6

1. Repeated eigenvalue $\lambda = -3$, eigenvector $\mathbf{v} = \begin{bmatrix} 1 & -1 \end{bmatrix}^T$; $x_1(t) = (c_1 + c_2 + c_2t)e^{-3t}$, $x_2(t) = (-c_1 - c_2t)e^{-3t}$



2. Repeated eigenvalue $\lambda = 2$, single eigenvector $\mathbf{v} = \begin{bmatrix} 1 & 1 \end{bmatrix}^T$; $x_1(t) = (c_1 + c_2 + c_2 t)e^{2t}, x_2(t) = (c_1 + c_2 t)e^{2t}$



- **3.** Repeated eigenvalue $\lambda = 3$, eigenvector $\mathbf{v} = \begin{bmatrix} -2 & 2 \end{bmatrix}^T$; $x_1(t) = (-2c_1 + c_2 - 2c_2t)e^{3t}, x_2(t) = (2c_1 + 2c_2t)e^{3t}$
- 4. Repeated eigenvalue $\lambda = 4$, single eigenvector $\mathbf{v} = \begin{bmatrix} -1 & 1 \end{bmatrix}^T$; $x_1(t) = (-c_1 + c_2 - c_2 t)e^{4t}$, $x_2(t) = (c_1 + c_2 t)e^{4t}$
- 5. Repeated eigenvalue $\lambda = 5$, eigenvector $\mathbf{v} = \begin{bmatrix} 2 & -4 \end{bmatrix}^T$; $x_1(t) = (2c_1 + c_2 + 2c_2t)e^{5t}$, $x_2(t) = (-4c_1 - 4c_2t)e^{5t}$



6. Repeated eigenvalue $\lambda = 5$, single eigenvector $\mathbf{v} = \begin{bmatrix} -4 & 4 \end{bmatrix}^T$; $x_1(t) = (-4c_1 + c_2 - 4c_2t)e^{5t}$, $x_2(t) = (4c_1 + 4c_2t)e^{5t}$



- 7. Eigenvalues $\lambda = 2, 2, 9$ with three linearly independent eigenvectors; $x_1(t) = c_1 e^{2t} + c_2 e^{2t}$, $x_2(t) = c_1 e^{2t} + c_3 e^{9t}$, $x_3(t) = c_2 e^{2t}$
- 8. Eigenvalues $\lambda = 7, 13, 13$ with three linearly independent eigenvectors; $x_1(t) = 2c_1e^{7t} - c_3e^{13t}$, $x_2(t) = -3c_1e^{7t} + c_3e^{13t}$, $x_3(t) = c_1e^{7t} + c_2e^{13t}$
- 9. Eigenvalues $\lambda = 5, 5, 9$ with three linearly independent eigenvectors; $x_1(t) = c_1 e^{5t} + 7c_2 e^{5t} + 3c_3 e^{9t}$, $x_2(t) = 2c_1 e^{5t}, x_3(t) = 2c_2 e^{5t} + c_3 e^{9t}$
- **10.** Eigenvalues $\lambda = 3, 3, 7$ with three linearly independent eigenvectors; $x_1(t) = 5c_1e^{3t} 3c_2e^{3t} + 2c_3e^{7t}$, $x_2(t) = 2c_1e^{3t} + c_3e^{7t}$, $x_3(t) = c_2e^{3t}$
- 11. Triple eigenvalue $\lambda = -1$ of defect 2; $x_1(t) = (-2c_2 + c_3 - 2c_3t)e^{-t}$, $x_2(t) = (c_1 - c_2 + c_2t - c_3t + \frac{1}{2}c_3t^2)e^{-t}$, $x_3(t) = (c_2 + c_3t)e^{-t}$
- 12. Triple eigenvalue $\lambda = -1$ of defect 2; $x_1(t) = e^{-t}(c_1 + c_3 + c_2t + \frac{1}{2}c_3t^2)$ $x_2(t) = e^{-t}(c_1 + c_2t + \frac{1}{2}c_3t^2),$ $x_3(t) = e^{-t}(c_2 + c_3t)$
- 13. Triple eigenvalue $\lambda = -1$ of defect 2; $x_1(t) = (c_1 + c_2 t + \frac{1}{2}c_3 t^2)e^{-t}$, $x_2(t) = (2c_2 + c_3 + 2c_3 t)e^{-t}$,
 - $x_3(t) = (c_2 + c_3 t)e^{-t}$
- 14. Triple eigenvalue $\lambda = -1$ of defect 2; $x_1(t) = e^{-t} (5c_1 + c_2 + c_3 + 5c_2t + c_3t + \frac{5}{2}c_3t^2),$ $x_2(t) = e^{-t} (-25c_1 - 5c_2 - 25c_2t - 5c_3t - \frac{25}{2}c_3t^2),$ $x_3(t) = e^{-t} (-5c_1 + 4c_2 - 5c_2t + 4c_3t - \frac{5}{2}c_3t^2)$

15. Triple eigenvalue
$$\lambda = 1$$
 of defect 1;
 $x_1(t) = (3c_1 + c_3 - 3c_3t)e^t$,
 $x_2(t) = (-c_1 + c_3t)e^t$, $x_3(t) = (c_2 + c_3t)e^t$

- **16.** Triple eigenvalue $\lambda = 1$ of defect 1; $x_1(t) = e^t (3c_1 + 3c_2 + c_3)$ $x_2(t) = e^t (-2c_1 - 2c_3t),$ $x_3(t) = e^t (-2c_2 + 2c_3t)$
- 17. Triple eigenvalue $\lambda = 1$ of defect 1; $x_1(t) = (2c_1 + c_2)e^t, x_2(t) = (-3c_2 + c_3 + 6c_3t)e^t,$ $x_3(t) = -9(c_1 + c_3t)e^t$
- **18.** Triple eigenvalue $\lambda = 1$ of defect 1; $x_1(t) = e^t (-c_1 - 2c_2 + c_3),$ $x_2(t) = e^t (c_2 + c_3t),$ $x_3(t) = e^t (c_1 - 2c_3t)$
- **19.** Double eigenvalues $\lambda = -1$ and $\lambda = 1$, with four linearly independent solutions;
 - $x_1(t) = c_1 e^{-t} + c_4 e^t$,
 - $x_2(t) = c_3 e^t$,
 - $x_3(t) = c_2 e^{-t} + 3c_4 e^t,$ $x_4(t) = c_1 e^{-t} - 2c_3 e^t$
- 20. Eigenvalue $\lambda = 2$ with multiplicity 4 and defect 3; $x_1(t) = (c_1 + c_3 + c_2t + c_4t + \frac{1}{2}c_3t^2 + \frac{1}{6}c_4t^3)e^{2t}$, $x_2(t) = (c_2 + c_3t + \frac{1}{2}c_4t^2)e^{2t}$,
- $x_3(t) = (c_3 + c_4t)e^{2\tilde{t}}, x_4(t) = c_4e^{2t}$ 21. Eigenvalue $\lambda = 1$ with multiplicity 4 and defect 2; $x_1(t) = (-2c_2 + c_3 - 2c_3t)e^t, x_2(t) = (c_2 + c_3t)e^t,$
- $x_1(t) = (-2c_2 + c_3 2c_3t)e^t, x_2(t) = (c_2 + c_3t)e^t,$ $x_3(t) = (c_2 + c_4 + c_3t)e^t, x_4(t) = (c_1 + c_2t + \frac{1}{2}c_3t^2)e^t$ 22. Eigenvalue $\lambda = 1$ with multiplicity 4 and defect 2;
- $\begin{aligned} x_1(t) &= (c_1 + 3c_2 + c_4 + c_2t + 3c_3t + \frac{1}{2}c_3t^2)e^t, \\ x_2(t) &= -(2c_2 c_3 + 2c_3t)e^t, \\ x_4(t) &= -(2c_1 + 6c_2 + 2c_2t + 6c_3t + c_3t^2)e^t \end{aligned}$
- 23. $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{-t} + (c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3) e^{3t}$ with $\mathbf{v}_1 = \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}^T, \mathbf{v}_2 = \begin{bmatrix} 4 & 0 & 9 \end{bmatrix}^T,$ $\mathbf{v}_3 = \begin{bmatrix} 0 & 2 & 1 \end{bmatrix}^T$
- **24.** $\mathbf{x}(t) = c_1 \mathbf{v}_1 e^{-t} + (c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3) e^{3t}$ with $\mathbf{v}_1 = \begin{bmatrix} 5 & 3 & -3 \end{bmatrix}^T, \mathbf{v}_2 = \begin{bmatrix} 4 & 0 & -1 \end{bmatrix}^T,$ $\mathbf{v}_3 = \begin{bmatrix} 2 & -1 & 0 \end{bmatrix}^T$
- 25. $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1t + \mathbf{v}_2) + c_3(\frac{1}{2}\mathbf{v}_1t^2 + \mathbf{v}_2t + \mathbf{v}_3)]e^{2t}$ with $\mathbf{v}_1 = [-1 \quad 0 \quad -1]^T$, $\mathbf{v}_2 = [-4 \quad -1 \quad 0]^T$, and $\mathbf{v}_3 = [1 \quad 0 \quad 0]^T$
- 26. $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1t + \mathbf{v}_2) + c_3(\frac{1}{2}\mathbf{v}_1t^2 + \mathbf{v}_2t + \mathbf{v}_3)]e^{3t}$ with $\mathbf{v}_1 = [0 \ 2 \ 2]^T$, $\mathbf{v}_3 = [2 \ 1 \ -3]^T$, and $\mathbf{v}_3 = [1 \ 0 \ 0]^T$
- **27.** $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1t + \mathbf{v}_2) + c_3\mathbf{v}_3]e^{2t}$ with $\mathbf{v}_1 = [-5 \ 3 \ 8]^T, \mathbf{v}_2 = [1 \ 0 \ 0]^T, \mathbf{v}_3 = [1 \ 1 \ 0]^T$
- **28.** $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1t + \mathbf{v}_2) + c_3(\frac{1}{2}\mathbf{v}_1t^2 + \mathbf{v}_2t + \mathbf{v}_3)]e^{2t}$ with $\mathbf{v}_1 = [119 - 289 \ 0]^T$, $\mathbf{v}_2 = [-17 \ 34 \ 17]^T$, and $\mathbf{v}_3 = [1 \ 0 \ 0]^T$
- **29.** $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1t + \mathbf{v}_2)]e^{-t} + [c_3\mathbf{v}_3 + c_4(\mathbf{v}_3t + \mathbf{v}_4)]e^{2t}$ with $\mathbf{v}_1 = \begin{bmatrix} 1 & -3 & -1 & -2 \end{bmatrix}^T$, $\mathbf{v}_2 = \begin{bmatrix} 0 & 1 & 0 & 0 \end{bmatrix}^T$, $\mathbf{v}_3 = \begin{bmatrix} 0 & -1 & 1 & 0 \end{bmatrix}^T$, $\mathbf{v}_4 = \begin{bmatrix} 0 & 0 & 2 & 1 \end{bmatrix}^T$
- **30.** $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1t + \mathbf{v}_2)]e^{-t} + [c_3\mathbf{v}_3 + c_4(\mathbf{v}_3t + \mathbf{v}_4)]e^{2t}$, with $\mathbf{v}_1 = \begin{bmatrix} 0 & 1 & -1 & -3 \end{bmatrix}^T$, $\mathbf{v}_2 = \begin{bmatrix} 0 & 0 & 1 & 2 \end{bmatrix}^T$, $\mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 0 & 0 \end{bmatrix}^T$, $\mathbf{v}_4 = \begin{bmatrix} 0 & 0 & 3 & 5 \end{bmatrix}^T$
- **31.** $\mathbf{x}(t) = [c_1\mathbf{v}_1 + c_2(\mathbf{v}_1 t + \mathbf{v}_2) + c_3(\frac{1}{2}\mathbf{v}_1 t^2 + \mathbf{v}_2 t + \mathbf{v}_3) + c_4\mathbf{v}_4]e^t$ with $\mathbf{v}_1 = [42 \ 7 \ -21 \ -42]^T$, $\mathbf{v}_2 = [34 \ 22 \ -10 \ -27]^T$, $\mathbf{v}_3 = [1 \ 0 \ 0 \ 0]^T$, $\mathbf{v}_4 = [0 \ 1 \ 3 \ 0]^T$

32.
$$\mathbf{x}(t) = (c_1\mathbf{v}_1 + c_2\mathbf{v}_2)e^{2t} + (c_3\mathbf{v}_3 + c_4\mathbf{v}_4 + c_5\mathbf{v}_5)e^{3t}$$
 with
 $\mathbf{v}_1 = \begin{bmatrix} 8 & 0 & -3 & 1 & 0 \end{bmatrix}^T, \mathbf{v}_2 = \begin{bmatrix} 1 & 0 & 0 & 0 & 3 \end{bmatrix}^T,$
 $\mathbf{v}_3 = \begin{bmatrix} 3 & -2 & -1 & 0 & 0 \end{bmatrix}^T,$
 $\mathbf{v}_4 = \begin{bmatrix} 2 & -2 & 0 & -3 & 0 \end{bmatrix}^T,$
 $\mathbf{v}_5 = \begin{bmatrix} 1 & -1 & 0 & 0 & 3 \end{bmatrix}^T$
33. $\mathbf{x}_1(t) = \begin{bmatrix} \cos 4t & \sin 4t & 0 & 0 \end{bmatrix}^T e^{3t},$
 $\mathbf{x}_2(t) = \begin{bmatrix} -\sin 4t & \cos 4t & 0 & 0 \end{bmatrix}^T e^{3t},$
 $\mathbf{x}_3(t) = \begin{bmatrix} t \cos 4t & t \sin 4t & \cos 4t & \sin 4t \end{bmatrix}^T e^{3t},$
 $\mathbf{x}_4(t) = \begin{bmatrix} -t \sin 4t & t \cos 4t & -\sin 4t & \cos 4t \end{bmatrix}^T e^{3t}$
34. $\mathbf{x}_1(t) = \begin{bmatrix} \sin 3t & 3t & 3t \\ 0 & \sin 3t \\ 3 \cos 3t - 3 \sin 3t \\ 0 & \sin 3t \end{bmatrix} e^{2t},$
 $\mathbf{x}_2(t) = \begin{bmatrix} 3\cos 3t & 3t & 3t \\ 0 & -\cos 3t \\ -\cos 3t \end{bmatrix} e^{2t},$
 $\mathbf{x}_3(t) = \begin{bmatrix} (3t - 10)\cos 3t - (3t + 9)\sin 3t \\ 0 & \sin 3t \\ t \sin 3t \end{bmatrix} e^{2t},$
 $\mathbf{x}_4(t) = \begin{bmatrix} -t\cos 3t + 3\sin 3t \\ (3t - 9)\cos 3t + (3t - 10)\sin 3t \\ -\cos 3t \\ -\cos 3t \end{bmatrix} e^{2t},$
 $\mathbf{x}_4(t) = \begin{bmatrix} -t\cos 3t + 3\sin 3t \\ (3t + 9)\cos 3t + (3t - 10)\sin 3t \\ -\cos 3t \\ -t\cos 3t \end{bmatrix} e^{2t}$

36.
$$x_1(t) = v_0(2 - 2e^{-t} - te^{-t}),$$

 $x_2(t) = v_0(2 - 2e^{-t} - te^{-t} - \frac{1}{2}t^2e^{-t}),$
 $\lim_{t \to \infty} x_1(t) = \lim_{t \to \infty} x_2(t) = 2v_0$

In Problems 37, 39, 41, 43, and 45 we give a nonsingular matrix **Q** and a Jordan-form matrix **J** such that $\mathbf{A} = \mathbf{QJQ}^{-1}$. Any scalar multiple of **Q** will do the same job.

37.
$$\mathbf{Q} = \begin{bmatrix} 1 & 4 & -2 \\ -1 & 0 & 9 \\ 2 & 9 & 0 \end{bmatrix}, \mathbf{J} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

39.
$$\mathbf{Q} = \begin{bmatrix} 1 & 4 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \mathbf{J} = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$$

41.
$$\mathbf{Q} = \begin{bmatrix} 0 & -5 & 0 \\ 1 & 3 & 0 \\ 1 & 8 & 1 \end{bmatrix}, \mathbf{J} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$$

43.
$$\mathbf{Q} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 3 & -1 & -1 & 2 \\ 1 & 0 & 1 & 0 \\ 2 & 0 & 0 & 1 \end{bmatrix}, \mathbf{J} = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

45.
$$\mathbf{Q} = \begin{bmatrix} -30 & 432 & 360 & 0 \\ -7 & 72 & 228 & 0 \\ 9 & -216 & -108 & 0 \\ 30 & -432 & -288 & 12 \end{bmatrix}, \mathbf{J} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Section 7.7

The format for the first eight answers is this: (x(t), y(t)) at t = 0.2 by the Euler method, by the improved Euler method, by the Runge-Kutta method, and finally the actual values.

- **1.** (0.8800, 2.5000), (0.9600, 2.6000), (1.0027, 2.6401), (1.0034, 2.6408)
- **2.** (0.8100, -0.8100), (0.8200, -0.8200), (0.8187, -0.8187), (0.8187, -0.8187)

ANSWERS TO SELECTED PROBLEM § 11

- **3.** (2.8100, 2.3100), (3.2200, 2.6200), (3.6481, 2.9407), (3.6775, 2.9628)
- **4.** (3.3100, -1.6200), (3.8200, -2.0400), (4.2274, -2.4060), (4.2427, -2.4205)
- **5.** (-0.5200, 2.9200), (-0.8400, 2.4400), (-0.5712, 2.4485), (-0.5793, 2.4488)
- **6.** (-1.7600, 4.6800), (-1.9200, 4.5600), (-1.9029, 4.4995), (-1.9025, 4.4999)
- **7.** (3.1200, 1.6800), (3.2400, 1.7600), (3.2816, 1.7899), (3.2820, 1.7902)
- **8.** (2.1600, -0.6300), (2.5200, -0.4600), (2.5320, -0.3867), (2.5270, -0.3889)
- 9. At t = 1 we obtain (x, y) = (3.99261, 6.21770) (h = 0.1) and (3.99234, 6.21768) (h = 0.05); the actual value is (3.99232, 6.21768).
- **10.** At t = 1 we obtain (x, y) = (1.31498, 1.02537) (h = 0.1) and (1.31501, 1.02538) (h = 0.05); the actual value is (1.31501, 1.02538).
- **11.** At t = 1 we obtain (x, y) = (-0.05832, 0.56664) (h = 0.1) and (-0.05832, 0.56665) (h = 0.05); the actual value is (-0.05832, 0.56665).
- 12. We solved x' = y, $y' = -x + \sin t$, x(0) = y(0) = 0. With h = 0.1 and also with h = 0.05 we obtain the actual value $x(1.0) \approx 0.15058$.
- **13.** Runge-Kutta, h = 0.1: about 1050 ft in about 7.7 s
- 14. Runge-Kutta, h = 0.1: about 1044 ft in about 7.8 s
- **15.** Runge-Kutta, h = 1.0: about 83.83 mi in about 168 s
- 16. At 40°: 5.0 s, 352.9 ft; at 45°: 5.4 s, 347.2 ft; at 50°: 5.8 s, 334.2 ft (all values approximate)
- **17.** At 39.0° the range is about 352.7 ft. At 39.5° it is 352.8; at 40°, 352.9; at 40.5°, 352.6; at 41.0°, 352.1.
- 18. Just under 57.5°
- 19. Approximately 253 ft/s
- **20.** Maximum height: about 1005 ft, attained in about 5.6 s; range: about 1880 ft; time aloft: about 11.6 s
- **21.** Runge-Kutta with h = 0.1 yields these results: (a) 21400 ft, 46 s, 518 ft/s; (b) 8970 ft, 17.5 s; (c) 368 ft/s (at $t \approx 23$).

Chapter 8

Section 8.1

1.
$$\Phi(t) = \begin{bmatrix} e^{t} & e^{3t} \\ -e^{t} & e^{3t} \end{bmatrix}, \mathbf{x}(t) = \frac{1}{2} \begin{bmatrix} 5e^{t} + e^{3t} \\ -5e^{t} + e^{3t} \end{bmatrix}$$
2.
$$\Phi(t) = \begin{bmatrix} 1 & e^{4t} \\ 2 & -2e^{4t} \end{bmatrix}, \mathbf{x}(t) = \frac{1}{4} \begin{bmatrix} 3 + 5e^{4t} \\ 6 - 10e^{4t} \end{bmatrix}$$
3.
$$\Phi(t) = \begin{bmatrix} 5\cos 4t & -5\sin 4t \\ 2\cos 4t + 4\sin 4t & 4\cos 4t - 2\sin 4t \end{bmatrix}, \mathbf{x}(t) = \frac{1}{4} \begin{bmatrix} -5\sin 4t \\ 4\cos 4t - 2\sin 4t \end{bmatrix}$$
4.
$$\Phi(t) = e^{2t} \begin{bmatrix} 1 & 1+t \\ 1 & t \end{bmatrix}, \mathbf{x}(t) = e^{2t} \begin{bmatrix} 1+t \\ t \end{bmatrix}$$
5.
$$\Phi(t) = \begin{bmatrix} 2\cos 3t & -2\sin 3t \\ -3\cos 3t + 3\sin 3t & 3\cos 3t + 3\sin 3t \end{bmatrix}, \mathbf{x}(t) = \frac{1}{3} \begin{bmatrix} 3\cos 3t - \sin 3t \\ -3\cos 3t + 6\sin 3t \end{bmatrix}$$
6.
$$\Phi(t) = e^{5t} \begin{bmatrix} \cos 4t - 2\sin 4t & 2\cos 4t + 2\sin 4t \\ 2\cos 4t & 2\sin 4t \end{bmatrix}, \mathbf{x}(t) = 2e^{5t} \begin{bmatrix} \cos 4t + \sin 4t \\ \sin 4t \end{bmatrix}$$

7.
$$\Phi(t) = \begin{bmatrix} 6 & 3e^{t} & 2e^{-t} \\ 2 & e^{t} & e^{-t} \\ 5 & 2e^{t} & 2e^{-t} \end{bmatrix},$$

$$\mathbf{x}(t) = \begin{bmatrix} -12 + 12e^{t} + 2e^{-t} \\ -4 + 4e^{t} + 2e^{-t} \\ -10 + 8e^{t} + 2e^{-t} \end{bmatrix}$$
8.
$$\Phi(t) = \begin{bmatrix} 0 & e^{t} & e^{3t} \\ e^{-2t} & -e^{t} & -e^{3t} \\ e^{-2t} & -e^{2t} & 0 & e^{3t} \end{bmatrix}, \mathbf{x}(t) = \begin{bmatrix} -e^{t} + e^{-2t} \\ -e^{-2t} \end{bmatrix}$$
9.
$$e^{At} = \begin{bmatrix} 2e^{3t} - e^{t} & -2e^{3t} + 2e^{t} \\ -2 + 2e^{2t} & 3 - 3e^{2t} \end{bmatrix}$$
10.
$$e^{At} = \begin{bmatrix} -2 + 3e^{2t} & 3 - 3e^{2t} \\ -2 + 2e^{2t} & 3 - 2e^{2t} \end{bmatrix}$$
11.
$$e^{At} = \begin{bmatrix} 3e^{3t} - 2e^{2t} & -3e^{3t} + 3e^{2t} \\ 2e^{3t} - 2e^{2t} & -2e^{3t} + 3e^{2t} \end{bmatrix}$$
12.
$$e^{At} = \begin{bmatrix} -3e^{t} + 4e^{2t} & 4e^{t} - 4e^{2t} \\ -3e^{t} + 3e^{2t} & 4e^{t} - 3e^{2t} \end{bmatrix}$$
13.
$$e^{At} = \begin{bmatrix} 4e^{3t} - 3e^{t} & -4e^{3t} + 4e^{t} \\ 3e^{3t} - 3e^{t} & -3e^{3t} + 4e^{t} \end{bmatrix}$$
14.
$$e^{At} = \begin{bmatrix} -3e^{t} + 9e^{2t} & 6e^{t} - 6e^{2t} \\ -12e^{t} + 12e^{2t} & 9e^{t} - 8e^{2t} \end{bmatrix}$$
15.
$$e^{At} = \begin{bmatrix} -9e^{t} + 10e^{2t} & 15e^{t} - 15e^{2t} \\ -12e^{t} + 22e^{t} & 6e^{t} - 9e^{2t} \end{bmatrix}$$
16.
$$e^{At} = \begin{bmatrix} -9e^{t} + 10e^{2t} & 15e^{t} - 15e^{2t} \\ -2e^{t} + 6e^{t} & -2t + 6e^{t} \end{bmatrix}$$
17.
$$e^{At} = \frac{1}{2} \begin{bmatrix} e^{2t} + e^{6t} \\ -e^{2t} + e^{6t} \\ e^{2t} + e^{2t} \end{bmatrix}$$
18.
$$e^{At} = \frac{1}{2} \begin{bmatrix} e^{2t} + e^{6t} \\ 2e^{10t} - 2e^{5t} \\ e^{10t} + 2e^{5t} \end{bmatrix}$$
19.
$$e^{At} = \frac{1}{5} \begin{bmatrix} 2e^{5t} + 4e^{15t} \\ -2e^{5t} + 2e^{15t} \\ 4e^{5t} + e^{15t} \end{bmatrix}$$
21.
$$e^{At} = \begin{bmatrix} 1 + t & -t \\ t & 1 -t \end{bmatrix}$$
22.
$$e^{At} = \begin{bmatrix} 1 + 6t & 4t \\ -9t & 1 - 6t \end{bmatrix}$$
23.
$$e^{At} = \begin{bmatrix} 1 + 5t & 4t^{2t} \\ 1 + t & -t \\ t \end{bmatrix}$$
24.
$$e^{At} = \begin{bmatrix} 1 + 3t & 0 & -3t \\ 5t + 18t^{2} & 1 & 7t - 18t^{2} \\ 3t & 0 & 1 - 3t \end{bmatrix}$$
25.
$$e^{At} = \begin{bmatrix} e^{2t} & 5te^{2t} \\ 0 & e^{2t} \\ 3t & 0 \end{bmatrix}$$
26.
$$e^{At} = \begin{bmatrix} e^{2t} & 5te^{2t} \\ 0 & e^{2t} \\ 0 & 0 \end{bmatrix}$$
27.
$$e^{At} = \begin{bmatrix} e^{2t} & 5te^{2t} \\ 0 & e^{2t} \\ 20t + 150t^{2} e^{5t} \end{bmatrix}$$
28.
$$e^{At} = \begin{bmatrix} e^{2t} & 5te^{2t} \\ 0 & e^{t} \\ 20t + 150t^{2} e^{5t} \end{bmatrix}$$
28.
$$e^{At} = \begin{bmatrix} e^{5t} & 0 & 0 \\ 10te^{5t} & e^{5t} \\ 0 & 0 \end{bmatrix}$$
28.
$$e^{At} = \begin{bmatrix} e^{4t} & 2e^{2t} \\ 0 & e^{4t} \\ 56 \end{bmatrix}$$
29.
$$e^{At} = \begin{bmatrix} e^{4t} & 2e^{4t} \\ 56 \end{bmatrix}$$

29.
$$e^{At} = \begin{bmatrix} 1 & 2t & 3t + 6t^{2} & 4t + 6t^{2} + 4t^{3} \\ 0 & 1 & 6t & 3t + 6t^{2} \\ 0 & 0 & 1 & 2t \\ 0 & 0 & 1 & 2t \end{bmatrix} e^{t},$$

$$\mathbf{x}(t) = e^{At} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$
30.
$$e^{At} = e^{3t} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 6t & 1 & 0 & 0 \\ 9t + 18t^{2} & 6t & 1 & 0 \\ 12t + 54t^{2} + 36t^{3} & 9t + 18t^{2} & 6t & 1 \end{bmatrix},$$

$$\mathbf{x}(t) = e^{At} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$
33.
$$\mathbf{x}(t) = \begin{bmatrix} c_{1} \cosh t + c_{2} \sinh t \\ c_{1} \sinh t + c_{2} \cosh t \end{bmatrix}$$
35.
$$e^{At} = \begin{bmatrix} e^{3t} & 4te^{3t} \\ 0 & e^{3t} \end{bmatrix}$$
36.
$$e^{At} = e^{t} \begin{bmatrix} 1 & 2t & 3t + 4t^{2} \\ 0 & 1 & 4t \\ 0 & 0 & 1 \end{bmatrix}$$
37.
$$e^{At} = \begin{bmatrix} e^{2t} & 3e^{2t} - 3e^{t} & 13e^{2t} - (13 + 9t)e^{t} \\ 0 & e^{t} & 3te^{t} \end{bmatrix}$$
38.
$$e^{At} = \begin{bmatrix} e^{5t} & 4e^{10t} - 4e^{5t} & 16e^{10t} - (16 + 50t)e^{5t} \\ 0 & 0 & e^{t} \end{bmatrix}$$
39.
$$e^{At} = \begin{bmatrix} e^{5t} & 4e^{10t} - 4e^{5t} & 16e^{10t} - (16 + 50t)e^{2t} \\ 0 & 0 & e^{2t} \end{bmatrix}$$
40.
$$e^{At} = \begin{bmatrix} e^{t^{2}} 3e^{2t} - 3e^{t} & 13e^{2t} - (12 + 9t)e^{t} & (51 + 18t)e^{t} - (51 - 36t)e^{2t} \\ 0 & 0 & e^{2t} & 4e^{2t} \end{bmatrix}$$
52.
$$\mathbf{x}(t) = \frac{1}{3}, \mathbf{y}(t) = -\frac{8}{3}$$
52.
$$\mathbf{x}(t) = \frac{1}{3}, \mathbf{y}(t) = -\frac{8}{3}$$
53.
$$\mathbf{x}(t) = \frac{1}{756} (84e^{-t} + 4e^{6t} - 868 + 840t - 504t^{2}), \mathbf{y}(t) = \frac{1}{756} (-864e^{-t} + 3e^{6t} + 861 - 882t + 378t^{2})$$
54.
$$\mathbf{x}(t) = \frac{1}{3t} (99e^{5t} + 8e^{-2t} - 63e^{t})$$
55.
$$\mathbf{x}(t) = \frac{1}{3t} (-12 - e^{-t} - 7te^{-t}), \mathbf{y}(t) = \frac{1}{3} (-6 - 7te^{-t})$$
65.
$$\mathbf{x}(t) = \frac{1}{3t} (1 + 16t), \mathbf{y}(t) = \frac{1}{3t} (3 \cosh t + 5 \sinh t), \mathbf{y}(t) = \frac{1}{4t} (3 \cosh t + 166e^{-9t} - 125 \cosh t - 150 \sinh t), \mathbf{y}(t) = \frac{1}{4t} (369e^{t} + 166e^{-9t} - 125 \cosh t - 150 \sinh t), \mathbf{y}(t) = \frac{1}{4t} (369e^{t} + 166e^{-9t} - 125 \cosh t - 150 \sinh t), \mathbf{y}(t) = \frac{1}{4t} (3 \cosh t + 2 \sinh t), \mathbf{y}(t) = \frac{1}{4t} (3 \cosh t + 2 \sinh t), \mathbf{y}(t) = \frac{1}{4t} (3 \cosh t + 2 \sinh t), \mathbf{y}(t) = \frac{1}{4t} \sin 2t$$
10.
$$\mathbf{x}(t) = \frac{1}{13} e^{t} (4 \cos t - 6 \sin t), \mathbf{y}(t) = \frac{1}{4} (-5 + 4t + e^{4t})$$

12.
$$x(t) = t^2$$
, $y(t) = -t^2$
13. $x(t) = \frac{1}{2}(1+5t)e^t$, $y(t) = -\frac{5}{2}te^t$

14.
$$x(t) = \frac{1}{8}(-2 + 4t - e^{4t} + 2te^{4t}), y(t) = \frac{1}{2}t(-2 + e^{4t})$$

ANSWERS TO SELECTED PROBLEM \$13

 $-5\sin 2t$ $2\cos 2t - \sin 2t$ $-5\sin 4t$ $4\cos 4t - 2\sin 4t$ $-2\sin 3t$ $3\cos 3t + 3\sin 3t$

 $-5\sin 2t$ $2\cos 2t + \sin 2t$ $-9\sin 3t$

 $3\cos 3t - 3\sin 3t$

 $-5\sin 4t$ $4\cos 4t - 2\sin 4t$

 e^{2t}

$$\begin{aligned} & 15, & (x), f(t) = 200(1 - e^{-r/(0)}, & (x) = 1 \\ & x(t) = 4 \\ & (t) = 1 \\ & x(t) = 4 \\ & (t) = 1 \\ & x(t) = 4 \\ & (t) = 1 \\ & x(t) = 4 \\ & (t) = 1 \\ & x(t) = 4 \\ & (t) = 1 \\ & x(t) = 4 \\ & (t) = 1 \\ & ($$

$$\begin{aligned} & 28. \ \mathrm{x}(t) = \begin{bmatrix} -2e^{7t} + 3e^{13t} & -2e^{7t} + 2e^{13t} & 0 \\ 3e^{7t} - 3e^{13t} & 3e^{7t} - 2e^{13t} & 0 \\ 3e^{7t} - 3e^{13t} & 3e^{7t} - 2e^{13t} & 0 \\ 3e^{7t} - 3e^{13t} & 3e^{7t} - 2e^{13t} & 0 \\ -e^{7t} + e^{13t} & -e^{13t} + e^{13t} & -2e^{13t} & 0 \\ 0 & e^{7t} & 0 & 0 \end{bmatrix} \\ & 29. \ \mathrm{x}(t) = \begin{bmatrix} 7e^{5t} - 6e^{9t} & -3e^{9t} + 3e^{9t} & -21e^{5t} + 21e^{9t} \\ 0 & 2e^{5t} - 2e^{9t} & -e^{5t} + e^{9t} & -6e^{5t} + 7e^{9t} \end{bmatrix} \\ & 30. \ \mathrm{x}(t) = \begin{bmatrix} 5e^{3t} - 4e^{7t} & -10e^{3t} + 10e^{7t} & 12e^{3t} - 12e^{7t} \\ 2e^{3t} - 2e^{7t} & -4e^{3t} + 5e^{7t} & 6e^{3t} - 6e^{7t} \\ 2e^{3t} - 2e^{7t} & -4e^{3t} + 5e^{7t} & 6e^{3t} - 6e^{7t} \\ 2e^{3t} - 2e^{7t} & -4e^{3t} + 5e^{7t} & 6e^{3t} - 6e^{7t} \\ 2e^{3t} - 2e^{7t} - 10e^{3t} + 10e^{7t} & 12e^{3t} - 12e^{7t} \\ 2e^{3t} - 2e^{7t} - 14e^{2t} + 15e^{3t} \end{bmatrix} \\ & 31. \ e^{At} = \frac{1}{4} \begin{bmatrix} -e^{-t} + 5e^{3t} & e^{-t} - e^{3t} \\ -e^{-t} - 14e^{2t} + 15e^{3t} \end{bmatrix} \\ & 31. \ e^{At} = \frac{1}{6e^{-t} - 14e^{2t} + 15e^{3t}} \end{bmatrix} \\ & 32. \ \text{With } e^{At} \ \text{as in Problem 31}, \\ & x(t) = \begin{bmatrix} -e^{-t} - 14e^{2t} + 16e^{2t} \\ 1 - 7 + 9t - 3t^{2t} + 11e^{-2t} + 11e^{-2t} + 11e^{-2t} + 11e^{-2t} + 11e^{-2t} + 11e^{-2t} + 1e^{-2t} + 1e^{12t} + 1t^{-2t} + 1e^{12t} \end{bmatrix} \\ & 33. \ e^{At} = \begin{bmatrix} 1+3t & -t \\ 9t & 1-3t \end{bmatrix}, \\ & x(t) = \begin{bmatrix} 3+11t + 8t^{2} \\ 5+17t + 24e^{2t} \end{bmatrix} \\ & 41. \ x(t) = \begin{bmatrix} -9e^{-t} + 11e^{0t} & -2e^{-t} + 2e^{0t} & 4e^{-t} + 4e^{3t} \\ -18e^{-t} + 18e^{3t} & -4e^{-t} + 4e^{3t} & 8e^{-t} - 7e^{3t} \end{bmatrix} \\ & 42. \ x(t) = \begin{bmatrix} -9e^{-t} + 18e^{3t} & -4e^{-t} + 2e^{3t} & 12e^{-2t} + 12e^{3t} \\ 3e^{-2t} + -3e^{3t} & 6e^{-2t} - 6e^{3t} & 12e^{-2t} + 12e^{3t} \end{bmatrix} \\ & 43. \ x(t) = \frac{1}{2}e^{2t} \begin{bmatrix} -1^{2} - 8t + 2 & 4t^{2} + 34t & t^{2} + 8t \\ -2t & 4t^{2} + 2t & t^{2} + 2t \end{bmatrix} \\ & 44. \ x(t) = \frac{1}{2}e^{2t} \begin{bmatrix} -1^{2} - 8t + 2 & 4t^{2} + 34t & t^{2} + 2t \\ -2t & 4t^{2} + 2t & t^{2} + 2t \end{bmatrix} \\ & 44. \ x(t) = \frac{1}{2}e^{2t} \begin{bmatrix} -1^{2} - 8t + 2 & -18t^{2} - 2tt & 2t \\ -2t^{2} - 2t^{2} - 2t^{2} - 2t^{2} & 2t^{2} \\ -2t^{2} - 2t^{2} - 2t^{2} & 2t^{2} \\ -2t^{2} - 2t^{2} - 2t^{2} & 2t^{2} \\ -2t^{2} - 2t^{2} - 2t^{2} & 2t^{2}$$

 $\begin{array}{c} -2te^{-t} \\ 1)e^{-t} + (2-t)e^{2t} \\ 2te^{-t} + te^{2t} \end{array}$ $te^{-t} + e^{2t}$ +60t+ 38*t* -18t48t+2 $= \begin{bmatrix} \cos 2t \\ \cos 2t \end{bmatrix} + i \begin{bmatrix} -\sin 4t \\ \sin 4t \end{bmatrix}$

There are two natural modes-one in which the two masses move in the same direction with frequency $\omega_1 = 1$ and with equal amplitudes, and one in which they move in opposite directions with frequency $\omega_2 = 3$ and with equal amplitudes.

48.
$$\mathbf{x}(t) = \begin{bmatrix} \cos t \\ \cos t \end{bmatrix} + i \begin{bmatrix} -\sin\sqrt{5}t \\ \sin\sqrt{5}t \end{bmatrix}$$

There are two natural modes-one in which the two masses move in the same direction with frequency $\omega_1 = 1$ and with equal amplitudes, and one in which they move in opposite directions with frequency $\omega_2 = \sqrt{5}$ and with equal amplitudes.

49.
$$\mathbf{x}(t) = \begin{bmatrix} \cos \sqrt{2}t \\ \cos \sqrt{2}t \end{bmatrix} + i \begin{bmatrix} -\sin 2t \\ \sin 2t \end{bmatrix}$$

There are two natural modes-one in which the two masses move in the same direction with frequency $\omega_1 = \sqrt{2}$ and with equal amplitudes, and one in which they move in opposite directions with frequency $\omega_2 = 2$ and with equal amplitudes.

There are two natural modes-one in which the two masses move in the same direction with frequency $\omega_1 = 2$ and with equal amplitudes, and one in which they move in opposite directions with frequency $\omega_2 = 4$ and with equal amplitudes.

Chapter 9

Section 9.1

1.	6.1.14	2. 6.1.16	3. 6.1.19	4. 6.1.13
5.	6.1.12	6. 6.1.18	7. 6.1.15	8. 6.1.17

9. Equilibrium solutions $x(t) \equiv 0, \pm 2$. The critical point (0, 0) in the phase plane looks like a center, whereas the points $(\pm 2, 0)$ look like saddle points.



10. Equilibrium solution $x(t) \equiv 0$. The critical point (0, 0) in the phase plane looks like a spiral sink.



- 11. Equilibrium solutions $x(t) \equiv \ldots, -2\pi, -\pi, 0, \pi, 2\pi, \ldots$ The phase portrait shown in the solutions manual suggests that the critical point $(n\pi, 0)$ in the phase plane is a spiral sink if *n* is even, but is a saddle point if *n* is odd.
- 12. Equilibrium solution $x(t) \equiv 0$. The critical point (0, 0) in the phase plane looks like a spiral source, with the solution curves emanating from this source spiraling outward toward a closed curve trajectory.



- 13. Solution $x(t) = x_0 e^{-2t}$, $y(t) = y_0 e^{-2t}$. The origin is a stable proper node similar to the one illustrated in Fig. 6.1.4.
- 14. Solution $x(t) = x_0 e^{2t}$, $y(t) = y_0 e^{-2t}$. The origin is an unsta-

ble saddle point.



15. Solution $x(t) = x_0 e^{-2t}$, $y(t) = y_0 e^{-t}$. The origin is a stable node.



16. Solution $x(t) = x_0 e^t$, $y(t) = y_0 e^{3t}$. The origin is an unstable improper node.



17. Solution $x(t) = A \cos t + B \sin t$, $y(t) = B \cos t - A \sin t$. The

716 ANSWERS TO SELECTED PROBLEMS

origin is a stable center.



- **18.** Solution $x(t) = A \cos 2t + B \sin 2t$, $y(t) = -2B \cos 2t + 2A \sin 2t$. The origin is a stable center.
- **19.** Solution $x(t) = A \cos 2t + B \sin 2t$, $y(t) = B \cos 2t A \sin 2t$. The origin is a stable center.
- 20. Solution $x(t) = e^{-2t} (A \cos t + B \sin t), y(t) = e^{-2t} [(-2A + B) \cos t (A + 2B) \sin t]$. The origin is a stable spiral point.



- **23.** The origin and the circles $x^2 + y^2 = C > 0$; the origin is a stable center.
- **24.** The origin and the hyperbolas $y^2 x^2 = C$; the origin is an unstable saddle point.



25. The origin and the ellipses $x^2 + 4y^2 = C > 0$; the origin is a

stable center.



26. The origin and the ovals of the form $x^4 + y^4 = C > 0$; the origin is a stable center.



Section 9.2

1. Asymptotically stable node



2. Unstable improper node

3. Unstable saddle point



- 4. Unstable saddle point
- 5. Asymptotically stable node



- 6. Unstable node
- 7. Unstable spiral point



9. Stable, but not asymptotically stable, center



- 10. Stable, but not asymptotically stable, center
- **11.** Asymptotically stable node: (2, 1)



- **12.** Unstable improper node: (2, -3)
- 13. Unstable saddle point: (2, 2)



14. Unstable saddle point: (3, 4)

8. Asymptotically stable spiral point

718 ANSWERS TO SELECTED PROBLEMS

15. Asymptotically stable spiral point: (1, 1)



16. Unstable spiral point: (3, 2)

17. Stable center: $(\frac{5}{2}, -\frac{1}{2})$



- **18.** Stable, but not asymptotically stable, center: (-2, -1)
- **19.** (0,0) is a stable node. Also, there is a saddle point at (0.67, 0.40).



20. (0,0) is an unstable node. Also, there is a saddle point at

(-1, -1) and a spiral sink at (-2.30, -1.70).



21. (0,0) is an unstable saddle point. Also, there is a spiral sink at (-0.51, -2.12).



22. (0, 0) is an unstable saddle point. Also, there are nodal sinks at $(\pm 0.82, \pm 5.06)$ and nodal sources at $(\pm 3.65, \mp 0.59)$.



23. (0,0) is a spiral sink. Also, there is a saddle point at

(-1.08, -0.68).



24. (0, 0) is a spiral source. No other critical points are visible.



25. Theorem 2 implies only that (0, 0) is a stable sink—either a node or a spiral point. The phase portrait for $-5 \le x, y \le 5$ also shows a saddle point at (0.74, -3.28) and spiral sink at (2.47, -0.46). The origin looks like a nodal sink in a second phase portrait for $-0.2 \le x, y \le 0.2$, which also reveals a second saddle point at (0.12, 0.07).





26. Theorem 2 implies only that (0,0) is an unstable source. The phase portrait for $-3 \le x, y \le 3$ also shows saddle points at (0.20, 0.25) and (-0.23, -1.50), as well as a nodal sink at (2.36, 0.58).



27. Theorem 2 implies only that (0,0) is a center or a spiral point, but does not establish its stability. The phase portrait for $-2 \le x, y \le 2$ also shows saddle points at (-0.25, -0.51) and (-1.56, 1.64), plus a nodal sink at (-1.07, -1.20). The origin looks like a likely center in a second phase portrait for $-0.6 \le x, y \le 0.6$.





28. Theorem 2 implies only that (0, 0) is a center or a spiral point, but does not establish its stability (though in the phase portrait it looks like a likely center). The phase portrait for $-0.25 \le x \le 0.25$, $-1 \le y \le 1$ also shows saddle points at (0.13, 0.63) and (-0.12, -0.47).

30. There is a saddle point at (1, 1) and a spiral sink at (-1, 1).



31. There is a saddle point at (1, 1) and a spiral sink at (-1, -1).



32. There is a saddle point at (2, 1) and a spiral sink at (-2, -1).



37. Note that the differential equation is homogeneous.



29. There is a saddle point at (0, 0). The other critical point (1, 1) is indeterminate, but looks like a center in the phase portrait.



Section 9.3

1. Linearization at (0,0): x' = 200x, y' = -150y; phase plane portrait:



3. Linearization at (75, 50): u' = -300v, v' = 100u; phase plane portrait:



- 5. The characteristic equation is $\lambda^2 + 45\lambda + 126 = 0$.
- 7. The characteristic equation is $(-24 \lambda)^2 2 \cdot (18)^2 = 0$. Phase plane portrait:



Phase plane portrait for the nonlinear system in Problems

4–7:



- 9. The characteristic equation is $\lambda^2 + 58\lambda 120 = 0$.
- 10. The characteristic equation is $(\lambda + 36)(\lambda + 18) 576 = 0$. Phase plane portrait:



Phase plane portrait for the nonlinear system in Problems 8–10:



12. The characteristic equation is $\lambda^2 + 2\lambda - 15 = 0$.

13. The characteristic equation is $\lambda^2 + 2\lambda + 6 = 0$. Phase plane

portrait:



- 15. The characteristic equation is $\lambda^2 + 2\lambda 24 = 0$.
- 17. The characteristic equation is $\lambda^2 4\lambda + 6 = 0$. Phase plane portrait:



19. The characteristic equation is $\lambda^2 + 10 = 0$. Phase plane portrait:



21. The characteristic equation is $\lambda^2 - \lambda - 6 = 0$.

22. The characteristic equation is $\lambda^2 - 5\lambda + 10 = 0$. Phase plane

portrait:



- 24. The characteristic equation is $\lambda^2 + 5\lambda 14 = 0$.
- **25.** The characteristic equation is $\lambda^2 + 5\lambda + 10 = 0$. Phase plane portrait:



- **26.** Naturally growing populations in competition Critical points: nodal source (0, 0) and saddle point (3, 2) Nonzero coexisting populations $x(t) \equiv 3$, $y(t) \equiv 2$
- **27.** Naturally declining populations in cooperation Critical points: nodal sink (0, 0) and saddle point (3, 2) Nonzero coexisting populations $x(t) \equiv 3$, $y(t) \equiv 2$



28. Naturally declining predator, naturally growing prey population Critical points: saddle point (0, 0) and apparent stable center (4, 8)

Nonzero coexisting populations $x(t) \equiv 4, y(t) \equiv 8$



29. Logistic and naturally growing populations in competition Critical points: nodal source (0, 0), nodal sink (3, 0), and saddle point (2, 2)

Nonzero coexisting populations $x(t) \equiv 2, y(t) \equiv 2$



30. Logistic and naturally declining populations in cooperation Critical points: saddle point (0, 0), nodal sink (3, 0), and saddle point (5, 4)

Nonzero coexisting populations $x(t) \equiv 5$, $y(t) \equiv 4$

31. Logistic prey, naturally declining predator population Critical points: saddle points (0, 0) and (3, 0), spiral sink (2, 4) Nonzero coexisting populations $x(t) \equiv 2, y(t) \equiv 4$



- **32.** Logistic populations in cooperation Critical points: nodal source (0,0), saddle points (10,0) and (0,20), nodal sink (30,60) Nonzero coexisting populations $x(t) \equiv 30$, $y(t) \equiv 60$
- **33.** Logistic prey and predator populations Critical points: nodal source (0,0), saddle points (30,0) and

(0, 20), nodal sink (4, 22) Nonzero coexisting populations $x(t) \equiv 4$, $y(t) \equiv 22$



34. Logistic prey and predator populations Critical points: nodal source (0,0), saddle points (15,0) and (0,5), spiral sink (10, 10) Nonzero coexisting populations $x(t) \equiv 10$, $y(t) \equiv 10$

Section 9.4

1. Eigenvalues: -2, -3; stable node



- 2. Eigenvalues: 1, 3; unstable node
- 3. Eigenvalues: -3, 5; unstable saddle point



724 ANSWERS TO SELECTED PROBLEMS

4. Eigenvalues: $-1 \pm 2i$; stable spiral point



5. Critical points: $(0, n\pi)$ where *n* is an integer; an unstable saddle point if *n* is even, a stable spiral point if *n* is odd



- 6. Critical points: (*n*, 0) where *n* is an integer; an unstable saddle point if *n* is even, a stable spiral point if *n* is odd
- Critical points: (nπ, nπ) where n is an integer; an unstable saddle point if n is even, a stable spiral point if n is odd



- Critical points: (nπ, 0) where n is an integer; an unstable node if n is even, an unstable saddle point if n is odd
- 9. If *n* is odd then $(n\pi, 0)$ is an unstable saddle point.
- 10. If *n* is odd then $(n\pi, 0)$ is a stable node.
- 11. $(n\pi, 0)$ is a stable spiral point.
- 12. Unstable saddle points at (2, 0) and (-2, 0), a stable center at (0, 0)
- 13. Unstable saddle points at (2, 0) and (-2, 0), a stable spiral point at (0, 0)
- 14. Stable centers at (2, 0) and (-2, 0), an unstable saddle point at (0, 0)

- 15. A stable center at (0, 0) and an unstable saddle point at (4, 0)
- 16. Stable centers at (2,0), (0,0), and (-2,0), unstable saddle points at (1,0) and (-1,0)
- **17.** (0, 0) is a spiral sink.



18. (0,0) is a spiral sink; the points $(\pm 2,0)$ are saddle points.



19. (0, 0) is a spiral sink.



20. $(n\pi, 0)$ is a spiral sink if *n* is even, a saddle point if *n* is odd.

Chapter 10

Section 10.1

- **1.** $1/s^2$, s > 0**2.** $\frac{1}{s}$ **3.** e/(s-3), s > 3**4.** $\frac{1}{s}$
- **2.** $2/s^3$, s > 0**4.** $s/(s^2 + 1)$, s > 0
- 5. $1/(s^2 1)$, s > 1
- 6. $\frac{1}{2}[1/s s/(s^2 + 4)], s > 0$
- 7. $(1 e^{-s})/s$, s > 08. $(e^{-s} - e^{-2s})/s$, s > 0
 - $(e^{-s} e^{-2s})/s, s > 0$ 9. $(1 e^{-s} se^{-s})/s^2, s > 0$

ANSWERS TO SELECTED PROBLEM \$25

10.
$$(s-1+e^{-s})/s^2$$
, $s > 0$
11. $\frac{1}{2}\sqrt{\pi s^{-3/2} + 3s^{-2}}$, $s > 0$
12. $(45\pi - 192s^{3/2})/(8s^{7/2})$, $s > 0$
13. $s^{-2} - 2(s-3)^{-1}$, $s > 3$
14. $3\sqrt{\pi}/(4s^{5/2}) + 1/(s+10)$, $s > 0$
15. $s^{-1} + s(s^2 - 25)^{-1}$, $s > 5$ 16. $(s+2)/(s^2+4)$, $s > 0$
17. $\cos^2 2t = \frac{1}{2}(1 + \cos 4t)$; $\frac{1}{2}[s^{-1} + s/(s^2 + 16)]$, $s > 0$
18. $3/(s^2 + 36)$, $s > 0$
19. $s^{-1} + 3s^{-2} + 6s^{-3} + 6s^{-4}$, $s > 0$
20. $1/(s-1)^2$, $s > 1$ 21. $(s^2 - 4)/(s^2 + 4)^2$, $s > 0$
22. $\frac{1}{2}[s/(s^2 - 36) - s^{-1}]$ 23. $\frac{1}{2}t^3$
24. $2\sqrt{t/\pi}$ 25. $1 - \frac{8}{3}t^{3/2}\pi^{-1/2}$
26. e^{-5t} 27. $3e^{4t}$
28. $3\cos 2t + \frac{1}{2}\sin 2t$ 29. $\frac{5}{3}\sin 3t - 3\cos 3t$
30. $-\cosh 2t - \frac{9}{2}\sinh 2t$ 31. $\frac{3}{5}\sinh 5t - 10\cosh 5t$
32. $2u(t-3)$

- **37.** f(t) = 1 u(t a). Your figure should indicate that the graph of *f* contains the point (a, 0), but not the point (a, 1).
- **38.** f(t) = u(t-a) u(t-b). Your figure should indicate that the graph of *f* contains the points (a, 1) and (b, 0), but not the points (a, 0) and (b, 1).
- **39.** Figure 10.2.8 shows the graph of the unit staircase function.

Section 10.2

1.
$$x(t) = 5\cos 2t$$

2. $x(t) = 3\cos 3t + \frac{4}{3}\sin 3t$
3. $x(t) = \frac{2}{3}(e^{2t} - e^{-t})$
4. $x(t) = \frac{1}{2}(7e^{-3t} - 3e^{-5t})$
5. $x(t) = \frac{1}{3}(2\sin t - \sin 2t)$
6. $x(t) = \frac{1}{3}(\cos t - \cos 2t)$
7. $x(t) = \frac{1}{8}(9\cos t - \cos 3t)$
8. $x(t) = \frac{1}{9}(1 - \cos 3t)$
9. $x(t) = \frac{1}{6}(2 - 3e^{-t} + e^{-3t})$
10. $x(t) = \frac{1}{4}(2t - 3 + 12e^{-t} - 9e^{-2t})$
11. $x(t) = 1, y(t) = -2$
12. $x(t) = \frac{2}{9}(e^{2t} - e^{-t} - 3te^{-t}), y(t) = \frac{1}{9}(e^{2t} - e^{-t} + 6te^{-t})$
13. $x(t) = -(2/\sqrt{3})\sinh(t/\sqrt{3}), y(t) = \cosh(t/\sqrt{3}) + (1/\sqrt{3})\sinh(t/\sqrt{3})$
14. $x(t) = \frac{1}{4}(2t - 3\sin 2t), y(t) = -\frac{1}{8}(2t + 3\sin 2t)$
15. $x(t) = \frac{1}{3}(2 + e^{-3t/2}[\cos(rt/2) + r\sin(rt/2)]), y(t) = \frac{1}{21}(28 - 9e^t + 2e^{-3t/2}[\cos(rt/2) + 4r\sin(rt/2)])$ where $r = \sqrt{3}$
16. $x(t) = \cos t + \sin t, y(t) = e^t - \cos t, z(t) = -2\sin t$
17. $f(t) = \frac{1}{3}(e^{3t} - 1)$
18. $f(t) = \frac{3}{5}(1 - e^{-5t})$
19. $f(t) = \frac{1}{9}(6\sin 3t - \cos 3t + 1)$
21. $f(t) = t - \sin t$
22. $f(t) = \frac{1}{9}(-1 + \cosh 3t)$
23. $f(t) = -t + \sinh t$
24. $f(t) = \frac{1}{2}(e^{-2t} - 2e^{-t} + 1)$

Section 10.3

1. $24/(s-\pi)^5$ 2. $\frac{3}{4}\sqrt{\pi}(s+4)^{-5/2}$ **3.** $3\pi/[(s+2)^2+9\pi^2]$ **4.** $\sqrt{2}(2s+5)/(4s^2+4s+17)$ 6. $(t-t^2)e^{-t}$ 5. $\frac{3}{2}e^{2t}$ 8. $e^{-2t} \cos t$ 7. te^{-2t} 9. $e^{3t} (3\cos 4t + \frac{7}{2}\sin 4t)$ 10. $\frac{1}{36}e^{2t/3} (8\cos \frac{4}{3}t - 5\sin \frac{4}{3}t)$ 12. $2 + 3e^{3t}$ 11. $\frac{1}{2} \sinh 2t$ 13. $3e^{-2t} - 5e^{-5t}$ 14. $2 + e^{2t} - 3e^{-t}$ 16. $\frac{1}{125}[e^{2t}(5t-2) + e^{-3t}(5t + 2)]$ 15. $\frac{1}{25}(e^{5t}-1-5t)$ 2)1 17. $\frac{1}{16}(\sinh 2t - \sin 2t)$ 18. $e^{4t} \left(1 + 12t + 24t^2 + \frac{32}{3}t^3\right)$ 19. $\frac{1}{3}(2\cos 2t + 2\sin 2t - 2\cos t - \sin t)$ **20.** $\frac{1}{32}[e^{2t}(2t-1)+e^{-2t}(2t+1)]$ 21. $\frac{1}{2}e^{-t}(5\sin t - 3t\cos t - 2t\sin t)$ 22. $\frac{1}{64}e^{t/2}[(4t+8)\cos t + (4-3t)\sin t]$ 27. $\frac{1}{4}e^{-3t}(8\cos 4t + 9\sin 4t)$ **28.** $\frac{1}{4}(1-2e^{2t}+e^{4t})$ **29.** $\frac{1}{8}(-6t + 3\sinh 2t)$ **30.** $\frac{1}{10}[2e^{-t} - e^{-2t}(2\cos 2t + \sin 2t)]$ **31.** $\frac{1}{15}(6e^{2t} - 5 - e^{-3t})$ **32.** $\frac{1}{2}(\cosh t + \cos t)$ 33. $x(t) = r(\cosh rt \sin rt - \sinh rt \cos rt)$ where $r = 1/\sqrt{2}$ **34.** $\frac{1}{2}\sin 2t + \frac{1}{3}\sin 3t$ **35.** $\frac{1}{16}(\sin 2t - 2t\cos 2t)$ **36.** $\frac{1}{50}[2e^{2t} + (10t-2)\cos t - (5t+14)\sin t]$ **37.** $\frac{1}{50}[(5t-1)e^{-t} + e^{-2t}(\cos 3t + 32\sin 3t)]$ **38.** $\frac{1}{510}e^{-3t}(489\cos 3t + 307\sin 3t) + \frac{1}{170}(7\cos 2t + 6\sin 2t)$ 39. 20 x = +t-20x =

Section 10.4

1.	$\frac{1}{2}t^2$	2.	$(e^{at} - at - 1)/a^2$
3.	$\frac{1}{2}(\sin t - t\cos t)$	4.	$2(t-\sin t)$
5.	te^{at}	6.	$(e^{at} - e^{bt})/(a - b)$
7.	$\frac{1}{3}(e^{3t}-1)$	8.	$\frac{1}{4}(1-\cos 2t)$
9.	$\frac{1}{54}(\sin 3t - 3t \cos 3t)$	10.	$(kt - \sin kt)/k^3$
11.	$\frac{1}{4}(\sin 2t + 2t \cos 2t)$	12.	$\frac{1}{5}[1 - e^{-2t}(\cos t +$
	+ ` `		$2\sin t$]
13.	$\frac{1}{10}(3e^{3t} - 3\cos t + \sin t)$	14.	$\frac{1}{3}(\cos t - \cos 2t)$
15.	$6s/(s^2+9)^2, s>0$	16.	$(2s^3 - 24s)/(s^2 + 4)^3$,
			s > 0
17.	$(s^2 - 4s - 5)/(s^2 - 4s + 13)^2$, s >	0
18	$\frac{2(3s^2+6s+7)}{s>0}$)	
10.	$(s+1)^2(s^2+2s+5)^2$, 	
19.	$\frac{1}{2}\pi - \arctan s = \arctan(1/s), s$	> 0	
20.	$\frac{1}{2}\ln(s^2+4) - \ln s, s > 0$	21.	$\ln s - \ln(s-3), s > 3$
22.	$\overline{\ln(s+1)} - \ln(s-1), s > 1$		
23.	$-(2\sinh 2t)/t$	24.	$2(\cos 2t - \cos t)/t$
25.	$e^{-2t} + e^{3t} - 2\cos t)/t$	26.	$(e^{-2t}\sin 3t)/t$
27.	$2(1-\cos t)/t$	28.	$\frac{1}{8}(t\sin t - t^2\cos t)$
29.	(s+1)X'(s) + 4X(s) = 0; x(s)	t) =	$Ct^3e^{-t}, C \neq 0$
30.	$X(s) = A/(s+3)^3$; $x(t) = C_1$	t^2e^-	$^{3t}, C \neq 0$

726 ANSWERS TO SELECTED PROBLEMS

- **31.** $(s-2)X'(s) + 3X(s) = 0; x(t) = Ct^2e^{2t}, C \neq 0$ **32.** $(s^2+2s)X'(s) + (4s+4)X(s) = 0; x(t) = C(1-t-e^{-2t}-te^{-2t}), C \neq 0$
- **33.** $(s^2 + 1)X'(s) + 4sX(s) = 0; x(t) = C(\sin t t\cos t), C \neq 0$
- 34. $x(t) = Ce^{-2t}(\sin 3t 3t \cos 3t), C \neq 0$

Section 10.5

1. $f(t) = u(t-3) \cdot (t-3)$



2.
$$f(t) = (t-1)u(t-1) - (t-3)u(t-3)$$











5.
$$f(t) = u(t - \pi) \cdot \sin(t - \pi) = -u(t - \pi) \sin t$$







7. $f(t) = \sin t - u(t - 2\pi)\sin(t - 2\pi) = [1 - u(t - 2\pi)]\sin t$



8. $f(t) = \cos \pi t - u(t-2) \cos \pi (t-2) = [1 - u(t-2)] \cos \pi t$



9. $f(t) = \cos \pi t + u(t-3) \cos \pi (t-3) = [1 - u(t-3)] \cos \pi t$



ANSWERS TO SELECTED PROBLEM \$27

34. x(t) = g(t) - u(t-1)[g(t-1) + h(t-1)], where g(t) =





11. $f(t) = 2[1 - u_3(t)]; F(s) = 2(1 - e^{-3s})/s$ 12. $F(s) = (e^{-s} - e^{-4s})/s$ 13. $F(s) = (1 - e^{-2\pi s})/(s^2 + 1)$ 14. $F(s) = s(1 - e^{-2s})/(s^2 + \pi^2)$ 15. $F(s) = (1 + e^{-3\pi s})/(s^2 + 1)$ 16. $F(s) = 2(e^{-\pi s} - e^{-2\pi s})/(s^2 + 4)$ 17. $F(s) = \pi(e^{-2s} + e^{-3s})/(s^2 + \pi^2)$ 18. $F(s) = 2\pi(e^{-3s} + e^{-5s})/(4s^2 + \pi^2)$ 19. $F(s) = e^{-s}(s^{-1} + s^{-2})$ 20. $F(s) = (1 - e^{-s})/s^2$ 21. $F(s) = (1 - 2e^{-s} + e^{-2s})/s^2$ 28. $F(s) = (1 - e^{-as} - ase^{-as})/[s^2(1 - e^{-2as})]$ 31. $x(t) = \frac{1}{2}[1 - u(t - \pi)]\sin^2 t$



32. x(t) = g(t) - u(t-2)g(t-2), where $g(t) = \frac{1}{12}(3-4e^{-t} + e^{-4t})$



33.
$$x(t) = \frac{1}{8} \left[1 - u(t - 2\pi) \right] \left(\sin t - \frac{1}{3} \sin 3t \right)$$







Chapter 11

Section 11.1

1.
$$y(x) = c_0 \left(1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \cdots \right) = c_0 e^x; \rho = +\infty$$

2. $y(x) = c_0 \left(1 + \frac{4x}{1!} + \frac{4^2 x^2}{2!} + \frac{4^3 x^3}{3!} + \frac{4^4 x^4}{4!} + \cdots \right) = c_0 e^{4x}; \rho = \infty$

3.
$$y(x) = c_0 \left(1 - \frac{3x}{2} + \frac{(3x)^2}{2!2^2} - \frac{(3x)^3}{3!2^3} + \frac{(3x)^4}{4!2^4} - \cdots \right) = c_0 e^{-3x^2}; \rho = +\infty$$
4.
$$y(x) = c_0 \left(1 - \frac{x^2}{1!} + \frac{x^4}{2!} - \frac{x^6}{3!} + \cdots \right) = c_0 e^{-x^2}; \rho = \infty$$
5.
$$y(x) = c_0 \left(1 + \frac{x^3}{3} + \frac{x^6}{2!3^2} + \frac{x^9}{3!3^3} + \cdots \right) = c_0 \exp\left(\frac{1}{3}x^3\right); \rho = +\infty$$
6.
$$y(x) = c_0 \left(1 + \frac{x}{2} + \frac{x^2}{4} + \frac{x^3}{8} + \frac{x^4}{16} + \cdots \right) = \frac{2c_0}{2-x}; \rho = \frac{1}{2}$$
7.
$$y(x) = c_0 \left(1 + 2x + 4x^2 + 8x^3 + \cdots \right) = \frac{c_0}{1-2x}; \rho = \frac{1}{2}$$
8.
$$y(x) = c_0 \left(1 + \frac{x}{2} - \frac{x^2}{8} + \frac{x^3}{16} - \frac{5x^4}{128} + \cdots \right) = c_0\sqrt{1+x}; \rho = 1$$
9.
$$y(x) = c_0 \left(1 + 2x + 3x^2 + 4x^3 + \cdots \right) = \frac{c_0}{(1-x)^2}; \rho = 1$$
10.
$$y(x) = c_0 \left(1 - \frac{3x}{2} + \frac{3x^2}{8} + \frac{x^3}{16} + \frac{3x^4}{128} + \cdots \right)$$

$$= c_0(1 - x)^{3/2}; \rho = 1$$
11.
$$y(x) = c_0 \left(1 - \frac{3x}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \cdots \right) + c_1 \left(x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \cdots \right)$$

$$= c_0 \cosh x + c_1 \sinh x; \rho = +\infty$$
12.
$$y(x) = c_0 \left(1 - \frac{(3x)^2}{2!} + \frac{(2x)^4}{4!} + \frac{(2x)^6}{6!} + \cdots \right) + \frac{c_1}{3!} \left(3x - \frac{(3x)^3}{3!} + \frac{(3x)^5}{5!} - \frac{(3x)^7}{7!} + \cdots \right)$$

$$= c_0 \cos 3x + \frac{1}{3} c_1 \sin 3x; \rho = +\infty$$
13.
$$y(x) = c_0 \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \right) + \frac{c_1}{3!} \left(3x - \frac{(3x)^3}{3!} + \frac{(3x)^5}{5!} - \frac{(3x)^7}{7!} + \cdots \right)$$

$$= x + c_0 \cos x + (c_1 - 1) \sin x; \rho = \infty$$
15.
$$(n + 1)c_n = 0 \text{ for all } n \ge 0, \text{ so } c_n = 0 \text{ for all } n \ge 0.$$
16.
$$2nc_n = c_n \text{ for all } n \ge 0, \text{ so } c_n = 0 \text{ for all } n \ge 0.$$
17.
$$c_0 = c_1 = 0 \text{ and } c_{n+1} = -nc_n \text{ for } n \ge 1.$$
18.
$$c_n = 0 \text{ for all } n \ge 0, \text{ so } c_n = 0 \text{ for all } n \ge 0.$$
19.
$$(n + 1)(n + 2)c_{n+2} = -4c_n;$$

$$y(x) = \frac{3}{2} \left[(2x) - \frac{(2x)^3}{2!} + \frac{(2x)^5}{5!} - \frac{(2x)^7}{7!} + \cdots \right] = \frac{3}{2} \sin 2x$$
20.
$$(n + 1)(n + 2)c_{n+2} = 4c_n;$$

$$y(x) = 2\left[1 + \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} + \frac{(2x)^6}{6!} + \cdots \right] = 2 \cosh 2x$$

21.
$$n(n+1)c_{n+1} = 2nc_n - c_{n-1};$$

 $y(x) = x + x^2 + \frac{x^3}{2!} + \frac{x^4}{3!} + \frac{x^5}{4!} + \dots = xe^x$

- 22. $n(n+1)c_{n+1} = -nc_n + 2c_{n-1}; y = e^{-2x}$
- 23. As $c_0 = c_1 = 0$ and $(n^2 n + 1)c_n + (n 1)c_{n-1} = 0$ for $n \ge 2, c_n = 0$ for all $n \ge 0$

Section 11.2

1.
$$c_{n+2} = c_n$$
;
 $y(x) = c_0 \sum_{n=0}^{\infty} x^{2n} + c_1 \sum_{n=0}^{\infty} x^{2n+1} = \frac{c_0 + c_1 x}{1 - x^2}$; $\rho = 1$
2. $c_{n+2} = -\frac{1}{2}c_n$; $\rho = 2$;
 $y(x) = c_0 \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^n} + c_1 \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2^n}$
3. $(n+2)c_{n+2} = -c_n$;
 $y(x) = c_0 \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{n!2^n} + c_1 \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!!}$; $\rho = +\infty$
4. $(n+2)c_{n+2} = -(n+4)c_n$; $\rho = 1$;
 $y(x) = c_0 \sum_{n=0}^{\infty} (-1)^n (n+1)x^{2n} + \frac{1}{3}c_1 \sum_{n=0}^{\infty} (-1)^n (2n+3)x^{2n+1}$

5.
$$3(n+2)c_{n+2} = nc_n; \ \rho = \sqrt{3};$$

 $y(x) = c_0 + c_1 \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)3^n}$

6.
$$(n+1)(n+2)c_{n+2} = (n-3)(n-4)c_n; \rho = \infty;$$

 $y(x) = c_0(1+6x^2+x^4) + c_1(x+x^3)$

7.
$$3(n+1)(n+2)c_{n+2} = -(n-4)^2 c_n;$$

 $y(x) = c_0 \left(1 - \frac{8x^2}{3} + \frac{8x^4}{27}\right) + c_1 \left(x - \frac{x^3}{2} + \frac{x^5}{120} + 9\sum_{n=3}^{\infty} \frac{(-1)^n [(2n-5)!!]^2 x^{2n+1}}{(2n+1)! 3^n}\right)$

8.
$$2(n+1)(n+2)c_{n+2} = (n-4)(n+4)c_n;$$

 $y(x) = c_0(1-4x^2+2x^4) +$
 $c_1\left(x - \frac{5x^3}{4} + \frac{7x^5}{32} + \sum_{n=3}^{\infty} \frac{(2n-5)!!(2n+3)!!x^{2n+1}}{(2n+1)!2^n}\right)$

9.
$$(n+1)(n+2)c_{n+2} = (n+3)(n+4)c_n; \rho = 1;$$

 $y(x) = c_0 \sum_{n=0}^{\infty} (n+1)(2n+1)x^{2n} + \frac{c_1}{3} \sum_{n=0}^{\infty} (n+1)(2n+3)x^{2n+1}$

10.
$$3(n+1)(n+2)c_{n+2} = -(n-4)c_n;$$

 $y(x) = c_0 \left(1 + \frac{2x^2}{3} + \frac{x^4}{27}\right) + c_1 \left(x + \frac{x^3}{6} + \frac{x^5}{360} + 3\sum_{n=3}^{\infty} \frac{(-1)^n (2n-5)!! x^{2n+1}}{(2n+1)! 3^n}\right)$

11.
$$5(n+1)(n+2)c_{n+2} = 2(n-5)c_n;$$

 $y(x) = c_1\left(x - \frac{4x^3}{15} + \frac{4x^5}{375}\right) + c_0\left(1 - x^2 + \frac{x^4}{10} + \frac{x^6}{750} + 15\sum_{n=4}^{\infty} \frac{(2n-7)!! 2^n x^{2n}}{(2n)! 5^n}\right)$

ANSWERS TO SELECTED PROBLEM 329

12.
$$c_2 = 0; (n+2)c_{n+3} = c_n;$$

 $y(x) = c_0 \left(1 + \sum_{n=1}^{\infty} \frac{x^{3n}}{2 \cdot 5 \cdots (3n-1)} \right) + c_1 \sum_{n=0}^{\infty} \frac{x^{3n+1}}{n! 3^n}$

13.
$$c_2 = 0; (n+3)c_{n+3} = -c_n;$$

 $y(x) = c_0 \sum_{n=0}^{\infty} \frac{(-1)^n x^{3n}}{n! 3^n} + c_1 \sum_{n=0}^{\infty} \frac{(-1)^n x^{3n+1}}{1 \cdot 4 \cdots (3n+1)}$

14.
$$c_2 = 0; (n+2)(n+3)c_{n+3} = -c_n;$$

 $y(x) = c_0 \left(1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^{3n}}{3^n \cdot n! \cdot 2 \cdot 5 \cdots (3n-1)} \right) + c_1 \sum_{n=0}^{\infty} \frac{(-1)^n x^{3n+1}}{3^n \cdot n! \cdot 1 \cdot 4 \cdots (3n+1)}$

15.
$$c_2 = c_3 = 0; (n+3)(n+4)c_{n+4} = -c_n;$$

 $y(x) = c_0 \left(1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^{4n}}{4^n \cdot n! \cdot 3 \cdot 7 \cdots (4n-1)} \right) + c_1 \sum_{n=0}^{\infty} \frac{(-1)^n x^{4n+1}}{4^n \cdot n! \cdot 5 \cdot 9 \cdots (4n+1)}$

16. y(x) = x

17.
$$y(x) = 1 + x^{2}$$

18. $y(x) = 2 \sum_{n=0}^{\infty} \frac{(-1)^{n} (x-1)^{2n}}{n! 2^{n}}$; converges for all x

19.
$$y(x) = \frac{1}{3} \sum_{n=0}^{\infty} (2n+3)(x-1)^{2n+1}$$
; converges if $0 < x < 2$

- **20.** $y(x) = 2 6(x 3)^2$; converges for all x
- **21.** $y(x) = 1 + 4(x + 2)^2$; converges for all x

22.
$$y(x) = 2x + 6$$

23.
$$2c_2 + c_0 = 0; (n+1)(n+2)c_{n+2} + c_n + c_{n-1} = 0$$
 for
 $n \ge 1; y_1(x) = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \cdots;$
 $y_2(x) = x - \frac{x^3}{6} - \frac{x^4}{12} + \cdots$

24.
$$y_1(x) = 1 + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^6}{45} + \cdots;$$

 $y_2(x) = x + \frac{x^3}{3} + \frac{x^4}{6} + \frac{x^5}{5} + \cdots$
25. $c_2 = c_3 = 0, (n+3)(n+4)c_{n+4} + (n+1)c_{n+1} + c_n = 0$

for
$$n \ge 0$$
; $y_1(x) = 1 - \frac{x^4}{12} + \frac{x^7}{126} + \cdots$;
 $y_2(x) = x - \frac{x^4}{12} - \frac{x^5}{20} + \cdots$

26.
$$y(x) = c_0 \left(1 - \frac{x^6}{30} + \frac{x^9}{72} + \cdots \right) + c_1 \left(x - \frac{x^7}{42} + \frac{x^{10}}{90} + \cdots \right)$$

27.
$$y(x) =$$

 $1 - x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{24} + \frac{x^5}{30} + \frac{29x^6}{720} + \frac{13x^7}{630} - \frac{143x^8}{40320} + \cdots;$
 $y(0.5) \approx 0.4156$
28. $y(x) =$

20.
$$y(x) = c_0 \left(1 - \frac{x^2}{2} + \frac{x^3}{6} + \cdots\right) + c_1 \left(x - \frac{x^3}{6} + \frac{x^4}{12} + \cdots\right)$$

29. $y_1(x) = 1 - \frac{1}{2}x^2 + \frac{1}{720}x^6 + \cdots;$
 $y_2(x) = x - \frac{1}{6}x^3 - \frac{1}{60}x^5 + \cdots$

- **30.** $y(x) = c_0 \left(1 \frac{x^2}{2} + \frac{x^3}{6} + \cdots \right) + c_1 \left(x \frac{x^2}{2} + \frac{x^4}{18} + \cdots \right)$
- **33.** The following figure shows the interlaced zeros of the 4th and 5th Hermite polynomials.



34. The figure below results when we use n = 40 terms in each summation. But with n = 50 we get the same picture as Fig. 8.2.3 in the text.



Section 11.3

- 1. Ordinary point
- **3.** Irregular singular point
- 4. Irregular singular point

2. Ordinary point

- 5. Regular singular point; $r_1 = 0, r_2 = -1$
- 6. Regular singular point; $r_1 = 1, r_2 = -2$
- 7. Regular singular point; r = -3, -3
- 8. Regular singular point; $r = \frac{1}{2}, -3$
- **9.** Regular singular point x = 1
- **10.** Regular singular point x = 1
- **11.** Regular singular points x = 1, -1
- **12.** Irregular singular point x = 2
- **13.** Regular singular points x = 2, -2
- 14. Irregular singular points x = 3, -3
- **15.** Regular singular point x = 2
- 16. Irregular singular point x = 0, regular singular point x = 1

17.
$$y_1(x) = \cos \sqrt{x}, y_2(x) = \sin \sqrt{x}$$

18.
$$y_1(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!(2n+1)!!},$$

 $y_2(x) = x^{-1/2} \sum_{n=0}^{\infty} \frac{x^n}{n!(2n-1)!!}$
19. $y_1(x) = x^{3/2} \left(1 + 3 \sum_{n=1}^{\infty} \frac{x^n}{n!(2n+3)!!} \right)$
 $y_2(x) = 1 - x - \sum_{n=2}^{\infty} \frac{x^n}{n!(2n-3)!!}$

20.
$$y_1(x) = x^{1/3} \sum_{n=0}^{\infty} \frac{(-1)^n 2^n x^n}{n! \cdot 4 \cdot 7 \cdots (3n+1)},$$

 $y_2(x) = \sum_{n=0}^{\infty} \frac{(-1)^n 2^n x^n}{n! \cdot 2 \cdot 5 \cdots (3n-1)}$

21.
$$y_1(x) = x \left(1 + \sum_{n=1}^{\infty} \frac{x^{2n}}{n! \cdot 7 \cdot 11 \cdots (4n+3)} \right),$$

 $y_2(x) = x^{-1/2} \left(1 + \sum_{n=1}^{\infty} \frac{x^{2n}}{n! \cdot 1 \cdot 5 \cdots (4n-3)} \right)$

22.
$$y_1(x) = x^{3/2} \left(1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{n! \cdot 9 \cdot 13 \cdots (4n+5)} \right),$$

 $y_2(x) = x^{-1} \left(1 + \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^{2n}}{n! \cdot 3 \cdot 7 \cdots (4n-1)} \right)$

23.
$$y_1(x) = x^{1/2} \left(1 + \sum_{n=1}^{\infty} \frac{x^{2n}}{2^n \cdot n! \cdot 19 \cdot 31 \cdots (12n+7)} \right)$$

 $y_2(x) = x^{-2/3} \left(1 + \sum_{n=1}^{\infty} \frac{x^{2n}}{2^n \cdot n! \cdot 5 \cdot 17 \cdots (12n-7)} \right)$

24.
$$y_1(x) = x^{1/3} \left(1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{2^n \cdot n! \cdot 7 \cdot 13 \cdots (6n+1)} \right)$$

 $y_2(x) = 1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{2^n \cdot n! \cdot 5 \cdot 11 \cdots (6n-1)}$

25.
$$y_1(x) = x^{1/2} \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{n! \cdot 2^n} = x^{1/2} e^{-x/2},$$

 $y_2(x) = 1 + \sum_{n=1}^{\infty} \frac{(-1)^n x^n}{(2n-1)!!}$

26.
$$y_1(x) = x^{1/2} \sum_{n=0}^{\infty} \frac{x^{2n}}{n! \cdot 2^n} = x^{1/2} \exp\left(\frac{1}{2}x^2\right)$$

 $y_2(x) = 1 + \sum_{n=1}^{\infty} \frac{2^n x^{2n}}{3 \cdot 7 \cdots (4n-1)}$

27.
$$y_1(x) = \frac{1}{x}\cos 3x, y_2(x) = \frac{1}{x}\sin 3x$$





30. $y_1(x) = \cos x^2$, $y_2(x) = \sin x^2$



31. $y_1(x) = x^{1/2} \cosh x$, $y_2(x) = x^{1/2} \sinh x$



34.
$$y_1(x) = x \left(1 - \frac{x^2}{42} + \frac{x^4}{1320} + \cdots \right),$$

 $y_2(x) = x^{-1/2} \left(1 - \frac{7x^2}{24} + \frac{19x^4}{3200} + \cdots \right)$

Section 11.4

5.
$$J_4(x) = \frac{1}{x^2}(x^2 - 24)J_0(x) + \frac{8}{x^3}(6 - x^2)J_1(x)$$

10. 3
11. $x^2J_1(x) + xJ_0(x) - \int J_0(x) dx + C$
12. $(x^3 - 4x)J_1(x) + 2x^2J_0(x) + C$
13. $(x^4 - 9x^2)J_1(x) + (3x^3 - 9x)J_0(x) + 9 \int J_0(x) dx + C$
14. $-xJ_1(x) + \int J_0(x) dx + C$
15. $2xJ_1(x) - x^2J_0(x) + C$
16. $3x^2J_1(x) + (3x - x^3)J_0(x) - 3 \int J_0(x) dx + C$
17. $(4x^3 - 16x)J_1(x) + (8x^2 - x^4)J_0(x) + C$
18. $-2J_1(x) + \int J_0(x) dx + C$
19. $y(x) = x[c_1J_0(x) + C_2Y_0(x)]$
20. $y(x) = \frac{1}{x}[c_1J_1(x) + c_2Y_1(x)]$
21. $y(x) = x[c_1J_{1/2}(3x^2) + c_2J_{-1/2}(3x^2)]$
22. $y(x) = x^3[c_1J_{2}(2x^{1/2}) + c_2Y_2(2x^{1/2})]$
23. $y(x) = x^{-1/3}[c_1J_{1/3}(\frac{1}{3}x^{3/2}) + c_2J_{-1/3}(\frac{1}{3}x^{3/2})]$
24. $y(x) = x^{-1/4}[c_1J_0(x) + c_2Y_0(x)]$
25. $y(x) = x^{-1}[c_1J_0(x) + c_2Y_0(x)]$
26. $y(x) = x^2[c_1J_1(4x^{1/2}) + c_2Y_1(4x^{1/2})]$
27. $y(x) = x^{1/2}[c_1J_{1/2}(2x^{3/2}) + c_2J_{-1/2}(2x^{3/2})]$
28. $y(x) = x^{-1/4}[c_1J_{3/2}(\frac{2}{5}x^{5/2}) + c_2J_{-3/2}(\frac{2}{5}x^{5/2})]$
29. $y(x) = x^{1/2}[c_1J_{1/6}(\frac{1}{3}x^3) + c_2J_{-1/6}(\frac{1}{3}x^3)]$
30. $y(x) = x^{1/2}[c_1J_{1/5}(\frac{4}{5}x^{5/2}) + c_2J_{-1/5}(\frac{4}{5}x^{5/2})]$

Appendix A

1.
$$y_0 = 3, y_1 = 3 + 3x, y_2 = 3 + 3x + \frac{3}{2}x^2, y_3 = 3 + 3x + \frac{3}{2}x^2 + \frac{1}{2}x^3, y_4 = 3 + 3x + \frac{3}{2}x^2 + \frac{1}{2}x^3 + \frac{1}{8}x^4; y(x) = 3e^x$$

3. $y_0 = 1, y_1 = 1 - x^2, y_2 = 1 - x^2 + \frac{1}{2}x^4, y_3 = 1 - x^2 + \frac{1}{2}x^4 - \frac{1}{6}x^6, y_4 = 1 - x^2 + \frac{1}{2}x^4 - \frac{1}{6}x^6 + \frac{1}{24}x^8; y(x) = \exp(-x^2)$
5. $y_0 = 0, y_1 = 2x, y_2 = 2x + 2x^2, y_3 = 2x + 2x^2 + \frac{4}{3}x^3 + \frac{2}{3}x^4; y(x) = e^{2x} - 1$
7. $y_0 = 0, y_1 = x^2, y_2 = x^2 + \frac{1}{2}x^4, y_3 = x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6, y_4 = x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6 + \frac{1}{24}x^8; y(x) = \exp(x^2) - 1$
9. $y_0 = 1, y_1 = (1 + x) + \frac{1}{2}x^2, y_2 = (1 + x + x^2) + \frac{1}{6}x^3, y_3 = (1 + x + x^2 + \frac{1}{3}x^3) + \frac{1}{24}x^4; y(x) = 2e^x - 1 - x = 1 + x + x^2 + \frac{1}{3}x^3 + \cdots$
11. $y_0 = 1, y_1 = 1 + x, y_2 = (1 + x + x^2) + \frac{1}{3}x^3, y_3 = (1 + x + x^2 + x^3) + \frac{2}{3}x^4 + \frac{1}{3}x^5 + \frac{1}{9}x^6 + \frac{1}{63}x^7; y(x) = \frac{1}{1 - x} = 1 + x + x^2 + x^3 + x^4 + x^5 + \cdots$
12. $y_0 = 1, y_1 = 1 + \frac{1}{2}x, y_2 = 1 + \frac{1}{2}x + \frac{3}{8}x^3 + \frac{1}{8}x^3 + \frac{1}{64}x^4, y_3 = 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \frac{13}{64}x^4 + \cdots; y(x) = (1 - x)^{-1/2}$
13. $\begin{bmatrix} x_0 \\ y_0 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 1 + 3t \\ -1 + 3t \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} = \begin{bmatrix} 1 + 3t + \frac{1}{2}t^2 \\ -1 + 5t - \frac{1}{2}t^2 + \frac{5}{6}t^3 \end{bmatrix}$
14. $\mathbf{x}(t) = \begin{bmatrix} e^t + te^t \\ e^t \end{bmatrix}$
16. $y_3(1) \approx 0.350185$

This page intentionally left blank

INDEX

Boldface page numbers indicate where terms are defined.

A

Abel's formula, 277, 289 Acceleration, 11 constant, 12 Addition of matrices. 164 associative law, 171 commutative law, 171 distributive law, 171 Addition of vectors, 212, 222 parallelogram law, 212, 223 triangle law, 212 Adjoint matrix, 199, 675 Air resistance, 20, 94 proportional to square of velocity, 97 proportional to velocity, 20, 95, 460 Airy equation, 625, 650 Airy function, 625 Algebra of inverse matrices, 178 Algebraic multiplicity, 354 Alligator population, 82 Almost linear system, 516 stability, 519 Amplification factor, 330 Amplitude, 305 Analytic function, 606 Apollo (satellite orbit), 462 Approximate logistic solution, 112 Argument (of complex number), 298 Arnold, David, 28 Artin, Emil, (1898-1962), 644 Associated eigenvector, 339 Associated homogeneous equation, 266, 279, 377 Asymptotic stability, 508 Augmented coefficient matrix. 148 Automobile: simplified model, 312 two-axle, 434 vibrations, 331, 336 Autonomous differential equation, 88 critical point, 88 equilibrium solution, 88

stable critical point, **89** unstable critical point, **89** Autonomous system, **503** linearized, 516 Auxiliary equation, *see* Characteristic equation Average error, 80

В

Back substitution, 140, 146 algorithm, 151 Basic unit vector, 218 Basis. 235 as maximal linearly independent set, 236, 241 as minimal spanning set, 242 as uniquely spanning set, 242 for **R**³, **218** for column space, 246 for row space, 244 for solution space, 240 orthogonal, 253 Batted baseball, 458 Beats, 329 Bernoulli equation, 61 Bessel, Friedrick W. (1784–1846), 642 Bessel equation, 590, 604, 635, 639, 642 Bessel function: identities, 647 order 1, 639 order $\frac{1}{2}$, 290, 645 12 order $\frac{3}{2}$, 650 32 order n, integral order, second kind, 646 order *p*, 642 order p, first kind, 645 order zero, first kind, 636 order zero, second kind, 646 solutions in. 649 Bifurcation diagram, 92 Bifurcation point, 92, 523

Binomial series, 605, 615 Birkhoff, Garrett (1911–1996), 118 Birth rate, 75 Bounded population growth, 76 Brachistochrone problem, 44 Brine tank examples, 366, 389, 393, 395, 496 Broughton Bridge, 330 Buoy problem, 311 Bus orbit, 462

С

Carbon-14, 36 Carrying capacity, 21, 78 Cart with flywheel, 327 Cascade of brine tanks, 54, 395, 483 Catenary problem, 44 Cauchy-Schwarz inequality, 251 Cayley-Hamilton theorem, 174, 361, 450, 477, 491, 498 Center: critical point, 412 of power series, 606 stable, 508 Central conic, 208 Chain (of generalized eigenvectors), 442, 444 Characteristic equation, 274, 292 of matrix, 340, 386 Characteristic value, see Eigenvalue Churchill, R. V. (1899-1987), 565, 592 Circular frequency, 305, 308 Clairaut equation, 71 Clarinet reed, 551 Clarke, Arthur (1917-2008), 17 Closed trajectory, 509 Clepsydra, 43 Coddington, E. A. (1920-1991), 633 Coefficient matrix, 147 augmented, 148 of a first-order system, 376 Coexistence of species, 532

Cofactor, 190 expansion, 190 matrix. 199. 675 Column (of a matrix), 147 Column rank, 244 and row rank, 247 Column space, 244 basis for, 246 Column vector, 148, 165, 169 Comet orbit, 467 Compartmental analysis, 388 Competing species, 529, 534 Competition situation, 80 Competition system, 529, 535 Complementary function, 287 Complete eigenvalue, 437 Complete independence of eigenvectors, 353 Complete set of eigenvectors, 437 Complex eigenvalue, 391, 410 with negative real part, 414 with positive real part, 415 Complex-valued function, real and imaginary parts, 296 Componentwise addition, 212, 222 multiplication of vector by scalar, 213.223 Compound interest, 36 Computational efficiency, 200 Conservation of energy, 303, 331 Constant acceleration, 12 Continuous dependence of solutions, 665 Convergence: of improper integral, 558 of power series, 605 Convolution (of functions), 587 Convolution property (of transforms), 588 Cooperation of species, 534 Coordinate plane, 214 space, 211 Corresponding eigenvalue, 339 Cramer's rule, 188, 197, 674 Criterion for diagonalizability, 349 Criterion for exactness, 66 Critical damping, 307, 308 Critical point (of system), 503 asymptotic stability, 508 center, 508 classification, 520 isolated, 514 node, 506 saddle point, 400, 507 spiral point, 509 stability, 507 Crossbow, 94, 98, 125, 134

Cumulative error, 110 Curvature, 71 Curve fitting, 203

D

Damped nonlinear vibrations, 542 Damped pendulum oscillations, 548 Damping constant, 302 Death rate, 75 Decay constant, 36 Defect (of eigenvalue), 439 Defective eigenvalue, 439 Dependence (linear), 214, 232, 270, 282, 378 Dependence on parameters, 91 Dependent variable missing, 68 Derivative (of matrix function), 375 Determinant, 188, 189, 190 2×2.188 $3 \times 3, 189$ $n \times n$, 190 and invertibility, 196, 672 and matrix product, 196, 672 and elementary row operations, 192, 669 by cofactor expansion, 190 by elimination method, 194 by permutations, 669 coefficient, 188 row and column properties, 192 Vandermonde, 204, 289 Diagonal (principal), 162 Diagonal matrix, 174 Diagonalizable matrix, 349, 352 Diagonalization, criterion for, 349 Differential equation, 1 autonomous, 88 Bernoulli, 61 Clairaut. 71 differential form, 65 Euler, 277 exact. 65 first-order, 7 general solution, 10, 11, 35, 271, 272, 286, 379, 382, 386 homogeneous, 59, 279 implicit solution, 35 linear, 47, 50, 265, 279 logistic, 21, 44, 76, 80 matrix, 469 normal form, 7 order. 6 order n, 6 ordinary, 7 partial, 7 particular solution, 10 Riccati, 70, 651

second-order, 11, 68 separable, 32 singular solution, 35 solution, 6 Differential equations and determinism, 313 Differential form, 65 Differential operator, 293 Differentiation of transforms, 589 Dimension and rank, 248 Dimension of a vector space, 237 Direction field, 18, 505 Direction of flow, 413 Displacement vector, 425 Distance between points, 252 Distinct real eigenvalues, 387, 399, 402 Dog problem, 71 Doomsday versus extinction, 81, 536 Dot product, 250 Drag coefficient, 96 Drug elimination, 37 Duplication (undetermined coefficients), 319 Dvnamic damper, 434 Dynamic phase plane graphics, 421

E

Earth-Moon satellite orbits, 461 Earthquake vibrations, 336, 435 Echelon form. 150 Echelon matrix, 150 reduced, 157 Eigenspace, 345 Eigenvalue, 339 complete, 437 complex, 391, 410, 414 defective, 439 distinct real, 351, 387, 399, 402 for matrix, 339, 386 geometric significance of, 412 imaginary, 411 multiplicity 2, 441 multiplicity k, 437 repeated, 399 zero, 403, 409 Eigenvalue matrix, 348 Eigenvalue solution, 386 Eigenvalue-eigenvector algorithm, 341 Eigenvector, 339, 386 associated with distinct eigenvalues, 351 chain of, 442 complex conjugate, 410 linearly independent, 386 rank r generalized, 442 Eigenvector equation, 386

Eigenvector matrix, 348 Element (of a matrix), 147 Elementary matrix, 180 and row operations, 180 Elementary (row) operation, 142, 149 effect on determinant, 193, 669 Elimination constant, 37 Elimination method: for linear systems, 140, 148 Gauss-Jordan, 157 Gaussian, 152 Elliptic integral, 548 Elliptical orbit, 412 Engineering functions, 603 Entry (of a matrix), 147 Equilibrium position, 302 Equilibrium solution, 88 of system, 504 Equivalent systems and matrices, 150 Error: in Euler's method, 117 in improved Euler method, 120 in Runge-Kutta method, 128 propagation, 112 Error function, 53 Escape velocity, 100 Euclidean space, inner product, 251 Euler, Leonhard (1707–1783), Euler (differential) equation, 277 Euler's formula, 296 Euler's method, 107 algorithm, 107 cumulative error, 110 error in, 117 for systems, 454 improved, 119 local error, 110 roundoff error, 111 Euler's theorem, 565 Exact equation, 65 Exactness, criterion for, 66 Existence, uniqueness of solutions: first-order equation, 23 linear system, 371, 662, 663 nth-order equation, 280 second-order equation, 268 Explosion-extinction equation, 81, 89 Exponential growth, see Natural growth Exponential matrix, 472 computation of, 478 Exponential order, 563 Exponential series, 606 Exponents (at a regular singular point), 632 External force. 303 periodic, 431

vector, 431 Extinction situation, 82, 359

F

Famous numbers, 117, 126, 136 Fibonacci sequence, 363, 615 Finite-dimensional vector space, 237 First-order equation, 7 First-order system, 366, 370, 376 FitzHugh, Richard (1922-2007), 555 FitzHugh-Nagumo equations, 555 Flight trajectories, 63 Flywheel on cart, 327 Folia of Descartes, 524 Forced motion, 303 Forced oscillations: and resonance, 431 damped, 332 undamped, 327 Forced vibrations, 266 Formal multiplication of series, 606 Fox-rabbit example, 357 Fps units, 13 Free motion, 303 damped, 303, 307 undamped, 303, 304 Free oscillations, 428 Free variables, 151, 239 Free vibrations, 266 Frequency, 305 resonance, 432 Frobenius, Georg (1848-1919), 631 Frobenius series, 631 Frobenius series solutions, 633 From the Earth to the Moon, 100 Function space, 259 Fundamental matrix, 469 Fundamental theorem of algebra, 292, 341 Funnel, 89

G

g (earth's surface gravitational acceleration), 13
G (universal gravitational constant), 99
Gallery of phase plane portraits, 417–418
Galvani, Luigi (1737–1798), 554
Gamma function, 559, 643
Gauss, hypergeometric equation, 640
Gaussian elimination algorithm, 152
Gauss-Jordan elimination algorithm, 157
General population equation, 76
General solution, 10, 11, 35, 271
of homogeneous equation, 272, 286

of homogeneous system, 379, 386 of nonhomogeneous equation, 287 of nonhomogeneous system, 382, 482 Generalized eigenvector, 442 Geometric multiplicity, 354 Geometric series, 605 Geometric significance of eigenvector, 412 Global existence of solutions, 659 Graphical method, 18

Н

Hailstone problem, 55 Half-life, 38 Halley's comet, 375, 467 Hard spring, 540 Harvesting a logistic population, 90 Harvesting and restocking, 127 Heaviside, Oliver (1850-1925), 565 Hermite equation, 624 Hermite polynomial, 624 Higher-order systems, 458 Hodgkin, A. L. (1914-1998), 554 Hodgkin-Huxley model, 555 Hole-through-Earth problem, 311 Homicide victim problem, 43 Homogeneous: first-order equation, 59 linear system of first-order equations, 370, 385 nth-order equation, 279, 291 second-order equation, 266 second-order system, 427, 496 Homogeneous linear (algebraic) system, 160 uniqueness of solution, 162 Hooke's law, 302, 425, 507, 539 Hopf bifurcation, 524 Huxley, A. F. (1917-2012), 554 Hypergeometric equation, series, 640

I

Identity matrix, **162**, 175 Identity principle: for polynomials, 260 for power series, 609 Imaginary eigenvalues, 411 Implicit solution, **34** Improper integral, 401, 558 Improper node, 401, **506** Improved Euler method, 119, 120 algorithm, 119 error in, 120 for systems, 455 Independence: see Linear independence Independent variable missing, 69 Indicial equation, 632 Indoor temperature oscillations, 56 Infinite-dimensional vector space, 237 Initial condition, 3, 7 Initial position, 12 Initial value problem, 7, 268, 280, 381 and elementary row operations, 380 Laplace transform solution, 568 order *n*, 280 Initial vector, 356 Initial velocity, 12 Inner product, 250 Integrating factor, 46 Integration of transforms, 591 Interpolating polynomial, 204 Intersection of subspaces, 228 Inverse Laplace transform, 561 uniqueness of, 565 Inverse matrix, 177 adjoint formula, 199, 676 algebra, 178 algorithm, 181 solution of linear system, 179 uniqueness, 177 Inverse of 2×2 matrix, 178 Inverse-square law of gravitation, 99, 373 Invertible matrix, 177 and determinants, 196 and row operations, 181 Irregular singular point, 629 Isolated critical point, 514

J

Jacobian matrix, **516** Joint proportion situation, 80 Jordan block (matrix), 449 Jordan normal form, **449** Jump, **562**

Κ

Kansas City, 330 Kepler, Johannes (1571–1630), 373 laws of planetary motion, 373, 466 Kinematic formula, 17 Kinetic energy, 303, 331 Kronecker delta, 492 Kutta, Wilhelm (1867–1944), 127

L

Lakes Erie, Huron, Ontario, 52 Laplace, Pierre Simon de (1749–1827), 565 Laplace transform, 558 and convolution, 588 and initial value problems, 568 and linear systems, 571 differentiation of, 589 existence, 563 integration of transforms, 591 inverse, 561 inverse transforms of series, 594 linearity, 560 notation, 561 of derivative, 568 of higher derivatives, 569 of integral, 574 of periodic function, 597 of square and triangular wave functions, 598 products of transforms, 587 translation on the s-axis, 579 translation on the *t*-axis, 594 uniqueness of inverse, 565 Law of cosines, 202 Leading entry, 150, 239 Leading variables, 151, 239 Legendre polynomial, 623 Legendre's equation, 290, 604, 622 Length of vector, 213 Leonardo Fibonacci (1175-1250?), 363 Limit cycle, 524 Limited environment situation, 80 Limiting population, 21, 78, 90, 358 Limiting velocity, 20 Linear approximation formula, 109 Linear combination, 167, 228 Linear differential equation: first-order, 47, 50 nth-order, 279 second-order, 265 Linear factor partial fractions, 579 Linear independence, 214, 216, 231, 270, 282 and orthogonality, 253 and unique linear combinations, 232 of vector-valued functions, 378 Linear independence and the determinant, 233, 234 Linear system (algebraic), 139

consistent, 139

equivalent, 150

homogeneous, 160

nonhomogeneous, 161

possibilities), 160

reduced echelon, 157

solution of. 139

number of solutions (three

inconsistent, 139

upper triangular form, 143 Linear system (differential equations): associated homogeneous system, 377 eigenspace of, 345 eigenvalue method, 386 first-order, 370, 376 general solution, 379, 386 homogeneous, 370 nonhomogeneous, 370, 382, 482 second-order, 426, 496 stability, 518 Linearity of Laplace transform, 560 Linearized system, 516 Lipschitz continuous, 658 Local error, 110 Local existence of solutions, 663 Logarithmic decrement, 313 Logistic equation, 21, 44, 76 competition situation, 80 joint proportion situation, 80 limited environment situation, 80 with harvesting, 90 Logistic populations, interactions of, 534 Logistic prey population, 536 Lower triangular matrix, 194 Lunar lander, 12, 99, 457

Μ

Maclaurin series, 606 Manchester (England), 330 Mass matrix, 425, 446 Mass-spring system, 266, 302, 309, 366, 425, 427, 432, 496, 507, 539, 570, 571, 580, 596 Mathematical model, modeling, 4 Matrix. 147 addition. 164 adjoint, 199, 675 augmented, 148 coefficient, 147 cofactor, 190, 675 column, 147 determinant, 188 diagonal, 174 diagonalizable, 349, 352 echelon, 150 eigenvalue, 330, 386 eigenvector, 339, 386 element, 147 elementary, 180 elementary row operations, 149 entry, 147 equality, 164 exponential, 472 fundamental. 469

identity, 162, 172, 175 inverse, **177**, 676 invertible, 177 Jacobian, 516 minor, 190 multiplication by scalar, 165 negative of, 165 nilpotent, 474 nonsingular, 183 norm, 658 orthogonal, 202 principal diagonal, 162 product, 168 projection matrices of, 491 reduced echelon, 157 row, 147 shape, 147 similar, 349 singular, 183 size. 147 spectral decomposition, 493 square, 161 stochastic, 357 trace. 347 transition. 356 triangular, 194 transpose, 195 with distinct eigenvalues, 352 zero, 172 Matrix algebra, 171 Matrix differential equation, 376, 469 Matrix equations, 182 Matrix exponential, 494, 499 Matrix exponential solution, 475 Matrix-valued function, 375 continuous, 375 differentiable. 375 derivative of. 375 Mechanical systems (modeling), 331 Method of elimination, 140 for determinants, 194 Method of Frobenius, 631 Method of successive approximations, 654 Method of undetermined coefficients, 314, 316, 320 for nonhomogeneous systems, 483 Mexico City, 330 Minimal spanning set, 242 Minor (of a matrix), 190 Mixture problems, 51 Mks units, 13 Modulus (of complex number), 298 Multiplication by an elementary matrix, 671 Multiplication of matrices, 168 and determinants, 196

associative law, 171 by scalars, **165** Multiplication of vector by scalar, **213** Multiplicity of eigenvalue, 354, 437 Multistory building, 435 Mutual extinction, 359

Ν

n-space, 222 *n*-tuple, 222 n-vector, 166, 222 Natural frequency, 328, 428 Natural growth and decay, 35 Natural growth equation, 37 Natural mode of oscillation, 428 Newton, Sir Isaac (1642-1727), 95, 373 Newton's law of cooling, 2, 38 Newton's law of gravitation, 99, 373 Newton's method, 97, 292 Newton's second law of motion, 11, 15, 94, 266, 302, 327, 365, 425 Nilpotent matrix, 474, 498 Nodal sink (or source), 401, 402, 406, . 408, 506 Node (improper or proper), 401, 506 Nonelementary function, 106 Nonhomogeneous equation, 266, 287 Nonhomogeneous system, 248, 370, 382, 482 Nonlinear pendulum, 544 Nonlinear spring, 539 Nonlinear vibrations, 542 Nonsingular matrix, 183 properties, 183 Null space, 255 and row space, 255

0

On-off function, 577 One-parameter family of solutions, 5 Operation (elementary), 149 Operator, polynomial differential, 293 Orbit (of comet), 467 Order of differential equation, 6 Ordinary differential equation, 7 Ordinary point, 616 Orthogonal basis, 253 Orthogonal complement, 254 properties of, 254 Orthogonal matrix, 202 Orthogonal vectors, 252 Orthogonality and linear independence, 253 Oscillating populations, 528 Oscillating springs:

hard, 540 soft, 541 Overdamping, 307

Ρ

Parallelogram law of addition, 212, 223 Parameter, 141 Parameters, variation of, 322 for linear systems, 485 Partial-fraction decomposition, 260, 579 Pendulum, 303, 311 nonlinear, 544 Period, 305, 547, 597 Periodic function, 597 Permutation, 668 Phase angle, 305 Phase diagram, 88 Phase plane, 369, 503 position-velocity, 539 Phase plane portrait, 369, 505 Physical units, 13 Picard, Emile (1856-1941), 654 Piecewise continuous function, 561 jump, 562 Piecewise smooth function, 568 Pivot columns of a matrix, 245 Polar form of a complex number, 298 Polking, John, 28, 513, 525 Polluted reservoir, 55 Polynomial, 203, 237, 259 interpolating, 204 Polynomial differential operator, 293 Population equation, 76 Population explosion, 76, 359 Population growth, 3, 35 Population vector, 356 Position function, 11 Position vector, 373 Potential energy, 303, 331 Power series. 604 Power series method, 604, 608 Power series representation, 605 Practical resonance, 333 Predation. 534 Predator-prey model (or situation or system), 357, 527, 535 Predictor-corrector methods, 120 Principal diagonal, 162 Principia Mathematica, 95, 373 Principle of superposition: for linear differential equations, 267.279 for systems, 377 Product: matrix. 168

matrix with number, **165** of determinants, 196 Projection matrix, 491, 498 Proper node, 401, **506** Proper subspace, 219 Pseudofrequency, 308 Pseudoperiod, 308 Pythagorean formula, 253 Python programming language, 115, 125, 135, 465

Q

Quadratic factor partial fraction, 579 Quadratic factors, repeated, 583

R

 \mathbf{R}^2 as a vector space, 214 \mathbf{R}^3 as a vector space, 214 Rabbit-fox example, 357 Radioactive decay, 36 Radius of convergence, 611 Railway car examples, 428, 446 Rank and dimension, 248 Rank r generalized eigenvector, 442 Rayleigh, Lord (John William Strutt, 1842-1919), 551 Recurrence relation, 609 many-term, 620 two-term, 620 Reduced echelon matrix, 157 uniqueness of, 158 Reducible second-order equation: dependent variable missing, 68 independent variable missing, 69 Reduction of order, 289 Regular singular point, 629 Repeated eigenvalue, 399 negative, 408 positive, 405 zero, 409 Resistance matrix, 446 Resonance, 330, 432 and repeated quadratic factors, 583 practical, 333 Riccati equation, 70, 651 River crossing, 15 Rocket propulsion equation, 104 Rodrigues' formula, 624 Rolling disk, 331 Roots of characteristic equation: complex, 297 distinct real, 274, 292 repeated complex, 299 repeated real, 275, 295, 405 Rota, Gian-Carlo (1932-1999), 118 Roundoff error, 111

Row (of a matrix), 147 Row equivalent matrices, 150 Row operations, 149 and determinants, 192 and elementary matrices, 180 and invertible matrices, 181 Row rank, 243 and column rank, 247 Row space, 243 and null space, 255 basis for, 244 of an echelon matrix, 243 of equivalent matrices, 243 Row vector, 166, 169 of a matrix, 169, 242 Runge, Carl (1856–1927), 127 Runge-Kutta method, 128 error in, 128 for systems, 456 variable step size methods, 460

S

Saddle point, 400, 507 Sawtooth function, 577 Second law of motion, 11, 13, 94, 95, 266, 302, 327, 365, 425 Second-order equation, 68 approximate solution, 456 Second-order system, 426, 496 Separable differential equation, 32 Separation of variables, 31 Separatrix, 532, 541 Series: binomial, 605 convergence, 605 exponential, 606 formal multiplication, 606 geometric, 605 hypergeometric, 640 identity principle, 609 Maclaurin, 606 power, 604 radius of convergence, 611 shift of index, 610 Taylor, 606 termwise addition, 606 termwise differentiation, 608 Shape (of a matrix), 147 Shift of index, 610 Similar matrices, 349 Simple harmonic motion, 305 Simple pendulum, 303 Sine integral function, 50 Singular matrix, 183 Singular point, 616 irregular, 629 regular, 629

Singular solution, 35 Sink, 401, 408, 507 Size (of a matrix), 147 Skydiver, 131, 136 Skywalk, 330 Slope field, 18, 505 applications of, 20 Snowplow problem, 43 Soft spring, 541 Soft touchdown, 12 Solar wind, 17 Solution: eigenvalue, 386 equilibrium, 49, 88, 504 existence, uniqueness, 23, 371, 662, 663 fundamental matrix, 470 general, 10, 11, 35, 271, 272, 286 implicit, 34 linear first-order, 47 of differential equation, 6 of linear (algebraic) system, 139 of nonhomogeneous equation, 287 of nonhomogeneous system, 382, 482 of second-order system, 427, 496 of system of differential equations, 365. 371. 376 one-parameter family, 5 particular, 10 singular, 35 straight-line, 403 trivial, 160 two-parameter family, 7 zero. 403 Solution curve, 18, 369 Solution near ordinary point, 617 Solution set, 139 Solution space: basis, 240 of a linear (algebraic) system, 226 of a linear differential equation, 261, 267 Solution subspace, 226 Solutions in Bessel functions, 649 Solve, an initial value problem, 7 Source, 402, 406, 507 Spacecraft landing, 464 Span, spanning set, 229, 238 unique, 232, 242 Spectral decomposition methods, 490 Spectral decomposition of a matrix, 493, 499 Spiral point (sink or source),415, 509 Spout, 89 Spring constant, **302** Square matrix, 161

Square wave function, 566, 577 Stability: of almost linear systems, 519 of linear systems, 518 Stable center, 508 Stable critical point, 89, 507 Staircase function, 566, 576 Standard basis for \mathbf{R}^n , 235 Standard unit vectors, 231 Star, see Proper node Static displacement, 303, 330 Static equilibrium position, 303 Steady periodic solution, 433 Step function, 562, 594 Step size, 106 Stiffness matrix, 425 Stirling's approximation, 55 Stochastic matrix, 357 Stokes' drag law, 313 Stonehenge, 38 Straight-line solution, 403 Subspace, 224 criterion, 225 intersection, 228 of **R**³, 219 proper, 226 solution, 226 sum, 228 zero, 226 Substitution method, 58 Sum: of matrices. 164 of subspaces, 228 of vectors, 212, 222 Superposition principle, see Principle of superposition Swimmer's problem, 14 Systems of dimension two, 398

Т

Taylor series, 606 Temperature oscillations, 56 Terminal speed, 96 Termwise addition of series, 606 Termwise differentiation of series, 608 Three possibilities (for linear systems), 139, 160 Threshold population, 81, 90 Time lag, 305 Time reversal, 402 Time-varying amplitude, 308 Torricelli's law, 2, 39 Trace (of a matrix), 347 Trajectory, 369, 503 closed, 509 Transform of derivative, 568 Transform of integral, 574 Transform of periodic function, 597 Transform perspective, 572 Transient solution, 332, 433 Transition matrix, 356 Translated series solutions, 619 Transpose (of a matrix), 195 Transposition, 668 Triangle inequality, 253 Triangle law of addition, 212 Triangular matrix, 194 Triangular wave function, 577 Trivial solution, 160 Two independent eigenvectors, 405

U

Undamped forced oscillations, 327 Undamped motion, 303 Underdamping, 308 Undetermined coefficients, 314, 483 case of duplication, 319 Unicycle model of car, 331 Uniform convergence, 659 Unique solution, 140 Uniqueness of linear combinations, 232 Uniqueness of solutions, see Existence Uniquely spanning set, 232, 242 Unit step function, 562, 594 Unit vector, 176, 218 standard, 231 Units (physical), 13 Unstable critical point, 89, 507 Upper triangular matrix, 194 U.S. population, 78, 86

V

van der Pol, Balthasar (1889–1959), 552 van der Pol's equation, 552 Vandermonde determinant, 204, 289 Vandermonde matrix, 204, 260 Variable gravitational acceleration, 99 Variables: free, **151**, 239 leading, **151**, 239 Variation of parameters, 322, 485 formula, 324, 486 Vector: 2-vector, 214 3-vector, 212 basic unit, 176, 218 column, 148, 169 component, 212 coordinates, 211 length, 213, 251 multiplication by scalar, 213, 223 *n*-vector, 222 norm, 658 orthogonal, 252 row, 166, 169 scalar product, 167 sum, 212, 222 unit, 176, **218** zero, 213, 223 Vector addition, 212, 222 Vector space, 223 basis of, 235 dimension, 237 finite-dimensional, 237 infinite dimensional, 237 of functions, 224, 259 subspace, 224 Velocity, 11 limiting, 20 Verhulst, Pierre-François (1804–1849), 78 Verne, Jules (1828–1905), 100, 103 Vertical motion with gravitational acceleration, 13 with air resistance, 95 Viscosity, 313 Volterra, Vito (1860-1940), 526

W

Watson, G. N. (1886–1965), 642
Weight, 13
Well-posed problems and mathematical models, 665
Without two independent eigenvectors, 406
World population, 37, 87, 205
Wronskian, 271, 283 of solutions, 272, 285, 289 of vector-valued functions, 378

Ζ

Zero matrix, **172** Zero subspace, 226 Zero vector, 213, 223

Table of Laplace Transforms

 s^{a+1}

This table summarizes the general properties of Laplace transforms and the Laplace transforms of particular functions derived in Chapter 10.

Function	Transform	Function	Transform
f(t)	F(s)	e ^{at}	$\frac{1}{s-a}$
af(t) + bg(t)	aF(s) + bG(s)	$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$
f'(t)	sF(s) - f(0)	$\cos k t$	$\frac{s}{s^2 + k^2}$
f''(t)	$s^2 F(s) - sf(0) - f'(0)$	sin k t	$\frac{k}{s^2 + k^2}$
$f^{(n)}(t)$	$s^{n}F(s) - s^{n-1}f(0) - \dots - f^{(n-1)}(0)$	$\cosh k t$	$\frac{s}{s^2 - k^2}$
$\int_0^t f(\tau) d\tau$	$\frac{F(s)}{s}$	sinh k t	$\frac{k}{s^2 - k^2}$
$e^{at}f(t)$	F(s-a)	$e^{at}\cos kt$	$\frac{s-a}{(s-a)^2+k^2}$
u(t-a)f(t-a)	$e^{-as}F(s)$	$e^{at}\sin kt$	$\frac{k}{(s-a)^2 + k^2}$
$\int_0^t f(\tau)g(t-\tau)d\tau$	F(s)G(s)	$\frac{1}{2k^3}(\sin kt - kt\cos kt)$	$\frac{1}{(s^2+k^2)^2}$
tf(t)	-F'(s)	$\frac{t}{2k}\sin kt$	$\frac{s}{(s^2+k^2)^2}$
$t^n f(t)$	$(-1)^n F^{(n)}(s)$	$\frac{1}{2k}(\sin kt + kt\cos kt)$	$\frac{s^2}{(s^2+k^2)^2}$
$\frac{f(t)}{t}$	$\int_{s}^{\infty} F(\sigma) d\sigma$	u(t-a)	$\frac{e^{-as}}{s}$
f(t), period p	$\frac{1}{1-e^{-ps}}\int_0^p e^{-st}f(t)dt$	$\delta(t-a)$	e^{-as}
1	$\frac{1}{s}$	$(-1)^{[t/a]}$ (square wave)	$\frac{1}{s} \tanh \frac{as}{2}$
t	$\frac{1}{s^2}$	$\left[\frac{t}{a}\right]$ (staircase)	$\frac{e^{-as}}{s(1-e^{-as})}$
t^n	$\frac{n!}{s^{n+1}}$		
$\frac{1}{\sqrt{\pi t}}$	$\frac{1}{\sqrt{s}}$		
t^a	$\frac{\Gamma(a+1)}{a+1}$		

Table of Integrals

ELEMENTARY FORMS

1.
$$\int u \, dv = uv - \int v \, du$$

2.
$$\int u^n \, du = \frac{1}{n+1}u^{n+1} + C \quad \text{if } n \neq -1$$

3.
$$\int \frac{du}{u} = \ln |u| + C$$

4.
$$\int e^u \, du = e^u + C$$

5.
$$\int a^u \, du = \frac{a^u}{\ln a} + C$$

6.
$$\int \sin u \, du = -\cos u + C$$

7.
$$\int \cos u \, du = \sin u + C$$

8.
$$\int \sec^2 u \, du = \tan u + C$$

9.
$$\int \csc^2 u \, du = -\cot u + C$$

10.
$$\int \sec u \, \tan u \, du = \sec u + C$$

11. $\int \csc u \, \cot u \, du = -\csc u + C$
12. $\int \tan u \, du = \ln |\sec u| + C$
13. $\int \cot u \, du = \ln |\sin u| + C$
14. $\int \sec u \, du = \ln |\sec u + \tan u| + C$
15. $\int \csc u \, du = \ln |\sec u - \cot u| + C$
16. $\int \frac{du}{\sqrt{a^2 - u^2}} = \sin^{-1} \frac{u}{a} + C$
17. $\int \frac{du}{a^2 + u^2} = \frac{1}{a} \tan^{-1} \frac{u}{a} + C$
18. $\int \frac{du}{a^2 - u^2} = \frac{1}{2a} \ln \left| \frac{u + a}{u - a} \right| + C$

r

TRIGONOMETRIC FORMS

 $19. \int \sin^2 u \, du = \frac{1}{2}u - \frac{1}{4}\sin 2u + C$ $23. \int \sin^3 u \, du = -\frac{1}{3}(2 + \sin^2 u)\cos u + C$ $20. \int \cos^2 u \, du = \frac{1}{2}u + \frac{1}{4}\sin 2u + C$ $24. \int \cos^3 u \, du = \frac{1}{3}(2 + \cos^2 u)\sin u + C$ $21. \int \tan^2 u \, du = \tan u - u + C$ $25. \int \tan^3 u \, du = \frac{1}{2}\tan^2 u + \ln|\cos u| + C$ $26. \int \cot^3 u \, du = -\frac{1}{2}\cot^2 u - \ln|\sin u| + C$ $27. \int \sec^3 u \, du = \frac{1}{2}\sec u \tan u + \frac{1}{2}\ln|\sec u + \tan u| + C$ $28. \int \csc^3 u \, du = -\frac{1}{2}\csc u \cot u + \frac{1}{2}\ln|\csc u - \cot u| + C$ $29. \int \sin au \sin bu \, du = \frac{\sin(a - b)u}{2(a - b)} - \frac{\sin(a + b)u}{2(a + b)} + C$ $if a^2 \neq b^2$

(Continued on Rear Endpaper)
Table of Integrals (cont.)

30.
$$\int \cos au \, \cos bu \, du = \frac{\sin(a-b)u}{2(a-b)} + \frac{\sin(a+b)u}{2(a+b)} + C \quad \text{if } a^2 \neq b^2$$
31.
$$\int \sin au \, \cos bu \, du = -\frac{\cos(a-b)u}{2(a-b)} - \frac{\cos(a+b)u}{2(a+b)} + C \quad \text{if } a^2 \neq b^2$$
32.
$$\int \sin^n u \, du = -\frac{1}{n} \sin^{n-1} u \, \cos u + \frac{n-1}{n} \int \sin^{n-2} u \, du$$
33.
$$\int \cos^n u \, du = \frac{1}{n} \cos^{n-1} u \, \sin u + \frac{n-1}{n} \int \cos^{n-2} u \, du$$
34.
$$\int \tan^n u \, du = \frac{1}{n-1} \tan^{n-1} u - \int \tan^{n-2} u \, du \quad \text{if } n \neq 1$$
35.
$$\int \cot^n u \, du = -\frac{1}{n-1} \cot^{n-1} u - \int \cot^{n-2} u \, du \quad \text{if } n \neq 1$$
36.
$$\int \sec^n u \, du = \frac{1}{n-1} \sec^{n-2} u \, \tan u + \frac{n-2}{n-1} \int \sec^{n-2} u \, du \quad \text{if } n \neq 1$$
37.
$$\int \csc^n u \, du = -\frac{1}{n-1} \csc^{n-2} u \, \cot u + \frac{n-2}{n-1} \int \csc^{n-2} u \, du \quad \text{if } n \neq 1$$
38.
$$\int u \sin u \, du = \sin u - u \cos u + C$$
39.
$$\int u \cos u \, du = \cos u + u \sin u + C$$
40.
$$\int u^n \sin u \, du = -u^n \cos u + n \int u^{n-1} \cos u \, du$$

FORMS INVOLVING $\sqrt{u^2 \pm a^2}$

42.
$$\int \sqrt{u^2 \pm a^2} \, du = \frac{u}{2} \sqrt{u^2 \pm a^2} \pm \frac{a^2}{2} \ln \left| u + \sqrt{u^2 \pm a^2} \right| + C$$

43.
$$\int \frac{du}{\sqrt{u^2 \pm a^2}} = \ln \left| u + \sqrt{u^2 \pm a^2} \right| + C$$

FORMS INVOLVING $\sqrt{a^2 - u^2}$

44.
$$\int \sqrt{a^2 - u^2} \, du = \frac{u}{2} \sqrt{a^2 - u^2} + \frac{a^2}{2} \sin^{-1} \frac{u}{a} + C$$

45.
$$\int \frac{\sqrt{a^2 - u^2}}{u} \, du = \sqrt{a^2 - u^2} - a \ln \left| \frac{a + \sqrt{a^2 - u^2}}{u} \right| + C$$

EXPONENTIAL AND LOGARITHMIC FORMS

$$46. \int ue^{u} du = (u-1)e^{u} + C$$

$$49. \int e^{au} \sin bu \, du = \frac{e^{au}}{a^{2} + b^{2}} (a \sin bu - b \cos bu) + C$$

$$47. \int u^{n}e^{u} du = u^{n}e^{u} - n \int u^{n-1}e^{u} \, du$$

$$50. \int_{C} e^{au} \cos bu \, du = \frac{e^{au}}{a^{2} + b^{2}} (a \cos bu + b \sin bu) + C$$

$$48. \int u^{n} \ln u \, du = \frac{u^{n+1}}{n+1} \ln u - \frac{u^{n+1}}{(n+1)^{2}} + C$$

INVERSE TRIGONOMETRIC FORMS

51.
$$\int \sin^{-1} u \, du = u \sin^{-1} u + \sqrt{1 - u^2} + C$$
52.
$$\int \tan^{-1} u \, du = u \tan^{-1} u - \frac{1}{2} \ln(1 + u^2) + C$$
53.
$$\int \sec^{-1} u \, du = u \sec^{-1} u - \ln \left| u + \sqrt{u^2 - 1} \right| + C$$
54.
$$\int u \sin^{-1} u \, du = \frac{1}{4} (2u^2 - 1) \sin^{-1} u + \frac{u}{4} \sqrt{1 - u^2} + C$$
55.
$$\int u \tan^{-1} u \, du = \frac{1}{2} (u^2 + 1) \tan^{-1} u - \frac{u}{2} + C$$
56.
$$\int u \sec^{-1} u \, du = \frac{u^2}{2} \sec^{-1} u - \frac{1}{2} \sqrt{u^2 - 1} + C$$
57.
$$\int u^n \sin^{-1} u \, du = \frac{u^{n+1}}{n+1} \sin^{-1} u - \frac{1}{n+1} \int \frac{u^{n+1}}{\sqrt{1 - u^2}} \, du \quad \text{if } n \neq -1$$
58.
$$\int u^n \tan^{-1} u \, du = \frac{u^{n+1}}{n+1} \tan^{-1} u - \frac{1}{n+1} \int \frac{u^{n+1}}{1 + u^2} \, du \quad \text{if } n \neq -1$$
59.
$$\int u^n \sec^{-1} u \, du = \frac{u^{n+1}}{n+1} \sec^{-1} u - \frac{1}{n+1} \int \frac{u^{n+1}}{\sqrt{u^2 - 1}} \, du \quad \text{if } n \neq -1$$

OTHER USEFUL FORMULAS

$$60. \int_{0}^{\infty} u^{n} e^{-u} du = \Gamma(n+1) = n! \quad (n \ge 0) \qquad 61. \int_{0}^{\infty} e^{-au^{2}} du = \frac{1}{2} \sqrt{\frac{\pi}{a}} \quad (a > 0)$$

$$62. \int_{0}^{\pi/2} \sin^{n} u du = \int_{0}^{\pi/2} \cos^{n} u du = \begin{cases} \frac{1 \cdot 3 \cdot 5 \cdots (n-1)}{2 \cdot 4 \cdot 6 \cdots n} \cdot \frac{\pi}{2} & \text{if } n \text{ is an even integer and } n \ge 2 \\ \frac{2 \cdot 4 \cdot 6 \cdots (n-1)}{3 \cdot 5 \cdot 7 \cdots n} & \text{if } n \text{ is an odd integer and } n \ge 3 \end{cases}$$